

# Measurement of Feynman- $x$ Spectra of Photons and Neutrons in the Very Forward Direction in DIS at HERA

---

**J. Olsson\* (on behalf of the H1 Collaboration)**

*DESY, Hamburg, Germany*

*E-mail: jan.olsson@desy.de*

H1  $ep$  scattering data taken at HERA in the years 2006-7 are used to measure the production of very forward directed photons and neutrons. Normalised cross sections are given as a function of Feynman- $x$  and of  $W$ , the centre-of-mass energy of the virtual photon-proton system. The phase space covered is defined by the ranges  $6 < Q^2 < 100 \text{ GeV}^2$ ,  $0.05 < y < 0.6$ ,  $70 < W < 245 \text{ GeV}$  and  $\eta > 7.9$ , where  $Q^2$  is the photon virtuality,  $y$  the inelasticity and  $\eta$  the pseudorapidity in the laboratory frame. The integrated luminosity is  $131 \text{ pb}^{-1}$ . Comparisons to the data are made with the predictions of several models used in cosmic ray physics for the simulation of hadronic interactions in the Earth atmosphere. The hypothesis of Feynman Scaling is tested, by studying the  $W$ -dependence of the cross sections differential in Feynman- $x$ . This is the first direct experimental test of Feynman Scaling for photon and neutron production in the very forward direction at HERA.

*The European Physical Society Conference on High Energy Physics  
22–29 July 2015  
Vienna, Austria*

---

\*Speaker.


## 1. Introduction

At the  $ep$  collider HERA the study of forward production of baryons (protons and neutrons) and photons has long been a subject of interest [1]. The term “forward” here implies that the produced particles have small polar angles with respect to the proton beam direction and carry a large fraction of the incoming proton longitudinal momentum. The early studies of forward baryon production at HERA [2, 3] showed that models of deep-inelastic scattering (DIS) are able to reproduce the data, but only if several production mechanisms are taken into account: string fragmentation, pion exchange, proton diffractive dissociation and elastic  $ep$  scattering. While the production mechanism for forward photons is almost solely the proton fragmentation, through the decay of neutral mesons from the fragmentation process, forward neutrons can also be produced in the colour singlet (pion) exchange process (Figure 1a,b). The fact that different production mechanisms have to be invoked for the simultaneous measurement of forward photons and neutrons is a challenge for the Monte Carlo (MC) simulation models. While it was found in a previous H1 analysis [3] that the forward neutron production can be well described as a sum of fragmentation and pion exchange processes, the forward photon production was in contrast found to be overestimated in the models, by large amounts (50-70%) [4]. Thus, very forward neutron and photon production offers a test of commonly used simulation models in an extreme corner of the DIS phase space.

Interest in the forward production of baryons and photons also arises in Cosmic Ray (CR) physics [5]. To estimate the energy of air shower initiating cosmic rays, extensive simulation of the hadronic interactions in the Earth atmosphere is needed, and several programs have been developed over the years for this purpose. For the calibration and tuning of such programs, forward scattering data obtained under controlled conditions, i.e. in high energy accelerator experiments, are needed. Such data are however surprisingly scarce. Thus, another motivation for the present study, and for the presentation of the data in terms of the Feynman- $x$  variable, was requests from the CR community.

The data presented here were taken in the last years of HERA operation, 2006-7, and consist of  $131 \text{ pb}^{-1}$ , taken with positron and proton beam energies of 27.6 and 920 GeV respectively, corresponding to the  $ep$  centre-of-mass (CM) energy  $\sqrt{s} = 319 \text{ GeV}$ . Neutrons and photons are detected in the H1 Forward Neutron Calorimeter (FNC) and to distinguish photons and neutrons the FNC Preshower is used. The latter is an electromagnetic calorimeter situated in front of the FNC Main Calorimeter. The aperture of the FNC, as given by the HERA beam optics elements, is limited to  $0.75 \text{ mrad}$  ( $\eta > 7.9$ , where  $\eta$  is the pseudorapidity in the laboratory frame).

The present analysis was recently published [6], as a continuation of the earlier analyses of forward neutrons [3] and forward photons [4]; thus, for details more than can be given here, please



**Figure 1:**  
Diagrams for neutron and photon forward production

NC DIS Selection	
$6 < Q^2 < 100 \text{ GeV}^2$	
$0.05 < y < 0.6$	
$70 < W < 245 \text{ GeV}$	
Forward photons	Forward neutrons
$\eta > 7.9$	$\eta > 7.9$
$0.1 < x_F < 0.7$	$0.1 < x_F < 0.94$
$0 < p_T^* < 0.4 \text{ GeV}$	$0 < p_T^* < 0.6 \text{ GeV}$
$W$ ranges for cross sections $\frac{1}{\sigma_{DIS}} \frac{d\sigma}{dx_F}$	
$70 < W < 130 \text{ GeV}$	
$130 < W < 190 \text{ GeV}$	
$190 < W < 245 \text{ GeV}$	

**Table 1:** Definition of the kinematic phase space of the measurements.

see these papers.

## 2. Data Sample

The data selection cuts are summarised in Table 1. Note that the ranges of  $x_F$ , the Feynman- $x$  variable, differ for neutrons and photons.  $x_F$  is defined as  $x_F = 2p_{\parallel}^*/W$ , where  $p_{\parallel}^*$  is the longitudinal momentum of the particle in the virtual photon-proton CM frame, with respect to the direction of the beam proton<sup>1</sup>. The lower value of the upper bound on  $x_F$  for photons is motivated by the wish to define the cross section for single photons; as known from MC simulations, at larger  $x_F$  values electromagnetic clusters in the FNC Preshower are mainly caused by unresolved double photons (from  $\pi^0$  decays). The common lower bound  $x_F = 0.1$  corresponds to lower values of about 92 GeV for the FNC cluster energies. The difference in upper bounds on the transverse momentum  $p_T^*$  is a consequence of the different upper bounds on  $x_F$ .

For the purpose of the test of the Feynman Scaling hypothesis, the  $W$  range is divided into three intervals. For the presentation of the results, the cross sections are normalised to the total DIS cross section.

The total sample, after all cuts, consists of  $\sim 230,000$  neutron and  $\sim 83,000$  photon events.

<sup>1</sup>In the kinematic range of this measurement the variable  $x_F$  is numerically almost equal to the longitudinal momentum fraction  $x_L$  used in previous publications. There,  $x_L$  was defined as  $x_L = E_{n,\gamma}/E_p$ , where  $E_p$ ,  $E_n$  and  $E_\gamma$  are the energies of the proton beam, the forward neutron and the forward photon in the laboratory frame, respectively.

### 3. Cosmic Ray Air Shower MC Models


The predictions of several models used for the simulation of cosmic ray air showers are compared to the measured cross sections: EPOS LHC [14], QGSJET 01 [15], QGSJET II-04 [16] and SIBYLL 2.1 [17]. These models have been developed over many years, and are based both on older (Regge theory, Gribov's Regge Calculus) and newer concepts, such as perturbative QCD. Central elements are mini-jets production and colour string fragmentation into hadrons. The programs differ in the internal treatment of these concepts, as well as in the treatment of fragmentation, saturation, multi-parton interactions, hadron remnant, etc..

Originally, these programs model hadron (proton, nuclei) interactions. For the purpose of comparison with the present data, all programs have been interfaced to the HERA  $ep$  scattering kinematics via the PHOJET [18] program. All program simulations used for the comparison with the data have been provided by the respective authors<sup>2</sup>; no adjustment of internal parameters was made from the H1 side.

## 4. Results

### 4.1 The $W$ -Dependence of the Cross Sections

Figure 2 shows the  $W$ -dependence of the normalised cross sections, for photons (Figure 2a) and neutrons (Figure 2b). The cross sections are within uncertainties constant, at values of  $\sim 0.027$  and  $\sim 0.083$ , respectively. The predictions of the CR Models for the photon rate are all about 30-40% too high and are not constant but show a falling  $W$ -dependence. Also in the case of the neutron data, none of the CR models describe the data rate, although EPOS LHC and QGSJET II-04 are closer than SIBYLL and QGSJET 01; a slight  $W$ -dependence is also here seen in all predictions, although weaker than in the photon case.



**Figure 2:** The fraction of DIS events with forward photons (a) and forward neutrons (b) as a function of  $W$ . The CR model predictions are compared to the data.


### 4.2 The $x_F$ -dependence of the cross sections

The normalised cross sections differential in  $x_F$  are shown for photons in Figure 3a and for neutrons in Figure 3b. Only the distributions in the  $W$ -interval 70 – 130 GeV are given here, the distributions in the two higher  $W$ -intervals are similar.

<sup>2</sup>Thanks to T.Pierog, R.Engel and S.Ostapchenko.

The CR model predictions for photons, also shown in Figure 3a, vary slightly for the  $x_F$ -dependence, but are in general above the data for low values of  $x_F$ . In the neutron  $x_F$ -dependence, Figure 3b, the CR model predictions vary greatly and again EPOS LHC and QGSJET II-04 are closer to the data than SIBYLL and QGSJET 01.

Since the CR models were concipated for hadronic interactions, a modified version of QGSJET 01, denoted “QGSJET 01 (no mi)” was also compared to the data; in this version the effects of multi-parton interactions are switched off, with the idea that this could be more appropriate for the  $ep$  DIS environment. However, while the “no mi” version shows almost no change in the case of the photon data, the description of the neutron  $x_F$ -data is not only not improved, but the dependence instead becomes significantly harder at large values of  $x_F$ .




**Figure 3:** Normalised cross sections of forward photon (a) and forward neutron (b) production in DIS, as a function of  $x_F$  in the  $W$ -interval 70 – 130 GeV. The CR model predictions are compared to the data.

### 4.3 Test of Feynman Scaling


The measured cross sections, differential in  $W$  and  $x_F$ , offer the possibility of performing a direct experimental test of the Feynman Scaling hypothesis [19]. For this purpose the  $d\sigma/dx_F$ -distributions in the two higher  $W$ -intervals, 130-190 and 190-245 GeV, are divided with the corresponding  $d\sigma/dx_F$ -distributions in the lowest  $W$ -interval, 70-130 GeV. The resulting ratios, as a function of  $x_F$ , are shown for photons in Figures 4a,b and for neutrons in Figures 5a,b. Within errors the ratios are compatible with unity, i.e. the  $x_F$ -distributions for both neutrons and photons stay practically unchanged with increasing  $W$ , confirming the approximate validity of the hypothesis. This is the first direct experimental test of Feynman Scaling for forwardly produced neutrons and photons at HERA.

The predictions of the CR models show here a mixed behaviour. All models, with exception of SIBYLL 2.1, are compatible with Feynman Scaling in the case of neutron production. However, in the case of photon production, all models violate the scaling hypothesis, showing decreasing ratios with increasing  $W$ . This effect is strongest for SIBYLL 2.1 and the QGSJET models, while EPOS LHC is closer to the data.

In previous analyses [3, 4] the neutron and photon data were shown to confirm the hypothesis of Limiting Fragmentation [20]. The hypotheses of Limiting Fragmentation and of Feynman Scaling both stem from the same time, late 1969, and both hypotheses were put forward independently. They are based on the same simple fact, namely the Lorentz contraction of the beam projectile, and both hypotheses aim at finding regularities in multi-particle production at high energies. While the hypothesis of Limiting Fragmentation states that single particle momentum distributions at high



**Figure 4:** Ratios of normalised cross sections of forward photon production in DIS corresponding to two different  $W$  intervals, as a function of  $x_F$ : respectively the ratio of the cross section in the  $130 < W < 190$  GeV (a) and  $190 < W < 245$  GeV (b) interval to the cross section in the  $70 < W < 130$  GeV interval. Predictions of the CR models are compared to the data.



**Figure 5:** Ratios of normalised cross sections of forward neutron production in DIS corresponding to two different  $W$  intervals, as a function of  $x_F$ : respectively the ratio of the cross section in the  $130 < W < 190$  GeV (a) and  $190 < W < 245$  GeV (b) interval to the cross section in the  $70 < W < 130$  GeV interval. Predictions of the CR models are compared to the data.

energy are limited by functions solely dependent on transverse momentum and rapidity, Feynman Scaling states independence of particle distributions at high energy when expressed as functions of transverse momentum and Feynman- $x$  only. Since the variables rapidity and Feynman- $x$  are related<sup>3</sup>, the two hypotheses are equivalent. It is therefore not unexpected that the same neutron and photon data in the present direct experimental test confirm the hypothesis of Feynman Scaling.

## 5. Question from the Audience

**Q:** Can this comparison of model calculations with data tell us something about the validity of the assumptions in the various models and help to improve the models ?

**A:** These models have all a long development history. See e.g. the recent overview given by T.Pierog [21]. In fact, the model which in the present comparison with data gives the best result, namely EPOS LHC, has recently been improved, using new data from the LHC experiments. In earlier versions, the program did not compare so well with data. Thus, experimental data at high energies are a great help in the development of these Cosmic Ray air shower models.

<sup>3</sup>The relation between  $x_F$  and rapidity  $y$  is  $x_F = 2\mu/W \sinh(y)$ ,  $\mu = \sqrt{p_t^2 + m^2}$ .

## 6. Acknowledgement

Many thanks to all colleagues in H1, in particular to A.Buniatyan and H.Zohrabyan, for providing the material in this report and for help given in its preparation. Warm thanks also to the Vienna team for the excellent organisation and the pleasant atmosphere of the conference.

## References

- [1] R. Engel, Nucl. Phys. Proc. Suppl. **75A** (1999) 62 [astro-ph/9811225].
- [2] C. Adloff *et al.* [H1 Collaboration], Eur. Phys. J. **C6** (1999) 587 [hep-ex/9811013];  
S. Chekanov *et al.* [ZEUS Collaboration], Nucl. Phys. **B637** 3 (2002) 3 [hep-ex/0205076];  
S. Chekanov *et al.* [ZEUS Collaboration], Nucl. Phys. **B776** (2007) 1 [hep-ex/0702028];  
S. Chekanov *et al.* [ZEUS Collaboration], JHEP **06** (2009) 074 [arXiv:0812.2416].
- [3] F. Aaron *et al.* [H1 Collaboration], Eur. Phys. J. **C68** (2010) 381 [arXiv:1001.0532].
- [4] F. Aaron *et al.* [H1 Collaboration], Eur. Phys. J. **C71** (2011) 1771 [arXiv:1106.5944].
- [5] A. Buniatyan *et al.*, "Cosmic Rays, HERA and the LHC: Working group summary." Proceedings of the Workshop on the Implications of HERA for LHC Physics, Geneva, Switzerland, 26-30 May 2008, p.566, DESY-PROC-2009-02 [arXiv:0903.3861].
- [6] V. Andreev *et al.* [H1 Collaboration], DESY 14-035 [arXiv:1404.0201].
- [7] K. Charchula, G. A. Schuler and H. Spiesberger, Comput. Phys. Commun. **81** (1994) 381.
- [8] G. Ingelman, A. Edin and J. Rathsmann, Comput. Phys. Commun. **101** (1997) 108 [hep-ph/9605286];  
A. Edin, G. Ingelman and J. Rathsmann, Phys. Lett. **B366** (1996) 371 [hep-ph/9508386].
- [9] L. Lönnblad, Comput. Phys. Commun. **71** (1992) 15.
- [10] H. Jung, Comp. Phys. Commun. **86** (1995) 147.
- [11] H. Holtmann *et al.*, Phys. Lett. **B338** (1994) 363.
- [12] A. Kwiatkowski, H. Spiesberger and H. J. Möhring, Comp. Phys. Commun. **69** (1992) 155.
- [13] B. Andersson *et al.*, Phys. Rep. **97** (1983) 31; T. Sjöstrand, [hep-ph/9508391].
- [14] K. Werner, F.-M. Liu and T. Pierog, Phys. Rev. **C74** (2006) 044902 [hep-ph/0506232];  
T. Pierog *et al.*, DESY-13-125 [arXiv:1306.0121].
- [15] N. N. Kalmykov and S. S. Ostapchenko, Phys. Atom. Nucl. **56** (1993) 346;  
N. N. Kalmykov, S. S. Ostapchenko and A. I. Pavlov, Nucl. Phys. Proc. Suppl. **52B** (1997) 17.
- [16] S. S. Ostapchenko, Phys. Rev. **D74** (2006) 014026 [hep-ph/0505259];  
S. S. Ostapchenko, Phys. Rev. **D83** (2011) 014018 [arXiv:1010.1869].
- [17] J. Engel *et al.*, Phys. Rev. **D46** (1992) 5013;  
E.-J. Ahn *et al.*, Phys. Rev. **D80** (2009) 094003 [arXiv:0906.4113].
- [18] R. Engel and J. Ranft, Phys. Rev. **D54** (1996) 4244 [hep-ph/9509373].
- [19] R. P. Feynman, Phys. Rev. Lett. **23** (1969) 1415.
- [20] J. Benecke *et al.*, Phys. Rev. **188** (1969) 2159;  
T. T. Chou and C.-N. Yang, Phys. Rev. **D50** (1994) 590.
- [21] T. Pierog, "Modelling hadronic interactions in cosmic ray Monte Carlo generators",  
Proceedings ISVHECRI 2014, <http://inspirehep.net/record/1389774>