

Double parton scattering: collider physics meets hadron structure

M. Diehl

Deutsches Elektronen-Synchrotron DESY

Berkeley Summer Program, 15 August 2013





Hadron-hadron collisions

- ▶ standard description based on **factorization formulae**

cross sect = parton distributions $\times$ parton-level cross sect

example: Z production

$$pp \rightarrow Z + X \rightarrow \ell^+ \ell^- + X$$



- ▶ factorization formulae are for **inclusive** cross sections $pp \rightarrow Y + X$ where Y = produced in parton-level scattering, specified in detail
 X = summed over, no details



Hadron-hadron collisions

- ▶ standard description based on **factorization formulae**

cross sect = parton distributions $\times$ parton-level cross sect

example: Z production

$$pp \rightarrow Z + X \rightarrow \ell^+ \ell^- + X$$



- ▶ factorization formulae are for **inclusive** cross sections $pp \rightarrow Y + X$ where Y = produced in parton-level scattering, specified in detail
 X = summed over, no details
- ▶ have also interactions between “spectator” partons
 their effects cancel in inclusive cross sections **thanks to unitarity**
but they affect the final state (**namely X**)

Multiparton interactions (MPI)




- ▶ secondary (and tertiary etc.) interactions generically take place in hadron-hadron collisions
- ▶ predominantly low- p_T scattering
 $\leadsto$ underlying event (UE)
- ▶ at high collision energy (Tevatron, LHC) can be hard
 $\leadsto$ multiple hard scattering
- ▶ many studies:
 - theory: phenomenology, theoretical foundations (1980s, recent activity)
 - experiment: ISR, SPS, HERA (photoproduction), Tevatron, LHC
 - Monte Carlo generators: Pythia, Herwig++, Sherpa
- ▶ expected to be important for many processes at LHC
 see e.g. workshops: <http://mpi11.desy.de>; MPI@LHC 2012, CERN

Relevance for LHC

example: $pp \rightarrow H + Z \rightarrow b\bar{b} + Z$

Del Fabbro, Treleani 1999


- multiple interactions contribute to signal and background



same for $pp \rightarrow H + W \rightarrow b\bar{b} + W$

study for Tevatron: Bandurin et al, 2010

Double vs. single hard scattering



- ▶ double hard scattering:
net p_T of produced system (Z or $b\bar{b}$ pair) $\ll$ hard scale Q (e.g. M_Z)
- ▶ single hard scattering:
 p_T distribution up to values $\sim Q$
- ▶ no generic suppression for transv. mom. $\ll Q$:


$$\frac{d\sigma_{\text{single}}}{d^2\mathbf{p}_Z d^2\mathbf{p}_{b\bar{b}}} \sim \frac{d\sigma_{\text{double}}}{d^2\mathbf{p}_Z d^2\mathbf{p}_{b\bar{b}}} \sim \frac{\Lambda^2}{Q^2}$$

but since single scattering populates larger phase space:

$$\sigma_{\text{single}} \sim \frac{1}{Q^2} \gg \sigma_{\text{double}} \sim \frac{\Lambda^2}{Q^4}$$


- ▶ in addition: double scattering enhanced at small x

Space-time structure



- ▶ transverse parton momenta **not** the same in amplitude $\mathcal{A}$ and in $\mathcal{A}^*$
 cross section $\propto \int d^2\mathbf{r} F(x_i, \mathbf{k}_i, \mathbf{r}) F(\bar{x}_i, \bar{\mathbf{k}}_i, -\mathbf{r})$
- ▶ Fourier trf to impact parameter: $F(x_i, \mathbf{k}_i, \mathbf{r}) \rightarrow F(x_i, \mathbf{k}_i, \mathbf{y})$
 cross section $\propto \int d^2\mathbf{y} F(x_i, \mathbf{k}_i, \mathbf{y}) F(\bar{x}_i, \bar{\mathbf{k}}_i, \mathbf{y})$
- ▶ interpretation: $\mathbf{y}$ = transv. dist. between two scattering partons
 = equal in both colliding protons

Cross section formula



$$\frac{d\sigma_{\text{double}}}{dx_1 d\bar{x}_1 dx_2 d\bar{x}_2} = \frac{1}{C} \hat{\sigma}_1 \hat{\sigma}_2 \int d^2\mathbf{y} F(x_1, x_2, \mathbf{y}) F(\bar{x}_1, \bar{x}_2, \mathbf{y})$$

C = combinatorial factor

$\hat{\sigma}_i$ = parton-level cross sections

$F(x_1, x_2, \mathbf{y})$ = double parton distribution (DPD)

$\mathbf{y}$ = transv. distance between partons

- ▶ follows from Feynman graphs and hard-scattering approximation
no semi-classical approximation required
- ▶ can make $\hat{\sigma}_i$ differential in further variables (e.g. for jet pairs)
- ▶ can extend $\hat{\sigma}_i$ to higher orders in α_s
get usual convolution integrals over x_i in $\hat{\sigma}_i$ and F

Paver, Treleani 1982, 1984; Mekhfi 1985, . . . , MD, Ostermeier, Schäfer 2012

Cross section formula

- ▶ for measured transv. momenta

$$\frac{d\sigma}{dx_1 d\bar{x}_1 d^2\mathbf{q}_1 dx_2 d\bar{x}_2 d^2\mathbf{q}_2} = \frac{1}{C} \hat{\sigma}_1 \hat{\sigma}_2$$


$$\times \left[\prod_{i=1}^2 \int d^2\mathbf{k}_i d^2\bar{\mathbf{k}}_i \delta(\mathbf{q}_i - \mathbf{k}_i - \bar{\mathbf{k}}_i) \right] \int d^2\mathbf{y} F(x_i, \mathbf{k}_i, \mathbf{y}) F(\bar{x}_i, \bar{\mathbf{k}}_i, \mathbf{y})$$

- ▶ $F(x_i, \mathbf{k}_i, \mathbf{y}) = k_T$ dependent two-parton distribution


- ▶ interpretation as **Wigner function**:

$\mathbf{k}_1, \mathbf{k}_2 =$ transv. parton momenta **averaged over $\mathcal{A}$ and $\mathcal{A}^*$**

$\mathbf{y} =$ transv. distance between partons **averaged over $\mathcal{A}$ and $\mathcal{A}^*$**



Cross section formula



- ▶ for measured transv. momenta

$$\frac{d\sigma}{dx_1 d\bar{x}_1 d^2\mathbf{q}_1 dx_2 d\bar{x}_2 d^2\mathbf{q}_2} = \frac{1}{C} \hat{\sigma}_1 \hat{\sigma}_2$$

$$\times \left[\prod_{i=1}^2 \int d^2\mathbf{k}_i d^2\bar{\mathbf{k}}_i \delta(\mathbf{q}_i - \mathbf{k}_i - \bar{\mathbf{k}}_i) \right] \int d^2\mathbf{y} F(x_i, \mathbf{k}_i, \mathbf{y}) F(\bar{x}_i, \bar{\mathbf{k}}_i, \mathbf{y})$$

- ▶ $F(x_i, \mathbf{k}_i, \mathbf{y}) = k_T$ dependent two-parton distribution
- ▶ operator definition as for TMDs

$$F(x_i, \mathbf{k}_i, \mathbf{y}) = \mathcal{FT}_{z_i \rightarrow (x_i, \mathbf{k}_i)} \langle p | \bar{q}(-\frac{1}{2}z_2) \Gamma_2 q(\frac{1}{2}z_2) \bar{q}(\mathbf{y} - \frac{1}{2}z_1) \Gamma_1 q(\mathbf{y} + \frac{1}{2}z_1) | p \rangle$$

- essential for studying factorization, scale dependence, etc.
- similar def for $F(x_i, \mathbf{y})$

bilinear op's $\bar{q} \Gamma_i q$ at different transv. positions

⇒ not a twist-four operator but product of two twist-two operators

Pocket formula

- ▶ if two-parton density factorizes as

$$F(x_1, x_2, \mathbf{y}) = f(x_1) f(x_2) G(\mathbf{y})$$

where $f(x_i)$ = usual PDF

- ▶ if assume same $G(\mathbf{y})$ for all parton types
then cross sect. formula turns into


$$\frac{d\sigma_{\text{double}}}{dx_1 d\bar{x}_1 dx_2 d\bar{x}_2} = \frac{1}{C} \frac{d\sigma_1}{dx_1 d\bar{x}_1} \frac{d\sigma_2}{x_2 \bar{x}_2} \frac{1}{\sigma_{\text{eff}}}$$

with $1/\sigma_{\text{eff}} = \int d^2\mathbf{y} G(\mathbf{y})^2$

↪ scatters are completely independent

- ▶ underlies bulk of phenomenological estimates
and implementation of MPI in Monte Carlo generators
- ▶ pocket formula **fails** if any of the above assumptions is invalid
and if further terms must be added to original expression of cross sect.

Experimental investigations



- ▶ also recent studies in double charm production ($c\bar{c}c\bar{c}$), LHCb data

Parton correlations

- ▶ if neglect correlations between two partons

$$F(x_1, x_2, \mathbf{y}) = \int d^2 \mathbf{y}' f(x_1, \mathbf{y}' + \mathbf{y}) f(x_2, \mathbf{y}')$$

where $f(x_i, \mathbf{y}) =$ impact parameter dependent single-parton density

and if neglect correlations between x and $\mathbf{y}$ of single parton

$$f(x_i, \mathbf{y}) = f(x_i) F(\mathbf{y})$$

with same $F(\mathbf{y})$ for all partons

then $G(\mathbf{y}) = \int d^2 \mathbf{y}' F(\mathbf{y}') F(\mathbf{y}' + \mathbf{y})$

$$\left| \begin{array}{c} \text{Diagram 1: Single vertex with } x_1, x_2 \text{ and } y \end{array} \right|^2 \approx \int d^2 b \left| \begin{array}{c} \text{Diagram 2: Two vertices with } x_1, x_2 \text{ and } y' \end{array} \right|^2 \times \left| \begin{array}{c} \text{Diagram 3: Two vertices with } x_1, x_2 \text{ and } y' + y \end{array} \right|^2$$

Parton correlations

- ▶ if neglect correlations between two partons

$$F(x_1, x_2, \mathbf{y}) = \int d^2 \mathbf{y}' f(x_1, \mathbf{y}' + \mathbf{y}) f(x_2, \mathbf{y}')$$

where $f(x_i, \mathbf{y}) =$ impact parameter dependent single-parton density

and if neglect correlations between x and $\mathbf{y}$ of single parton

$$f(x_i, \mathbf{y}) = f(x_i) F(\mathbf{y})$$

with same $F(\mathbf{y})$ for all partons

then $G(\mathbf{y}) = \int d^2 \mathbf{y}' F(\mathbf{y}') F(\mathbf{y}' + \mathbf{y})$

- ▶ for Gaussian $F(\mathbf{y})$ with average $\langle \mathbf{y}^2 \rangle$

$$\sigma_{\text{eff}} = 4\pi \langle \mathbf{y}^2 \rangle = 41 \text{ mb} \times \langle \mathbf{y}^2 \rangle / (0.57 \text{ fm})^2$$

determinations of $\langle \mathbf{y}^2 \rangle$ from GPDs and form factors: $(0.57 \text{ fm} - 0.67 \text{ fm})^2$

is $\gg \sigma_{\text{eff}} \sim 10$ to 20 mb from experimental extractions

if $F(\mathbf{y})$ is Fourier trf. of dipole then $41 \text{ mb} \rightarrow 36 \text{ mb}$

complete independence between two partons is disfavored


or something is seriously wrong with σ_{eff}

cf. Calucci, Treleani 1999; Frankfurt, Strikman, Weiss 2003-04; Blok et al 2013

Correlations involving x

- ▶ $F(x_1, x_2, \mathbf{y}) = f(x_1) f(x_2) G(\mathbf{y})$ cannot hold for all x_1, x_2
- ▶ most obvious: energy conservation $\Rightarrow x_1 + x_2 \leq 1$
often used: $F(x_1, x_2, \mathbf{y}) = f(x_1) f(x_2) (1 - x_1 - x_2)^n G(\mathbf{y})$
to suppress region of large $x_1 + x_2$
- ▶ significant $x_1 - x_2$ correlations found in constituent quark model

Rinaldi, Scopetta, Vento: [arXiv:1302.6462](https://arxiv.org/abs/1302.6462)



plot shows $\int d^2\mathbf{y} F_{uu}(x_1, x_2, \mathbf{y}) / f_u(x_2)$
is x_2 independent if factorization holds

- ▶ unknown: size of correlations when one or both of x_1, x_2 small

Correlations involving x and $\mathbf{y}$

- ▶ single-parton distribution $f(x, \mathbf{y})$ is Fourier trf. of generalized parton distributions at zero skewness

↪ information from exclusive processes and theory

- ▶ HERA results on $\gamma p \rightarrow J/\Psi p$ give

$$\langle \mathbf{y}^2 \rangle \propto \text{const} + 4\alpha' \log(1/x)$$

with $\alpha' \approx 0.15 \text{ GeV}^{-2} = (0.08 \text{ fm})^2$ for gluons at $x \sim 10^{-3}$

- ▶ lattice simulations $\rightarrow$ strong decrease of $\langle \mathbf{y}^2 \rangle$ with x above ~ 0.1 seen by comparing moments $\int dx x^{n-1} f(x, \mathbf{y})$ for $n = 0, 1, 2$
- ▶ expect similar correlations between x_i and $\mathbf{y}$ in two-parton dist's even if factorization $F(x_1, x_2, \mathbf{y}) = f(x_1, \mathbf{y}) f(x_2, \mathbf{y})$ does not hold
- ▶ in double parton scattering $\mathbf{y}$ unobservable:
$$d\sigma \propto \int d^2\mathbf{y} F(x_i, \mathbf{y}) F(\bar{x}_i, \mathbf{y})$$
in $f(x, \mathbf{y})$ impact parameter is Fourier conjugate to measurable momentum transfer

Correlations involving x and y

- ▶ single-parton distribution $f(x, \mathbf{y})$ is Fourier trf. of generalized parton distributions at zero skewness

↪ information from exclusive processes and theory

- ▶ HERA results on $\gamma p \rightarrow J/\Psi p$ give


$$\langle \mathbf{y}^2 \rangle \propto \text{const} + 4\alpha' \log(1/x)$$

with $\alpha' \approx 0.15 \text{ GeV}^{-2} = (0.08 \text{ fm})^2$ for gluons at $x \sim 10^{-3}$

- ▶ lattice simulations → strong decrease of $\langle \mathbf{y}^2 \rangle$ with x above ~ 0.1 seen by comparing moments $\int dx x^{n-1} f(x, \mathbf{y})$ for $n = 0, 1, 2$
- ▶ expect similar correlations between x_i and $\mathbf{y}$ in two-parton dist's even if factorization $F(x_1, x_2, \mathbf{y}) = f(x_1, \mathbf{y}) f(x_2, \mathbf{y})$ does not hold


Consequence for multiple interactions:

- ▶ if interaction 1 produces high-mass system
 - have large $x_1, \bar{x}_1$
 - smaller $\mathbf{y}$, more central collision
 - secondary interactions enhanced



Frankfurt, Strikman, Weiss 2003, study in Pythia: Corke, Sjöstrand 2011

Spin correlations



- ▶ polarizations of two partons can be correlated even in unpolarized target already pointed out by Mekhfi (1985)
 - ▶ quarks: longitudinal and transverse pol.
 - ▶ gluons: longitudinal and linear pol.

have eight (k_T integrated leading-twist) distributions for each combination qq , qg , gg

- ▶ fulfill positivity constraints analogous to Soffer bound for usual PDFs, e.g.


$$F_{qq} - F_{\Delta q \Delta q} \geq 2|F_{\delta q \delta q}|$$

$q = \text{unpol.}$, $\Delta q = \text{long.}$, $\delta q = \text{transv.}$; schematic notation

MD, Kasemets 2013

- ▶ in general not suppressed in hard scattering consequences for rate and distributions

Spin and angular distributions



- ▶ detailed calc'n for gauge boson pair production followed by leptonic decay

T. Kasemets, MD 2012; see also A. Manohar, W. Waalewijn 2011


- ▶ longitudinal quark spin correlations
 - ↪ overall rate and distribution in lepton rapidities
- ▶ transverse quark spin correlations
 - ↪ azimuthal correlation between lepton planes
 - ↪ two hard scatters are not independent
- ▶ expect similar effects for gluon initiated processes (esp. for jets)
 - linear gluon pol. ↪ azimuthal correlation between scattering planes
- ▶ **note:** independent scattering planes sometimes assumed as **criterion** to characterize double parton scattering

Spin correlations

- how important are spin correlations?

large effects expected in valence quark region

study in bag model: Chang, Manohar, Waalewijn: [arXiv:1211:3132](https://arxiv.org/abs/1211.3132)



plots show $F(x_1, x_2 = 0.4, k_\perp)$ for different pol. combinations

$k_\perp$ = Fourier conjugate to $\mathbf{y}$


- unknown: size of correlations when one or both of x_1, x_2 small

Color structure

- quark lines in amplitude and its conjugate can couple to color singlet or octet:

$$^1F \rightarrow (\bar{q}_2 \mathbb{1} q_2) (\bar{q}_1 \mathbb{1} q_1)$$

$$^8F \rightarrow (\bar{q}_2 t^a q_2) (\bar{q}_1 t^a q_1)$$




- 8F describes color correlation between quarks 1 and 2
is essentially unknown (no probability interpretation as a guide)
- for two-gluon dist's more color structures: 1, 8_S , 8_A , 10, $\overline{10}$, 27
- for k_T integrated distributions:
color correlations suppressed by Sudakov logarithms

Mekhfi 1988


... but not necessarily negligible
for moderately hard scales

Manohar, Waalewijn arXiv:1202:3794

U = Sudakov factor, Q = hard scale



Scale and energy dependence



$$\frac{s d\sigma}{\prod_{i=1}^2 dx_i d\bar{x}_i d^2q_i} \quad \frac{s d\sigma}{\prod_{i=1}^2 dx_i d\bar{x}_i}$$

$$\frac{1}{\Lambda^2 Q^2}$$

$$1$$

$$\frac{1}{\Lambda^2 Q^2}$$

$$\frac{\Lambda^2}{Q^2}$$

$$\frac{1}{\Lambda^2 Q^2}$$

$$\frac{\Lambda^2}{Q^2}$$

- ▶ interference between single and double scattering:
 - leading power when differential in $\mathbf{q}_i$
 - power suppressed when $\int d^2\mathbf{q}_i$, **twist-three parton distributions**


- ▶ at small $x_1 \sim x_2 \sim x$ expect

- single scattering $\propto x^{-\lambda}$
- double scattering $\propto x^{-2\lambda}$
- interference? how do three-particle correlators behave for small x ?

$$\text{with } xf(x) \sim x^{-\lambda}$$

Behavior at small interparton distance

- ▶ for $y \ll 1/\Lambda$ in perturbative region $F(x_1, x_2, y)$ dominated by graphs with splitting of single parton



- ▶ find **strong** correlations in x_1, x_2 , spin and color between two partons
e.g. 100% correlation for longitudinal pol. of q and $\bar{q}$
- ▶ can compute short-distance behavior:

$$F(x_1, x_2, y) \sim \frac{1}{y^2} \text{ splitting fct} \otimes \text{usual PDF}$$

Scale evolution for distributions without color correlation

- ▶ if define two-parton distributions as operator matrix elements in analogy with usual PDFs

$$F(x_1, x_2, \mathbf{y}; \mu) \sim \langle p | \mathcal{O}_1(\mathbf{0}; \mu) \mathcal{O}_2(\mathbf{y}; \mu) | p \rangle \quad f(x; \mu) \sim \langle p | \mathcal{O}(\mathbf{0}; \mu) | p \rangle$$

where $\mathcal{O}(\mathbf{y}; \mu)$ = twist-two operator renormalized at scale μ

- ▶ $F(x_i, \mathbf{y})$ for $\mathbf{y} \neq \mathbf{0}$:

separate DGLAP evolution for partons 1 and 2

$$\frac{d}{d \log \mu} F(x_i, \mathbf{y}) = P \otimes_{x_1} F + P \otimes_{x_2} F$$

two independent parton cascades


- ▶ $\int d^2 \mathbf{y} F(x_i, \mathbf{y})$:

extra term from $2 \rightarrow 4$ parton transition

since $F(x_i, \mathbf{y}) \sim 1/\mathbf{y}^2$


Kirschner 1979; Shelest, Snigirev, Zinovev 1982

Gaunt, Stirling 2009; Ceccopieri 2011



- ▶ which evolution eq. is relevant for double hard scattering?

Deeper problems with the splitting graphs





- ▶ contribution from splitting graphs in cross section
gives **divergent** integrals $\int d^2\mathbf{y} F(x_1, x_2, \mathbf{y}) F(\bar{x}_1, \bar{x}_2, \mathbf{y}) \sim \int d\mathbf{y}^2 / \mathbf{y}^4$
- ▶ **double counting** problem between double scattering with splitting and single scattering at loop level

MD, Ostermeier, Schäfer 2011; Gaunt, Stirling 2011; Gaunt 2012
 Blok, Dokshitzer, Frankfurt, Strikman 2011; Ryskin, Snigirev 2011, 2012
 same problem for jets: Cacciari, Salam, Sapeta 2009


- ▶ possible solution:
 subtract splitting contribution from two-parton dist's when $\mathbf{y}$ is small
 will also modify their scale evolution; remains to be worked out

Deeper problems with the splitting graphs



- ▶ contribution from splitting graphs in cross section gives **divergent** integrals $\int d^2\mathbf{y} F(x_1, x_2, \mathbf{y}) F(\bar{x}_1, \bar{x}_2, \mathbf{y}) \sim \int d\mathbf{y}^2 / \mathbf{y}^4$
- ▶ also have graphs with single PDF for one proton and double PDF for other


What is double parton scattering?



Blok et al, 2011-13


Sudakov factors

- ▶ for k_T dependent distributions, i.e. measured q_i :
Sudakov logarithms for **all** color channels
close relation with physics of parton showers
- ▶ for double Drell-Yan process
can adapt **Collins-Soper-Sterman** formalism for single Drell-Yan
 $\rightsquigarrow$ include and resum Sudakov logs in k_T dependent parton dist's
MD, D Ostermeier, A Schäfer 2011
for jet production inherit problems of usual TMD factorization
- ▶ at leading double log accuracy: singlet and octet dist's 1F and 8F
have **same** Sudakov factor as in single scattering
- ▶ beyond double log: Sudakov factors mix singlet and octet dist's



Factorization?

- ▶ open problem (for TMD and collinear formulations):
exchange of gluons in Glauber region



Not discussed in this talk:

- ▶ multiparton interactions in pA collisions
- ▶ small- x approach
connection with diffraction, AGK rules
ridge effect in pp and pA

Bartels, Salvadore, Vacca 2008

Dumitru et al 2011; ...

Conclusions

- ▶ multiple hard scattering is **not** generically suppressed in sufficiently differential cross sections
- ▶ current phenomenology relies on strong simplifications
- ▶ have several elements for a formulation of factorization but important open questions still unsolved
 - ▶ crosstalk with single hard scattering at small distances
 - ▶ Glauber gluon exchange
- ▶ double hard scattering depends on detailed **hadron structure** including correlation and interference effects
 - ▶ corresponding nucleon matrix elements largely unknown theoretical activity only started
 - ▶ transverse distance between partons essential
 $\rightsquigarrow$ (yet another aspect of) $2 + 1$ dim. proton structure
 - ▶ parton imaging from *ep* scattering can provide quantitative baseline for **uncorrelated** parton distributions
- ▶ LHC has potential for studying double hard scattering in great detail