

# PDF Constraints and $\alpha_s$ Results from CMS

Engin Eren (DESY)  
on behalf of CMS Collaboration


*DIS 2016, DESY  
Hamburg*

# Particle production in proton-proton collisions

proton structure

hard interaction

Factorization: proton structure  $\otimes$  sub-process ME



$$\sigma(s) = \sum_{i,j} \int_{\tau_0}^1 \left[ \frac{d\tau}{\tau} \cdot \frac{dL_{ij}(\mu_F^2)}{d\tau} \right] \cdot \hat{s} \cdot \hat{\sigma}_{ij}$$

Ingredients for SM predictions for pp@LHC:

- partonic cross section calculable in pQCD
- parton luminosity:

$$\tau \cdot \frac{dL_{ij}}{d\tau} \propto \int_0^1 dx_1 dx_2 (x_1 f_i(x_1, \mu_F^2) \cdot x_2 f_j(x_2, \mu_F^2)) + (1 \leftrightarrow 2) \delta(\tau - x_1 x_2)$$

Parton Distribution Functions (PDFs)

universal functions of partonic fraction  $x$  of proton momentum  
and energy scale  $Q$  of the process

***Precision of PDFs essential for interpretation of the LHC measurements***


***Precise LHC data are used to improve PDF constraints***

# How are PDFs determined ?

PDF for flavor  $i$ :  $f_i = f_i(x, Q^2)$ :

$Q^2$  dependence predicted by QCD

$x$ -dependence determined from data




**LHC provides important constraints in particular at large  $x$**

**Example of PDF determination:**



- parameterize  $x$ -shape at a scale  $Q^2_0$ :  

$$f(x) = Ax^B(1-x)^C(1+Dx+Ex^2)$$
- evolve these PDFs to  $Q^2 > Q^2_0$  (e.g. DGLAP)
- construct expected cross sections
- $\chi^2$ - fit to the experimental data



# Asymmetry in $W^\pm$ production at CMS

- $W$  production probes bi-linear quark combinations



Muon asymmetry in  $W$  production  
probes quark distributions  
at  $10^{-3} < x < 10^{-1}$

$$A(\eta) = \frac{\sigma_\eta^+ - \sigma_\eta^-}{\sigma_\eta^+ + \sigma_\eta^-} \approx \frac{u_v - d_v}{u_v + d_v + 2u_{sea}}$$

# QCD analyses of W charge asymmetry data

QCD analysis at NNLO, parton evolution in  $Q^2$  via DGLAP implemented in QCDNUM.

xFitter version 1.1.1 is used.

## Data in the QCD analysis:

- HERA I+II combined inclusive DIS data, Charged and Neutral Current [[Eur. Phys. J. C 75 \(2015\) 2604](#)]
- Investigated CMS data :
  - W charge asymmetry measurement with muons @ 8 TeV [[arXiv:1603.01803](#)]

PDF Uncertainties : Quadratic sum of experimental, model and parametrisation errors.

Experimental uncertainties: originate from uncertainties of the data

- Hessian error estimate: criterion  $\Delta\chi^2=1$  is applied

Model uncertainties: originate from variations of model input parameters:

- Fraction of strange quarks in the sea  $f_s=0.31\pm0.08$
- Values of charm and beauty quark masses.
- $Q^2$  cut on inclusive DIS data.

Parametrisation uncertainties:

- Originate from variations on assumed parametrization

# PDF parametrization and $\chi^2$

Basic parametrization at the starting scale  $Q^2_0=1.9 \text{ GeV}^2$  :

## W Charge Asymmetry (13p)

$$xg(x) = A_g x^{B_g} \cdot (1-x)^{C_g} (1 + D_g x)$$

$$xu_v(x) = A_{u_v} x^{B_{u_v}} \cdot (1-x)^{C_{u_v}} \cdot (1 + E_{u_v} x^2)$$

$$xd_v(x) = A_{d_v} x^{B_{d_v}} \cdot (1-x)^{C_{d_v}}$$

$$x\bar{U}(x) = A_{\bar{U}} x^{B_{\bar{U}}} \cdot (1-x)^{C_{\bar{U}}} (1 + E_{\bar{U}} x^2)$$

$$x\bar{D}(x) = A_{\bar{D}} x^{B_{\bar{D}}} \cdot (1-x)^{C_{\bar{D}}}$$

| Data sets                                                                             | Partial $\chi^2/n_{\text{dp}}$ |
|---------------------------------------------------------------------------------------|--------------------------------|
| HERA1+2 neutral current, $e^+p$ , $E_p = 920 \text{ GeV}$                             | 440/377                        |
| HERA1+2 neutral current, $e^+p$ , $E_p = 820 \text{ GeV}$                             | 69/70                          |
| HERA1+2 neutral current, $e^+p$ , $E_p = 575 \text{ GeV}$                             | 214/254                        |
| HERA1+2 neutral current, $e^+p$ , $E_p = 460 \text{ GeV}$                             | 210/204                        |
| HERA1+2 neutral current, $e^-p$ , $E_p = 920 \text{ GeV}$                             | 218/159                        |
| HERA1+2 charged current, $e^+p$ , $E_p = 920 \text{ GeV}$                             | 46/39                          |
| HERA1+2 charged current, $e^-p$ , $E_p = 920 \text{ GeV}$                             | 50/42                          |
| CMS $W^\pm$ muon charge asymmetry $\mathcal{A}(\eta\mu)$ , $\sqrt{s} = 8 \text{ TeV}$ | 3/11                           |
| Correlated $\chi^2$                                                                   | 141                            |
| Global $\chi^2/n_{\text{dof}}$                                                        | 1391/1143                      |

Normalization parameters  $A$  are determined by QCD sum rules

$B$ : define low- $x$  behaviour,  $C$ : high- $x$  shape

## Parametrization uncertainties :

originate from variations on assumed parametrization, in which additional parameters are added one by-one in the functional form of the parametrization;  
additional variation of 1.  $5 < Q^2_0 < 2.5 \text{ GeV}^2$

Largest difference of resulting PDFs to the central result (envelope) is assigned as uncertainty

# Asymmetry in $W^\pm$ production at CMS in PDF studies



## QCD analysis at NNLO

**Data:** Combined HERA I+II DIS data [[Eur. Phys. J. C 75 \(2015\) 2604](#)]  
+ CMS muon charge asymmetry measurement at 8-TeV [[arXiv:1603.01803](#)]

**Theory for  $A_W$ :** NLO prediction with MCFM, interfaced via APPLGRID

K-factors, which are calculated by FEWZ, are applied


QCD scales  $\mu_r = \mu_f = m_W$ , strong coupling  $\alpha_s(m_Z)=0.118$ ;




*Improvement in the uncertainty of the valence-quark distributions*

# Probing QCD with Jet Cross Sections

- Jet production sensitive to quark and gluon distributions, and to  $\alpha_s$




Strong correlation between inclusive jet cross section and gluon at high-middle  $x$




Eur. Phys. J. C 75 (2015) 288

Double differential inclusive jet cross section measurement (Paolo's Talk)



cross-section ratios for different  $\sqrt{s}$  are also provided!

# Determination of strong coupling $\alpha_s$ : inclusive jets @ 8TeV



CMS jet cross section measurement  
 $\sqrt{s}=8\text{TeV}$  (19.7 fb<sup>-1</sup>) CMS-PAS-SMP-14-001

compared to NLO QCD ⊗ PDFs  
in each bin of p<sub>T</sub> and y  
different sets of PDFs used  
each set has its α<sub>s</sub> - dependence


In each y bin, for each PDF, α<sub>s</sub> is determined by minimizing χ<sup>2</sup> between data and NLO

Similar results obtained with different PDFs



Using CT10NLO :

$$\alpha_s(M_Z) = 0.1164^{+0.0025}_{-0.0029}(\text{PDF})^{+0.0053}_{-0.0028}(\text{Scale}) \\ \pm 0.0001(\text{NP})^{+0.0014}_{-0.0015}(\text{Exp})$$

Analysis performed in 6 p<sub>T</sub> bins  
⇒ running of α<sub>s</sub>=α<sub>s</sub>(Q<sup>2</sup>)




# Determination of strong coupling $\alpha_s$ : inclusive jets @ 8TeV



Using CT10NLO :

$$\alpha_s(M_Z) = 0.1164^{+0.0025}_{-0.0029}(\text{PDF})^{+0.0053}_{-0.0028}(\text{Scale}) \pm 0.0001(\text{NP})^{+0.0014}_{-0.0015}(\text{Exp})$$

Analysis performed in 6  $p_T$  bins  
 $\Rightarrow$  running of  $\alpha_s = \alpha_s(Q^2)$



# QCD analyses of Inclusive Jets at 8 TeV.

QCD analysis at NLO, parton evolution in  $Q^2$  via DGLAP implemented in QCDNUM.

xFitter version 1.1.1 is used.

Data in the QCD analysis:

- HERA I+II combined inclusive DIS data, Charged and Neutral Current [Eur. Phys. J. C 75 (2015) 2604]
- Investigated CMS data :
  - Inclusive Jets @ 8 TeV [CMS-PAS-SMP-14-001]

PDF Uncertainties : Quadratic sum of experimental, model and parametrisation errors.

Experimental uncertainties: originate from uncertainties of the data

- Hessian error estimate: criterion  $\Delta\chi^2=1$  is applied
- MC Method (as a cross check)

Model uncertainties: originate from variations of model input parameters:

- Fraction of strange quarks in the sea  $f_s=0.31\pm0.08$
- Values of charm and beauty quark masses.
- $Q^2$  cut on inclusive DIS data.

Parametrisation uncertainties:

- Originate from variations on assumed parametrization

# QCD Analysis of Inclusive Jet Measurements at 8 TeV

Basic parametrization at the starting scale  $Q^2_0=1.9 \text{ GeV}^2$  :

Inclusive Jets @8TeV (18p)

$$xg(x) = A_g x^{B_g} \cdot (1-x)^{C_g} (1 + E_g x^2) - A'_g x^{B'_g} (1-x)^{C'_g}$$

$$xu_v(x) = A_{u_v} x^{B_{u_v}} \cdot (1-x)^{C_{u_v}} \cdot (1 + D_{u_v} x)$$

$$xd_v(x) = A_{d_v} x^{B_{d_v}} \cdot (1-x)^{C_{d_v}} \cdot (1 + D_{d_v} x)$$

$$x\overline{U}(x) = A_{\overline{U}} x^{B_{\overline{U}}} \cdot (1-x)^{C_{\overline{U}}} (1 + D_{\overline{U}} x)$$

$$x\overline{D}(x) = A_{\overline{D}} x^{B_{\overline{D}}} \cdot (1-x)^{C_{\overline{D}}} (1 + E_{\overline{D}} x^2)$$

| Data sets                      |                               | Partial $\chi^2/n_{\text{dp}}$ |
|--------------------------------|-------------------------------|--------------------------------|
| HERA1+2 Neutral Current        | $e^+p, E_p = 920 \text{ GeV}$ | 376/332                        |
| HERA1+2 Neutral Current        | $e^+p, E_p = 820 \text{ GeV}$ | 61/63                          |
| HERA1+2 Neutral Current        | $e^+p, E_p = 575 \text{ GeV}$ | 197/234                        |
| HERA1+2 Neutral Current        | $e^+p, E_p = 460 \text{ GeV}$ | 204/187                        |
| HERA1+2 Neutral Current        | $e^-p$                        | 219/159                        |
| HERA1+2 Charged Current        | $e^+p$                        | 41/39                          |
| HERA1+2 Charged Current        | $e^-p$                        | 50/42                          |
| CMS inclusive jets 8 TeV       | $0 < y < 0.5$                 | 53/36                          |
|                                | $0.5 < y < 1.0$               | 34/36                          |
|                                | $1.0 < y < 1.5$               | 35/35                          |
|                                | $1.5 < y < 2.0$               | 52/29                          |
|                                | $2.0 < y < 2.5$               | 49/24                          |
|                                | $2.5 < y < 3.0$               | 4.9/18                         |
| Correlated $\chi^2$            |                               | 94                             |
| Global $\chi^2/n_{\text{dof}}$ |                               | 1471/1216                      |

Normalization parameters  $A$  are determined by QCD sum rules

$B$ : define low- $x$  behaviour,  $C$ : high- $x$  shape

# Impact of the CMS jet measurements on PDFs



## QCD analysis at NLO

**Data:** combined HERA I+II DIS [Eur. Phys. J. C 75 (2015) 2604]

+ CMS inclusive jet production at 8 TeV,  $L = 19.71 \text{ fb}^{-1}$

**Theory for jet production in  $pp$ :** NLOJET++ version 4.1.3, interfaced via fastNLO


QCD scales  $\mu_r = \mu_f = p_{T \text{ jet}}$ , strong coupling  $\alpha_s(m_Z)=0.1180$ ;



*Improvement in the uncertainty of the gluon distributions at high-x*

# Estimation of PDF Uncertainty with MC Method

- 200 of replicas allowing the central values of cross-sections to fluctuate within their systematic and statistical uncertainties.
- For each replica, **NLO QCD** fit is performed. Errors on the PDFs are estimated from the RMS of the spread of the curves.



*Similar reduction of uncertainties observed as in hessian estimate*

# Summary

- W Charge Asymmetry (Muon) Data :
  - ★ Precision of the valence quark distributions improves.
- CMS Inclusive Jet Data :
  - ★ Precision of the gluon distributions improves.
- Prospects :
  - ★ Both data can be used in global QCD analysis by PDF collaborations.

# Backup Slides

# Framework for CMS QCD analyses in this talk

QCD analysis at NLO and NNLO, parton evolution in  $Q^2$  via DGLAP implemented in QCDNUM

Data in the QCD analysis:

- HERA I+II combined inclusive DIS data, Charged and Neutral Current [[arXiv:1506.06042](https://arxiv.org/abs/1506.06042)]
- Different CMS data sets (W charge asymmetry and Inclusive Jets @ 8TeV)

Experimental uncertainties: originate from uncertainties of the data, criterion  $\Delta\chi^2=1$  is applied

Model input:

- Theory calculations at NLO and NNLO appropriate for each data set
- Starting scale of PDF evolution  $Q_0^2 = 1.9 \text{ GeV}^2$
- Heavy quark treatment: general mass variable flavor number scheme by Thorne-Roberts (TR)
- Heavy quark masses:
  - NLO :  $m_c = 1.47 \text{ GeV}$ ,  $m_b = 4.50 \text{ GeV}$ .
  - NNLO :  $m_c = 1.43 \text{ GeV}$ ,  $m_b = 4.50 \text{ GeV}$ .

Model uncertainties: originate from variations of model input parameters:

- NLO (Incl. Jets) :  $1.41 \text{ GeV} < m_c < 1.53 \text{ GeV}$  and  $5 \text{ GeV}^2 < Q_{\min}^2 < 10 \text{ GeV}^2$
- NNLO (W asymmetry) :  $1.37 \text{ GeV} < m_c < 1.49 \text{ GeV}$  and  $2.5 \text{ GeV}^2 < Q_{\min}^2 < 5 \text{ GeV}^2$
- $4.25 \text{ GeV} < m_b < 4.75 \text{ GeV}$  same for both.

fraction of strange quarks in the sea  $f_s = 0.31 \pm 0.08$

# K-Factors and $\chi^2$ definitions


$$K = \frac{\hat{\sigma}_{NNLO} \otimes PDF(NNLO)}{\hat{\sigma}_{NLO} \otimes PDF(NNLO)}$$

$$\chi^2(m, b) = \sum_i \frac{[\mu_i - m_i(1 - \sum_j \gamma_j^i b_j)]^2}{\delta_{i,unc}^2 m_i^2 + \delta_{i,stat} \mu_i m_i (1 - \sum_j \gamma_j^i b_j)} + \sum_j b_j^2$$


$$\chi^2(m) = \sum_{i,k} (m_i - \mu_i) C_{ik}^{-1} (m_k - \mu_k)$$

# Probing QCD with Jet Cross Sections

Strong correlation between inclusive jet cross section and gluon at high-middle  $x$





Strong correlation between inclusive jet cross section and quark at high  $x$





The potential impact of the CMS inclusive jet data can be illustrated by the correlation between the inclusive jet cross section  $\sigma_{\text{jet}}(Q)$  and the PDF  $xf(x, Q^2)$  for any parton flavour  $f$ . The NNPDF Collaboration [63] provides PDF sets in the form of an ensemble of replicas  $i$ , which sample variations in the PDF parameter space within allowed uncertainties. The correlation coefficient  $\varrho_f(x, Q)$  between a cross section and the PDF for flavour  $f$  at a point  $(x, Q)$  can be computed by evaluating means and standard deviations from an ensemble of  $N$  replicas as

$$\varrho_f(x, Q) = \frac{N}{(N-1)} \frac{\langle \sigma_{\text{jet}}(Q)_i \cdot xf(x, Q^2)_i \rangle - \langle \sigma_{\text{jet}}(Q)_i \rangle \cdot \langle xf(x, Q^2)_i \rangle}{\Delta_{\sigma_{\text{jet}}(Q)} \Delta_{xf(x, Q^2)}}. \quad (12)$$

# Hessian vs MC



# u and d valence PDFs from inclusive jet @8TeV



# Parametrisation Uncertainty



- Get the largest difference
- Construct envelope

