

Higgs boson mass and new physics

Fedor Bezrukov,^{a,b} Mikhail Yu. Kalmykov,^c Bernd A. Kniehl^c and
Mikhail Shaposhnikov^d

^a*Physics Department, University of Connecticut,
Storrs, CT 06269-3046, U.S.A.*

^b*RIKEN-BNL Research Center, Brookhaven National Laboratory,
Upton, NY 11973, U.S.A.*

^c*II. Institut für Theoretische Physik, Universität Hamburg,
Luruper Chaussee 149, 22761, Hamburg, Germany*

^d*Institut de Théorie des Phénomènes Physiques, École Polytechnique Fédérale de Lausanne,
CH-1015 Lausanne, Switzerland*

E-mail: Fedor.Bezrukov@uconn.edu, mikhail.kalmykov@desy.de,
kniehl@desy.de, Mikhail.Shaposhnikov@epfl.ch

ABSTRACT: We discuss the lower Higgs boson mass bounds which come from the absolute stability of the Standard Model (SM) vacuum and from the Higgs inflation, as well as the prediction of the Higgs boson mass coming from the asymptotic safety of the SM. We account for the three-loop renormalization group evolution of the couplings of the SM and for a part of the two-loop corrections that involve the QCD coupling α_s to the initial conditions for their running. This is one step beyond the current state-of-the-art procedure (“one-loop matching–two-loop running”). This results in a reduction of the *theoretical* uncertainties in the Higgs boson mass bounds and predictions, associated with the SM physics, to 1–2 GeV. We find that with the account of existing experimental uncertainties in the mass of the top quark and α_s (taken at the 2σ level) the bound reads $M_H \geq M_{\min}$ (equality corresponds to the asymptotic-safety prediction), where $M_{\min} = (129 \pm 6)$ GeV. We argue that the discovery of the SM Higgs boson in this range would be in agreement with the hypothesis of the absence of new energy scales between the Fermi and Planck scales, whereas the coincidence of M_H with M_{\min} would suggest that the electroweak scale is determined by Planck physics. In order to clarify the relation between the Fermi and Planck scales a construction of an electron-positron or muon collider with a center-of-mass energy $\sim (200 + 200)$ GeV (Higgs and t-quark factory) would be needed.

KEYWORDS: Standard Model, Higgs Physics, Renormalization Group

ARXIV EPRINT: [1205.2893](https://arxiv.org/abs/1205.2893)

Contents

1	Introduction	1
2	The stability bound	4
2.1	The benchmark mass	4
2.2	Value of M_{\min}	5
3	M_{\min} and BSM physics	9
3.1	Asymptotic safety	10
3.2	M_{\min} and cosmology	12
3.3	Higgs inflation	13
4	Conclusions	14
A	$O(\alpha\alpha_s)$ electroweak corrections to the top Yukawa and Higgs self couplings in the SM	18
A.1	$O(\alpha\alpha_s)$ corrections to the relation between the on-shell and $\overline{\text{MS}}$ Fermi constants	18
A.2	$O(\alpha\alpha_s)$ corrections to the relation between the $\overline{\text{MS}}$ and pole masses of the top quark	22
A.3	$O(\alpha\alpha_s)$ corrections to the relation between the $\overline{\text{MS}}$ and pole masses of the Higgs boson	22
A.4	$O(\alpha\alpha_s)$ corrections to the top Yukawa and Higgs self-couplings	25
B	Beta functions	27

1 Introduction

The mass M_H of the Higgs boson in the SM is an important indicator of the presence of new energy scales in particle physics. It is well known that if $M_{\min}^{\text{meta}} < M_H < M_{\max}^{\text{Landau}}$ then the SM is a consistent effective field theory all the way from the Fermi scale up to the (reduced) Planck scale $M_P = 2.44 \times 10^{18}$ GeV. The upper limit comes from the requirement that the Landau pole in the scalar self-coupling¹ must not appear at energies below M_P [1–3]. The lower limit comes from the requirement of the stability of the SM vacuum against tunneling to the states with the Higgs field ϕ exceeding substantially the electroweak value 250 GeV [4–6] (see figure 1).

The estimates of M_{\max}^{Landau} give a number around 175 GeV [1–3, 7] which is in the M_H range excluded (at least in the range 129 – 525 GeV) by the searches for the SM Higgs

¹To be more precise, the scalar self-coupling is infinite in the one-loop approximation only. If higher-order terms are included, it may not become infinite, but move away from the region of the weak coupling.

boson at the LHC and the Tevatron [8, 9]. In other words, we already know that the SM is a weakly coupled theory up to the Planck scale.

One can distinguish between two types of the stability bounds. If $M_H > M_{\min}^{\text{stability}}$, the electroweak vacuum is absolutely stable, whereas if $M_{\min}^{\text{meta}} < M_H < M_{\min}^{\text{stability}}$, then it is metastable with a life-time exceeding that of the Universe. Numerically, $M_{\min}^{\text{meta}} \simeq 111 \text{ GeV}$ [10]. The existence of the Higgs boson with a mass smaller than M_{\min}^{meta} would provide an undisputed argument in favor of the existence of new physics between the Fermi and Planck scales. However, already since LEP we know that this is not the case.

The Higgs mass $M_{\min}^{\text{stability}}$ is not at all special from the point of view of the validity of the SM up to the Planck scale. The value of $M_{\min}^{\text{stability}}$, however, plays a crucial role if the SM is embedded to a bigger picture which includes gravity. First, only if $M_H > M_{\min}^{\text{stability}}$, the Higgs boson of the SM can play the role of the inflaton and make the Universe flat, homogeneous and isotropic, and produce a necessary spectrum of perturbations needed for structure formation [11, 12]. Second, $M_H = M_{\min}^{\text{stability}}$ is a prediction of an asymptotically safe scenario of the SM [13], making it consistent up to an arbitrarily large scale.

Thus, we will focus on the upgrade of existing computations of $M_{\min}^{\text{stability}}$ and on the discussion of the significance of the relation between the Higgs boson (to be discovered yet) mass M_H and $M_{\min}^{\text{stability}}$ for beyond-the-SM (BSM) physics.

The computation of $M_{\min}^{\text{stability}}$ has been already done in a large number of papers [10, 12, 14–17]. It is divided into two parts. The first one is the determination of the $\overline{\text{MS}}$ parameters from the physical observables and the second one is the renormalization group running of the $\overline{\text{MS}}$ constants from the electroweak to a high-energy scale. The most advanced recent works [10, 12, 17, 18] use the so-called “one-loop-matching–two-loop-running” procedure. It can determine the Higgs boson mass bounds with a theoretical accuracy of 2–5 GeV (see the discussion of uncertainties in [12] and below). Meanwhile, the most important terms in the three-loop running of the gauge and Higgs coupling constants were computed in [19, 20] (we thank K. Chetyrkin and M. Zoller for sharing these results with us prior to publication). The present work accounts for $O(\alpha\alpha_s)$ corrections in the $\overline{\text{MS}}$ -pole matching procedure, which were not known previously. This allows us to decrease the theoretical uncertainties in the Higgs boson mass prediction/bounds associated with SM physics down to 1–2 GeV. This is a new result, based on a superior partial “two-loop-matching–three-loop-running” procedure. These findings are described in section 2.2. We will see that the experimental errors in the mass of the top quark and in the value of the strong coupling constant are too large to settle the question of the stability of the electroweak vacuum, even if the LHC will confirm the evidence for the Higgs signal presented by the ATLAS and CMS collaborations [8, 9] in the region $M_H = 124 - 126 \text{ GeV}$.

In section 3 we will discuss the significance of the relationship between the true Higgs boson mass M_H and $M_{\min}^{\text{stability}}$ for BSM physics. We will argue that if $M_H = M_{\min}^{\text{stability}}$ then the electroweak symmetry breaking is likely to be determined by Planck physics and that this would indicate the absence of new energy scales between the Fermi and gravitational scales. We will also address here the significance of $M_{\min}^{\text{stability}}$ for the SM with gravity included. Of course, this can only be done under certain assumptions. Specifically, we will discuss the non-minimal coupling of the Higgs field to the Ricci scalar (relevant for Higgs-inflation [11, 12, 21]) and the asymptotic-safety scenario of the SM [13].




Figure 1. Higgs self-coupling in the SM as a function of the energy scale. The top plot depicts possible behaviors for the whole Higgs boson mass range — Landau pole, stable, or unstable electroweak vacuum. The lower plots show the detailed behaviors for low Higgs boson masses, with dashed (dotted) line corresponding to the experimental uncertainty in the top mass M_t (strong coupling constant α_s), and the shaded yellow (pink) regions correspond to the total experimental error and the theoretical uncertainty, with the latter estimated as 1.2 GeV (2.5 GeV), see section 2 for a detailed discussion.

In section 4 we present our conclusions. We will argue that if only the Higgs boson with a mass around $M_{\min}^{\text{stability}}$ and nothing else will be found at the LHC, the next step in high-energy physics should be the construction a new electron-positron (or muon) collider — the Higgs and t-factory. It will not only be able to investigate in detail the Higgs and top physics, but also elucidate the possible connection of the Fermi and Planck scales.

Appendix A contains a full set of formulas required for the determination of the $\overline{\text{MS}}$ coupling constants from the pole masses of the SM particles, including the corrections of the orders of up to $O(\alpha_s^3)$, $O(\alpha)$, and $O(\alpha\alpha_s)$. The computer code for the matching is made publicly available at <http://www.inr.ac.ru/~fedor/SM/>.




Figure 2. Schematic depiction of the SM effective potential V for the Higgs field for $M_H > M_{\min}^{\text{stability}}$ (left) and $M_H < M_{\min}^{\text{stability}}$ (right).

2 The stability bound

The stability bound will be found in the “canonical” SM, without any new degrees of freedom or any extra higher-dimensional operators added, see figure 2.

2.1 The benchmark mass

It will be convenient for computations to introduce yet another parameter, the “benchmark mass”, which we will call M_{\min} (without any superscript). Suppose that all parameters of the SM, except for the Higgs boson mass, are exactly known. Then M_{\min} , together with the normalisation point μ_0 , are found from the solution of the two equations:

$$\lambda(\mu_0) = 0, \quad \beta_\lambda(\lambda(\mu_0)) = 0, \quad (2.1)$$

where β_λ is the β -function governing the renormalisation group (RG) running of λ . Here we define all the couplings of the SM in the $\overline{\text{MS}}$ renormalisation scheme, which is used de-facto in the most of the higher-loop computations. Clearly, if any other renormalization scheme is used, the equations $\lambda = \beta_\lambda = 0$ will give another benchmark mass, since the definition of all the couplings are scheme dependent.

The procedure of computing M_{\min} is very clean and transparent. Take the standard $\overline{\text{MS}}$ definition of all coupling constants of the SM, fix all of them at the Fermi scale given the experimentally known parameters such as the mass of the top quark, the QCD coupling, etc., and consider the running Higgs self-coupling $\lambda(\mu)$ depending on the standard ’t Hooft-Veltman parameter μ . Then, adjust M_{\min} in such a way that eq. (2.1) are satisfied at some μ_0 .

For the stability bound one should find the effective potential $V(\phi)$ and solve the equations

$$V(\phi_{SM}) = V(\phi_1), \quad V'(\phi_{SM}) = V'(\phi_1) = 0, \quad (2.2)$$

where ϕ_{SM} corresponds to the SM Higgs vacuum, and ϕ_1 correspond to the extra vacuum states at large values of the scalar field. Though the effective potential and the field ϕ

are both gauge and scheme dependent, the solution for the Higgs boson mass to these equations is gauge and scheme invariant.

In fact, $M_{\min}^{\text{stability}}$ is very close to M_{\min} . Numerically, the difference between them is much smaller, than the current theoretical and experimental precisions for M_{\min} , see below. The following well known argument explains why this is the case. The RG improved effective potential for large ϕ can be written as [15, 16, 22]

$$V(\phi) \propto \lambda(\phi)\phi^4 \left[1 + O\left(\frac{a}{4\pi} \log(M_i/M_j)\right) \right], \quad (2.3)$$

where a is a common name for all the SM coupling constants (which are rather small at Planck scale), and M_i are the masses of different particles in the background of the Higgs field. If $O(\alpha)$ corrections are neglected, the eq. (2.2) coincides with eq. (2.1), meaning that $M_{\min} \simeq M_{\min}^{\text{stability}}$. The numerical evaluation for the one loop effective potential gives $\Delta m^{\text{stability}} \equiv M_{\min}^{\text{stability}} - M_{\min} \simeq -0.15 \text{ GeV}$, which can be neglected in view of the uncertainties discussed below.

Note that in many papers the stability bound is shown as a function of the cutoff scale Λ (the energy scale up to which the SM can be considered as a valid effective field theory). It is required that $V(\phi) > V(\phi_{SM})$ for all $\phi < \Lambda$. This can be reformulated as $\lambda(\mu) > 0$ for all $\mu < \Lambda$ with pretty good accuracy. Interestingly, if $\Lambda = M_P$, this bound is very close to the stability bound following from eq. (2.2), having nothing to do with the Planck scale (see also below). Note also that the uncertainties in experimental determinations of M_t and α_s together with theoretical uncertainties, described in the next section, lead to significant changes in the scale Λ . Figure 1 illustrates that for Higgs boson masses 124 – 127 GeV this scale may vary from 10^8 GeV up to infinity within currently available precisions.

2.2 Value of M_{\min}

The state-of-the-art computation of M_{\min} has contained up to now the so-called one-loop $\overline{\text{MS}}$ -pole matching, relating the experimentally measured physical parameters to the parameters of the SM in the $\overline{\text{MS}}$ scheme (to be more precise, the two-loop α_s corrections to the top pole mass– $\overline{\text{MS}}$ mass relation has been included [10]). Then the results of the first step are plugged into two-loop RG equations and solved numerically.

Before discussing the upgrade of the one-loop-matching–two-loop-running procedure, we will remind of the results already known and their uncertainties. We will make use of our computations of M_{\min} presented in [12].² Somewhat later papers [17, 18, 23] contain exactly the same numbers for $M_{\min}^{\text{stability}}$ (note, however, that the theoretical uncertainties were not discussed in [23]). See also earlier computations in [7, 10, 14–16, 18].

In [12] we found:

$$M_{\min} = \left[126.3 + \frac{M_t - 171.2 \text{ GeV}}{2.1 \text{ GeV}} \times 4.1 - \frac{\alpha_s - 0.1176}{0.002} \times 1.5 \right] \text{ GeV}, \quad (2.4)$$

²The main interest in this paper was the lower bound on the Higgs boson mass in the Higgs inflation, see below. However, M_{\min} has been estimated as well as a by-product of the computation.

Source of uncertainty	Nature of estimate	$\Delta_{\text{theor}} M_{\text{min}}, \text{ GeV}$
2-loop matching λ	Sensitivity to μ	1.7
2-loop matching y_t	Sensitivity to μ	0.6
3-loop α_s to y_t	known	1.4
4-loop α_s to y_t	educated guess [27, 28]	0.4
confinement, y_t	educated guess $\sim \Lambda_{QCD}$	0.5
3-loop running $M_W \rightarrow M_P$	educated guess	0.8
total uncertainty	sum of squares	2.5
total uncertainty	linear sum	5.4
Corrections to [12]		$\Delta M_{\text{min}}, \text{ GeV}$
Typos in the code used in [12]	error	+0.2
Extra δ_t^{QED} in (A.5) of [10]	error	+0.4
“Exact” formula instead of approximation (2.20) in [32]	clarification	+0.1
Total correction to (7.1) of [12]		+0.7
Total shift to be applied to (7.1) of [12] for comparison		+0.7

Table 1. Theoretical uncertainties and mistakes in the M_{min} evaluation in [12].

and estimated the theoretical uncertainties as summarized in table 1 (see also [17]).³ While repeating this analysis, we found some numerical errors which are given at the bottom section of this table (see a detailed discussion below). In total, they shift the value given in eq. (2.4) up by 0.7 GeV. As for the uncertainties, they were estimated as follows. The one-loop matching formulas can be used directly at $\mu = m_t$, or at some other energy scale, e.g. at $\mu = M_Z$, and then the coupling constants at $\mu = m_t$ can be derived using RG running. The difference between the procedures gives an estimate of the two-loop effects in the matching procedure. This is presented by the first two lines in table 1 (in fact, we underestimated before the uncertainty from the λ matching — previously we had here 1.2 GeV and now 1.7 GeV). The next two lines are associated with three- and four-loop corrections to the top Yukawa coupling y_t . The three-loop corrections were computed in [24–26] and the four-loop α_s contribution to the top mass was estimated to be of the order $\delta y_t(m_t)/y_t(m_t) \simeq 0.001$ in [27, 28]. The non-perturbative QCD effects in the top pole mass– $\overline{\text{MS}}$ mass matching are expected to be at the same level [29–31]. For three-loop running we put the typical coefficients in front of the largest couplings α_s and y_t . If these uncertainties are not correlated and can be summed up in squares, the theoretical uncertainty is 2.5 GeV. If they are summed up linearly, then the theoretical error can be as large as ~ 5 GeV.

Now, this computation can be considerably improved. First, in [19] the three-loop corrections to the running of all gauge couplings has been calculated. Second, in [20] the

³The reader not interested in details of the comparison with the previous results can skip this discussion and restart with the following paragraph.

Contribution	ΔM_{\min} , GeV
Three loop beta functions	-0.23
$\delta y_t \propto O(\alpha_s^3)$	-1.15
$\delta y_t \propto O(\alpha\alpha_s)$	-0.13
$\delta\lambda \propto O(\alpha\alpha_s)$	0.62

Table 2. Contributions to the value of the M_{\min} .

leading contributions (containing the top Yukawa coupling and α_s) to the running of the top quark Yukawa and the Higgs boson self-coupling have been determined. This removes the uncertainty related to the three-loop RG running. In addition, in the present paper, we determine the two-loop corrections of the order $O(\alpha\alpha_s)$ to the matching of the pole masses and the top quark Yukawa and Higgs boson self coupling constants. Also, the known [24–26] three-loop QCD correction to the top quark mass relation of the order $O(\alpha_s^3)$ can be included (previously it had been used for estimates of uncertainties). All this considerably decreases the theoretical uncertainties in M_{\min} .

The individual contributions of the various new corrections on top of the previous result are summarized in table 2. It is clearly seen that there are two new significant contributions — one is the three-loop pure QCD correction to the top quark mass [24–26], and another is the two-loop correction of $O(\alpha\alpha_s)$ to the Higgs boson mass, found in the present paper. Together the new contributions add -0.89 GeV to the overall shift of the previous prediction [12], giving the result⁴

$$M_{\min} = \left[128.95 + \frac{M_t - 172.9 \text{ GeV}}{1.1 \text{ GeV}} \times 2.2 - \frac{\alpha_s - 0.1184}{0.0007} \times 0.56 \right] \text{ GeV}. \quad (2.5)$$

The new result (2.5) is less than 0.2 GeV away from the old one (2.4) if the same central values for M_t and α_s are inserted. This coincidence is the result of some magic. In the old evaluation several mistakes were present, summarized in table 1. The largest one was the double counting of δ_t^{QED} in eq. (A.5) of [10], as compared to the original result [32]. Also, there were minor typos in the computer code for the matching of the Higgs coupling constant, and finally there was a small correction coming from the use of an approximate rather than exact one-loop formula for the $O(\alpha)$ corrections from [32]. These corrections add 0.7 GeV to the original number in [12]. By chance this almost exactly canceled the -0.89 GeV contribution from the higher loops, table 2, nearly leading to a coincidence of eqs. (2.5) and (2.4).

Table 3 summarizes the uncertainties in the new computation. It contains fewer lines. Now we can ignore safely the error from higher-order (four-loop) RG corrections for the running up to the Planck scale. The first two lines were derived in the same manner as previously. For the Higgs boson self-coupling we can use the matching formulas (A.35) to get the value of $\lambda(\mu)$ at scale $\mu = M_t$ directly, or to get the value $\lambda(M_Z)$ and then

⁴Note, that this value is the benchmark mass defined by eq. (2.1). The mass corresponding to the metastability bound (2.2) is ~ 0.1 – 0.2 GeV smaller.

Source of uncertainty	Nature of estimate	$\Delta_{\text{theor}} M_{\text{min}}, \text{ GeV}$
3-loop matching λ	Sensitivity to μ	1.0
3-loop matching y_t	Sensitivity to μ	0.2
4-loop α_s to y_t	educated guess [27, 28]	0.4
confinement, y_t	educated guess $\sim \Lambda_{QCD}$	0.5
4-loop running $M_W \rightarrow M_P$	educated guess	< 0.2
total uncertainty	sum of squares	1.2
total uncertainty	linear sum	2.3

Table 3. Theoretical uncertainties in the present M_{min} evaluation.

evolve the constants to the scale $\mu = M_t$ with the RG equations. The obtained difference $\delta\lambda(M_t)/\lambda(M_t) \simeq 0.016$ (cf. figure 12) corresponds to the error $\delta m \sim 1.0 \text{ GeV}$. A similar procedure of comparing the evolution between M_t and M_Z using RG equations and direct matching formulas to the order $O(\alpha_s^3, \alpha, \alpha\alpha_s)$ leads for the change in the top quark Yukawa coupling $\delta y_t/y_t \sim 0.0005$ (cf. figure 12), leading to $\delta M_{\text{min}} \sim 0.2 \text{ GeV}$. Note, however, that strictly speaking this test verifies the error of the μ dependent terms in the matching formulas, while the constant ones may lead to larger contributions. We also do not estimate now the contributions of the order $O(\alpha^2)$, where numerically one order in α may correspond to y_t^4 . Thus, this estimate should be better considered as a lower estimate of the error. The four-loop matching and confinement contributions are the same as before.

As an indication of the dependence on the matching point we present figure 3, where the reference Higgs boson mass M_{min} was obtained using the matching formulas at scale μ_0 varying between the Z-boson and top quark masses. One can see that the overall change of the Higgs boson mass is about 1 GeV.

If we assume that these uncertainties are not correlated and symmetric we get a theoretical error in the determination of the critical Higgs boson mass $\delta M_{\text{min}} \simeq 1.2 \text{ GeV}$. If they are summed up linearly, we get an error of 2.4 GeV. We leave it to the reader to decide which estimate of the uncertainties is more adequate. The precision of the theoretical value of M_{min} can be further increased by computing the $O(\alpha^2)$ two-loop corrections to the matching procedure. Numerically, the most important terms are those when α corresponds to y_t^2 and λ .

The result (2.5) is visualized by figure 4. The experimentally allowed regions for the top mass M_t and strong coupling α_s are adopted from the PDG 2010 edition [33].⁵ On top of these allowed regions the bands corresponding to the reference values of the Higgs boson mass M_{min} being equal to 124, 125, 126, 127 GeV are shown, with the dashed and dotted lines corresponding to quadratically or linearly added estimates of the theoretical uncertainties.

One can see that the accuracy of the theoretical computations and of the experimental measurements of the top and the Higgs boson masses does not allow yet to conclude with

⁵Note, however, that the current experimental error estimate is based on averaging over different experimental approaches. In some methods quite a different central values are obtained. See e.g. [34–36] about the α_s determination and [37–39] about M_t .




Figure 3. The dependence of the reference Higgs boson mass M_{\min} on the matching scale μ_0 (the $\overline{\text{MS}}$ constants are obtained by matching formulas at the scale μ_0 and then used for the solution of eq. (2.1)). The thick solid line corresponds to the full matching formulas $\lambda \sim O(\alpha, \alpha\alpha_s)$, $y_t \sim O(\alpha_s^3, \alpha, \alpha\alpha_s)$; the thin lines correspond to using matching formulas of lower order. The thick dashed line corresponds to using additionally the two-loop electroweak contributions to the Higgs coupling constant in the gauge-less limit, eq. (48) of [119], see the discussion in the “Note added”. Here $M_t = 172.9 \text{ GeV}$ and $\alpha_s = 0.1184$.

confidence whether the discovery of the Higgs boson with a mass $124 - 127 \text{ GeV}$ would indicate stability or metastability of the SM vacuum. All these reference values of Higgs masses are compatible within 2σ with current observations.

3 M_{\min} and BSM physics

Our definition of the “benchmark” Higgs boson mass consists of the solution of the two equations (2.1) and gives, in addition to M_{\min} , the value of the scale μ_0 at which the scalar self-coupling and its β -function vanish simultaneously. The central value of μ_0 is $2.9 \times 10^{18} \text{ GeV}$ and is quite stable if m_t and α_s are varied in their confidence intervals (see figure 5). One can see that there is a remarkable coincidence between μ_0 and the (reduced) Planck scale $M_P = 2.44 \times 10^{18} \text{ GeV}$. The physics input in the computation of μ_0 includes the parameters of the SM only, while the result gives the gravity scale. A possible explanation may be related to the asymptotic safety of the SM, see [13] and below.⁶ It remains to be seen if this is just the random play of the numbers or a profound indication that the electroweak symmetry breaking is related to Planck physics. If real, this coincidence indicates that there should be no new energy scales between the Planck and Fermi scales, as they would remove this coincidence unless some conspiracy is taking place.

We will discuss below two possible minimal embeddings of the SM into the theory of gravity and discuss the significance of M_{\min} in them.

⁶Other explanation include the “multiple point principle” of [40, 41] and requiring the degeneracy between the SM vacuum and an extra one appearing at the Planck scale, the inflation from false vacuum decay [42–45], etc.




Figure 4. The values of the strong coupling constant α_s and top mass M_t corresponding to several minimal Higgs boson masses M_{\min} . The 68% and 95% experimentally allowed regions for α_s and M_t are given by shaded areas. The dashed (dotted) lines correspond to 1.2 GeV (2.45 GeV) uncertainty in the M_{\min} theoretical determination.

3.1 Asymptotic safety

The asymptotic safety of the SM [13], associated with the asymptotic safety of gravity [46], is strongly related to the value of the Higgs boson mass. Though General Relativity is non-renormalizable by perturbative methods, it may exist as a field theory non-perturbatively, exhibiting a non-trivial ultraviolet fixed point (for a review see [47]). If true, all other coupling of the SM (including the Higgs self-interaction) should exhibit asymptotically safe behaviors with the gravity contribution to the RG running included.

The prediction of the Higgs boson mass from the requirement of asymptotic safety of the SM is found as follows [13]. Consider the SM running of the coupling constants and add to the β -functions extra terms coming from gravity, deriving their structure from




Figure 5. The scale μ_0 (solution of (2.1)) depending on the top mass M_t . The dashed lines correspond to 1σ uncertainty in the α_s . The yellow shaded region corresponds to adding the α_s experimental error and the theoretical uncertainty in the matching of the top Yukawa coupling y_t and top pole mass.

dimensional analysis:

$$\beta_h^{\text{grav}} = \frac{a_h}{8\pi} \frac{\mu^2}{M_P^2(\mu)} h, \quad (3.1)$$

where a_1 , a_2 , a_3 , a_y , and a_λ are some constants (anomalous dimensions) corresponding to the gauge couplings of the SM g , g' , g_s , the top Yukawa coupling y_t , and the Higgs self-coupling λ . In addition,

$$M_P^2(\mu) \simeq M_P^2 + 2\xi_0\mu^2 \quad (3.2)$$

is the running Planck mass with $\xi_0 \approx 0.024$ following from numerical solutions of functional RG equations [48–50]. Now, require that the solution for all coupling constants is finite for all μ and that λ is always positive. The SM can only be asymptotically safe if a_1 , a_2 , a_3 , a_y are all negative, leading to asymptotically safe behaviors of the gauge and Yukawa couplings. For $a_\lambda < 0$ we are getting the interval of admissible Higgs boson masses, $M_{\min}^{\text{safety}} < M_H < M_{\max}^{\text{safety}}$. However, if $a_\lambda > 0$, as follows from the computations of [49, 50], only one value of the Higgs boson mass $M_H = M_{\min}^{\text{safety}}$ lead to an asymptotically safe behavior of λ . As is explained in [13], this behavior is only possible provided $\lambda(M_P) \approx 0$ and $\beta_\lambda(\lambda(M_P)) \approx 0$. And, due to a miraculous coincidence of μ_0 and M_P , the difference $\Delta m^{\text{safety}} \equiv M_{\min}^{\text{safety}} - M_{\min}$ is extremely small, of the order 0.1 GeV. The evolution of the Higgs self-coupling for the case of $a_h < 0$ is shown in figure 6, and for the case $a_h > 0$ in figure 7.

In fact, in the discussion of the asymptotic safety of the SM one can consider a more general situation, replacing the Planck mass in eq. (3.2) by some cutoff scale $\Lambda = \kappa M_P$.



Figure 6. Schematic depiction of the behavior of the scalar self-coupling if $a_h < 0$ for $M_{\min}^{\text{safety}} < M_H < M_{\max}^{\text{safety}}$ (left) and $M_H < M_{\min}^{\text{safety}}$ (right). In both cases gravity leads to an asymptotically free behavior of the scalar self-coupling. Negative λ leads to instability and is thus excluded.




Figure 7. Schematic depiction of the behavior of the scalar self-coupling if $a_h > 0$ for $M_H > M_{\min}^{\text{safety}}$, leading to a Landau-pole behavior (left), $M_H > M_{\min}^{\text{safety}}$, leading to instability (right) and $M_H = M_{\min}^{\text{safety}}$, leading to an asymptotically safe behavior (middle). Only this choice is admissible.

Indeed, if the Higgs field has a non-minimal coupling with gravity (see below), the behavior of the SM coupling may start to change at energies smaller than M_P by a factor $1/\xi$, leading to an expectation for the range of κ as $1/\xi \lesssim \kappa \lesssim 1$. Still, the difference between M_{\min} and M_{\min}^{safety} remains small even for $\kappa \sim 10^{-4}$, where $M_{\min}^{\text{safety}} \simeq 128.4 \text{ GeV}$, making the prediction $M_H \simeq M_{\min}$ sufficiently stable against specific details of Planck physics within the asymptotic safety scenario.

3.2 M_{\min} and cosmology

It is important to note that if the mass of the Higgs boson is smaller than the stability bound M_{\min} , this does not invalidate the SM. Indeed, if the time-life of the metastable SM vacuum exceeds the age of the Universe (this is the case when $M_H > M_{\text{meta}}$, with

$M_{\text{meta}} \simeq 111 \text{ GeV}$ [10]) then finding a Higgs boson in the mass interval $M_{\text{meta}} < M_H < M_{\text{min}}$ would simply mean that we live in a metastable state with a very long lifetime. Of course, if the Higgs boson were discovered with a mass below M_{meta} , this would prove that there *must be* new physics between the Fermi and Planck scales, stabilizing the SM vacuum state. However, the latest LEP results, confirmed recently by LHC, tell us that in fact $M_H > M_{\text{meta}}$, and, therefore, that the presence of a new energy scale is not required, if only the metastability argument is used.

The bound $M_H > M_{\text{meta}}$ can be strengthened if thermal cosmological evolution is considered [10]. After inflation the universe should find itself in the vicinity of the SM vacuum and stay there till present. As the probability of the vacuum decay is temperature dependent, the improved Higgs boson mass bound is controlled by the reheating temperature after inflation (or maximal temperature of the Big Bang). The latter is model dependent, leading to the impossibility to get a robust bound much better than M_{meta} . For example, in R^2 inflation [51, 52] the reheating temperature is rather low, $T \sim 10^9 \text{ GeV}$ [52], leading to the lower bound 116 GeV [53] on the Higgs boson mass, which exceeds M_{meta} only by 4 GeV .

However, if no new degrees of freedom besides those already present in the SM are introduced and the Higgs boson plays the role of the inflaton, the bound $M_H \gtrsim M_{\text{min}}$ reappears, as is discussed below.

3.3 Higgs inflation

The inclusion of a non-minimal interaction of the Higgs field with gravity, given by the Lagrangian $\xi|\phi|^2 R$, where R is the Ricci scalar, changes drastically the behavior of the Higgs potential in the region of large Higgs fields $\phi > M_{\text{inflation}} \simeq M_P/\sqrt{\xi}$ [11]. Basically, the potential becomes flat at $\phi > M_{\text{inflation}}$, keeping the value it acquired at $\phi \simeq M_P/\sqrt{\xi}$. This feature leads to a possibility of Higgs-inflation: if the parameter ξ is sufficiently large, $700 < \xi < 10^5$, [12] the Higgs boson of the SM can make the Universe flat, homogeneous and isotropic, and can produce the necessary spectrum of primordial fluctuations. The possibility of the Higgs inflation is also strongly related to the value of the Higgs boson mass: successful inflation can only occur if $M_{\text{min}}^{\text{inflation}} < M_H < M_{\text{max}}^{\text{inflation}}$. The upper limit $M_{\text{max}}^{\text{inflation}}$ comes from the requirement of the validity of the SM up to the inflation scale $M_{\text{inflation}}$. Near $M_{\text{min}}^{\text{inflation}}$ the behavior of the effective potential in the Einstein frame changes as shown in figure 8: if $M_H < M_{\text{min}}^{\text{inflation}}$ the “bump” in the Higgs potential prevents the system to go to the SM vacuum state. As in the previous case, these bounds can be formulated with the use of the Higgs self-coupling λ . Basically, it must be perturbative and positive for all energy scales below $M_{\text{inflation}}$. Though any Higgs boson mass in the interval $M_{\text{min}}^{\text{inflation}} < M_H < M_{\text{max}}^{\text{inflation}}$ can lead to successful inflation, the value $M_{\text{min}}^{\text{inflation}}$ is somewhat special. For the lower part of the admitted interval the value of the non-minimal coupling ξ reaches its minimal value $\xi \simeq 700$, extending the region of the applicability of perturbation theory [12, 54, 55].

The computation of the lower bound on the Higgs boson mass from inflation is more complicated. It is described in detail in [12, 21]. Basically, one has to compute the Higgs potential in the chiral electroweak theory associated with large values of the Higgs field




Figure 8. Schematic depiction of the effective potential V for the Higgs field in the Higgs-inflationary theory in the Einstein frame for $M_H > M_{\min}^{\text{inflation}}$ (left) and $M_H < M_{\min}^{\text{inflation}}$ (right).

and find when the slow-roll inflation in this potential can give the large-scale perturbations observed by the COBE satellite. The outcome of these computations, however, can be formulated in quite simple terms: for inflationary bound find $M_{\min}^{\text{inflation}}$ from the condition $\lambda(\mu) > 0$ for all $\mu < M_{\text{inflation}}$ [12]. A priori, the inflationary bound could have been very different from M_{\min} and thus from $M_{\min}^{\text{stability}}$. Indeed, both M_{\min} and $M_{\min}^{\text{stability}}$ know *nothing* about the Planck scale and are defined entirely within the SM, whereas the inflationary bound does use M_P . However, the remarkable numerical coincidence, between μ_0 and M_P , makes M_{\min} and $M_{\min}^{\text{inflation}}$ practically the same. The coupling constant λ evolves very slowly near the Planck scale, so that the regions for the Higgs boson mass following from the conditions $\lambda(\mu) > 0$ for $\mu < M_P$ and $\mu < M_{\text{inflation}}$ are almost identical. This leads to the result that $\Delta m^{\text{inflation}} \equiv M_{\min}^{\text{inflation}} - M_{\min} \simeq -0.1 - 0.2 \text{ GeV}$. This number is derived within the SM without addition of any higher-dimensional operators.

One must note that simple scale analysis leads to unitarity violation in the Higgs inflation below the Planck energy scale [56–59]. This means, that calculations in Higgs inflationary models should be done with some additional assumptions about the high-energy physics, formulated in [54], specifically the approximate scale invariance at high field backgrounds. As is explained in [54], adding to the SM higher-dimensional operators with a Higgs-field dependent cutoff modifies the lower bound on the Higgs boson mass in Higgs inflation. If these operators are coming with “natural” power counting coefficients (for an exact definition see [54]) the sensitivity of the Higgs boson mass bound to unknown details of ultraviolet physics is rather small $\Delta M_{\min}^{\text{inflation}} \simeq 0.6 \text{ GeV}$ [54]. At the same time, it is certainly not excluded that the change of $M_{\min}^{\text{inflation}}$ can be larger.

4 Conclusions

If the SM Higgs boson will be discovered at LHC in the remaining mass interval $115.5 < M_H < 127 \text{ GeV}$ not excluded at 95% [8, 9], there is no necessity for a new energy scale between the Fermi and Planck scales. The electroweak theory remains in a weakly coupled

region all the way up to M_P , whereas the SM vacuum state lives longer than the age of the Universe. If the SM Higgs boson mass will be found to *coincide* with M_{\min} given by eq. (2.5), this would provide a strong argument in favor of the *absence* of such a scale and indicate that the electroweak symmetry breaking may be associated with the physics at the Planck scale.

The experimental precision in the Higgs boson mass measurements at the LHC can eventually reach 200 MeV and thus be much smaller than the present theoretical ($\sim 1\text{--}2\text{ GeV}$) and experimental ($\sim 5\text{ GeV}$, 2σ) uncertainties in the determination of M_{\min} . The largest uncertainty comes from the measurement of the mass of the top quark. It does not look likely that the LHC will substantially reduce the error in the top quark mass determination. Therefore, to clarify the relation between the Fermi and Planck scales the construction of an electron-positron or muon collider with a center-of-mass energy of $\sim (200 + 200)\text{ GeV}$ (Higgs and t-quark factory) would be needed. This would be decisive for setting up the question about the necessity for a new energy scale besides the two ones already known — the Fermi and the Planck scales. In addition, this will allow to study in detail the properties of the two heaviest particles of the SM, potentially most sensitive to any types on new physics.

Surely, even if the SM is a valid effective field theory all the way up the the Planck scale, it cannot be complete as it contradicts a number of observations. We would like to use this opportunity to underline once more that the confirmed observational signals in favor of BSM physics which were not discussed in this paper (neutrino masses and oscillations, dark matter and baryon asymmetry of the Universe) can be associated with new physics *below* the electroweak scale, for reviews see [60, 61] and references therein.⁷ The minimal model, νMSM , contains, in addition to the SM particles, three relatively light singlet Majorana fermions. These fermions could be responsible for neutrino masses, dark matter and the baryon asymmetry of the Universe. The νMSM predicts that the LHC will continue to confirm the SM and see no deviations from it. At the same time, new experiments at the high-intensity frontier, discussed in [64], may be needed to uncover the new physics below the Fermi scale. In addition, new observations in astrophysics, discussed in [61], may shed light to the nature of dark matter. As the running of the couplings in the νMSM coincides with that in the SM, all results of the present paper are equally applicable to the νMSM .

Acknowledgments

The work of M.S. was supported by the Swiss National Science Foundation. The work of M.Yu.K. and B.A.K. was supported in part by the German Federal Ministry for Education and Research BMBF through Grants No. 05H09GUE, and No. 05H12GUE by the German Research Foundation DFG through the Collaborative Research Centre No. 676 *Particles, Strings and the Early Universe — The Structure of Matter and Space Time*, and by the Helmholtz Association HGF through the Helmholtz Alliance Ha 101 *Physics at the Terascale*. We thank A. Boyarsky, K. Chetyrkin, D. Gorbunov, F. Jegerlehner, G. Passarino,

⁷As for the dark energy, it may be related to a massless dilaton realizing spontaneously broken scale invariance [62, 63].

O. Ruchayskiy, and M. Zoller for helpful discussions, collaboration and interest to our work. M.Yu.K. is indebted to Fred Jegerlehner for a fruitful long-time collaboration on studies of the $\overline{\text{MS}}$ scheme beyond the one-loop order in the framework of quantum field theory models with spontaneously symmetry breaking, and in particular for collaboration on [65], the results and methods of which were heavily used in the present work.

Note added. After our paper was submitted to the electronic preprint archive arXiv (on May 13) a number of events happened, which require its update. First, extra corrections to the matching procedure at low energy scale, not computed in our work, were found in [119] (May 29).⁸ Ref. [119] finds agreement with our results on the $\alpha\alpha_s$ order in the Higgs self-coupling constant. In addition, this paper computed a part of the $O(\alpha^2)$ corrections to the top Yukawa and Higgs coupling constants in the “gauge-less” limit of the Standard model, i.e. the two-loop terms containing the top Yukawa and scalar self-coupling were accounted for. The overall effect of these terms happened to be quite small. The corrections shift the benchmark Higgs mass up by 0.2 GeV, and reduce the sensitivity of the results to the normalisation point from 1.2 GeV to 0.8 GeV, decreasing somewhat the theoretical error-bars, see figure 3.

Second, the discovery of the Higgs-like resonance was announced by the ATLAS and CMS collaborations [66, 67] at CERN. According to CMS,

$$M_H = 125.3 \pm 0.4(\text{stat}) \pm 0.5(\text{syst}) \text{ GeV}, \quad (4.1)$$

while ATLAS gives the slightly higher value

$$M_H = 126 \pm 0.4(\text{stat}) \pm 0.4(\text{syst}) \text{ GeV}. \quad (4.2)$$

And, finally, updated results on the mass of the top quark were announced at ICHEP 2012 (July 9), see [38, 68]. The combination of the Tevatron results reads:

$$m_t = 173.2 \pm 0.6(\text{stat}) \pm 0.8(\text{syst}) \text{ GeV} = 173.2 \pm 1.0 \text{ GeV}, \quad (4.3)$$

whereas the present LHC value is

$$m_t = 173.3 \pm 0.5(\text{stat}) \pm 1.3(\text{syst}) \text{ GeV} = 173.3 \pm 1.4 \text{ GeV}. \quad (4.4)$$

The central values of the top mass are somewhat higher (by 0.3–0.4 GeV) than those which were given by the Particle Data Group at the time we were writing our paper.

In figure 9, which is an update of figure 4, we show the changes due to the experimental shift in the top mass (we take the Tevatron result as having smaller errors) and due to the additional two-loop corrections found in [119] for the ATLAS and CMS values of the Higgs mass. The value of the benchmark Higgs mass is inside the 2σ contours in the m_t, α_s plane for the ATLAS and CMS values of M_H , entering the 1σ contours if theoretical uncertainties are incorporated. The red curves show the shrinking in the size of the uncertainties in m_t

⁸Preliminary results of ref. [119] were presented in ref. [118]. However, the $O(\alpha\alpha_s)$ threshold correction to the Higgs self-coupling is not specified in ref. [118], but is presented here for the first time. Moreover, the conclusions reached in ref. [118] disagree with ours.




Figure 9. Update of figure 4. The values of the strong coupling constant α_s and top mass M_t corresponding to a minimal Higgs boson mass M_{\min} coinciding with the recent CMS and ATLAS values 125.3 GeV and 126 GeV, and to the Higgs mass values shifted by approximately $1\sigma \sim 0.6$ GeV. The latest Tevatron 68% and 95% experimentally allowed regions for α_s and M_t are given by shaded areas. The dashed (dotted) lines correspond to a 1 GeV (2 GeV) uncertainty in the M_{\min} theoretical determination. The red lines in the center correspond to the expected precision from an e^+e^- collider.

and α_s (1 and 2σ) that can be achieved with an e^+e^- collider operated as a $t\bar{t}$ factory (this estimate is taken from [69–76] and [77] discussing the ILC physics).

It is also important to note, that it is difficult to determine which renormalization scheme corresponds to the numerical value of m_t quoted by experiments. The m_t determinations by the Tevatron and LHC collaborations [68, 78] are based on Monte Carlo event generators implemented with LO hard-scattering matrix elements. Although it is plausible and likely that the experimental m_t values thus extracted are close to the pole mass of the top quark, this is by no means guaranteed with the due theoretical rigor [79]. In fact, rigorous determinations of the top-quark pole mass from total production cross sections yield somewhat smaller values, albeit with larger errors [30, 31, 120]. For the time being,

we adopt the working hypothesis that the experimental values of m_t [68, 78] correspond to the pole mass, bearing in mind that this is probably subject to change once a proper NLO treatment of the resonating top-quark propagators is implemented.

Thus, all our considerations remain in force and call for further improvement of the theoretical computations, which should account for all $O(\alpha^2)$ corrections to the mapping procedure, and for a construction of a $t\bar{t}$ factory to pin down α_s and m_t . A recent paper [120], where uncertainties in determination m_t and α_s have been analyzed, reached exactly the same conclusion.

A $O(\alpha\alpha_s)$ electroweak corrections to the top Yukawa and Higgs self couplings in the SM

The evaluation of radiative corrections to the relations between $\overline{\text{MS}}$ parameters (coupling constants) and masses of particles includes two steps: the evaluation of radiative corrections between the Fermi constant G_F and its $\overline{\text{MS}}$ counterpart [80] (see [81–83] for recent reviews) and the evaluation of the radiative corrections between $\overline{\text{MS}}$ and pole masses.

The one-loop electroweak correction of $\mathcal{O}(\alpha)$ to the relation between the self-coupling $\lambda(\mu^2)$ and the pole mass of the Higgs boson was obtained in [84] and the one to the relation between the Yukawa coupling y_t and the pole mass of top quark was found in [32]. The corresponding ingredients for the two-loop mixed electroweak-QCD corrections were evaluated in [65, 85–89], but have never been assembled. We performed independent (re)calculations of all $O(\alpha)$ and $O(\alpha\alpha_s)$ contributions. In the following we will denote the on-shell masses by capital M and the $\overline{\text{MS}}$ masses by lowercase m .

The numerical results for the values of the $\overline{\text{MS}}$ constants at scale M_t obtained up to different loop order is presented by the figures 10, 11.

A.1 $O(\alpha\alpha_s)$ corrections to the relation between the on-shell and $\overline{\text{MS}}$ Fermi constants

The relation between the Fermi coupling constant and the bare parameters is as follows [80]:

$$\frac{G_F}{\sqrt{2}} = \frac{g_0^2}{8m_{W,0}^2} \{1 + \Delta R_0\}, \quad (\text{A.1})$$

where ΔR_0 includes unrenormalized electroweak corrections and $g_0, m_{W,0}^2$ are the SU(2) coupling constant and the bare W boson mass (see for details [81–83]). After performing $\overline{\text{MS}}$ renormalization this relation has the following form:

$$\frac{G_F}{\sqrt{2}} = \frac{G_F(\mu^2)}{\sqrt{2}} (1 + \Delta_{G_F,\alpha} + \Delta_{G_F,\alpha\alpha_s} + \dots), \quad (\text{A.2})$$

where on the r.h.s. all masses and coupling constants are taken in the $\overline{\text{MS}}$ renormalization scheme. The one-loop coefficient, $\Delta_{G_F,\alpha}$, is known from [80] and for $N_c = 3$, $C_F = 4/3$




Figure 10. Contributions to the top Yukawa constant through orders $O(\alpha_s^2)$, $O(\alpha_s^3)$, $O(\alpha_s^3) + O(\alpha)$ and $O(\alpha_s^3) + O(\alpha) + O(\alpha\alpha_s^2)$. One parameter is varied, while the two others are chosen from $M_t = 172.9$ GeV, $\alpha_s = 0.1184$, $M_h = 125$ GeV.

and $m_b = 0$ has the following form:

$$\begin{aligned}
 \Delta_{G_F, \alpha} = \frac{g^2}{16\pi^2} & \left\{ \frac{m_t^4}{m_W^2 m_H^2} \left(6 - 6 \ln \frac{m_t^2}{\mu^2} \right) + \frac{m_t^2}{m_W^2} \left(-\frac{3}{4} + \frac{3}{2} \ln \frac{m_t^2}{\mu^2} \right) + \frac{m_H^2}{m_W^2} \left(-\frac{7}{8} + \frac{3}{4} \ln \frac{m_H^2}{\mu^2} \right) \right. \\
 & + \frac{m_Z^4}{m_H^2 m_W^2} \left(-\frac{1}{2} + \frac{3}{2} \ln \frac{m_Z^2}{\mu^2} \right) + \frac{m_W^2}{m_H^2} \left(-1 + 3 \ln \frac{m_W^2}{\mu^2} \right) + \frac{3}{4} \frac{m_W^2}{m_H^2 - m_W^2} \ln \left(\frac{m_W^2}{m_H^2} \right) \\
 & + \frac{m_Z^2}{m_W^2} \left(\frac{5}{8} + \frac{17}{4} \ln \frac{m_W^2}{\mu^2} - 5 \ln \frac{m_Z^2}{\mu^2} \right) - \frac{3}{4} \ln \frac{m_W^2}{\mu^2} - \frac{3}{4} \ln \frac{m_H^2}{\mu^2} \\
 & \left. - \frac{17}{4} \frac{m_Z^2}{m_W^2 \sin^2 \theta_W} \ln \left(\frac{m_W^2}{m_Z^2} \right) + \frac{5}{4} + \frac{7}{2 \sin^2 \theta_W} \ln \left(\frac{m_W^2}{m_Z^2} \right) \right\}. \quad (\text{A.3})
 \end{aligned}$$

Here, $\sin^2 \theta_W$ is defined in the $\overline{\text{MS}}$ scheme as

$$\sin^2 \theta_W \equiv \sin^2 \theta_W^{\overline{\text{MS}}}(\mu^2) = \frac{g'^2(\mu^2)}{g^2(\mu^2) + g'^2(\mu^2)} = 1 - \frac{m_W^2(\mu^2)}{m_Z^2(\mu^2)}, \quad (\text{A.4})$$




Figure 11. Contributions to the Higgs self coupling constant of the orders $O(\alpha)$ and $O(\alpha) + O(\alpha\alpha_s)$. One parameter is varied, while the two others are chosen from $M_t = 172.9 \text{ GeV}$, $\alpha_s = 0.1184$, $M_h = 125 \text{ GeV}$.

where $g'(\mu^2)$ and $g(\mu^2)$ are the U(1) and SU(2) $\overline{\text{MS}}$ gauge coupling constants, respectively. The matching conditions between the $\overline{\text{MS}}$ parameter, defined by eq. (A.4), and its on-shell version [80] follows from the identification

$$\sin^2 \theta_W^{OS} = 1 - \frac{M_W^2}{M_Z^2}, \quad (\text{A.5})$$

where M_Z and M_W are the pole masses of the gauge bosons (see detailed discussion in [90–92]). The evaluation of the mixed QCD-electroweak coefficient, $\Delta_{G_F, \alpha\alpha_s}$, is reduced to the evaluation of the $O(\alpha\alpha_s)$ corrections to the W boson self-energy at zero momenta transfer

and may be written in the following way [93–97]:

$$\begin{aligned}\Delta_{G_F, \alpha\alpha_s} &\equiv 2g_R^2 Z_{g, \alpha\alpha_s} - \left[Z_{W, \alpha\alpha_s} - Z_{m_t^2, \alpha_s} m_t^2 \frac{\partial}{\partial m_t^2} \frac{\Pi_{WW, \alpha}(0)}{m_W^2(\mu^2)} - \frac{\Pi_{WW, \alpha\alpha_s}(0)}{m_W^2(\mu^2)} \right] \\ &= C_f N_c \frac{g^2 g_s^2}{(16\pi^2)^2} \frac{m_t^2}{m_W^2} \left[20 \frac{m_t^2}{m_H^2} - \frac{13}{8} + \zeta_2 \right. \\ &\quad \left. + \left(1 - 20 \frac{m_t^2}{m_H^2} \right) \ln \left(\frac{m_t^2}{\mu^2} \right) - \left(\frac{3}{2} - 12 \frac{m_t^2}{m_H^2} \right) \ln^2 \left(\frac{m_t^2}{\mu^2} \right) \right],\end{aligned}\quad (\text{A.6})$$

where for $Z_{g, \alpha\alpha_s}$, $Z_{g, \alpha\alpha_s}$ and $Z_{W, \alpha\alpha_s}$ we used the results⁹ of [92] and n_F is the number of fermion families ($n_F = 3$ in the SM).

Using the fact, that G_F is RG invariant, i.e. $\mu^2 \frac{d}{d\mu^2} G_F = 0$, the μ -dependent terms in eq. (A.6) can be evaluated explicitly from the one-loop correction and the explicit knowledge of the anomalous dimension γ_{G_F} . As was shown in [90–92, 98, 99], the anomalous dimension γ_{G_F} can be extracted (i) via the beta-function β_λ of the scalar self-coupling and the anomalous dimension of the mass parameter m^2 (in the unbroken phase) or (ii) via the β -function of the SU(2) gauge coupling g and the anomalous dimension of the W boson (in the broken phase):

$$\gamma_{G_F} \equiv \mu^2 \frac{\partial}{\partial \mu^2} \ln G_F(\mu^2) = \frac{\beta_\lambda}{\lambda} - \gamma_{m^2} = 2 \frac{\beta_g}{g} - \gamma_W. \quad (\text{A.7})$$

Eq. (A.2) can be written as

$$\begin{aligned}\frac{G_F}{G_F(\mu^2)} &= 1 - \frac{g^2}{16\pi^2} \left[\gamma_{G_F, \alpha} L - \Delta X_{G_F, \alpha}^{(1)} \right] \\ &\quad + \frac{g^2 g_s^2}{(16\pi)^2} \left[\Delta X_{G_F, \alpha\alpha_s}^{(2)} + C_{G_F, \alpha\alpha_s}^{(2,2)} L^2 - C_{G_F, \alpha\alpha_s}^{(2,1)} L \right],\end{aligned}\quad (\text{A.8})$$

where $L = \ln \frac{\mu^2}{m_t^2}$ and the coefficients $C_{G_F, \alpha\alpha_s}^{(2,2)}$ and $C_{G_F, \alpha\alpha_s}^{(2,1)}$ are defined via the RG equations:

$$2C_{G_F, \alpha\alpha_s}^{(2,2)} = Z_{m_t^2, \alpha_s} \frac{\partial}{\partial m_t^2} \gamma_{G_F, \alpha} = -6C_f \left[\frac{3}{2} \frac{m_t^2}{m_W^2} - 12 \frac{m_t^4}{m_W^2 m_H^2} \right] \Big|_{N_c=3}, \quad (\text{A.9})$$

$$C_{G_F, \alpha\alpha_s}^{(2,1)} = \gamma_{G_F, \alpha\alpha_s} + Z_{m_t^2, \alpha_s} m_t^2 \frac{\partial}{\partial m_t^2} \Delta X_{G_F, \alpha}^{(1)} + Z_{m_t^2, \alpha_s} \gamma_{G_F, \alpha}, \quad (\text{A.10})$$

⁹There are typos in eq. (4.41) of [92]: in all $\overline{\text{MS}}$ renormalization constants, $Z_W^{\alpha_s}$ and $Z_Z^{\alpha_s}$, “ m_t^2/m_H^2 ” should be replaced by “ m_t^2/m_W^2 ”, i.e.

$$\begin{aligned}Z_W^{\alpha_s} &= 1 + \frac{g^2}{(16\pi^2)} \frac{\alpha_s}{4\pi} N_c C_f \left[\frac{1}{\varepsilon} \left(4 \frac{m_t^4}{m_H^2 m_W^2} - \frac{5}{4} \frac{m_t^2}{m_W^2} + \frac{1}{2} n_F \right) + \frac{1}{\varepsilon^2} \left(-12 \frac{m_t^4}{m_H^2 m_W^2} + \frac{3}{2} \frac{m_t^2}{m_W^2} \right) \right] \\ Z_Z^{\alpha_s} &= 1 + \frac{g^2}{(16\pi^2)} \frac{\alpha_s}{4\pi} N_c C_f \left[\frac{1}{\varepsilon} \left(4 \frac{m_t^4}{m_H^2 m_W^2} - \frac{5}{4} \frac{m_t^2}{m_W^2} + \frac{10}{9} n_F \frac{m_W^2}{m_Z^2} + \frac{11}{18} n_F \frac{m_Z^2}{m_W^2} - \frac{11}{9} n_F \right) \right. \\ &\quad \left. + \frac{1}{\varepsilon^2} \left(-12 \frac{m_t^4}{m_H^2 m_W^2} + \frac{3}{2} \frac{m_t^2}{m_W^2} \right) \right],\end{aligned}$$

with

$$\gamma_{G_F, \alpha\alpha_s} = \left[2 \frac{\beta_{g, \alpha\alpha_s}}{g} - 2Z_{W, \alpha\alpha_s} \right] = \frac{1}{2} N_c C_f \left[5 \frac{m_t^2}{m_W^2} - 16 \frac{m_t^4}{m_W^2 m_H^2} \right]. \quad (\text{A.11})$$

Collecting all terms in eq. (A.10) we get

$$C_{G_F, \alpha\alpha_s}^{(2,1)} \Big|_{N_c=3, C_f=\frac{4}{3}} = 4 \frac{m_t^2}{m_W^2} - 80 \frac{m_t^4}{m_W^2 m_H^2}. \quad (\text{A.12})$$

At the end of this section we again point out that the anomalous dimension of the vacuum expectation value $v^2(\mu^2) = 1/(\sqrt{2}G_F(\mu^2))$ within the diagram technique is defined by eq. (A.7) and that it is not equal to the anomalous dimension of the scalar field as in the effective-potential approach [100]. Another important property of eq. (A.7) is the appearance of an inverse power of the coupling constant λ due to the explicit inclusion of the tadpole contribution. As consequence, the limit of zero Higgs mass, $m_H^2 = 0$, does not exist within the perturbative approach. The importance of the inclusion of the tadpole contribution to restore gauge invariance of on-shell counterterms was recognized a long time ago [101] and was explicitly included in the one-loop electroweak corrections to the matching conditions [32, 84]. The RG equations for the mass parameters were discussed in [90–92, 98, 99].

A.2 $O(\alpha\alpha_s)$ corrections to the relation between the $\overline{\text{MS}}$ and pole masses of the top quark

A detailed discussion and an explicit evaluation¹⁰ have been presented in [89] (the results of [89] were also used for the analysis of the convergence of the series of the set of Feynman diagrams evaluated in [103, 104]). For our analysis is enough to write the following symbolic relation between the $\overline{\text{MS}}$ and pole masses of the top quark:

$$\frac{m_t(\mu^2)}{M_t} = 1 + \sigma_\alpha + \sigma_{\alpha_s} + \sigma_{\alpha_s^2} + \sigma_{\alpha_s^3} + \sigma_{\alpha\alpha_s} + \dots, \quad (\text{A.13})$$

where σ_α and $\sigma_{\alpha\alpha_s}$ are defined by eq. (5.54) or eq. (5.57) of [89].

The pure QCD corrections can be found in [24–26] (only the value of $\sigma_{\alpha_s}(M_t)$ is given there, but the expression for other μ values can be readily reconstructed from the beta functions).

A.3 $O(\alpha\alpha_s)$ corrections to the relation between the $\overline{\text{MS}}$ and pole masses of the Higgs boson

At the two-loop level the relation between the pole and $\overline{\text{MS}}$ masses is defined as follows:

$$\begin{aligned} s_P &= m_0^2 - \Pi_0^{(1)} - \Pi_0^{(2)} - \Pi_0^{(1)} \Pi_0^{(1)'} - \left[\sum_j (\delta m_{j,0}^2)^{(1)} \frac{\partial}{\partial m_{j,0}^2} + \sum_j (\delta g_{j,0})^{(1)} \frac{\partial}{\partial g_{j,0}} \right] \Pi_0^{(1)} \\ &= m_a^2 - \left\{ \Pi_a^{(1)} \right\}_{\overline{\text{MS}}} - \left\{ \Pi_a^{(2)} + \Pi_a^{(1)} \Pi_a^{(1)'} \right\}_{\overline{\text{MS}}}, \end{aligned} \quad (\text{A.14})$$

¹⁰There is typo in eq. (4.46) of [89]: the common factor C_f was lost. The correct result is

$$= \frac{\alpha_s}{4\pi} \frac{e^2}{16\pi^2 \sin^2 \theta_W} C_f \left(\frac{1}{C_f} C_{\alpha\alpha_s}^{(2,2)} \ln^2 \frac{m_t^2}{\mu^2} + \frac{1}{C_f} C_{\alpha\alpha_s}^{(2,1)} \ln \frac{m_t^2}{\mu^2} + \text{without modifications} \right).$$

However, all plots, eq. (5.57) and the Maple program [102] are correct.

where the sum runs over all species of particles, $g_j = \alpha, g_s, (\delta g_{j,0})^{(1)}$ and $(\delta m_{j,0}^2)^{(1)}$ are the one-loop counterterms for the charges and physical masses in the $\overline{\text{MS}}$ scheme and after differentiation we put all parameters equal to their on-shell values. The derivatives in eq. (A.14) correspond to the subtraction of sub-divergences. The genuine two-loop mass counterterm comes from the shift of the m_0^2 term. The relation between the bare and $\overline{\text{MS}}$ masses of the Higgs boson has the form

$$(m_H^B)^2 = (m_H^R(\mu^2))^2 \left[1 + \frac{g^2}{16\pi^2\varepsilon} Z_{H,\alpha} + \frac{g^4}{(16\pi^2)^2} \left(\frac{1}{\varepsilon} Z_{H,\alpha^2}^{(2,1)} + \frac{1}{\varepsilon^2} Z_{H,\alpha^2}^{(2,2)} \right) + \frac{g_s^2 g^2}{(16\pi^2)^2} \left(\frac{1}{\varepsilon} Z_{H,\alpha\alpha_s}^{(2,1)} + \frac{1}{\varepsilon^2} Z_{H,\alpha\alpha_s}^{(2,2)} \right) \right], \quad (\text{A.15})$$

where g is the $\text{SU}(2)$ $\overline{\text{MS}}$ renormalized coupling constant.

The exact analytical result for the $O(\alpha\alpha_s)$ two-loop quark contribution to the Higgs-boson self-energy was calculated in [88, 95, 95]. The bare two-loop mixed QCD-electroweak contribution (with explicit inclusion of the tadpole) for the quark with mass m_q reads:

$$\begin{aligned} \Pi_{0,m_H^2,\alpha\alpha_s,q}^{(2)} = & \frac{g^2 g_s^2}{(16\pi^2)^2} N_c C_f \frac{m_q^2}{m_W^2} \left\{ -J_{0qq}(1, 1, 1; m_H^2)(n-3) \right. \\ & + J_{0qq}(1, 1, 2; m_H^2) [m_H^2 - 4m_q^2] \frac{(n^2 - 5n + 8)}{(n-4)(n-3)} \\ & + A_0(m_q^2) B_0(m_q^2, m_q^2; m_H^2) \frac{(n-2)}{(n-3)(n-4)} \\ & \times \left[(n^3 - 8n^2 + 19n - 16) + \frac{m_H^2}{2m_q^2} (n^2 - 5n + 8) \right] \\ & + [B_0(m_q^2, m_q^2; m_H^2)]^2 \left[m_H^2 \frac{(n-2)^2}{2(n-4)} - m_q^2 \frac{2(n^2 - 4n + 2)}{(n-4)} \right] \\ & \left. + [A_0(m_q^2)]^2 \frac{(n-2)(n^2 - 5n + 8)}{2m_q^2(n-4)(n-3)} + [A_0(m_q^2)]^2 \frac{3}{2} \frac{(n-1)(n-2)^2}{m_q^2(n-3)} \right\}, \end{aligned} \quad (\text{A.16})$$

where the last terms come from the tadpole, n is the dimension of space-time [105] and

$$\begin{aligned} J_{0qq}(a, b, c; m^2) &= \int \frac{d^n(k_1 k_2)}{[(k_1 + k_2 - p)^2]^a [k_1^2 + m_q^2]^b [k_2^2 + m_q^2]^c} \Big|_{p^2 = -m^2}, \\ B_0(m_1^2, m_2^2, m^2) &= \int \frac{d^n k_1}{[k_1^2 + m_1^2][(k_1 - p)^2 + m_2^2]} \Big|_{p^2 = -m^2}, \\ A_0(m^2) &= \int \frac{d^n k_1}{k_1^2 + m^2} \equiv \frac{4(m^2)^{\frac{n}{2}-1}}{(n-2)(n-4)}. \end{aligned} \quad (\text{A.17})$$

In accordance with eq. (A.14), the coefficient $\Delta_{m_H^2,\alpha\alpha_s,q}$ of order $O(\alpha\alpha_s)$ relating the pole

and $\overline{\text{MS}}$ masses of the Higgs boson, $s_p - m_H^2$, can be written as

$$\Delta_{m_H^2, \alpha\alpha_s, q} = \lim_{\varepsilon \rightarrow 0} \left(\frac{g_s^2 g_s^2}{(16\pi^2)^2} \left[\frac{1}{\varepsilon} Z_{H, \alpha\alpha_s, q}^{(2,1)} + \frac{1}{\varepsilon^2} Z_{H, \alpha\alpha_s, q}^{(2,2)} \right] - \frac{g_s^2}{16\pi^2} \frac{1}{\varepsilon} Z_{m_q^2, \alpha_s} m_q^2 \frac{\partial}{\partial m_q^2} \Pi_{0, H, \alpha}^{(1)} - \Pi_{0, m_H^2, \alpha\alpha_s, q}^{(2)} \right), \quad (\text{A.18})$$

where

$$\frac{\partial}{\partial m_q^2} \Pi_{0, H, \alpha}^{(1)} = \frac{N_c}{m_W^2} \frac{g^2}{16\pi^2} \left\{ B_0(m_q^2, m_q^2, m_H^2) \left[\frac{m_H^2 - 2m_q^2(n+1)}{2} \right] - \frac{(3n-2)}{2} A_0(m_q^2) \right\}. \quad (\text{A.19})$$

As result of our calculation we find:

$$Z_{H, \alpha\alpha_s, q}^{(2,1)} = \frac{g_s^2 g_s^2}{(16\pi^2)^2} N_c C_f \frac{5}{4} \frac{m_q^2}{m_W^2}, \quad Z_{H, \alpha\alpha_s, q}^{(2,2)} = -\frac{g_s^2 g_s^2}{(16\pi^2)^2} N_c C_f \frac{3}{2} \frac{m_q^2}{m_W^2}. \quad (\text{A.20})$$

The contributions of other quarks with non-zero mass are additive. Exploring the ε expansion for the master integral J_{0qq} from [106], we have for the t -quark contribution ($q = t$):

$$\begin{aligned} \Delta_{m_H^2, \alpha_s \alpha} \equiv \Delta_{m_H^2, \alpha_s \alpha, t} &= \frac{g_s^2 g_s^2}{(16\pi^2)^2} N_c C_f \frac{m_t^4}{m_W^2} \left\{ \frac{4(z-2)(z-4)}{z} F(y) - \frac{4(1+y)^3}{y(1-y)} G(y) \right. \\ &\quad + \frac{3+20y+16y^2-4y^3-9y^4}{2y(1-y)^2} \ln^2 y + \frac{(1+y)(17+88y+17y^2)}{(1-y)2y} \ln y \\ &\quad \left. + \frac{(131+258y+131y^2)}{8y} - 6\zeta_3 \frac{(1+y)^2(1+y^2)}{y(1-y)^2} \right\} \\ &\quad + C_{H, \alpha\alpha_s}^{(2,2)} \ln^2 \left(\frac{m_t^2}{\mu^2} \right) + C_{H, \alpha\alpha_s}^{(2,1)} \ln \left(\frac{m_t^2}{\mu^2} \right), \end{aligned} \quad (\text{A.21})$$

where

$$z = \frac{m_H^2}{m_t^2}, \quad y = \frac{1 - \sqrt{\frac{z}{z-4}}}{1 + \sqrt{\frac{z}{z-4}}}, \quad z = -\frac{(1-y)^2}{y}, \quad 4m_t^2 - m_H^2 = m_t^2 \frac{(1+y)^2}{y}, \quad (\text{A.22})$$

and we have introduced the two functions $F(y)$ and $G(y)$ (see also [87, 88]) defined as¹¹

$$\begin{aligned} F(y) &= 3 [\text{Li}_3(y) + 2\text{Li}_3(-y)] - 2 \ln y [\text{Li}_2(y) + 2\text{Li}_2(-y)] \\ &\quad - \frac{1}{2} \ln^2 y [\ln(1-y) + 2 \ln(1+y)], \\ G(y) &= [\text{Li}_2(y) + 2\text{Li}_2(-y)] + \ln y [\ln(1-y) + 2 \ln(1+y)], \end{aligned} \quad (\text{A.24})$$

¹¹We cross checked, that eq. (A.21) minus the tadpole contribution coincides with results of refs. [87, 95–97] after the following substitutions:

$$\begin{aligned} r &= \frac{z}{4}, & 1-r &= \frac{(1+y)^2}{4y}, & r_+ &= 1/\sqrt{y}, & r_- &= \sqrt{y}, \\ f &= -\frac{1}{2} \ln y, & g &= \ln(1-y) - 1/2 \ln y, & h &= \ln(1+y) - 1/2 \ln y. \end{aligned} \quad (\text{A.23})$$

and

$$\ln\left(\frac{m_H^2}{m_t^2}\right) = 2\ln(1-y) - \ln y + i\pi. \quad (\text{A.25})$$

In eq. (A.21) we explicitly factorized the RG logarithms, $C_{H,\alpha\alpha_s}^{(2,2)}$ and $C_{H,\alpha\alpha_s}^{(2,1)}$, which may be calculated also from the one-loop result and the mass anomalous dimensions (see [98, 99] for the general case). From the parametrization

$$\begin{aligned} M_H^2 &= m_H^2 + \frac{g^2}{16\pi^2} \left[\Delta X_{H,\alpha}^{(1)} - C_{H,\alpha}^{(1)} L \right] + \frac{g^2 g_s^2}{(16\pi^2)^2} \left[\Delta X_{H,\alpha\alpha_s}^{(2)} + C_{H,\alpha\alpha_s}^{(2,2)} L^2 - C_{H,\alpha\alpha_s}^{(2,1)} L \right] \\ &= m_H^2 + \Delta_{m_H^2,\alpha} + \Delta_{m_H^2,\alpha\alpha_s}, \end{aligned} \quad (\text{A.26})$$

where $L = \ln \frac{\mu^2}{m_t^2}$, and using the fact that pole mass is RG invariant, we have:

$$C_{H,\alpha}^{(1)} = m_H^2 Z_{H,\alpha}, \quad \gamma_{m_t^2,\alpha_s} = Z_{m_t^2,\alpha_s} = -6C_f, \quad (\text{A.27})$$

$$2C_{H,\alpha\alpha_s}^{(2,2)} = Z_{m_t^2,\alpha_s} m_t^2 \frac{\partial}{\partial m_t^2} m_H^2 Z_{H,\alpha} = -3m_H^2 C_f N_c \frac{m_t^2}{m_W^2}, \quad (\text{A.28})$$

$$C_{H,\alpha\alpha_s}^{(2,1)} = m_H^2 \gamma_{H,\alpha\alpha_s} + Z_{m_t^2,\alpha_s} C_{H,\alpha}^{(1)} + Z_{m_t^2,\alpha_s} m_t^2 \frac{\partial}{\partial m_t^2} \Delta X_{H,\alpha}^{(1)}, \quad (\text{A.29})$$

where

$$Z_{H,\alpha} = -\frac{3}{2} - \frac{3}{4} \frac{m_Z^2}{m_W^2} + \frac{3}{4} \frac{m_H^2}{m_W^2} + \sum_{\text{lepton}} \frac{1}{2} \frac{m_l^2}{m_W^2} + N_c \sum_u \frac{1}{2} \frac{m_u^2}{m_W^2} + N_c \sum_d \frac{1}{2} \frac{m_d^2}{m_W^2}. \quad (\text{A.30})$$

In terms of the variable y , defined by eq. (A.22), the final result reads:

$$C_{H,\alpha\alpha_s}^{(2,1)} = -C_f N_c \frac{m_t^4}{m_W^2} \left[3 \frac{(1+y)(1+8y+y^2)}{y(1-y)} \ln y + \frac{(17+38y+17y^2)}{2y} \right]. \quad (\text{A.31})$$

A.4 $O(\alpha\alpha_s)$ corrections to the top Yukawa and Higgs self-couplings

The relation between the top Yukawa (Higgs) coupling and the Fermi constant G_F is obtained from eqs. (A.2), (A.13) and (A.26) as:

$$\frac{y_t^2(\mu^2)}{2\sqrt{2}G_F M_t^2} = \frac{m_t^2(\mu^2)}{M_t^2} \frac{G_F(\mu^2)}{G_F}, \quad (\text{A.32})$$

$$\frac{\lambda(\mu^2)}{\sqrt{2}G_F M_H^2} = \frac{m_H^2(\mu^2)}{M_H^2} \frac{G_F(\mu^2)}{G_F}, \quad (\text{A.33})$$

and the relation between the Higgs coupling constant $\lambda \equiv h_{\text{Sirlin}}$ used in [84] and the parametrization of [89–92] follows from the comparison of the RG functions: $h_{\text{Sirlin}} = \lambda_{\text{Jegerlehner}}(\mu^2)/6$.

The $O(\alpha\alpha_s)$ result for the top Yukawa coupling reads (see eq. (21) in [65] and [107])

$$\begin{aligned}
 & \sqrt{\frac{y_t^2(\mu^2)}{2\sqrt{2}G_F M_t^2}} - 1 = (1 + \sigma_\alpha + \sigma_{\alpha_s} + \sigma_{\alpha\alpha_s}) \\
 & \times \left(1 - \Delta_{G_F, \alpha} - \Delta_{G_F, \alpha\alpha_s} - \sum_f [m_f^2 - M_f^2]_{\alpha_s} \frac{\partial}{\partial m_f^2} \Delta_{G_F, \alpha} \right)^{\frac{1}{2}} \bigg|_{m_J^2 = M_J^2, e^2 = e_{OS}^2} - 1 \\
 & = \left(\sigma_\alpha - \frac{1}{2} \Delta_{G_F, \alpha} + \sigma_{\alpha_s} \right) \bigg|_{m_J^2 = M_J^2, e^2 = e_{OS}^2} \\
 & + \left(\sigma_{\alpha\alpha_s} - \frac{1}{2} \Delta_{G_F, \alpha\alpha_s} - \frac{1}{2} \sigma_{\alpha_s} \Delta_{G_F, \alpha} - \frac{1}{2} \sum_f [m_f^2 - M_f^2]_{\alpha_s} \frac{\partial}{\partial m_f^2} \Delta_{G_F, \alpha}(m_t^2) \right) \bigg|_{m_J^2 = M_J^2, e^2 = e_{OS}^2},
 \end{aligned} \tag{A.34}$$

where σ_X are defined in eq. (A.13). The $O(\alpha\alpha_s)$ result for the Higgs coupling is

$$\begin{aligned}
 & \frac{\lambda(\mu^2)}{\sqrt{2}G_F M_H^2} - 1 = + \left(-\Delta_{G_F, \alpha} - \frac{\Delta_{m_H^2, \alpha}}{M_H^2} \right) \bigg|_{m_J^2 = M_J^2, e^2 = e_{OS}^2} \\
 & + \left(-\Delta_{G_F, \alpha\alpha_s} - \frac{\Delta_{m_H^2, \alpha\alpha_s}}{M_H^2} - [m_t^2 - M_t^2]_{\alpha_s} \frac{\partial}{\partial m_t^2} \left[\Delta_{G_F, \alpha} + \frac{\Delta_{m_H^2, \alpha}}{M_H^2} \right] \right) \bigg|_{m_J^2 = M_J^2, e^2 = e_{OS}^2},
 \end{aligned} \tag{A.35}$$

where

$$[m_t^2 - M_t^2]_{\alpha_s} = -2M_t^2 C_f \frac{g_s^2}{16\pi^2} \left(4 - 3 \ln \frac{M_t^2}{\mu^2} \right),$$

and the sum runs over all quarks.

For completeness, we present also the explicit expressions for the derivatives (for $N_c = 3$, $C_F = 4/3$ and $m_b = 0$):

$$m_t^2 \frac{\partial}{\partial m_t^2} \Delta_{m_H^2, \alpha} = \frac{g^2}{16\pi^2} \frac{3m_t^4}{m_H^2 m_W^2} \left[\frac{1+4y+y^2}{y} + \frac{(1+y)(1+8y+y^2)}{2y(1-y)} \ln y + \frac{1}{2} \frac{m_H^2}{m_t^2} \ln \left(\frac{m_t^2}{\mu^2} \right) \right], \tag{A.36}$$

$$m_t^2 \frac{\partial}{\partial m_t^2} \Delta_{G_F, \alpha} = \frac{g^2}{16\pi^2} \left\{ \frac{m_t^4}{m_W^2 m_H^2} \left(6 - 12 \ln \frac{m_t^2}{\mu^2} \right) + \frac{m_t^2}{m_W^2} \left(\frac{3}{4} + \frac{3}{2} \ln \frac{m_t^2}{\mu^2} \right) \right\}. \tag{A.37}$$

The terms of the orders $O(\alpha)$ and $O(\alpha_s)$ in eq. (A.34) and eq. (A.35) correspond to [32] and [84], respectively. Terms of the order $O(\alpha\alpha_s)$ in eq. (A.34) and eq. (A.35) are the mixed QCD-electroweak corrections and $\Delta_{G_F, \alpha\alpha_s}$, $\sigma_{\alpha\alpha_s}$, $\Delta_{m_H^2, \alpha\alpha_s}$ are defined by eq. (A.6), eq. (A.21), and eq. (5.54) or eq. (5.57) of [89].

For completeness, we present also the coefficient $\Delta_{m_H^2, \alpha}$. We divide all corrections into bosonic (diagrams without any fermions) and fermionic (diagrams exhibiting a fermion loop) ones: $\Delta_{m_H^2, \alpha} = \frac{g^2}{16\pi^2} m_H^2 \left(\Delta_{m_H^2, \alpha, \text{boson}} + \Delta_{m_H^2, \alpha, \text{fermion}} \right)$. Using the nota-

tions of [90–92], we may write the one-loop corrections in the following form

$$\begin{aligned}\Delta_{m_H^2, \alpha, \text{boson}} = & \frac{1}{2} - \frac{1}{2} \ln \frac{m_W^2}{\mu^2} - B(m_W^2, m_W^2; m_H^2) \\ & + \frac{m_H^2}{m_W^2} \left(-\frac{3}{2} + \frac{9}{8} \frac{\pi}{\sqrt{3}} + \frac{3}{8} \ln \frac{m_H^2}{\mu^2} + \frac{1}{4} B(m_W^2, m_W^2; m_H^2) + \frac{1}{8} B(m_Z^2, m_Z^2; m_H^2) \right) \\ & + \frac{m_Z^2}{m_W^2} \left(\frac{1}{4} - \frac{1}{4} \ln \frac{m_Z^2}{\mu^2} - \frac{1}{2} B(m_Z^2, m_Z; m_H^2) \right) \\ & + \frac{m_W^2}{m_H^2} \left(3 - 3 \ln \frac{m_W^2}{\mu^2} + 3 B(m_W^2, m_W; m_H^2) \right) \\ & + \frac{m_Z^4}{m_W^2 m_H^2} \left(\frac{3}{2} - \frac{3}{2} \ln \frac{M_Z^2}{\mu^2} + \frac{3}{2} B(m_Z^2, m_Z; m_H^2) \right),\end{aligned}\tag{A.38}$$

$$\begin{aligned}\Delta_{m_H^2, \alpha, \text{fermion}} = & \frac{1}{2} \frac{m_l^2}{m_W^2} \sum_{\text{lepton}} \left[B_0(m_l^2, m_l^2; m_H^2) \left(1 - 4 \frac{m_l^2}{m_H^2} \right) - 4 \frac{m_l^2}{m_H^2} \left(1 - \ln \frac{m_l^2}{\mu^2} \right) \right] \\ & + \frac{1}{2} \frac{m_q^2}{m_W^2} N_c \sum_{\text{quark}} \left[B_0(m_q^2, m_q^2; m_H^2) \left(1 - 4 \frac{m_q^2}{m_H^2} \right) - 4 \frac{m_q^2}{m_H^2} \left(1 - \ln \frac{m_q^2}{\mu^2} \right) \right],\end{aligned}\tag{A.39}$$

where (see eq. (E.6) in [106])

$$\begin{aligned}B(m^2, m^2; m_H^2) &= \int_0^1 dx \ln \left(\frac{m^2}{\mu^2} x + \frac{m^2}{\mu^2} (1-x) - \frac{m_H^2}{\mu^2} x(1-x) - i0 \right) \\ &= \ln \frac{m^2}{\mu^2} - 2 - \frac{1+Y}{1-Y} \ln Y,\end{aligned}\tag{A.40}$$

with

$$Y = \frac{1 - \sqrt{\frac{r}{r-4}}}{1 + \sqrt{\frac{r}{r-4}}}, \quad r = \frac{m_H^2}{m^2}.$$

All results are collected in the Maple code of [102].

B Beta functions

Two-loop SM beta functions above the top mass are collected in [10] (see [108–117] for original works). The three-loop beta functions can be read off [19, 20].

Below the top mass the one loop beta functions for the gauge couplings were used to evolve the PDG values from M_Z to M_t . For example, for $\alpha(\mu)$ we have

$$\alpha(\mu) = \frac{\alpha(M_Z)}{1 + \frac{11}{6\pi} \alpha(M_Z) \log(\frac{\mu}{m_Z})}.\tag{B.1}$$

The higher-loop corrections are not important numerically for the electroweak constants for the small energy range between M_Z and M_t .




Figure 12. Top Yukawa (left) and Higgs coupling (right) at scale M_t . The constants are extracted by using matching formulas at the scale μ and then evolving to the scale M_t by RG equations. The dotted, dashed and solid lines correspond to matching through orders $O(\alpha_s^2)+O(\alpha)$, $O(\alpha_s^3)+O(\alpha)$ and $O(\alpha_s^3)+O(\alpha)+O(\alpha\alpha_s^2)$, respectively.

For the strong coupling $\alpha_s \equiv g_s^2/(4\pi)$ the RG equation up to order $O(\alpha_s^3)$ is used

$$\frac{d\alpha_s}{d\log\mu} = -(11 - \frac{2}{3}N_f)\frac{\alpha_s^2}{2\pi} - (51 - \frac{19}{3}N_f)\frac{\alpha_s^3}{4\pi^2} - (2857 - \frac{5033}{9}N_f + \frac{325}{27}N_f^2)\frac{\alpha_s^4}{64\pi^3}, \quad (\text{B.2})$$

where $N_f = 5$ is the number of flavors below the top quark. Strictly speaking, the value of $\alpha_s(M_t)$ obtained from this equation should also be shifted to the six-quark value as

$$\alpha_{S,N_f=6}(M_t) = \alpha_{S,N_f=5}(M_t) - \frac{11}{72\pi^2}\alpha_{S,N_f=5}^3(M_t), \quad (\text{B.3})$$

but this introduces a negligible effect (< 0.1 GeV) for the Higgs mass.

In all the formulas of appendix A we use the values of α and α_s at the matching scale μ .

References

- [1] L. Maiani, G. Parisi and R. Petronzio, *Bounds on the Number and Masses of Quarks and Leptons*, *Nucl. Phys.* **B 136** (1978) 115 [[INSPIRE](#)].
- [2] N. Cabibbo, L. Maiani, G. Parisi and R. Petronzio, *Bounds on the Fermions and Higgs Boson Masses in Grand Unified Theories*, *Nucl. Phys.* **B 158** (1979) 295 [[INSPIRE](#)].
- [3] M. Lindner, *Implications of Triviality for the Standard Model*, *Zeit. Phys.* **C31** (1986) 295.
- [4] N. Krasnikov, *Restriction of the Fermion Mass in Gauge Theories of Weak and Electromagnetic Interactions*, *Yad. Fiz.* **28** (1978) 549 [[INSPIRE](#)].
- [5] P.Q. Hung, *Vacuum Instability and New Constraints on Fermion Masses*, *Phys. Rev. Lett.* **42** (1979) 873 [[INSPIRE](#)].
- [6] H.D. Politzer and S. Wolfram, *Bounds on Particle Masses in the Weinberg-Salam Model*, *Phys. Lett.* **B 82** (1979) 242 [Erratum *ibid.* **83B** (1979) 421] [[INSPIRE](#)].
- [7] T. Hambye and K. Riessellmann, *Matching conditions and Higgs mass upper bounds revisited*, *Phys. Rev.* **D 55** (1997) 7255 [[hep-ph/9610272](#)] [[INSPIRE](#)].
- [8] CMS collaboration, S. Chatrchyan et al., *Combined results of searches for the standard model Higgs boson in pp collisions at $\sqrt{s} = 7$ TeV*, *Phys. Lett.* **B 710** (2012) 26 [[arXiv:1202.1488](#)] [[INSPIRE](#)].

- [9] ATLAS collaboration, G. Aad et al., *Combined search for the Standard Model Higgs boson using up to 4.9fb^{-1} of pp collision data at $\sqrt{s}=7\text{TeV}$ with the ATLAS detector at the LHC*, *Phys. Lett. B* **710** (2012) 49 [[arXiv:1202.1408](#)] [[INSPIRE](#)].
- [10] J. Espinosa, G. Giudice and A. Riotto, *Cosmological implications of the Higgs mass measurement*, *JCAP* **05** (2008) 002 [[arXiv:0710.2484](#)] [[INSPIRE](#)].
- [11] F. Bezrukov and M. Shaposhnikov, *The Standard Model Higgs boson as the inflaton*, *Phys. Lett. B* **659** (2008) 703 [[arXiv:0710.3755](#)] [[INSPIRE](#)].
- [12] F. Bezrukov and M. Shaposhnikov, *Standard Model Higgs boson mass from inflation: Two loop analysis*, *JHEP* **07** (2009) 089 [[arXiv:0904.1537](#)] [[INSPIRE](#)].
- [13] M. Shaposhnikov and C. Wetterich, *Asymptotic safety of gravity and the Higgs boson mass*, *Phys. Lett. B* **683** (2010) 196 [[arXiv:0912.0208](#)] [[INSPIRE](#)].
- [14] G. Altarelli and G. Isidori, *Lower limit on the Higgs mass in the standard model: An Update*, *Phys. Lett. B* **337** (1994) 141 [[INSPIRE](#)].
- [15] J. Casas, J. Espinosa and M. Quirós, *Improved Higgs mass stability bound in the standard model and implications for supersymmetry*, *Phys. Lett. B* **342** (1995) 171 [[hep-ph/9409458](#)] [[INSPIRE](#)].
- [16] J. Casas, J. Espinosa and M. Quirós, *Standard model stability bounds for new physics within LHC reach*, *Phys. Lett. B* **382** (1996) 374 [[hep-ph/9603227](#)] [[INSPIRE](#)].
- [17] M. Holthausen, K.S. Lim and M. Lindner, *Planck scale Boundary Conditions and the Higgs Mass*, *JHEP* **02** (2012) 037 [[arXiv:1112.2415](#)] [[INSPIRE](#)].
- [18] J. Elias-Miro et al., *Higgs mass implications on the stability of the electroweak vacuum*, *Phys. Lett. B* **709** (2012) 222 [[arXiv:1112.3022](#)] [[INSPIRE](#)].
- [19] L.N. Mihaila, J. Salomon and M. Steinhauser, *Gauge Coupling β -functions in the Standard Model to Three Loops*, *Phys. Rev. Lett.* **108** (2012) 151602 [[arXiv:1201.5868](#)] [[INSPIRE](#)].
- [20] K. Chetyrkin and M. Zoller, *Three-loop β -functions for top-Yukawa and the Higgs self-interaction in the Standard Model*, *JHEP* **06** (2012) 033 [[arXiv:1205.2892](#)] [[INSPIRE](#)].
- [21] F.L. Bezrukov, A. Magnin and M. Shaposhnikov, *Standard Model Higgs boson mass from inflation*, *Phys. Lett. B* **675** (2009) 88 [[arXiv:0812.4950](#)] [[INSPIRE](#)].
- [22] S.R. Coleman and E.J. Weinberg, *Radiative Corrections as the Origin of Spontaneous Symmetry Breaking*, *Phys. Rev. D* **7** (1973) 1888 [[INSPIRE](#)].
- [23] J. Ellis, J. Espinosa, G. Giudice, A. Hoecker and A. Riotto, *The Probable Fate of the Standard Model*, *Phys. Lett. B* **679** (2009) 369 [[arXiv:0906.0954](#)] [[INSPIRE](#)].
- [24] K. Chetyrkin and M. Steinhauser, *Short distance mass of a heavy quark at order α_s^3* , *Phys. Rev. Lett.* **83** (1999) 4001 [[hep-ph/9907509](#)] [[INSPIRE](#)].
- [25] K. Chetyrkin and M. Steinhauser, *The Relation between the $\overline{\text{MS}}$ and the on-shell quark mass at order α_s^3* , *Nucl. Phys. B* **573** (2000) 617 [[hep-ph/9911434](#)] [[INSPIRE](#)].
- [26] K. Melnikov and T.v. Ritbergen, *The Three loop relation between the $\overline{\text{MS}}$ and the pole quark masses*, *Phys. Lett. B* **482** (2000) 99 [[hep-ph/9912391](#)] [[INSPIRE](#)].
- [27] A. Kataev and V. Kim, *Uncertainties of QCD predictions for Higgs boson decay into bottom quarks at NNLO and beyond*, *PoS(ACAT08)004* [[arXiv:0902.1442](#)] [[INSPIRE](#)].
- [28] A. Kataev and V. Kim, *Peculiar features of the relations between pole and running heavy quark masses and estimates of the $O(\alpha_s^4)$ contributions*, *Phys. Part. Nucl.* **41** (2010) 946 [[arXiv:1001.4207](#)] [[INSPIRE](#)].

- [29] M.C. Smith and S.S. Willenbrock, *Top quark pole mass*, *Phys. Rev. Lett.* **79** (1997) 3825 [[hep-ph/9612329](#)] [[INSPIRE](#)].
- [30] A.H. Hoang, A. Jain, I. Scimemi and I.W. Stewart, *Infrared Renormalization Group Flow for Heavy Quark Masses*, *Phys. Rev. Lett.* **101** (2008) 151602 [[arXiv:0803.4214](#)] [[INSPIRE](#)].
- [31] A.H. Hoang, A. Jain, I. Scimemi and I.W. Stewart, *R-evolution: Improving perturbative QCD*, *Phys. Rev. D* **82** (2010) 011501 [[arXiv:0908.3189](#)] [[INSPIRE](#)].
- [32] R. Hempfling and B.A. Kniehl, *On the relation between the fermion pole mass and \overline{MS} Yukawa coupling in the standard model*, *Phys. Rev. D* **51** (1995) 1386 [[hep-ph/9408313](#)] [[INSPIRE](#)].
- [33] PARTICLE DATA GROUP collaboration, K. Nakamura et al., *Review of particle physics*, *J. Phys. G* **37** (2010) 075021 [[INSPIRE](#)].
- [34] S. Alekhin, J. Blumlein and S. Moch, *Parton Distribution Functions and Benchmark Cross sections at NNLO*, *Phys. Rev. D* **86** (2012) 054009 [[arXiv:1202.2281](#)] [[INSPIRE](#)].
- [35] G. Watt, *Parton distribution function dependence of benchmark Standard Model total cross sections at the 7 TeV LHC*, *JHEP* **09** (2011) 069 [[arXiv:1106.5788](#)] [[INSPIRE](#)].
- [36] G. Watt, *MSTW PDFs and impact of PDFs on cross sections at Tevatron and LHC*, *Nucl. Phys. Proc. Suppl.* **222-224** (2012) 61 [[arXiv:1201.1295](#)] [[INSPIRE](#)].
- [37] U. Langenfeld, S. Moch and P. Uwer, *Measuring the running top-quark mass*, *Phys. Rev. D* **80** (2009) 054009 [[arXiv:0906.5273](#)] [[INSPIRE](#)].
- [38] TEVATRON ELECTROWEAK WORKING GROUP, CDF and D0 collaborations, *Combination of CDF and D0 results on the mass of the top quark using up to 5.8fb^{-1} of data*, [arXiv:1107.5255](#) [[INSPIRE](#)].
- [39] D0 collaboration, V.M. Abazov et al., *Determination of the pole and \overline{MS} masses of the top quark from the $t\bar{t}$ cross section*, *Phys. Lett. B* **703** (2011) 422 [[arXiv:1104.2887](#)] [[INSPIRE](#)].
- [40] D. Bennett and H.B. Nielsen, *Predictions for nonAbelian fine structure constants from multicriticality*, *Int. J. Mod. Phys. A* **9** (1994) 5155 [[hep-ph/9311321](#)] [[INSPIRE](#)].
- [41] C. Froggatt and H.B. Nielsen, *Standard model criticality prediction: Top mass $173 \pm 5\text{ GeV}$ and Higgs mass $135 \pm 9\text{ GeV}$* , *Phys. Lett. B* **368** (1996) 96 [[hep-ph/9511371](#)] [[INSPIRE](#)].
- [42] G. Isidori, V.S. Rychkov, A. Strumia and N. Tetradis, *Gravitational corrections to standard model vacuum decay*, *Phys. Rev. D* **77** (2008) 025034 [[arXiv:0712.0242](#)] [[INSPIRE](#)].
- [43] I. Masina and A. Notari, *The Higgs mass range from Standard Model false vacuum Inflation in scalar-tensor gravity*, *Phys. Rev. D* **85** (2012) 123506 [[arXiv:1112.2659](#)] [[INSPIRE](#)].
- [44] I. Masina and A. Notari, *Standard Model False Vacuum Inflation: Correlating the Tensor-to-Scalar Ratio to the Top Quark and Higgs Boson masses*, *Phys. Rev. Lett.* **108** (2012) 191302 [[arXiv:1112.5430](#)] [[INSPIRE](#)].
- [45] I. Masina and A. Notari, *Inflation from the Higgs field false vacuum with hybrid potential*, [arXiv:1204.4155](#) [[INSPIRE](#)].
- [46] S. Weinberg, *Ultraviolet divergences in quantum theories of gravity*, in *General Relativity: An Einstein centenary survey*, S.W. Hawking and W. Israel eds., Cambridge University Press, Cambridge, U.K. (1979), pg. 790.
- [47] M. Niedermaier and M. Reuter, *The Asymptotic Safety Scenario in Quantum Gravity*, *Living Rev. Rel.* **9** (2006) 5.

- [48] M. Reuter, *Nonperturbative evolution equation for quantum gravity*, *Phys. Rev. D* **57** (1998) 971 [[hep-th/9605030](#)] [[INSPIRE](#)].
- [49] R. Percacci and D. Perini, *Asymptotic safety of gravity coupled to matter*, *Phys. Rev. D* **68** (2003) 044018 [[hep-th/0304222](#)] [[INSPIRE](#)].
- [50] G. Narain and R. Percacci, *Renormalization Group Flow in Scalar-Tensor Theories. I*, *Class. Quant. Grav.* **27** (2010) 075001 [[arXiv:0911.0386](#)] [[INSPIRE](#)].
- [51] A.A. Starobinsky, *A New Type of Isotropic Cosmological Models Without Singularity*, *Phys. Lett. B* **91** (1980) 99 [[INSPIRE](#)].
- [52] D. Gorbunov and A. Panin, *Scalaron the mighty: producing dark matter and baryon asymmetry at reheating*, *Phys. Lett. B* **700** (2011) 157 [[arXiv:1009.2448](#)] [[INSPIRE](#)].
- [53] F. Bezrukov and D. Gorbunov, *Distinguishing between R^2 -inflation and Higgs-inflation*, *Phys. Lett. B* **713** (2012) 365 [[arXiv:1111.4397](#)] [[INSPIRE](#)].
- [54] F. Bezrukov, A. Magnin, M. Shaposhnikov and S. Sibiryakov, *Higgs inflation: consistency and generalisations*, *JHEP* **01** (2011) 016 [[arXiv:1008.5157](#)] [[INSPIRE](#)].
- [55] A. Barvinsky, A.Y. Kamenshchik, C. Kiefer, A. Starobinsky and C. Steinwachs, *Asymptotic freedom in inflationary cosmology with a non-minimally coupled Higgs field*, *JCAP* **12** (2009) 003 [[arXiv:0904.1698](#)] [[INSPIRE](#)].
- [56] C. Burgess, H.M. Lee and M. Trott, *Power-counting and the Validity of the Classical Approximation During Inflation*, *JHEP* **09** (2009) 103 [[arXiv:0902.4465](#)] [[INSPIRE](#)].
- [57] J. Barbon and J. Espinosa, *On the Naturalness of Higgs Inflation*, *Phys. Rev. D* **79** (2009) 081302 [[arXiv:0903.0355](#)] [[INSPIRE](#)].
- [58] C. Burgess, H.M. Lee and M. Trott, *Comment on Higgs Inflation and Naturalness*, *JHEP* **07** (2010) 007 [[arXiv:1002.2730](#)] [[INSPIRE](#)].
- [59] M.P. Hertzberg, *On Inflation with Non-minimal Coupling*, *JHEP* **11** (2010) 023 [[arXiv:1002.2995](#)] [[INSPIRE](#)].
- [60] M. Shaposhnikov, *Is there a new physics between electroweak and Planck scales?*, [arXiv:0708.3550](#) [[INSPIRE](#)].
- [61] A. Boyarsky, O. Ruchayskiy and M. Shaposhnikov, *The Role of sterile neutrinos in cosmology and astrophysics*, *Ann. Rev. Nucl. Part. Sci.* **59** (2009) 191 [[arXiv:0901.0011](#)] [[INSPIRE](#)].
- [62] M. Shaposhnikov and D. Zenhausern, *Scale invariance, unimodular gravity and dark energy*, *Phys. Lett. B* **671** (2009) 187 [[arXiv:0809.3395](#)] [[INSPIRE](#)].
- [63] M. Shaposhnikov and D. Zenhausern, *Quantum scale invariance, cosmological constant and hierarchy problem*, *Phys. Lett. B* **671** (2009) 162 [[arXiv:0809.3406](#)] [[INSPIRE](#)].
- [64] D. Gorbunov and M. Shaposhnikov, *How to find neutral leptons of the ν MSM?*, *JHEP* **10** (2007) 015 [[arXiv:0705.1729](#)] [[INSPIRE](#)].
- [65] F. Jegerlehner and M. Kalmykov, *$O(\alpha\alpha_s)$ relation between pole- and $\overline{\text{MS}}$ mass of the t quark*, *Acta Phys. Polon. B* **34** (2003) 5335 [[hep-ph/0310361](#)] [[INSPIRE](#)].
- [66] ATLAS collaboration, G. Aad et al., *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, *Phys. Lett. B* **716** (2012) 1 [[arXiv:1207.7214](#)] [[INSPIRE](#)].
- [67] CMS collaboration, S. Chatrchyan et al., *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, *Phys. Lett. B* **716** (2012) 30 [[arXiv:1207.7235](#)] [[INSPIRE](#)].

- [68] T. Muller, *New results from the top quark*, in *ICHEP 2012 Conference*, Melbourne (2012) [<http://indico.cern.ch/getFile.py/access?contribId=7&sessionId=17&resId=1&materialId=slides&confId=181298>].
- [69] ECFA/DESY LC PHYSICS WORKING GROUP collaboration, E. Accomando et al., *Physics with e^+e^- linear colliders*, *Phys. Rept.* **299** (1998) 1 [[hep-ph/9705442](#)] [[INSPIRE](#)].
- [70] ECFA/DESY LC PHYSICS WORKING GROUP collaboration, J. Aguilar-Saavedra et al., *TESLA: The Superconducting electron positron linear collider with an integrated x-ray laser laboratory. Technical design report. Part 3. Physics at an e^+e^- linear collider*, [hep-ph/0106315](#) [[INSPIRE](#)].
- [71] AMERICAN LINEAR COLLIDER WORKING GROUP collaboration, T. Abe et al., *Linear Collider Physics Resource Book for Snowmass 2001 - Part 1: Introduction*, [hep-ex/0106055](#).
- [72] AMERICAN LINEAR COLLIDER WORKING GROUP collaboration, T. Abe et al., *Linear Collider Physics Resource Book for Snowmass 2001 - Part 2: Higgs and Supersymmetry Studies*, [hep-ex/0106056](#) [[INSPIRE](#)].
- [73] T. Abe, *Linear Collider Physics Resource Book for Snowmass 2001 - Part 3: Studies of Exotic and Standard Model Physics*, [hep-ex/0106057](#).
- [74] AMERICAN LINEAR COLLIDER WORKING GROUP collaboration, T. Abe et al., *Linear Collider Physics Resource Book for Snowmass 2001 - Part 4: Theoretical, Accelerator and Experimental Options*, [hep-ex/0106058](#).
- [75] ACFA LINEAR COLLIDER WORKING GROUP collaboration, K. Abe et al., *Particle physics experiments at JLC*, [hep-ph/0109166](#) [[INSPIRE](#)].
- [76] ILC collaboration, G. Aarons et al., *International Linear Collider Reference Design Report Volume 2: Physics at the ILC*, [arXiv:0709.1893](#) [[INSPIRE](#)].
- [77] M. Winter, *Determination of the Strong Coupling Constant at Giga-Z*, ILC note [PHSM-2001-016](#).
- [78] ATLAS AND CMS collaborations, S. Blyweert, *Top-quark mass measurements at the LHC*, [arXiv:1205.2175](#) [[INSPIRE](#)].
- [79] C. Schwanenberger, *Measurements of the inclusive cross section and of differential distributions in top quark pair production ($D0$)*, in *ICHEP 2012 Conference*, Melbourne (2012), [<https://indico.cern.ch/getFile.py/access?contribId=534&sessionId=67&resId=0&materialId=slides&confId=181298>].
- [80] A. Sirlin, *Radiative Corrections in the $SU(2)_L \times U(1)$ Theory: A Simple Renormalization Framework*, *Phys. Rev.* **D 22** (1980) 971 [[INSPIRE](#)].
- [81] W. Marciano and A. Sirlin, *Radiative Corrections to Neutrino Induced Neutral Current Phenomena in the $SU(2)_L \times U(1)$ Theory*, *Phys. Rev.* **D 22** (1980) 2695 [Erratum *ibid.* **D 31** (1985) 213] [[INSPIRE](#)].
- [82] G. Burgers and F. Jegerlehner, *Δr or the relation between the electroweak couplings and the weak vector boson masses*, *Conf.Proc.* **C8902201** (1989) 55.
- [83] S. Actis and G. Passarino, *Two-Loop Renormalization in the Standard Model Part III: Renormalization Equations and their Solutions*, *Nucl. Phys.* **B 777** (2007) 100 [[hep-ph/0612124](#)] [[INSPIRE](#)].
- [84] A. Sirlin and R. Zucchini, *Dependence of the quartic coupling $\bar{h}_{\overline{MS}}(M)$ on m_H and the possible onset of new physics in the higgs sector of the standard model*, *Nucl. Phys.* **B 266** (1986) 389 [[INSPIRE](#)].

- [85] T. Chang, K. Gaemers and W. van Neerven, *QCD Corrections to the Mass and Width of the Intermediate Vector Bosons*, *Nucl. Phys. B* **202** (1982) 407 [INSPIRE].
- [86] A. Djouadi, *$O(\alpha_s)$ Vacuum Polarization Functions of the Standard Model Gauge Bosons*, *Nuovo Cim. A* **100** (1988) 357 [INSPIRE].
- [87] B.A. Kniehl, *Two Loop Corrections to the Vacuum Polarizations in Perturbative QCD*, *Nucl. Phys. B* **347** (1990) 86 [INSPIRE].
- [88] A. Djouadi and P. Gambino, *QCD corrections to Higgs boson selfenergies and fermionic decay widths*, *Phys. Rev. D* **51** (1995) 218 [Erratum *ibid.* **D 53** (1996) 4111] [[hep-ph/9406431](#)] [INSPIRE].
- [89] F. Jegerlehner and M.Y. Kalmykov, *$O(\alpha_s)$ correction to the pole mass of the t quark within the standard model*, *Nucl. Phys. B* **676** (2004) 365 [[hep-ph/0308216](#)] [INSPIRE].
- [90] F. Jegerlehner, M.Y. Kalmykov and O. Veretin, *\overline{MS} versus pole masses of gauge bosons: Electroweak bosonic two loop corrections*, *Nucl. Phys. B* **641** (2002) 285 [[hep-ph/0105304](#)] [INSPIRE].
- [91] F. Jegerlehner, M.Y. Kalmykov and O. Veretin, *Full two loop electroweak corrections to the pole masses of gauge bosons*, *Nucl. Phys. Proc. Suppl.* **116** (2003) 382 [[hep-ph/0212003](#)] [INSPIRE].
- [92] F. Jegerlehner, M.Y. Kalmykov and O. Veretin, *\overline{MS} versus pole masses of gauge bosons. 2. Two loop electroweak fermion corrections*, *Nucl. Phys. B* **658** (2003) 49 [[hep-ph/0212319](#)] [INSPIRE].
- [93] F. Halzen and B.A. Kniehl, *Δr beyond one loop*, *Nucl. Phys. B* **353** (1991) 567 [INSPIRE].
- [94] S. Fanchiotti, B.A. Kniehl and A. Sirlin, *Incorporation of QCD effects in basic corrections of the electroweak theory*, *Phys. Rev. D* **48** (1993) 307 [[hep-ph/9212285](#)] [INSPIRE].
- [95] B.A. Kniehl and A. Sirlin, *Comparative analysis of three methods to evaluate vacuum polarization functions*, *Phys. Lett. B* **318** (1993) 367 [INSPIRE].
- [96] B.A. Kniehl, *Two loop $O(\alpha_s G_F M_t^2)$ corrections to the fermionic decay rates of the standard model Higgs boson*, *Phys. Rev. D* **50** (1994) 3314 [[hep-ph/9405299](#)] [INSPIRE].
- [97] A. Djouadi and P. Gambino, *Electroweak gauge bosons selfenergies: Complete QCD corrections*, *Phys. Rev. D* **49** (1994) 3499 [Erratum *ibid.* **D 53** (1996) 4111] [[hep-ph/9309298](#)] [INSPIRE].
- [98] F. Jegerlehner, M.Y. Kalmykov and O. Veretin, *Pole masses of gauge bosons: 2 loop electroweak corrections*, *Nucl. Instrum. Meth. A* **502** (2003) 618 [INSPIRE].
- [99] F. Jegerlehner and M.Y. Kalmykov, *Steps towards full two-loop calculations for 2 fermion to 2 fermion processes: Running versus pole masses schemes*, *Nucl. Instrum. Meth. A* **534** (2004) 299 [[hep-ph/0404213](#)] [INSPIRE].
- [100] H. Arason et al., *Renormalization group study of the standard model and its extensions. 1. The Standard model*, *Phys. Rev. D* **46** (1992) 3945 [INSPIRE].
- [101] J. Fleischer and F. Jegerlehner, *Radiative Corrections to Higgs Decays in the Extended Weinberg-Salam Model*, *Phys. Rev. D* **23** (1981) 2001 [INSPIRE].
- [102] M.Y. Kalmykov, <http://theor.jinr.ru/~kalmykov/pole/pole.html>.
- [103] M. Faisst, J.H. Kuhn and O. Veretin, *Pole versus MS mass definitions in the electroweak theory*, *Phys. Lett. B* **589** (2004) 35 [[hep-ph/0403026](#)] [INSPIRE].

- [104] D. Eiras and M. Steinhauser, *Two-loop $O(\alpha\alpha_s)$ corrections to the on-shell fermion propagator in the standard model*, *JHEP* **02** (2006) 010 [[hep-ph/0512099](#)] [[INSPIRE](#)].
- [105] G. 't Hooft and M. Veltman, *Regularization and Renormalization of Gauge Fields*, *Nucl. Phys. B* **44** (1972) 189 [[INSPIRE](#)].
- [106] A.I. Davydychev and M.Y. Kalmykov, *Massive Feynman diagrams and inverse binomial sums*, *Nucl. Phys. B* **699** (2004) 3 [[hep-th/0303162](#)] [[INSPIRE](#)].
- [107] B.A. Kniehl, J.H. Piclum and M. Steinhauser, *Relation between bottom-quark $\overline{\text{MS}}$ Yukawa coupling and pole mass*, *Nucl. Phys. B* **695** (2004) 199 [[hep-ph/0406254](#)] [[INSPIRE](#)].
- [108] M. Fischler and J. Oliensis, *Two Loop Corrections to the Evolution of the Higgs-Yukawa Coupling Constant*, *Phys. Lett. B* **119** (1982) 385 [[INSPIRE](#)].
- [109] M. Fischler and J. Oliensis, *Detailed Calculation of the Complete Two Loop Higgs-Yukawa β -function in an Arbitrary alpha Gauge*, *Phys. Rev. D* **28** (1983) 2027 [[INSPIRE](#)].
- [110] C. Ford, D. Jones, P. Stephenson and M. Einhorn, *The Effective potential and the renormalization group*, *Nucl. Phys. B* **395** (1993) 17 [[hep-lat/9210033](#)] [[INSPIRE](#)].
- [111] I. Jack, *Two loop background field calculations for gauge theories with scalar fields*, *J. Phys. A* **16** (1983) 1083 [[INSPIRE](#)].
- [112] I. Jack and H. Osborn, *General two loop β -functions for gauge theories with arbitrary scalar fields*, *J. Phys. A* **16** (1983) 1101 [[INSPIRE](#)].
- [113] I. Jack, *two loop β -functions for supersymmetric gauge theories*, *Phys. Lett. B* **147** (1984) 405 [[INSPIRE](#)].
- [114] I. Jack and H. Osborn, *general background field calculations with fermion fields*, *Nucl. Phys. B* **249** (1985) 472 [[INSPIRE](#)].
- [115] M.E. Machacek and M.T. Vaughn, *Two Loop Renormalization Group Equations in a General Quantum Field Theory. 1. Wave Function Renormalization*, *Nucl. Phys. B* **222** (1983) 83 [[INSPIRE](#)].
- [116] M.E. Machacek and M.T. Vaughn, *Two-loop renormalization group equations in a general quantum field theory (II). Yukawa couplings*, *Nucl. Phys. B* **236** (1984) 221.
- [117] M.E. Machacek and M.T. Vaughn, *Two-loop renormalization group equations in a general quantum field theory (III). Scalar quartic couplings*, *Nucl. Phys. B* **249** (1985) 70.
- [118] A. Strumia, *Higgs weights 125 GeV! Now what?*, talk given at *Implications of LHC results for TeV-scale physics*, CERN, Geneva, 28 March 2012 [<http://indico.cern.ch/materialDisplay.py?contribId=47&sessionId=20&materialId=slides&confId=162621>].
- [119] G. Degrand et al., *Higgs mass and vacuum stability in the Standard Model at NNLO*, *JHEP* **08** (2012) 098 [[arXiv:1205.6497](#)] [[INSPIRE](#)].
- [120] S. Alekhin, A. Djouadi and S. Moch, *The top quark and Higgs boson masses and the stability of the electroweak vacuum*, *Phys. Lett. B* **716** (2012) 214 [[arXiv:1207.0980](#)] [[INSPIRE](#)].