

Hidden SUSY from precision gauge unification

Sven Krippendorff^{1a}, Hans Peter Nilles^{2a}, Michael Ratz^{3b},
Martin Wolfgang Winkler^{4c}

^a *Bethe Center for Theoretical Physics
and
Physikalisches Institut der Universität Bonn,
Nussallee 12, 53115 Bonn, Germany*

^b *Physik-Department T30, Technische Universität München,
James-Frank-Straße, 85748 Garching, Germany*

^c *Deutsches Elektronen-Synchrotron DESY, Notkestrasse 85, 22607 Hamburg, Germany.*

Abstract

We revisit the implications of naturalness and gauge unification in the MSSM. We find that precision unification of the couplings in connection with a small μ parameter requires a highly compressed gaugino pattern as it is realized in mirage mediation. Due to the small mass difference between gluino and LSP, collider limits on the gluino mass are drastically relaxed. Without further assumptions, the relic density of the LSP is very close to the observed dark matter density due to coannihilation effects.

¹Email: krippendorf@th.physik.uni-bonn.de

²Email: nilles@th.physik.uni-bonn.de

³Email: michael.ratz@tum.de

⁴Email: martin.winkler@desy.de

Contents

1	Introduction	1
2	The effective SUSY threshold scale	2
3	Gauge coupling unification in MSSM models	4
3.1	Uncompressed gaugino mass spectra	5
3.2	Compressed gaugino mass spectra	6
4	Implications for the LHC and dark matter	7
4.1	LHC discovery potential	7
4.2	Dark Matter	12
5	Conclusions	14
A	Dependence of PGU on $\tan\beta$ and $m_{3/2}$	16

1 Introduction

Two key motivations for studying low-energy supersymmetry are the possibility for a natural solution to the electroweak hierarchy problem [1] and gauge coupling unification [2]. Given this input of a natural solution to the electroweak hierarchy problem and gauge coupling unification, we analyze in this article their effect on the spectrum of supersymmetric soft masses. Here we concentrate on the minimal supersymmetric extension of the standard model (MSSM).

If one ponders about the question whether or not gauge couplings really unify in the MSSM, it turns out that there is a non-negligible dependence on the pattern of supersymmetry breaking. In particular, in some of the most popular scenarios, such as the CMSSM, gauge coupling unification is not precise given soft masses around the weak scale. This refers to the fact that the strong coupling $\alpha_3 = g_3^2/(4\pi)$ turns out to be about 3 % smaller than α_1 and α_2 at M_{GUT} , which is defined as the scale where α_1 and α_2 unify.

Of course, this discrepancy of the couplings at M_{GUT} may originate from some thresholds at the high scale [3] (see also [4, S. Raby, ‘Grand unified theories’]). However, here we would like to discuss precision gauge coupling unification (PGU) in the absence of high-scale threshold corrections¹ and determine the consequences for the spectrum of soft masses.

Several examples of spectra achieving PGU have been discussed in the literature [5–7]. Most of these are now excluded by the non-observation of superpartners at the LHC or by the relatively large mass of the Higgs boson. The consideration of naturalness leads us to the choice of a small μ parameter which (further) narrows down viable soft

¹The size of these threshold corrections is ultimately set by the UV theory of choice. For a bottom-up analysis we take it as a UV parameter which we here assume to be vanishing.

mass spectra. Remarkably, we are led to a highly compressed pattern of gaugino and higgsino masses at or below the TeV scale, while the scalar superpartners can be out of reach for the LHC experiment. As the LHC sensitivity is drastically reduced in case of a small mass splitting between gluino and the lightest supersymmetric particle (LSP), PGU provides an attractive explanation on why SUSY has not been discovered so far.

The required spectrum of gaugino masses can be accommodated in mirage mediation and as such is directly connected to UV scenarios arising within string theory [8–12]. Fixing the ratio of gaugino masses with PGU opens interesting windows towards explicit UV realizations of mirage mediation.

Further, the problem of dark matter overproduction — usually arising in SUSY models with a bino LSP — can naturally be solved by a compressed gaugino spectrum. We find that in mirage mediation, PGU is intimately linked to the occurrence of coannihilations. These allow for a consistent explanation of dark matter in terms of the lightest neutralino.

This article is organized as follows: in section 2 we perform a model-independent analytical discussion of gauge coupling unification. In section 3, we study whether or not PGU can be achieved in realistic SUSY models taking the CMSSM and mirage mediation as examples. Then, we turn to the LHC and dark matter phenomenology of models with successful PGU in section 4, before concluding in section 5.

2 The effective SUSY threshold scale

The MSSM gauge couplings evaluated at the scale of grand unification can be written as [13]

$$\frac{1}{g_i^2(M_{\text{GUT}})} = \frac{1}{g_i^2(M_Z)} - \frac{b_i^{\text{MSSM}}}{8\pi^2} \ln \left(\frac{M_{\text{GUT}}}{M_Z} \right) + \frac{1}{g_{i,\text{Thr}}^2} + \Delta_i \quad (1)$$

where the b_i^{MSSM} denote the standard MSSM beta function coefficients, $(b_1, b_2, b_3) = (33/5, 1, -3)$, while Δ_i includes two-loop effects, threshold corrections related to the heavy standard model fields as well as a possible term required for the transition between renormalization schemes — the gauge couplings $g_i^2(M_Z)$ are typically given in the $\overline{\text{MS}}$ rather than the $\overline{\text{DR}}$ scheme. The threshold corrections within the MSSM read [13]

$$\frac{1}{g_{i,\text{Thr}}^2} = \sum_{\eta} \frac{b_i^{\eta}}{8\pi^2} \ln \left(\frac{M_{\eta}}{M_Z} \right), \quad (2)$$

where the sum runs over all sparticles and heavy Higgs fields with mass M_{η} evaluated at the low scale and b_i^{η} denoting the contribution of the particle η to the i^{th} β -function coefficient (cf. [14] for the exact expressions of b_i^{η} in the MSSM).²

To arrive at a consistent picture of unification, the g_i are required to meet in a single point. Assuming the absence of threshold corrections induced by heavy fields of a grand

²Here we neglect two-loop contributions to the thresholds.

unified theory (GUT), this implies that the masses of the MSSM fields must arrange such that their overall threshold to the g_i allows for PGU.

To get a first impression on the effect of soft masses on gauge coupling unification, let us make the ad hoc assumption that all sparticles as well as the heavy Higgs doublet have a common mass T_{SUSY} which we call the SUSY threshold scale. In this case, the overall threshold correction (2) to the gauge couplings takes a very simple form. We find

$$\frac{1}{g_{i,\text{Thr}}^2} = \frac{b_i^{\text{MSSM}} - b_i^{\text{SM}}}{8\pi^2} \ln \left(\frac{T_{\text{SUSY}}}{M_Z} \right), \quad (3)$$

where b_i^{SM} stands for the beta function coefficients of the standard model. Now we can determine the SUSY threshold scale for which the gauge couplings meet exactly. To quantify the deviation from PGU, we follow [7, 15] and introduce

$$\epsilon_3 = \frac{g_3^2(M_{\text{GUT}}) - g_{1,2}^2(M_{\text{GUT}})}{g_{1,2}^2(M_{\text{GUT}})}, \quad (4)$$

where we define M_{GUT} as the scale at which g_1 and g_2 meet.

In figure 1 we depict ϵ_3 as a function of the SUSY threshold scale. For the determination of ϵ_3 , we have used Softsusy (version 3.3.2) [16], where we set the strong coupling strength to $\alpha_s(M_Z) = 0.1184 \pm 0.0007$ [17] and $\tan \beta = 10$. It can be seen that the three gauge couplings would exactly meet in a single point if all superpartners as well as the heavy Higgs doublet had a common mass $T_{\text{SUSY}} \simeq 2 \text{ TeV}$. Note that the required value of T_{SUSY} has a very mild dependence on $\tan \beta$ which only affects the gauge coupling at the two-loop level.




Figure 1: Deviation from PGU as a function of T_{SUSY} . The width of the band corresponds to the 1σ experimental error in $\alpha_s(M_Z)$.

Turning to realistic SUSY models, despite the fact that not all superpartners have a common mass, it is still possible to define an effective SUSY threshold scale T_{SUSY}

through the relation

$$\frac{b_i^{\text{MSSM}} - b_i^{\text{SM}}}{8\pi^2} \ln \left(\frac{T_{\text{SUSY}}}{M_Z} \right) \equiv \sum_{\eta} \frac{b_i^{\eta}}{8\pi^2} \ln \left(\frac{M_{\eta}}{M_Z} \right) + (c_1 + c_2 b_i^{\text{MSSM}}) \quad (5)$$

with the same $(T_{\text{SUSY}}, c_1, c_2)$ for all three standard model gauge groups. Note that the term c_1 accounts for an overall shift of all $g_i(M_{\text{GUT}})$, while the c_2 term results in a change of M_{GUT} .

Therefore T_{SUSY} has a very simple interpretation: a SUSY spectrum with an effective SUSY threshold scale T_{SUSY} has the same effect on gauge coupling unification as a degenerate spectrum with all superpartners having this common mass, up to a change of M_{GUT} and/or the unified gauge coupling.

The effective SUSY threshold scale takes the form³

$$T_{\text{SUSY}} = \frac{m_{\widetilde{W}}^{32/19} m_{\widetilde{h}}^{12/19} m_H^{3/19}}{m_{\widetilde{g}}^{28/19}} X_{\text{sfermion}} , \quad (6)$$

where $m_{\widetilde{W}}$, $m_{\widetilde{h}}$, m_H and $m_{\widetilde{g}}$ denote the mass of the wino, the light and heavy MSSM Higgs and gluino, respectively. The sfermion contribution reads

$$X_{\text{sfermion}} = \prod_{i=1\dots 3} \left(\frac{m_{\widetilde{L}^{(i)}}^{3/19}}{m_{\widetilde{D}^{(i)}}^{3/19}} \right) \left(\frac{m_{\widetilde{Q}_L^{(i)}}^{7/19}}{m_{\widetilde{E}^{(i)}}^{2/19} m_{\widetilde{U}^{(i)}}^{5/19}} \right) , \quad (7)$$

where the masses of the sfermions appear in a self-explanatory notation. As we will show explicitly in the next section, $X_{\text{sfermion}} \simeq 1$ whenever the sfermion masses are universal within a $\text{SU}(5)$ multiplet at the high scale which is expected to hold in a GUT model. The mass splitting of sleptons and squarks through renormalization group running does not affect this conclusion. Hence, the sfermions leave no direct effect on PGU.

Note that the insensitivity of gauge coupling unification to the scale of sfermion masses is due to the completeness of these GUT multiplets. Only split GUT multiplets give non-trivial contributions to T_{SUSY} (cf. equation (6)). This is also utilized in models of split supersymmetry [18]. It is also interesting that the decoupling of complete generations of soft scalar masses, e.g. the first and second generation, does not affect PGU, allowing for a realization of “natural supersymmetry” with a light third generation of soft scalar masses.

3 Gauge coupling unification in MSSM models

As can be seen from equation (6), T_{SUSY} mainly depends on the higgsino mass and on the mass ratio $m_{\widetilde{W}} : m_{\widetilde{g}}$. In the following, after discussing the case of uncompressed gaugino masses, we shall show how a change in this mass ratio can lead to PGU.

³A simplified version of this formula was discussed in [13].

3.1 Uncompressed gaugino mass spectra

In models with gravity mediated SUSY breaking, the gaugino masses are typically expected to unify at the GUT scale M_{GUT} . Taking $m_{1/2}$ to be the universal gaugino mass at the GUT scale, the physical mass of wino and gluino can be approximated as

$$m_{\tilde{W}} \simeq 0.9 m_{1/2} \quad \text{and} \quad m_{\tilde{g}} \simeq 2.5 m_{1/2} . \quad (8)$$

In turn the effective SUSY threshold scale reads

$$T_{\text{SUSY}} \simeq 0.3 \left(m_{\tilde{h}}^{12} m_{1/2}^4 m_H^3 \right)^{1/19} X_{\text{sfermion}} . \quad (9)$$

In the limit where the sfermions are substantially heavier than the gauginos, sfermion masses are only weakly affected by RGE running. As long as the sfermion masses are universal within SU(5) multiplets at the high scale, we find $X_{\text{sfermion}} \simeq 1$. Even if gaugino masses are non-negligible and the masses of squarks and sleptons are split by RGE running, their effect on gauge coupling unification remains small. This is illustrated in figure 2 where we depict X_{sfermion} in the CMSSM for a fixed gaugino mass $m_{1/2}$ and a varying scalar mass m_0 . It can be seen that X_{sfermion} never deviates by more than a few per cent from one.




Figure 2: Effect of sfermions on gauge coupling unification in the CMSSM for varying m_0 .

Taking $X_{\text{sfermion}} = 1$, we can determine the higgsino mass which is required to obtain PGU, i.e. $T_{\text{SUSY}} \simeq 2 \text{ TeV}$. We find

$$m_{\tilde{h}} \simeq 20 \text{ TeV} \times \left(\frac{\text{TeV}}{m_{1/2}} \right)^{1/3} \left(\frac{\text{TeV}}{m_H} \right)^{1/4} . \quad (10)$$

Due to the small exponents of the last two terms, higgsino masses as large as $m_{\tilde{h}} = \mathcal{O}(10 \text{ TeV})$ are required for gauge coupling unification, even if we take $m_{1/2}$ and m_H in the multi-TeV range. As is well known, such heavy higgsinos lead to ‘unnaturally’ large fine-tuning as can be seen from the mass of the Z boson which — at tree level — can

be written as $M_Z \simeq -2m_{h_u}^2 - 2|\mu|^2$ for $\tan\beta \gg 1$. In the case of a large higgsino mass, severe cancellations between m_{h_u} and μ are required.

We will therefore now turn to a class of models in which PGU can be accommodated more naturally.

3.2 Compressed gaugino mass spectra

The effective T_{SUSY} can be enhanced if one increases the wino mass relative to the gluino mass. Specifically, we now discuss the possibility of realizing PGU by non-universal gaugino masses which can be accommodated in soft mass spectra arising from UV completions of the MSSM. To obtain non-universal gaugino masses, an obvious possibility is to use anomaly mediated contributions. While in pure anomaly mediation the hierarchy between the gluino and wino is even larger than in gravity mediation, more compressed gaugino spectra can be obtained in schemes with mixed gravity/anomaly mediation, also referred to as mirage scheme.

At first sight, having gravity and anomaly mediation at similar strength appears as an ad hoc assumption. But note that this naturally occurs in a wide class of string models due to the interplay between moduli stabilization and SUSY breaking. Examples of such mixed mediation schemes have been discussed in the context of type IIB string theory with moduli stabilization following KKLT [8, 9, 11, 19–21], in the context of heterotic string theory [12, 22–24], and in the context of G_2 compactifications [25].

In mirage mediation, the gaugino masses at the high scale can be written as

$$M_i = \frac{m_{3/2}}{16\pi^2} (\varrho + b_i^{\text{MSSM}} g^2) , \quad (11)$$

where $m_{3/2}$ denotes the gravitino mass and ρ parametrizes the gravity mediated contribution to the gaugino masses, in the limit of vanishing ϱ pure anomaly mediation is recovered. Notice that $|M_3| < |M_2|$ at the GUT scale unless for $\varrho \lesssim 0.5$. The gaugino masses at the low scale can be approximated as

$$m_{\tilde{B}} \simeq 0.45 (\varrho + 3.3) \frac{m_{3/2}}{16\pi^2} , \quad (12a)$$

$$m_{\tilde{W}} \simeq 0.9 (\varrho + 0.5) \frac{m_{3/2}}{16\pi^2} , \quad (12b)$$

$$m_{\tilde{g}} \simeq 2.4 (\varrho - 1.5) \frac{m_{3/2}}{16\pi^2} . \quad (12c)$$

The gaugino spectrum is thus typically more compressed than in models with gaugino mass unification. In the regime $0.8 \lesssim \varrho \lesssim 2.5$ the gluino is even the lightest gaugino.

In the context of mirage mediation, the scalar masses and trilinear terms are model dependent. The former are of the order of the gravitino mass unless in case of sequestering. The trilinear soft terms are typically suppressed compared to the gravitino mass and can thus be neglected. In this study we assume sfermion masses in the multi-TeV regime as suggested by the KKLT and heterotic models with F -term uplifting [11, 12, 26, 27].

In order to avoid a naturalness problem, we should require that $m_{h_{u,d}} = m_{\tilde{Q}_L^{(3)}} = m_{\tilde{U}^{(3)}}$ approximately holds at the GUT scale.⁴ In this case, the RGE trajectory of m_{h_u} exhibits the well known focus point [28, 29] such that multi-TeV scalars are not unnatural. In the focus point scenario m_{h_u} is driven to a very small value at the low scale, while m_{h_d} is not considerably affected by RGE running. This implies $m_H \simeq m_{h_d}$ for the physical mass of the heavy MSSM Higgs as well as a suppressed μ term [28]. Therefore, the above assumption is equivalent to $m_H = m_{\tilde{Q}_L^{(3)}} = m_{\tilde{U}^{(3)}} \gg \mu$ up to sub-leading corrections. As the running of the gauge couplings depends on μ rather than the Higgs soft masses this form of the input is more suitable for our purposes. To be specific we set

$$m_H = m_i \equiv m_0, \quad (13)$$

where m_i denotes the mass of the sfermion i at the high scale. We are thus left with the five free parameters $m_{3/2}$, ϱ , m_0 , μ and $\tan\beta$. To avoid excessive fine-tuning we restrict our attention to $\mu \leq 2$ TeV. Further, for any point in the parameter space, we eliminate m_0 by requiring that the mass of the light MSSM Higgs is $m_h = 126$ GeV, which typically leads to $m_0 \sim 15$ TeV. We should point out that our conclusions are insensitive to the assumptions in the scalar sector as the latter hardly affect gauge coupling unification. Thus our discussions are also valid in schemes where some of the complete sfermion families receive a mass $\sim m_{3/2}$ which typically is larger than m_0 .

We now can ask what constraints on the parameter space arise from imposing PGU. In figure 3, we present the parameter region consistent with PGU for fixed $m_{3/2}$ and $\tan\beta$. Precision unification clearly favors a compressed spectrum, i.e. a spectrum where the mass difference between gluino and LSP is small. For $1.1 < \varrho < 2.4$, the gluino is the LSP, unless the higgsino becomes even lighter (shrinking of the red band at the left side in figure 3). The two regions in parameter space consistent with PGU correspond to the two regions where the mass ratio of wino and gluino gives rise to the required effective SUSY threshold scale.

There is only a weak dependence on the overall scale of gaugino masses which is set by $m_{3/2}$. Therefore, our results remain qualitatively unchanged for different choices of $m_{3/2}$. Similarly, PGU only shows a weak dependence on $\tan\beta$. For illustration, we have included scans with varying $m_{3/2}$ and $\tan\beta$ in appendix A.

4 Implications for the LHC and dark matter

4.1 LHC discovery potential

We can now turn to the implications of PGU for SUSY searches at the LHC. As described in the previous section, this naturally leads us to models with mirage mediation where PGU can naturally be accommodated with a small μ .

Within the mirage scheme, solid predictions on the superpartner spectrum can be made. While the scalar superpartners are typically outside the reach of the LHC, the

⁴This is a very reasonable assumption as e.g. in the heterotic models, these fields have the same localization properties (see discussion in [22, 23]).




Figure 3: Parameter scan in the (μ, ρ) -plane for fixed $m_{3/2} = 50$ TeV. The region where the gauge couplings unify precisely within the experimental error on the strong coupling are shown in green. The red region exhibits a gluino LSP. The dotted contour indicates where the mass ratio between gluino and LSP becomes two.

gauginos and higgsinos remain relatively light. We cannot predict the hierarchy among gauginos and higgsinos, but there is a general trend that PGU prefers a spectrum in which the mass difference between the gluino and the lightest neutralino is small.

Such compressed SUSY models are much more difficult to detect at the LHC than ordinary SUSY models like the CMSSM (see e.g. [30–32]). While there is still a strong gluino pair production, the subsequent decay of gluinos yields only a small amount of visible energy. In the experimental searches, a sizable fraction of the potential signal events is rejected due to too soft jets. In the case of extreme compression, only events with initial state radiation can pass the event cuts, and the signal acceptance is typically well below 1% [31].

If the gluino is the next-to-lightest supersymmetric particle, the relevant gluino decay modes read

$$\tilde{g} \rightarrow q\bar{q} + \tilde{\chi}_1, \quad (14a)$$

$$\tilde{g} \rightarrow g + \tilde{\chi}_1. \quad (14b)$$

where $\tilde{\chi}_1$ denotes the lightest neutralino. The first process proceeds via an off-shell squark, the second via a quark / squark loop. Additional decay modes open up if further neutralinos or charginos are lighter than the gluino.

It is instructive to determine the regime of gluino and (lightest) neutralino masses preferred by PGU. For this purpose, we have generated a large data sample with random

choice of input parameters in the intervals

$$\varrho = 0.5 - 30, \quad m_{3/2} = \frac{40 - 200 \text{ TeV}}{\varrho}, \quad \mu = 0.1 - 2 \text{ TeV}, \quad \tan \beta = 10 - 50. \quad (15)$$

Note that the considered range of $m_{3/2}$ corresponds to a gravity mediated contribution to the gaugino masses of $0.25 - 1.25 \text{ TeV}$.

In figure 4, we provide a scatter plot of those parameter sets which lead to successful PGU in the $(m_{\tilde{g}}, m_{\tilde{\chi}_1})$ -plane. To guide the eye, we have also included the present limits on the gluino mass from the ATLAS search for jets plus missing energy at $\int dt L = 5.8 \text{ fb}^{-1}$ [33] and from the CMS search for b -jets plus missing energy at $\int dt L = 19.4 \text{ fb}^{-1}$ [34].⁵ Both analyses are based on a simplified model with just the gluino and the LSP assuming 100% branching of the process (14a). CMS additionally requires the final state quarks to be bottoms. Both limits are not strictly applicable as the considered decay modes do not necessarily dominate in the set-up we consider. In particular, we may encounter longer decay chains if the gluino decay to heavier neutralinos / charginos is kinematically accessible, leading to events with more and softer jets than in the simplified model. As too soft jets typically fail the cuts performed in the ATLAS and CMS analyses, this can clearly affect the constraints. However, very soft jets would mainly arise in parameter regions with a strongly compressed spectrum where initial state radiation is anyway required to pass the event trigger, the latter being insensitive to the gluino decay. Therefore, we believe that at least the ATLAS limit can still be used as a reasonable estimate. The stronger CMS limit should not be applied, as we do not find a preference for b -jets in the final state compared to light quark or gluon jets. We have, nevertheless, included it in figure 4 as the LHC gluon searches without b -tagging should reach a similar sensitivity once the full data sample collected at $\sqrt{s} = 8 \text{ TeV}$ is analyzed. Therefore, the CMS constraint can be seen as a projection of the current sensitivity of the LHC to gluinos.

A small fraction of the benchmark points features a gluino LSP. Apart from the fact that a gluino LSP is very unfavorable from a cosmological perspective, stable gluinos with $m_{\tilde{g}} < 985 \text{ GeV}$ are excluded by the ATLAS search for R-hadrons [36]. This constraint is also depicted in figure 4.

We observe that more than 90% of the benchmark points with PGU fulfill $m_{\tilde{\chi}_1} > 0.5 m_{\tilde{g}}$. The strong compression of the spectrum implies that the limits on the gluino mass are considerably weaker than in ordinary SUSY schemes like the CMSSM. In particular we find that a large fraction of the benchmark points with $m_{\tilde{g}} = 500 - 1000 \text{ GeV}$ is not excluded by the LHC.

We should also point out that the gluino can be rather long-lived in the scenario we consider. This is partly because of the phase space suppression due to the small mass difference between gluino and neutralino, and partly because of the large mass of the squarks which mediate the gluino decay. The gluino decay rates in case of $m_{\tilde{\chi}_1} \sim m_{\tilde{g}}$

⁵See [35] for a similar analysis by ATLAS.




Figure 4: Parameter points with successful gauge coupling unification (gray). Also shown are the constraints on the gluino mass for the decay modes $\tilde{g} \rightarrow q\bar{q} + \tilde{\chi}_1$ and $\tilde{g} \rightarrow b\bar{b} + \tilde{\chi}_1$ by ATLAS and CMS (see text). The yellow region features a gluino LSP which is constrained by the ATLAS search for stable R-hadrons.

scale as (see e.g. [37])

$$\Gamma(\tilde{g} \rightarrow q\bar{q} + \tilde{\chi}_1) \propto \text{gaugino fraction} \times \frac{(m_{\tilde{g}} - m_{\tilde{\chi}_1})^5}{m_{\tilde{q}}^4}, \quad (16a)$$

$$\Gamma(\tilde{g} \rightarrow g + \tilde{\chi}_1) \propto \text{higgsino fraction} \times \frac{(m_{\tilde{g}} - m_{\tilde{\chi}_1})^3 m_t^2}{m_{\tilde{q}}^4}, \quad (16b)$$

where m_t denotes the top mass and $m_{\tilde{q}}$ the mass of the squark in the intermediate state. In the set-up we consider, the squarks are in the range $m_{\tilde{q}} \sim 15$ TeV as implied by the requirement $m_h = 126$ GeV (see section 3.2). Note that the decay of the gluino into two quarks scales with the gaugino fraction of $\tilde{\chi}_1$, as the higgsino coupling to light quarks is suppressed, while top quarks in the final state are kinematically inaccessible for $m_{\tilde{\chi}_1} \sim m_{\tilde{g}}$. The decay into gluon and neutralino, on the other hand, scales with the higgsino component of $\tilde{\chi}_1$. In our set-up the mass splitting between left- and right-handed squarks is typically small, i.e. parity is approximately preserved in the squark sector. Therefore, the decay of the gluino into gaugino and gluon — which requires parity violation [38] — can be neglected.

We have systematically determined the total gluino decay rate $\Gamma_{\tilde{g}}$ for the benchmark sample using the tool SDECAY 1.3 [39], the corresponding distribution is shown in figure 5.⁶ We find that for slightly more than 10% of the benchmark points the decay length $c/\Gamma_{\tilde{g}}$ exceeds $10 \mu\text{m}$ which roughly corresponds to the LHC resolution. This

⁶The distribution was slightly smoothed.

suggests that, in a non-negligible fraction of the parameter space, one might be able to observe displaced vertex signatures. Note that, in our approach, we assumed universal squark masses at the high scale and fixed $m_{\tilde{q}}$ such that the correct Higgs mass is obtained. There exist, however, theoretical uncertainties in the calculation of the Higgs mass at the level of a few GeV [40]. As the squark mass enters the loop corrections to m_h only logarithmically this translates into an $\mathcal{O}(1)$ uncertainty on the squark mass.⁷ Given that $\Gamma_{\tilde{g}} \propto m_{\tilde{q}}^{-4}$, our calculation of the gluino decay rate can, thus, only be seen as an order of magnitude estimate. For many benchmark points $c/\Gamma_{\tilde{g}}$ is at least close to the LHC position resolution. Therefore, we believe that there is indeed a realistic chance to observe displaced vertices. Particularly long gluino life-times are obtained if the LSP is gaugino-like. In this case $\Gamma_{\tilde{g}}$ is suppressed by the fifth power of the mass splitting (or the decay must proceed through the subleading higgsino admixture of $\tilde{\chi}_1$).




Figure 5: Distribution of the gluino decay rate within the benchmark sample. The vertical dashed lines indicate where the gluino decay length $c/\Gamma_{\tilde{g}}$ reaches $10\mu\text{m}$ (roughly the LHC resolution) and 1mm respectively.

The gluino will, however, almost certainly decay within the inner pixel detectors of ATLAS / CMS. It is unlikely that the gluino decay vertex is more than a few mm from the initial collision vertex (see figure 5). While dedicated searches for gluino decays with such tiny displacements do currently not exist, it seems feasible to employ similar methods as for the identification of b -jets. In particular one could study the distribution of the transverse impact parameter in events with jets and missing energy. One might worry that the jets arising from the gluino decays are too soft to be detected. However, we find that a decay length above the LHC resolution can be obtained for mass splittings $m_{\tilde{g}} - m_{\tilde{\chi}_1}$ as large as about 100GeV (depending on the composition of $\tilde{\chi}_1$). We thus want to emphasize the importance to develop search strategies for gluinos with decays slightly displaced from the collision vertex. It is also important to determine how b -tagging is

⁷Note also that in the benchmark sample we have set $\tan\beta = 10 - 50$. For $\tan\beta < 10$, larger squark masses are required in order to obtain $m_h = 126\text{GeV}$, i.e. the life-time of the gluino would be further enhanced.

affected if the superpartner spectrum contains gluinos with a decay length similar as that of b -mesons.

Even if gluinos are too short-lived to give rise to displaced vertices, the study of gluino decays is of high interest. This is because details of the superpartner sector are encrypted in the gluino decay pattern. It will be difficult to distinguish the multi-jet decay modes of the gluino. However, as was pointed out in [41], it is feasible to identify the radiative mode $\tilde{g} \rightarrow g + \tilde{\chi}_1$ which typically gives rise to events with a low jet multiplicity. This particular decay mode is very sensitive to the higgsino component of the lightest neutralino (cf. (16)). A large branching fraction in this mode would thus hint at a relatively small μ term. Of course, it will be challenging to determine the full set of high scale parameters just from the gluino decay. But in any case it is remarkable that gluino decays can in principle be used to learn about the spectrum of superpartners which are, in principle, outside the reach of the LHC.

4.2 Dark Matter

As it is well known, the lightest neutralino is a very good candidate for the dark matter in the universe. However, in the simplest realizations of gravity mediation, the lightest neutralino is either bino-like or higgsino-like. Considerable neutralino mixing does only occur in some narrow regions of parameter space. Due to its small annihilation cross section, the bino density from thermal production typically exceeds the observed dark matter density Ω_{DM} by far. The latter has recently been determined at high accuracy by the Planck collaboration [42],

$$\Omega_{\text{DM}} h^2 = 0.1196 \pm 0.0031 . \quad (17)$$

Higgsinos, on the other hand, undergo efficient annihilations into third generation quarks or gauge bosons, coannihilations with the charged higgsinos further enhance their cross section. Hence, the relic density of a higgsino LSP is typically well below the dark matter density. In both cases one has to invoke a non-standard cosmological history if the LSP is to account for the observed dark matter. In case of the higgsino, non-thermal production e.g. by the decay of a heavy gravitino or modulus field is a viable option (see e.g. [43–45]). For a bino LSP the situation is more unfavorable as, if its abundance is diluted by the decay of a heavy field, binos are typically regenerated by the same decay, reintroducing precisely the same problem [46, 47].

In mirage mediation, the gaugino masses are non-universal at the high scale due to the anomaly-mediated contributions. We have seen that if we require PGU, we are drawn into a region of parameter space where the gaugino spectrum is highly compressed. This turns out to be very favorable for the dark matter density. We have determined the neutralino relic density $\Omega_\chi h^2$ for a large benchmark sample using the boundaries (15) with the tool MicrOMEGAs 2.4.5 [48]. In figure 6 we compare the distribution of $\Omega_\chi h^2$ among the sample points with and without imposing PGU.⁸

⁸For this figure we have disregarded benchmark points with a gluino LSP. The distribution was again slightly smoothed.




Figure 6: Distribution of the thermal neutralino relic density for the benchmark sample with (solid) and without (dashed) requiring PGU. The observed dark matter density is indicated by the gray contour.

We find that without the requirement of PGU, the lightest neutralino is typically bino- or higgsino-like, similar as in the models with gaugino mass unification. The distribution of the relic density among the benchmark points has two peaks corresponding to the bino and the higgsino case respectively.

If we include the requirement of PGU, the picture changes dramatically. Remarkably, as most benchmark points with PGU feature a compressed spectrum, the distribution is now peaked close to observed dark matter density. The reason is that in the parameter regions with precision unification, the mass difference between bino and wino is typically $\sim 10\%$, suggesting the occurrence of coannihilations. In particular in case of a bino LSP, wino pair processes would still dominate the annihilation cross section. These would be suppressed by a Boltzmann factor

$$B = \exp\left(-2 \frac{m_{\tilde{W}} - m_{\tilde{B}}}{T_F}\right) \quad (18)$$

with the freeze-out temperature $T_F \sim m_{\tilde{B}}/20$. Given a splitting of $\sim 10\%$ one finds $B \sim 0.01$. This factor compensates for the large wino pair-annihilation cross section, yielding a relic dark matter density consistent with observation. Another possibility to obtain the correct relic density is by mixing effects in the neutralino sector.

Turning to direct dark matter detection, we find that the cross section of the lightest neutralino with nucleons is dominated by exchange of the light Higgs (as the other scalars are in the multi-TeV regime). The coupling of the lightest neutralino to the Higgs scales with the gaugino / higgsino mixing angle, it vanishes in the limit of a pure state. The neutralino proton cross section σ_p for the benchmark points with PGU is shown in figure 7. For the determination of σ_p we used MicrOMEGAs 2.4.5, but took the updated nuclear quark form factors from [49] suggested by recent lattice QCD calculations (see also [50]). Note that these form factors suggest a smaller strange quark contribution to the nucleon mass compared to earlier computations [51]. They thus tend to give a smaller σ_p .




Figure 7: Neutralino proton cross section for the benchmark points with successful gauge coupling unification (gray). The current limit from the XENON100 direct dark matter search is also shown. The latter is only applicable if the lightest neutralino accounts for all dark matter in the universe.

We find that σ_p scatters between 10^{-48} cm² and 10^{-43} cm². While the XENON100 experiment [52] has just begun to probe this regime of cross sections, the next generation of direct detection experiments will be able to test a significant fraction of the depicted benchmark points. Still, there exist benchmark points with a bino-like LSP which have a strongly suppressed σ_p . These may even be consistent with thermal dark matter production due to coannihilations. Consequently, there is no guarantee for a signal in direct dark matter detection. One should also have in mind that the lightest neutralino must not necessarily account for the dark matter in the universe such that direct detection experiments would be doomed to fail.

5 Conclusions

This analysis is based on the hypothesis that precision gauge unification (PGU) is not an accident. In general, imposing PGU reduces the dimension of the soft supersymmetry breaking parameters by one. Within the more ‘traditional’ patterns of soft masses, imposing PGU implies typically a large μ parameter, which is in conflict with usual ‘naturalness’ arguments. On the other hand, a particularly attractive pattern of soft masses, consistent with PGU, is the one of mirage mediation where the gaugino spectrum is compressed and the μ parameter can be small. Such a spectrum can be realized in various string constructions which exhibit a ‘competition’ between gravity- and anomaly-mediated contributions to soft masses. The similarity of gaugino masses at low energies has profound implications for collider and dark matter searches.

By requiring PGU we are driven in a corner of the MSSM parameter space with a non-standard collider phenomenology. The small mass difference between gluino and neutralino typically leads to events with a reduced amount of visible energy. The detection of gluinos becomes far more challenging than in most of the standard SUSY models. In the case of extreme compression, initial state radiation is required to trigger on the SUSY events and the detection efficiency is strongly reduced. The gluino is typically rather long-lived, its decay length may exceed the LHC resolution and lead to displaced vertices at the level of μm to mm . Even if displaced vertices are absent, interesting information on the superpartner spectrum can be obtained by studying the gluino decay modes.

Another virtue of compressed gaugino masses is that they generically lead to coannihilation effects in the early universe. Cosmological problems related to the overproduction of dark matter which are present e.g. in the CMSSM can easily be solved. Indeed, we find that in the parameter space with PGU, the relic neutralino density is expected to be close to the observed dark matter density.

Finally, it is needless to say that measuring superpartner spectra that support PGU will provide us with invaluable information on how the standard model is completed in the ultraviolet.

Acknowledgments

We would like to thank Jamie Tattersall and Jan Hajer for useful discussions. This work was supported by the SFB-Transregio TR33 “The Dark Universe” (Deutsche Forschungsgemeinschaft), the European Union 7th network program “Unification in the LHC era” (PITN-GA-2009-237920), the DFG cluster of excellence Origin and Structure of the Universe by Deutsche Forschungsgemeinschaft (DFG) and the ERC Advanced Grant project “FLAVOUR” (267104).

A Dependence of PGU on $\tan \beta$ and $m_{3/2}$

Figures 8 and 9 illustrate the dependence of PGU on $\tan \beta$ and $m_{3/2}$.




Figure 8: Same as figure 3 for different choices of $\tan \beta$.




Figure 9: Same as figure 3 for different choices of $m_{3/2}$.

References

- [1] E. Witten, “Dynamical Breaking of Supersymmetry,” *Nucl.Phys.* **B188** (1981) 513.
- [2] S. Dimopoulos, S. Raby, and F. Wilczek, “Supersymmetry and the Scale of Unification,” *Phys.Rev.* **D24** (1981) 1681–1683.
- [3] M. L. Alciati, F. Feruglio, Y. Lin, and A. Varagnolo, “Proton lifetime from SU(5) unification in extra dimensions,” *JHEP* **0503** (2005) 054, [arXiv:hep-ph/0501086 \[hep-ph\]](#).
- [4] Particle Data Group, C. Amsler *et al.*, “Review of Particle Physics,” *Phys.Lett.* **B667** (2008) 1–1340.
- [5] M. S. Carena, L. Clavelli, D. Matalliotakis, H. P. Nilles, and C. Wagner, “Light gluinos and unification of couplings,” *Phys.Lett.* **B317** (1993) 346–353, [arXiv:hep-ph/9307320 \[hep-ph\]](#).
- [6] L. Roszkowski and M. A. Shifman, “Reconciling supersymmetric grand unification with $\alpha^{-1}_s(m(Z))$ approximates 0.11,” *Phys.Rev.* **D53** (1996) 404–412, [arXiv:hep-ph/9503358 \[hep-ph\]](#).
- [7] S. Raby, M. Ratz, and K. Schmidt-Hoberg, “Precision gauge unification in the MSSM,” *Phys.Lett.* **B687** (2010) 342–348, [arXiv:0911.4249 \[hep-ph\]](#).
- [8] K. Choi, A. Falkowski, H. P. Nilles, and M. Olechowski, “Soft supersymmetry breaking in KKLT flux compactification,” *Nucl.Phys.* **B718** (2005) 113–133, [arXiv:hep-th/0503216 \[hep-th\]](#).
- [9] K. Choi, K. S. Jeong, and K.-i. Okumura, “Phenomenology of mixed modulus-anomaly mediation in fluxed string compactifications and brane models,” *JHEP* **0509** (2005) 039, [arXiv:hep-ph/0504037 \[hep-ph\]](#).
- [10] A. Falkowski, O. Lebedev, and Y. Mambrini, “SUSY phenomenology of KKLT flux compactifications,” *JHEP* **0511** (2005) 034, [arXiv:hep-ph/0507110 \[hep-ph\]](#).
- [11] O. Lebedev, H. P. Nilles, and M. Ratz, “De Sitter vacua from matter superpotentials,” *Phys.Lett.* **B636** (2006) 126–131, [arXiv:hep-th/0603047 \[hep-th\]](#).
- [12] V. Löwen and H. P. Nilles, “Mirage Pattern from the Heterotic String,” *Phys.Rev.* **D77** (2008) 106007, [arXiv:0802.1137 \[hep-ph\]](#).
- [13] M. S. Carena, S. Pokorski, and C. Wagner, “On the unification of couplings in the minimal supersymmetric Standard Model,” *Nucl.Phys.* **B406** (1993) 59–89, [arXiv:hep-ph/9303202 \[hep-ph\]](#).

- [14] A. Lahanas and K. Tamvakis, “Low-energy thresholds and the renormalization group in the MSSM,” *Phys.Lett.* **B348** (1995) 451–456, [arXiv:hep-ph/9412281](#) [[hep-ph](#)].
- [15] A. Anandakrishnan and S. Raby, “Yukawa Unification Predictions with effective “Mirage” Mediation,” [arXiv:1303.5125](#) [[hep-ph](#)].
- [16] B. Allanach, “SOFTSUSY: a program for calculating supersymmetric spectra,” *Comput.Phys.Commun.* **143** (2002) 305–331, [arXiv:hep-ph/0104145](#) [[hep-ph](#)].
- [17] S. Bethke, “World Summary of α_s (2012),” *Nucl.Phys.Proc.Suppl.* **234** (2013) 229–234, [arXiv:1210.0325](#) [[hep-ex](#)].
- [18] N. Arkani-Hamed and S. Dimopoulos, “Supersymmetric unification without low energy supersymmetry and signatures for fine-tuning at the LHC,” *JHEP* **0506** (2005) 073, [arXiv:hep-th/0405159](#) [[hep-th](#)].
- [19] S. Kachru, R. Kallosh, A. D. Linde, and S. P. Trivedi, “De Sitter vacua in string theory,” *Phys.Rev.* **D68** (2003) 046005, [arXiv:hep-th/0301240](#) [[hep-th](#)].
- [20] K. Choi, A. Falkowski, H. P. Nilles, M. Olechowski, and S. Pokorski, “Stability of flux compactifications and the pattern of supersymmetry breaking,” *JHEP* **0411** (2004) 076, [arXiv:hep-th/0411066](#) [[hep-th](#)].
- [21] E. Dudas, A. Linde, Y. Mambrini, A. Mustafayev, and K. A. Olive, “Strong moduli stabilization and phenomenology,” *Eur.Phys.J.* **C73** (2013) 2268, [arXiv:1209.0499](#) [[hep-ph](#)].
- [22] S. Krippendorff, H. P. Nilles, M. Ratz, and M. W. Winkler, “The heterotic string yields natural supersymmetry,” *Phys.Lett.* **B712** (2012) 87–92, [arXiv:1201.4857](#) [[hep-ph](#)].
- [23] M. Badziak, S. Krippendorff, H. P. Nilles, and M. W. Winkler, “The heterotic MiniLandscape and the 126 GeV Higgs boson,” *JHEP* **1303** (2013) 094, [arXiv:1212.0854](#) [[hep-ph](#)].
- [24] B. L. Kaufman, B. D. Nelson, and M. K. Gaillard, “Mirage Models Confront the LHC: I. Kahler-Stabilized Heterotic String Theory,” [arXiv:1303.6575](#) [[hep-ph](#)].
- [25] B. S. Acharya, G. Kane, and P. Kumar, “Compactified String Theories – Generic Predictions for Particle Physics,” *Int.J.Mod.Phys.* **A27** (2012) 1230012, [arXiv:1204.2795](#) [[hep-ph](#)].
- [26] H. Abe, T. Higaki, T. Kobayashi, and Y. Omura, “Moduli stabilization, F-term uplifting and soft supersymmetry breaking terms,” *Phys.Rev.* **D75** (2007) 025019, [arXiv:hep-th/0611024](#) [[hep-th](#)].

- [27] R. Kallosh and A. D. Linde, “O’KKLT,” *JHEP* **0702** (2007) 002, [arXiv:hep-th/0611183](#) [hep-th].
- [28] J. L. Feng, K. T. Matchev, and T. Moroi, “Multi - TeV scalars are natural in minimal supergravity,” *Phys.Rev.Lett.* **84** (2000) 2322–2325, [arXiv:hep-ph/9908309](#) [hep-ph].
- [29] J. L. Feng, K. T. Matchev, and T. Moroi, “Focus points and naturalness in supersymmetry,” *Phys.Rev.* **D61** (2000) 075005, [arXiv:hep-ph/9909334](#) [hep-ph].
- [30] T. J. LeCompte and S. P. Martin, “Large Hadron Collider reach for supersymmetric models with compressed mass spectra,” *Phys.Rev.* **D84** (2011) 015004, [arXiv:1105.4304](#) [hep-ph].
- [31] T. J. LeCompte and S. P. Martin, “Compressed supersymmetry after 1/fb at the Large Hadron Collider,” *Phys.Rev.* **D85** (2012) 035023, [arXiv:1111.6897](#) [hep-ph].
- [32] H. K. Dreiner, M. Kramer, and J. Tattersall, “How low can SUSY go? Matching, monojets and compressed spectra,” *Europhys.Lett.* **99** (2012) 61001, [arXiv:1207.1613](#) [hep-ph].
- [33] ATLAS Collaboration, “Search for squarks and gluinos with the ATLAS detector using final states with jets and missing transverse momentum and 5.8 fb⁻¹ of \sqrt{s} =8 TeV proton-proton collision data,” *ATLAS-CONF-2012-109* (2012) .
- [34] CMS Collaboration, “Search for Supersymmetry in pp collisions at 8 TeV in events with a single lepton, multiple jets and b-tags,”.
- [35] ATLAS Collaboration, “Search for gluino pair production in final states with missing transverse momentum and at least three b-jets using 12.8 fb-1 of pp collisions at \sqrt{s} = 8 TeV with the ATLAS Detector.,”.
- [36] ATLAS Collaboration, G. Aad *et al.*, “Searches for heavy long-lived sleptons and R-Hadrons with the ATLAS detector in *pp* collisions at \sqrt{s} = 7 TeV,” *Phys.Lett.* **B720** (2013) 277–308, [arXiv:1211.1597](#) [hep-ex].
- [37] R. Barbieri, G. Gamberini, G. F. Giudice, and G. Ridolfi, “Constraining Supergravity Models from Gluino Production,” *Nucl.Phys.* **B301** (1988) 15.
- [38] H. E. Haber and G. L. Kane, “Gluino Decays and Experimental Signatures,” *Nucl.Phys.* **B232** (1984) 333.
- [39] M. Mühlleitner, A. Djouadi, and Y. Mambrini, “SDECAY: A Fortran code for the decays of the supersymmetric particles in the MSSM,” *Comput.Phys.Commun.* **168** (2005) 46–70, [arXiv:hep-ph/0311167](#) [hep-ph].

- [40] P. Draper, J. Feng, P. Kant, S. Profumo, and D. Sanford, “Dark Matter Detection in Focus Point Supersymmetry,” [arXiv:1304.1159](#) [hep-ph].
- [41] R. Sato, S. Shirai, and K. Tobioka, “Gluino Decay as a Probe of High Scale Supersymmetry Breaking,” *JHEP* **1211** (2012) 041, [arXiv:1207.3608](#) [hep-ph].
- [42] Planck Collaboration, P. Ade *et al.*, “Planck 2013 results. XVI. Cosmological parameters,” [arXiv:1303.5076](#) [astro-ph.CO].
- [43] T. Moroi and L. Randall, “Wino cold dark matter from anomaly mediated SUSY breaking,” *Nucl.Phys.* **B570** (2000) 455–472, [arXiv:hep-ph/9906527](#) [hep-ph].
- [44] M. Fujii and K. Hamaguchi, “Higgsino and wino dark matter from Q ball decay,” *Phys.Lett.* **B525** (2002) 143–149, [arXiv:hep-ph/0110072](#) [hep-ph].
- [45] R. Kitano, H. Murayama, and M. Ratz, “Unified origin of baryons and dark matter,” *Phys.Lett.* **B669** (2008) 145–149, [arXiv:0807.4313](#) [hep-ph].
- [46] T. Moroi, M. Yamaguchi, and T. Yanagida, “On the solution to the Polonyi problem with 0 (10-TeV) gravitino mass in supergravity,” *Phys.Lett.* **B342** (1995) 105–110, [arXiv:hep-ph/9409367](#) [hep-ph].
- [47] S. Nakamura and M. Yamaguchi, “Gravitino production from heavy moduli decay and cosmological moduli problem revived,” *Phys.Lett.* **B638** (2006) 389–395, [arXiv:hep-ph/0602081](#) [hep-ph].
- [48] G. Belanger, F. Boudjema, A. Pukhov, and A. Semenov, “MicrOMEGAs 2.0: A Program to calculate the relic density of dark matter in a generic model,” *Comput.Phys.Comm.* **176** (2007) 367–382, [arXiv:hep-ph/0607059](#) [hep-ph].
- [49] G. Belanger, F. Boudjema, A. Pukhov, and A. Semenov, “micrOMEGAs3.1 : a program for calculating dark matter observables,” [arXiv:1305.0237](#) [hep-ph].
- [50] A. Thomas, P. Shanahan, and R. Young, “Strangeness in the nucleon: what have we learned?,” *Nuovo Cim.* **C035N04** (2012) 3–10, [arXiv:1202.6407](#) [nucl-th].
- [51] M. Pavan, I. Strakovsky, R. Workman, and R. Arndt, “The Pion nucleon Sigma term is definitely large: Results from a G.W.U. analysis of pi nucleon scattering data,” *PiN Newslett.* **16** (2002) 110–115, [arXiv:hep-ph/0111066](#) [hep-ph].
- [52] XENON100 Collaboration, E. Aprile *et al.*, “Dark Matter Results from 225 Live Days of XENON100 Data,” *Phys.Rev.Lett.* **109** (2012) 181301, [arXiv:1207.5988](#) [astro-ph.CO].