DETERMINATION OF α_s AND m_c IN DEEP-INELASTIC SCATTERING

SERGEY ALEKHIN* and JOHANNES BLÜMLEIN†

*Deutsches Elektronen-Synchrotron DESY, Platanenallee 6,
Zeuthen, D15738, Germany*

**sergey.alekhin@desy.de*

†johannes.bluemlein@desy.de

SVEN-OLAF MOCH

*II. Institut für Theoretische Physik, Universität Hamburg,
Luruper Chaussee 149, Hamburg, D-22761, Germany*

sven-olaf.moch@desy.de

Received 4 July 2013

Accepted 4 July 2013

Published 13 August 2013

We describe the determination of the strong coupling constant $\alpha_s(M_Z^2)$ and of the charm-quark mass $m_c(m_c)$ in the $\overline{\text{MS}}$ -scheme, based on the QCD analysis of the unpolarized World deep-inelastic scattering data. At NNLO the values of $\alpha_s(M_Z^2) = 0.1134 \pm 0.0011(\text{exp})$ and $m_c(m_c) = 1.24 \pm 0.03(\text{exp})_{-0.02}^{+0.03}(\text{scale})_{-0.07}^{+0.00}(\text{th})$ are obtained and are compared with other determinations, and also clarifying discrepancies.

Keywords: Strong coupling constant; charm quark mass; parton distributions.

PACS Nos.: 12.38.Qk, 12.38.Bx

1. Introduction

The process of lepton–nucleon deep-inelastic scattering (DIS) is a clean source of basic information about the hadron substructure in terms of the parton model. Moreover, the QCD corrections to the parton model provide connection of the DIS structure functions with the parameters of the QCD Lagrangian, in particular to the strong coupling α_s and the heavy-quark masses. The higher order QCD corrections are manifest in the scaling violations of the structure functions w.r.t. the virtual photon momentum transfer Q^2 . This phenomenon was observed shortly after the discovery of the partonic structure of the nucleon and provided one of the first

*Permanent address: Institute for High Energy Physics, Pobeda 1, Protvino 142280, Russia.

constraints on α_s . With the dramatic improvement in the accuracy of the lepton–nucleon DIS data and the progress in the theoretical calculations, the value of α_s can in principle be determined with an accuracy of $O(1\%)$.^a The scaling violations, however, are also sensitive to the parton distribution functions (PDFs). Therefore the determination of α_s has to be performed simultaneously with the nucleon PDFs in global fits. Furthermore, an elaborate theoretical description of the QCD scaling violations is available for the leading-twist terms only. In practice this requires a careful isolation of the higher-twist effects and/or their independent phenomenological parametrization. Another important aspect of the DIS phenomenology is related to the c - and b -quark contributions. The heavy-quark production cross-section is sensitive to the heavy-quark masses. Therefore the DIS data provide a constraint on the c - and b -quark masses, $m_{c,b}$. The structure functions of the semi-inclusive process with the heavy quark in the final state are particularly useful for this purpose, although the data on inclusive structure functions are competitive with the semi-inclusive ones due to a much better accuracy. A major pitfall arising in the analysis of heavy-quark production is related to the account of the higher-order QCD corrections. Due to the two scales appearing in the problem, the calculations are quite involved. Therefore the NNLO corrections to the heavy-quark lepto-production are known in partial form only at present.² The problem of the high-order corrections is bypassed in the so-called variable-flavor-number (VFN) scheme assuming zero mass for the c - and b -quark. In this approximation the available high-order massless DIS Wilson coefficients can be employed for the calculation of the heavy-quark lepto-production rates. As is well known, the VFN approximation is obviously inapplicable at scales $Q^2 \sim m_{c,b}^2$ and it is commonly supplemented by *modeling* of the Wilson coefficient at low Q^2 in order to arrive in this way at a general-mass VFN (GMVFN) scheme. In this paper, we essentially focus on the determination of α_s and m_c based on the fixed-flavor-number (FFN) scheme. Here the mass effects are taken into account on field-theoretic grounds, free of model ambiguities appearing in the GMVFN schemes. Moreover, we employ the massive Wilson coefficients derived using the running-mass definition, which provide improved perturbative stability of the heavy-quark production rate.³

The paper is organized as follows. In Sec. 2, we outline the theoretical basis of the analysis and describe the data used. Sections 3 and 4 contain our results on the determination of α_s and the c -quark mass, respectively. In Sec. 5, we compare the FFN and VFN schemes with particular emphasis on the uncertainties in the determination of α_s and m_c .

2. Theoretical and Experimental Ingredients of the Analysis

Our determination of α_s and m_c is based on the QCD analysis of the DIS data obtained in fixed-target experiments and at the HERA collider. Only the proton-

^aFor a recent overview on precision determinations of $\alpha_s(M_Z^2)$ see Ref. 1.

and deuteron-target samples are selected which allows to minimize the impact of the nuclear corrections on the results.^b The DIS data are combined with the ones on the fixed-target Drell–Yan process, providing a supplementary constraint on the PDFs and facilitating the separation of the valence and sea quark distributions. The main version of our analysis is performed at NNLO using the three-loop anomalous dimensions in the PDF evolution and corresponding Wilson coefficients for the light-flavor DIS structure functions and the Drell–Yan process. For the neutral-current (NC) heavy-quark contribution we employ the approximate NNLO Wilson coefficients.² These terms were derived combining results obtained with the soft-gluon resummation technique and the high-energy limit of the DIS structure functions.⁵ These two approaches provide a good approximation at kinematics close to threshold of the heavy-quark production and far beyond the threshold, respectively. Between these two regimes the constraint coming from the available NNLO massive operator-matrix-element (OME) Mellin moments⁶ are employed. A remaining uncertainty in the NNLO Wilson coefficient obtained in this way is quantified by its margins, A and B . To find the best shape of the NNLO term preferred by the data, we use a linear interpolation between these margins

$$c_2^{(2)} = (1 - d_N) c_2^{(2),A} + d_N c_2^{(2),B} \quad (1)$$

and fit the interpolation parameter d_N to the data simultaneously with the PDF parameters, α_s and m_c . The charged-current (CC) heavy-quark production is calculated with account of the NLO corrections,^{7–9} which are the highest-order ones presently available. Both NC and CC massive Wilson coefficients are used in our analysis within the $\overline{\text{MS}}$ -definition of the heavy-quark mass. If compared to the case using the pole-mass, the present choice is perturbatively more stable³ since the mass can be defined at the typical renormalization/factorization scale of the process considered.

The leading-twist terms provided by the QCD-improved parton model are not sufficient at small Q^2 and/or final-state hadronic mass W , where parton correlations cannot be neglected. To account for these we add on the top of the leading-twist term the twist-4 contribution to the DIS structure functions $F_{2,T}$, parametrized in a model-independent form using spline interpolation.^c The twist-4 spline coefficients are fitted to the data together with other parameters. This is particularly important for the case of α_s in view of its strong correlation with the higher twist terms.

3. Strong Coupling Constant

The strong coupling constant α_s can be determined by comparing the Q^2 -dependence of the DIS cross-section measurements with the predictions based on the

^bFor a detailed description of the data set used, cf. Ref. 4.

^cThe twist-6 terms were also checked in the fit and found comparable to zero within errors applying the cut of $Q^2 > 2.5 \text{ GeV}^2$ used in the present analysis.

Fig. 1. The χ^2 -profile vs. the value of $\alpha_s(M_Z^2)$, for the separate data subsets, all obtained in variants of the ABM11 analysis with the value of α_s fixed and all other parameters fitted (solid lines: NNLO fit, dashes: NLO fit); from Ref. 4.

QCD-improved parton model. In our analysis the value of α_s is obtained simultaneously with the nucleon PDFs and the twist-4 terms. This allows to take into account the correlation of α_s with other parameters affecting the data Q^2 -dependence. The central value of α_s obtained in this way depends on the perturbative order and reduces from NLO to NNLO. In particular, the ABM11 fit⁴ yields

$$\begin{aligned}\alpha_s(M_Z^2) &= 0.1180 \pm 0.0012 \text{ (exp)} & \text{NLO}, \\ \alpha_s(M_Z^2) &= 0.1134 \pm 0.0011 \text{ (exp)} & \text{NNLO}.\end{aligned}\tag{2}$$

The values of α_s preferred by each particular DIS data set are demonstrated in Fig. 1 by means of the χ^2 -profiles obtained in the variants of the ABM11 fit with α_s fixed at the values in the range of 0.104–0.130. The HERA and BCDMS data sets have a similar χ^2 -shape with minima around the values Eq. (2), while the SLAC and NMC data pull the value of α_s somewhat up and down, respectively. Note that the two latter sets are sensitive to higher-twist terms due to substantial small- Q^2 contributions in these samples. In contrast, the HERA and BCDMS data are far less sensitive to higher twist terms and it is worth noting that in the variant of the ABM11 fit excluding the SLAC and NMC data and setting the higher twist terms to zero we find $\alpha_s(M_Z^2) = 0.1133 \pm 0.0011 \text{ (exp.)}$ at NNLO. The good agreement of this value with Eq. (2) substantiates the consistency between different data sets

Fig. 2. The values of $\alpha_s(M_Z^2)$ obtained in the ABM fit⁴ in comparison to the ones obtained by JR,¹² CTEQ,¹³ MSTW,¹⁴ and NNPDF¹⁵ from the analysis of the DIS data (squares) and from the combination of the DIS and jet Tevatron data^{16,17} (circles).

in our analysis once the higher twist terms are taken into account. Moreover, this cross-check confirms that the combination of the BCDMS and HERA data can be used for accurate determination of $\alpha_s(M_Z^2)$ since these two data sets provide complimentary constraints on the PDFs,¹⁰ see also Ref. 11.

The NNLO value of α_s , Eq. (2), is in a good agreement with the results of the JR analysis¹² and the recent CTEQ determination,¹³ while the MSTW¹⁴ and NNPDF¹⁵ groups report substantially bigger values, cf. Fig. 2. The discrepancy with MSTW can be explained in part by impact of the jet Tevatron data, which pull the value of $\alpha_s(M_Z^2)$ up by 0.001–0.002, depending on the fit details. Furthermore, changing our fit ansatz in direction of the MSTW and NNPDF, we obtain their value of α_s . In particular, dropping the higher twist terms simultaneously with an additional cut of $W^2 > 12.5 \text{ GeV}^2$ imposed by MSTW and NNPDF, we obtain $\alpha_s(M_Z^2) = 0.1191 \pm 0.0006(\text{exp.})$. In a similar way, disregarding the error correlation in the HERA and NMC data, like in the MSTW analysis, we obtain $\alpha_s(M_Z^2)$ shifted by +0.0026, in the direction of the MSTW value. Note that in this context the recently updated JR analysis¹² treats the higher twist properly. The CTEQ analysis¹³ seems to be less sensitive to the impact of the higher twist contributions, compared to that by MSTW and NNPDF due to a more stringent cut on Q^2 .

4. The Mass of the Charm Quark

The sensitivity to the charm quark mass m_c in our analysis appears essentially due to the data on the NC and CC inclusive DIS,¹⁹ and the CC semi-inclusive charm production in DIS^{20,21} with the most essential experimental constraint on m_c

Fig. 3. The combined HERA data on the reduced cross-section for the open charm production¹⁸ vs. x at different values of Q^2 in comparison with the analysis of Ref. 22 at NLO (dashed line) and NNLO (solid line) together with a fit variant based on the option $(A+B)/2$ of the NNLO Wilson coefficients of Ref. 2, cf. Eq. (1) (dotted line); from Ref. 22.

coming from the semi-inclusive charm-production HERA data.¹⁸ The latter sample comprises the statistics of the H1 and ZEUS experiments obtained for different c -quark decay channels. The combination was performed similarly to the case of the inclusive HERA data¹⁹ and allows to reduce the systematic errors of each experiment due to cross-calibration of the experiments. The FFN scheme provides a good description of the semi-inclusive HERA data up to the largest Q^2 -values covered, cf. Fig. 3, with a value of $\chi^2/\text{NDP} = 61/52$ at NNLO. Here NDP denotes the number of data points. The $\overline{\text{MS}}$ -values of m_c found in the analysis of Ref. 22 are

$$m_c(m_c) = 1.15 \pm 0.04 (\text{exp})_{-0.00}^{+0.04} (\text{scale}) \quad \text{NLO}, \quad (3)$$

$$m_c(m_c) = 1.24 \pm 0.03 (\text{exp})_{-0.02}^{+0.03} (\text{scale})_{-0.07}^{+0.00} (\text{th}) \quad \text{NNLO}_{\text{approx}}, \quad (4)$$

at NLO and NNLO, respectively, see also Ref. 23. The experimental accuracy of 30 MeV obtained at NNLO is quite competitive with other determinations of m_c

Fig. 4. The values of $m_c(m_c)$ obtained in the NLO and NNLO variants of our analysis with the value of $\alpha_s(M_Z^2)$ fixed. The position of the star displays the result with the value of $\alpha_s(M_Z^2)$ fitted⁴; from Ref. 22.

based on the e^+e^- data and the central value Eq. (4) is in a good agreement with the world average.²⁴ The scale error in Eqs. (3) and (4) is obtained varying the factorization scale by a factor of 1/2 and 2 around the nominal value of $\sqrt{m_c^2 + \kappa Q^2}$, where $\kappa = 4$ for NC and $\kappa = 1$ for CC heavy-quark production, respectively. In the NNLO case an additional error related to the uncertainty in the massive Wilson coefficients contributes. The value of Eq. (4) is obtained for the interpolation parameter $d_N = -0.1$ being preferred by the fit, roughly corresponding to option *A* of the Wilson coefficients of Ref. 2. Meanwhile, option *B* is clearly excluded by the data with $\chi^2/\text{NDP} = 115/52$. Therefore the uncertainty due to the missing NNLO massive terms is estimated as a variation between options *A* and $(A+B)/2$ which yields the value of 70 MeV, Eq. (4). The value of $m_c(m_c)$ obtained in our analysis demonstrates remarkable stability w.r.t. $\alpha_s(M_Z^2)$. Performing variants of our analysis with the values of α_s fixed in the wide range around the best one preferred by the data, we find a variation of $m_c(m_c)$ in the range of 10–20 MeV, depending on the order, cf. Fig 4.

The NLO value of $m_c(m_c)$ obtained in our analysis is somewhat lower than the one of $m_c(m_c) = 1.26 \pm 0.05(\text{exp})$ GeV from the analysis based on the HERA data only.¹⁸ To understand the difference we checked the cases of a cut on $Q^2 = 3.5 \text{ GeV}^2$, likewise in the HERA fit, and no semi-inclusive Tevatron data^{20,21} included. As a result we obtain shifts in $m_c(m_c)$ by +30 MeV and +40 MeV, respectively. The remaining discrepancy should be attributed to the particularities of the HERA PDFs. The value of $m_c(m_c)$ was also recently determined by the CTEQ collaboration.²⁵ In contrast to our case this determination is based on the S-ACOT- χ prescription as GMVFN scheme. Furthermore, the $\overline{\text{MS}}$ coefficients of

Ref. 25 are obtained by straightforward substitution of the pole- and running-mass matching relation into the pole-mass coefficients. The expressions obtained in such a way correspond to a mixed order in α_s . Moreover, the advantage of this approach is not evident in view of the poor perturbative convergence of the mass matching relation. The central CTEQ value of $m_c(m_c) = 1.12^{+0.11}_{-0.17}$ GeV is lower than the world average, while the CTEQ errors are much larger than those in Eqs. (3) and (4) due to the impact of the uncertainty in the GMVFN scheme modeling.

5. VFN Uncertainties

The choice of the factorization scheme plays an essential role in the analysis of the existing DIS data due to important constraints coming from the small- x region, where the heavy quark contribution is numerically large. While our analysis is based on the FFN scheme, many other groups employ different variants of the GMVFN scheme, which differ by *modeling* of the low- Q^2 region. The spread between these variants is rather substantial and thus implies a corresponding uncertainty in the basic parameters determined in these GMVFN fits. In particular, the value of m_c determined from a combination of the inclusive and semi-inclusive HERA data with the different versions of the ACOT and RT prescriptions for the VFN scheme demonstrate a spread of 400 MeV.¹⁸ There are also sources of the VFN scheme uncertainties, which are common for all these prescriptions. Firstly, the matching of heavy-quark PDFs is commonly performed at the factorization scale μ equal to the $\mu_0 = m_c$, respectively m_b . Clearly, at these scales neither the heavy flavors can be dealt with as massless. The matching point μ_0 is not fixed by theory and in principle it can vary in a wide range being an artefact of the description not contributing to the observables according to the renormalization group equations. Additional uncertainty emerge for the 4(5)-flavor PDFs in the NNLO analysis. They are commonly matched with the 3(4)-flavor ones using *NLO matching condition*, since the NNLO OMEs are not yet known in the complete form.^d However, to provide consistency with the NNLO Wilson coefficients the evolution of these PDFs is performed in the NNLO approximation that introduces an additional uncertainty due to missing higher-order corrections into the analysis. At the same time, the evolution of the 4(5)-flavor PDFs in the VFN scheme leads to a resummation of the terms $\sim \ln(\mu^2)$ which in part reproduce the higher-order corrections, being known, however, not to be dominant. Relations between the resummation effects and the VFN evolution uncertainties are illustrated in Fig. 5 by comparison of the μ -derivatives for the c -quark distribution being calculated in different ways. In one case the distributions are matched at the scale of μ_0 using the fixed-order-perturbation-theory (FOPT) matching conditions and then evolved starting from μ_0 with the massless splitting functions. In another case they are calculated with the FOPT matching conditions at all scales. The difference between these two

^dFor progress in this field, cf. Refs. 26–29.

Fig. 5. The difference between the c -quark PDFs derivatives $\dot{c}(x, \mu^2) \equiv \frac{dc(x, \mu^2)}{d \ln \mu^2}$ calculated with the FOPT matching condition and with the massless 4-flavor evolution starting at the matching point $\mu_0 = m_c = 1.4$ GeV vs. the factorization scale μ^2 at different values of x in the LO, NLO, and NNLO* approximations. The arrows display upper margin of the HERA collider kinematics with the collision c.m.s. energy squared $s = 10^5$ GeV² and the vertical lines correspond to the matching point position μ_0 .

cases do not demonstrate a significant rise with μ . The only exclusion is observed at $x \lesssim 0.0001$ and at scales outside the kinematics being probed in experiment. Therefore it cannot be attributed to the impact of the log-term resummation. In contrast, there is a substantial difference in the derivatives calculated in the NLO and NNLO*, i.e. the combination of the NLO matching with the NNLO evolution. This difference yields an estimate of the uncertainty in the VFN scheme due to the missing higher-orders, which is obviously larger than the resummation effects. Checking the impact of this uncertainty w.r.t. the value of $\alpha_s(M_Z^2)$ in combination with the variation of the matching point for the 4-flavor PDFs in the range of 1.2–1.5 GeV we find a value of ± 0.001 . This is comparable to the experimental uncertainty and makes the VFN schemes incompetent with the FFN one in the precision determination of $\alpha_s(M_Z^2)$.

Acknowledgments

We thank P. Jimenez-Delgado and E. Reya for discussions. This work has been supported in part by Helmholtz Gemeinschaft under contract VH-HA-101 (*Alliance Physics at the Terascale*), DFG Sonderforschungsbereich/Transregio 9 and by the European Commission through contract PITN-GA-2010-264564 (*LHCPhenoNet*).

References

1. S. Bethke *et al.*, arXiv:1110.0016.
2. H. Kawamura, N. A. Lo Presti, S. Moch and A. Vogt, *Nucl. Phys. B* **864**, 399 (2012).

3. S. Alekhin and S. Moch, *Phys. Lett. B* **699**, 345 (2011).
4. S. Alekhin, J. Blümlein and S. Moch, *Phys. Rev. D* **86**, 054009 (2012).
5. S. Catani, M. Ciafaloni and F. Hautmann, *Nucl. Phys. B* **366**, 135 (1991).
6. I. Bierenbaum, J. Blümlein and S. Klein, *Nucl. Phys. B* **820**, 417 (2009).
7. T. Gottschalk, *Phys. Rev. D* **23**, 56 (1981).
8. M. Glück, S. Kretzer and E. Reya, *Phys. Lett. B* **380**, 171 (1996) [Erratum-*ibid.* **405**, 391 (1997)].
9. J. Blümlein, A. Hasselhuhn, P. Kovacikova and S. Moch, *Phys. Lett. B* **700**, 294 (2011).
10. H1 Collab. (C. Adloff *et al.*), *Eur. Phys. J. C* **21**, 33 (2001).
11. S. Alekhin, J. Blümlein and S. Moch, arXiv:1303.1073.
12. P. Jimenez-Delgado and E. Reya, private communication.
13. J. Gao *et al.*, arXiv:1302.6246.
14. A. Martin, W. J. Stirling, R. Thorne and G. Watt, *Eur. Phys. J. C* **64**, 653 (2009).
15. R. D. Ball *et al.*, *Phys. Lett. B* **707**, 66 (2012).
16. A. Abulencia *et al.*, *Phys. Rev. D* **75**, 092006 (2007).
17. V. Abazov *et al.*, *Phys. Rev. Lett.* **101**, 062001 (2008).
18. H1 and ZEUS Collabs. (H. Abramowicz *et al.*), *Eur. Phys. J. C* **73**, 2311 (2013).
19. H1 and ZEUS Collabs. (F. D. Aaron *et al.*), *JHEP* **1001**, 109 (2010).
20. CCFR Collab. (A. O. Bazarko *et al.*), *Z. Phys. C* **65**, 189 (1995).
21. NuTeV Collab. (M. Goncharov *et al.*), *Phys. Rev. D* **64**, 112006 (2001).
22. S. Alekhin *et al.*, *Phys. Lett. B* **720**, 172 (2013).
23. S. Alekhin, K. Daum, K. Lipka and S. Moch, *Phys. Lett. B* **718**, 550 (2012).
24. Particle Data Group Collab. (J. Beringer *et al.*), *Phys. Rev. D* **86**, 010001 (2012).
25. J. Gao, M. Guzzi and P. M. Nadolsky, arXiv:1304.3494.
26. J. Ablinger *et al.*, *PoS LL* **2012**, 033 (2012).
27. J. Ablinger *et al.*, *Nucl. Phys. B* **864**, 52 (2012).
28. J. Ablinger *et al.*, *Nucl. Phys. B* **844**, 26 (2011).
29. J. Blümlein, A. Hasselhuhn, S. Klein and C. Schneider, *Nucl. Phys. B* **866**, 196 (2013).