


Transverse and longitudinal lambda polarization at HERMES



Gevorg Karyan

on behalf of the HERMES Collaboration

**Alikhanyan National Science Laboratory
Yerevan, Armenia**

Why Λ polarization is important

Self-analyzing power through
the parity-violating weak decay

Λ (uds)




Spin structure of the
lightest hyperon

Why Λ polarization is important

Sensitivity to the strange
quark polarization

Λ (uds)


A diagram illustrating the internal structure of a Lambda baryon (Λ), which consists of two up quarks (u) and one down quark (d). The quarks are represented by colored spheres (red for up, blue for down) with arrows indicating their spin directions. A green circular arrow around the leftmost quark indicates its transverse spin. A solid red arrow points from the center of the baryon towards the right, representing the polarization of the baryon. A dashed blue arrow points from the center towards the right, representing the polarization of the strange quark (s) within the baryon.

Access to the quarks
transverse spin distribution

How to access Λ polarization




$$\frac{dN}{d\Omega_p} \simeq 1 + \alpha \vec{P}_\Lambda \cdot \hat{\mathbf{k}}_p$$

$\Lambda(u d s)$



Decay protons prefer to follow the spin direction of Λ

Experiment



Beam : e^-/e^+ 27.6 GeV


Target : H, D, He^3 , Ne, Kr, Xe pure gaseous

Good momentum resolution : $\frac{\delta p}{p} < 2 \%$

Excellent particle identification

Longitudinal Λ polarization

$$e + N \rightarrow e + \Lambda + X$$




Longitudinal Λ polarization

$$\vec{e} + N \rightarrow e + \vec{\Lambda} + X$$

Longitudinally polarized

Unpolarized



Longitudinal Λ polarization


$$\vec{e} + N \rightarrow e + \vec{\Lambda} + X$$

Longitudinally polarized

Unpolarized

$$D(y) \approx (1 - (1 - y)^2) / (1 + (1 - y)^2)$$

$$y = \nu / E_b$$



Longitudinal Λ polarization


$$\vec{e} + N \rightarrow e + \vec{\Lambda} + X$$

Longitudinally polarized

Unpolarized

$$D(y) \approx (1 - (1-y)^2) / (1 + (1-y)^2)$$


$$y = \nu / E_b$$



Longitudinal Λ polarization

Phys. Rev. D74 (2006) 072004

(all targets except Xe)



$$D_{LL'} = 0.11 \pm 0.10 \pm 0.03$$


Small value for $D_{LL'}$ is observed.

Dominance of scattering from u and d quarks.

Strong rise at high z in models: polarized $s \rightarrow \Lambda$ (no hyperon decay in these models).

Transverse Λ polarization

$e + N \rightarrow \Lambda + X$ (quasi-real photo-production)



Transverse Λ polarization




Parity

conservation

Unpolarized

direction of spontaneous polarization



$$P_B P_T = 0.0000 \pm 0.0005$$


P_B - beam polarization

P_T - target polarization

Transverse Λ polarization



$$\frac{dN}{d\Omega_p} = \frac{dN_0}{d\Omega_p} (1 + \alpha |\vec{P}_\Lambda| \cos \theta_p)$$




$$\vec{P}_\Lambda = |\vec{P}_\Lambda| \cdot \hat{n}, \quad \hat{n} = \frac{\vec{p}_e \times \vec{p}_\Lambda}{|\vec{p}_e \times \vec{p}_\Lambda|}$$
$$\alpha = 0.642 \pm 0.013$$

Transverse Λ polarization

Phys. Rev. D76 (2007) 092008

(all targets except Xe)



$$P_n^\Lambda = 0.078 \pm 0.006 \pm 0.012$$


$$P_n^{\bar{\Lambda}} = -0.025 \pm 0.015 \pm 0.018$$

Λ polarization is found to be positive (opposite sign compared to pion and proton beam data).

Transverse Λ polarization

Phys. Rev. D76 (2007) 092008

(all targets except Xe)



$$P_n^\Lambda = 0.078 \pm 0.006 \pm 0.012$$
$$P_n^{\bar{\Lambda}} = -0.025 \pm 0.015 \pm 0.018$$


$$\zeta = \frac{(E_\Lambda + p_{z\Lambda})}{E_e + p_e}$$

correlated with x_F


Transverse Λ polarization

Phys. Rev. D76 (2007) 092008

(all targets except Xe)




$$P_n^\Lambda = 0.078 \pm 0.006 \pm 0.012$$
$$P_n^{\bar{\Lambda}} = -0.025 \pm 0.015 \pm 0.018$$



Transverse Λ polarization

Phys. Rev. D76 (2007) 092008

(all targets except Xe)




$$P_n^\Lambda = 0.078 \pm 0.006 \pm 0.012$$

$$P_n^{\bar{\Lambda}} = -0.025 \pm 0.015 \pm 0.018$$

Different magnitudes for Λ polarization in the "backward" ($\zeta < 0.25$) and "forward" ($\zeta > 0.25$) kinematic regions.

Transverse Λ polarization



Λ polarization rises linearly with p_T in both kinematic regions.

Transverse Λ polarization on nuclei

Atomic mass
dependence of
 Λ polarization

H

D

^4He


Ne

Kr

Xe

Transverse Λ polarization on nuclei

Phys. Rev. D90 (2014) 072007




Positive polarization for light nuclei.

Compatible with zero polarization for heavy nuclei.

Transverse Λ polarization on nuclei

Phys. Rev. D90 (2014) 072007



H+D data : polarization increases linearly with p_T at small ζ (backward region).

H+D data : polarization is substantially smaller in forward region ($\zeta > 0.3$) with very little dependence on p_T .

Kr+Xe data : polarization is compatible with zero within experimental uncertainties.

- **Small polarization transfer from a polarized beam to the lambda.**
- **Positive sign for the lambda transverse polarization.**
- **Different transverse polarizations for the lambda in “backward” and “forward” regions.**
- **Linear rise for the lambda transverse polarization with its transverse momentum.**
- **Positive transverse polarization for light nuclei and compatible with zero polarization for heavy nuclei.**