

Elucidating SUSY in the Interplay of LHC and ILC - $\tilde{\tau}$ co-annihilation models

M. Berggren, J. List,

I. Melzer-Pellmann, A. Cakir, B. Samani, C. Seitz, D. Krücker, S.
Wayand

¹DESY, Hamburg

9th Annual Meeting of the Helmholtz Alliance "Physics at the
Terascale"



Outline

- 1 Introduction: Our model
- 2 The STCx benchmark @ LHC
- 3 The ILC
- 4 The STCx benchmark @ ILC
 - STCx @ 500 GeV: Globaly
 - STCx @ 500 GeV: $\tilde{e}, \tilde{\mu}, \tilde{\tau}_1$
- 5 Summary

For all the details, see arXiv:1508.04383 (submitted to EPJC)

Studying SUSY in rich models

Remember, apart from stabilising the Higgs mass (naturalness) SUSY can address:

- Anomaly in $g - 2$ of the μ : Would prefer a not-too-heavy smuon.
- Dark matter : A WIMP of ~ 100 GeV would be required. And a process not to over-produce it, eg. by co-annihilation by a nearby NLSP.
- EW symmetry breaking, coupling constant unification: points to NP at or below 1 TeV
- To suppress the SUSY flavour problem (FCNC:s etc): Heavy 1:st & 2:nd generation squarks would be nice ...
- Need to meet Low-energy constrains : $b \rightarrow s\gamma$, $b \rightarrow \mu\mu$, ρ -parameter, $\Gamma(Z)$...
- All points to SUSY with a rich spectrum to explore for LHC14 and ILC!

But: Can we still get all this, without contradicting LHC8 limits ?!

Studying SUSY in rich models

Remember, apart from stabilising the Higgs mass (naturalness) SUSY can address:

- Anomaly in $g - 2$ of the μ : Would prefer a not-too-heavy smuon.
- Dark matter : A WIMP of ~ 100 GeV would be required. And a process not to over-produce it, eg. by co-annihilation by a nearby NLSP.
- EW symmetry breaking, coupling constant unification: points to NP at or below 1 TeV
- To suppress the SUSY flavour problem (FCNC:s etc): Heavy 1:st & 2:nd generation squarks would be nice ...
- Need to meet Low-energy constrains : $b \rightarrow s\gamma$, $b \rightarrow \mu\mu$, ρ -parameter, $\Gamma(Z)$...
- All points to SUSY with a rich spectrum to explore for LHC14 and ILC!

But: Can we still get all this, without contradicting LHC8 limits ?!

Studying SUSY in rich models

Remember, apart from stabilising the Higgs mass (naturalness) SUSY can address:

- Anomaly in $g - 2$ of the μ : Would prefer a not-too-heavy smuon.
- Dark matter : A WIMP of ~ 100 GeV would be required. And a process not to over-produce it, eg. by co-annihilation by a nearby NLSP.
- EW symmetry breaking, coupling constant unification: points to NP at or below 1 TeV
- To suppress the SUSY flavour problem (FCNC:s etc): Heavy 1:st & 2:nd generation squarks would be nice ...
- Need to meet Low-energy constrains : $b \rightarrow s\gamma$, $b \rightarrow \mu\mu$, ρ -parameter, $\Gamma(Z)$...
- All points to SUSY with a rich spectrum to explore for LHC14 and ILC!

But: Can we still get all this, without contradicting LHC8 limits ?!

Studying SUSY in rich models

Remember, apart from stabilising the Higgs mass (naturalness) SUSY can address:

- Anomaly in $g - 2$ of the μ : Would prefer a not-too-heavy smuon.
- Dark matter : A WIMP of ~ 100 GeV would be required. And a process not to over-produce it, eg. by co-annihilation by a nearby NLSP.
- EW symmetry breaking, coupling constant unification: points to NP at or below 1 TeV
- To suppress the SUSY flavour problem (FCNC:s etc): Heavy 1:st & 2:nd generation squarks would be nice ...
- Need to meet Low-energy constrains : $b \rightarrow s\gamma$, $b \rightarrow \mu\mu$, ρ -parameter, $\Gamma(Z)$...
- All points to SUSY with a rich spectrum to explore for LHC14 and ILC!

But: Can we still get all this, without contradicting LHC8 limits ?!

Studying SUSY in rich models

Remember, apart from stabilising the Higgs mass (naturalness) SUSY can address:

- Anomaly in $g - 2$ of the μ : Would prefer a not-too-heavy smuon.
- Dark matter : A WIMP of ~ 100 GeV would be required. And a process not to over-produce it, eg. by co-annihilation by a nearby NLSP.
- EW symmetry breaking, coupling constant unification: points to NP at or below 1 TeV
- To suppress the SUSY flavour problem (FCNC:s etc): Heavy 1:st & 2:nd generation squarks would be nice ...
- Need to meet Low-energy constrains : $b \rightarrow s\gamma$, $b \rightarrow \mu\mu$, ρ -parameter, $\Gamma(Z)$...
- All points to SUSY with a rich spectrum to explore for LHC14 and ILC!


But: Can we still get all this, without contradicting LHC8 limits ?!

Yes! Matercode global 10-parameter fit

pMSSM10 prediction: best-fit masses




[2015]



- ⇒ high colored masses
- ⇒ relatively low electroweak masses
 - partially with not too large ranges
- ⇒ clear prediction for ILC and CLIC

Our model: STCx

High mass squarks+gluino




Well-tempered higgs, bosino
and slepton sector

Varying 3-gen squarks

Our model: STCx

High mass squarks+gluino




Well-tempered higgs, bosino
and slepton sector

Varying 3-gen squarks

Our model: STCx

High mass squarks+gluino



Well-tempered higgs, bosino
and slepton sector

Varying 3-gen squarks

STCx @ LHC14

- Main features at LHC 14 TeV:

- Cross-sections:

- $\tilde{\chi}_k^0 \tilde{\chi}_l^\pm \sim 1 \text{ pb} > \tilde{\chi}_k^\pm \tilde{\chi}_l^\pm > \tilde{\tau}\tilde{\tau} > \tilde{\ell}\tilde{\ell} > \tilde{t}\tilde{t} > \tilde{b}\tilde{b} > \tilde{q}\tilde{q} > \tilde{\chi}_k^0 \tilde{\chi}_l^0 > \tilde{g}\tilde{g} \sim 1 \text{ fb}$

- $\tilde{\chi}$ cascade-decays to τ :s + the LSP in 75 % of the cases, often together with a boson (Z , W or h).

- For $\tilde{\chi}_k^0$, the rest is either only bosons, or "nothing" (ie. neutrinos).

- For $\tilde{\chi}_k^\pm$ the rest is other leptons.

- The τ :s mostly come from $\tilde{\tau}_1 \rightarrow \tau \tilde{\chi}_0^0$, where the mass difference is only 10 GeV \Rightarrow little missing energy.

- \tilde{b} mostly decays to $b \tilde{\chi}_0^0$: > 50 % to $b \tilde{\chi}_1^0$.

- \tilde{t} always goes to $t \tilde{\chi}_0^0$, but rarely to $t \tilde{\chi}_1^0$ (~ 10 %).

- The right-handed gen1 and 2 squarks almost always decay directly to quark+LSP.

STCx @ LHC14

- Main features at LHC 14 TeV:

- Cross-sections:

- $\tilde{\chi}_k^0 \tilde{\chi}_l^\pm \sim 1 \text{ pb} > \tilde{\chi}_k^\pm \tilde{\chi}_l^\pm > \tilde{\tau}\tilde{\tau} > \tilde{\ell}\tilde{\ell} > \tilde{t}\tilde{t} > \tilde{b}\tilde{b} > \tilde{q}\tilde{q} > \tilde{\chi}_k^0 \tilde{\chi}_l^0 > \tilde{g}\tilde{g} \sim 1 \text{ fb}$

- $\tilde{\chi}$ cascade-decays to τ :s + the LSP in 75 % of the cases, often together with a boson (Z , W or h).

- For $\tilde{\chi}_k^0$, the rest is either only bosons, or "nothing" (ie. neutrinos).

- For $\tilde{\chi}_k^\pm$ the rest is other leptons.

- The τ :s mostly come from $\tilde{\tau}_1 \rightarrow \tau \tilde{\chi}_0^0$, where the mass difference is only 10 GeV \Rightarrow little missing energy.

- \tilde{b} mostly decays to $b \tilde{\chi}_0^0$: $> 50\%$ to $b \tilde{\chi}_1^0$.

- \tilde{t} always goes to $t \tilde{\chi}_0^0$, but rarely to $t \tilde{\chi}_1^0$ ($\sim 10\%$).

- The right-handed gen1 and 2 squarks almost always decay directly to quark+LSP.

STCx @ LHC14

- Main features at LHC 14 TeV:
 - Cross-sections:
 - $\tilde{\chi}_k^0 \tilde{\chi}_l^\pm \sim 1 \text{ pb} > \tilde{\chi}_k^\pm \tilde{\chi}_l^\pm > \tilde{\tau} \tilde{\tau} > \tilde{\ell} \tilde{\ell} > \tilde{t} \tilde{t} > \tilde{b} \tilde{b} > \tilde{q} \tilde{q} > \tilde{\chi}_k^0 \tilde{\chi}_l^0 > \tilde{g} \tilde{g} \sim 1 \text{ fb}$
 - $\tilde{\chi}$ cascade-decays to τ :s + the LSP in 75 % of the cases, often together with a boson (Z , W or h).
 - For $\tilde{\chi}^0$, the rest is either only bosons, or "nothing" (ie. neutrinos).
 - For $\tilde{\chi}^\pm$ the rest is other leptons.
 - The τ :s mostly come from $\tilde{\tau}_1 \rightarrow \tau \tilde{\chi}_0^0$, where the mass difference is only 10 GeV \Rightarrow little missing energy.
 - \tilde{b} mostly decays to $b \tilde{\chi}_1^0$: $> 50\%$ to $b \tilde{\chi}_1^0$.
 - \tilde{t} always goes to $t \tilde{\chi}_1^0$, but rarely to $t \tilde{\chi}_1^0$ ($\sim 10\%$).
 - The right-handed gen1 and 2 squarks almost always decay directly to quark+LSP.

STCx @ LHC14

- Main features at LHC 14 TeV:

- Cross-sections:

- $\tilde{\chi}_k^0 \tilde{\chi}_l^\pm \sim 1 \text{ pb} > \tilde{\chi}_k^\pm \tilde{\chi}_l^\pm > \tilde{\tau}\tilde{\tau} > \tilde{\ell}\tilde{\ell} > \tilde{t}\tilde{t} > \tilde{b}\tilde{b} > \tilde{q}\tilde{q} > \tilde{\chi}_k^0 \tilde{\chi}_l^0 > \tilde{g}\tilde{g} \sim 1 \text{ fb}$

- $\tilde{\chi}$ cascade-decays to τ :s + the LSP in 75 % of the cases, often together with a boson (Z , W or h).

- For $\tilde{\chi}^0$, the rest is either only bosons, or "nothing" (ie. neutrinos).
 - For $\tilde{\chi}^\pm$ the rest is other leptons.

- The τ :s mostly come from $\tilde{\tau}_1 \rightarrow \tau \tilde{\chi}_0^0$, where the mass difference is only 10 GeV \Rightarrow little missing energy.

- \tilde{b} mostly decays to $b \tilde{\chi}_0^0$: $> 50\%$ to $b \tilde{\chi}_1^0$.

- \tilde{t} always goes to $t \tilde{\chi}_0^0$, but rarely to $t \tilde{\chi}_1^0$ ($\sim 10\%$).

- The right-handed gen1 and 2 squarks almost always decay directly to quark+LSP.

STCx @ LHC14

- Main features at LHC 14 TeV:
 - Cross-sections:
 - $\tilde{\chi}_k^0 \tilde{\chi}_l^\pm \sim 1 \text{ pb} > \tilde{\chi}_k^\pm \tilde{\chi}_l^\pm > \tilde{\tau}\tilde{\tau} > \tilde{\ell}\tilde{\ell} > \tilde{t}\tilde{t} > \tilde{b}\tilde{b} > \tilde{q}\tilde{q} > \tilde{\chi}_k^0 \tilde{\chi}_l^0 > \tilde{g}\tilde{g} \sim 1 \text{ fb}$
 - $\tilde{\chi}$ cascade-decays to τ :s + the LSP in 75 % of the cases, often together with a boson (Z , W or h).
 - For $\tilde{\chi}^0$, the rest is either only bosons, or "nothing" (ie. neutrinos).
 - For $\tilde{\chi}^\pm$ the rest is other leptons.
 - The τ :s mostly come from $\tilde{\tau}_1 \rightarrow \tau \tilde{\chi}_0^0$, where the mass difference is only 10 GeV \Rightarrow little missing energy.
 - \tilde{b} mostly decays to $b \tilde{\chi}_1^0$: $> 50\%$ to $b \tilde{\chi}_1^0$.
 - \tilde{t} always goes to $t \tilde{\chi}_1^0$, but rarely to $t \tilde{\chi}_1^0$ ($\sim 10\%$).
 - The right-handed gen1 and 2 squarks almost always decay directly to quark+LSP.

STCx @ LHC14

- Main features at LHC 14 TeV:
 - Cross-sections:
 - $\tilde{\chi}_k^0 \tilde{\chi}_l^\pm \sim 1 \text{ pb} > \tilde{\chi}_k^\pm \tilde{\chi}_l^\pm > \tilde{\tau}\tilde{\tau} > \tilde{\ell}\tilde{\ell} > \tilde{t}\tilde{t} > \tilde{b}\tilde{b} > \tilde{q}\tilde{q} > \tilde{\chi}_k^0 \tilde{\chi}_l^0 > \tilde{g}\tilde{g} \sim 1 \text{ fb}$
 - $\tilde{\chi}$ cascade-decays to τ :s + the LSP in 75 % of the cases, often together with a boson (Z , W or h).
 - For $\tilde{\chi}^0$, the rest is either only bosons, or "nothing" (ie. neutrinos).
 - For $\tilde{\chi}^\pm$ the rest is other leptons.
 - The τ :s mostly come from $\tilde{\tau}_1 \rightarrow \tau \tilde{\chi}_0^0$, where the mass difference is only 10 GeV \Rightarrow little missing energy.
 - \tilde{b} mostly decays to $b \tilde{\chi}_1^0$: $> 50\%$ to $b \tilde{\chi}_1^0$.
 - \tilde{t} always goes to $t \tilde{\chi}_1^0$, but rarely to $t \tilde{\chi}_1^0$ ($\sim 10\%$).
 - The right-handed gen1 and 2 squarks almost always decay directly to quark+LSP.

STCx @ LHC14

- Main features at LHC 14 TeV:

⇒ LHC expectations

- Despite the high cross-section, the low amount of missing E_T and the long decay chains will make **direct bosino and slepton observations hard**.
- The simple decay-chains and very high missing E_T will make **first- and second-generation squark** production easy to detect. However, the **cross-section is low** so that it is still **challenging**.
- **Third generation squark** production constitute a good compromise between cross-section and visibility, and will be the **most powerful discovery channel**. The lower cross-section in STC10 is compensated by higher visibility.

STCx @ LHC14

- Main features at LHC 14 TeV:

⇒ LHC expectations

- Despite the high cross-section, the low amount of missing E_T and the long decay chains will make **direct bosino and slepton observations hard**.
- The simple decay-chains and very high missing E_T will make **first- and second-generation squark** production easy to detect. However, the **cross-section is low** so that it is still **challenging**.
- **Third generation squark** production constitute a good compromise between cross-section and visibility, and will be the **most powerful discovery channel**. The lower cross-section in STC10 is compensated by higher visibility.

STCx @ LHC14

- Main features at LHC 14 TeV:

⇒ LHC expectations


- Despite the high cross-section, the low amount of missing E_T and the long decay chains will make **direct bosino and slepton observations hard**.
- The simple decay-chains and very high missing E_T will make **first- and second-generation squark** production easy to detect. However, the **cross-section is low** so that it is still **challenging**.
- **Third generation squark** production constitute a good compromise between cross-section and visibility, and will be the **most powerful discovery channel**. The lower cross-section in STC10 is compensated by higher visibility.

STCx @ LHC14: Full-hadronic search

- Select events with large amount of E_T in central, high E jets (HT) \Rightarrow hard interaction.
- Further select large *missing* E_T (MHT) \Rightarrow Partly invisible decay of heavy particle to a (much) lighter one.
- Avoid “signals” from bad measurements: Angle between any jet and the MHT direction large. Veto isolated leptons.
- HT distribution.
- Same, but enhance in \tilde{b} and $\tilde{\tau}$ by demanding two b-tagged jets.


STCx @ LHC14: Full-hadronic search

- Select events with large amount of E_T in central, high E jets (HT) \Rightarrow hard interaction.
- Further select large *missing* E_T (MHT) \Rightarrow Partly invisible decay of heavy particle to a (much) lighter one.
- Avoid “signals” from bad measurements: Angle between any jet and the MHT direction large. Veto isolated leptons.
- HT distribution.
- Same, but enhance in \tilde{b} and $\tilde{\tau}$ by demanding two b-tagged jets.



STCx @ LHC14: Full-hadronic search with btag

- Select events with large amount of E_T in central, high E jets (HT) \Rightarrow hard interaction.
- Further select large *missing* E_T (MHT) \Rightarrow Partly invisible decay of heavy particle to a (much) lighter one.
- Avoid “signals” from bad measurements: Angle between any jet and the MHT direction large. Veto isolated leptons.
- HT distribution.
- Same, but enhance in \tilde{b} and \tilde{t} by demanding two b-tagged jets.




STCx @ LHC14: Search for $\tilde{b} \rightarrow b\tilde{\chi}_1^0$

- Select events with **exactly two** central, high E_T , b-tagged jets.
- No other jet with significant E_T allowed, veto leptons.
- Require high missing E_T and high transverse mass.
- The “contransverse mass” is an observable that is predicted to have an edge at $(M_{\tilde{b}}^2 - M_{\tilde{\chi}_1^0}^2)/M_{\tilde{b}} \approx M_{\tilde{\tau}}$.


STCx @ LHC14: Search for $\tilde{b} \rightarrow b\tilde{\chi}_1^0$

- Select events with **exactly two** central, high E_T , b-tagged jets.
- No other jet with significant E_T allowed, veto leptons.
- Require high missing E_T and high transverse mass.
- The “contransverse mass” is an observable that is predicted to have an edge at $(M_{\tilde{b}}^2 - M_{\tilde{\chi}_0^0}^2)/M_{\tilde{b}} \approx M_{\tilde{b}}$.



STCx @ LHC14: Search for $\tilde{b} \rightarrow b\tilde{\chi}_1^0$

- Select events with **exactly two** central, high E_T , b-tagged jets.
 - No other jet with significant E_T allowed, veto leptons.
 - Require high missing E_T and high m_{CT} .
 - The analysis is optimized to have a high discovery reach (up to $M_{\tilde{b}} \sim 1$ TeV).
 - Quite pure sample.
 - Limits on $M_{\tilde{b}}$ (or measurement if $M_{\tilde{\chi}_1^0}$ known).
- $$(M_{\tilde{b}}^2 - M_{\tilde{\chi}_1^0}^2) / M_{\tilde{b}} \approx M_{\tilde{b}}$$




STCx @ LHC14: Search for \tilde{t} in single-lepton channel

- Low BR $\tilde{t} \rightarrow t\tilde{\chi}_1^0$ in the model. \Rightarrow Look for $\tilde{t} \rightarrow t\tilde{\chi}_{2,3,4}^0$, with bosinos going to $W \rightarrow \ell\nu$ on one side, to $W \rightarrow qq'$ on the other.
- One (and only one) **isolated lepton** centrally + “Nothing”
- **Many central jets** (≥ 5), with one or two b-tagged.
- Transverse mass m_T of missing momentum and lepton > 260 GeV.
- **Select events where the angle between leading jets and missing momentum is large**


STCx @ LHC14: Search for \tilde{t} in single-lepton channel

- Low BR $\tilde{t} \rightarrow t\tilde{\chi}_1^0$ in the model. \Rightarrow Look for $\tilde{t} \rightarrow t\tilde{\chi}_{2,3,4}^0$, with bosinos going to $W \rightarrow \ell\nu$ on one side, to $W \rightarrow qq'$ on the other.
- One (and only one) **isolated lepton** centrally + “Nothing”
- **Many central jets** (≥ 5), with one or two b-tagged.
- Transverse mass m_T of missing momentum and lepton > 260 GeV.
- Select events where the angle between leading jets and missing momentum is large



STCx @ LHC14: Search for \tilde{t} in single-lepton channel

- Low BR $\tilde{t} \rightarrow t\tilde{\chi}_1^0$ in the model. \Rightarrow Look for $\tilde{t} \rightarrow t\tilde{\chi}_{2,3,4}^0$, with bosinos going to $W \rightarrow \ell\nu$ on one side, to $W \rightarrow qq'$ on the other.
- One (and only one) **isolated lepton** central
- **Many** two b-
- Transverse momentum and lepton > 200 GeV.
- Select events where the angle between leading jets and missing momentum is large



STCx @ LHC14: Search for bosinos

- *Dominating* SUSY channel at LHC is $\tilde{\chi}_2^0 \tilde{\chi}_1^\pm$ production: > 1 pb.
- \Rightarrow search for direct *un-coloured* bosino production (“whiteinos”) !
- Golden channel: \geq three leptons.
- Plethora of channels, and of backgrounds (WZ, $t\bar{t}$, Drell-Yan)
- \Rightarrow 45 different searches binned in
 - $m_{\tilde{g}}, m_{\tilde{t}},$ and E_T^{miss}
- No single bin highly significant, but together, $> 5\sigma$ @ $< 200 \text{ fb}^{-1}$.
- Signal is a mainly $\tilde{\chi}_2^0 \tilde{\chi}_1^\pm$, followed by $\tilde{\chi}_2^\pm \chi_x$.

Ch	Total SM \pm unc.	STCs	$Z_{\tilde{g}}$	$\sqrt{s_{\tilde{g}}}$	$\sqrt{s_{\tilde{t}}}$	Other EWK	No EWK
$m_{\tilde{g}} < 75\text{GeV}$							
1	10900 \pm 3300	88	<-0.5	76	7	3	2
2	5900 \pm 960	130	<-0.5	110	10	10	0
3	1390 \pm 340	140	<-0.5	110	20	10	0
4	1290 \pm 210	26	<-0.5	16	7	2	2
5	348 \pm 121	19	<-0.5	10	6	2	1
6	45.1 \pm 32.4	8.5	<-0.5	2.3	4.5	0.5	1.2
7	469 \pm 125	26	<-0.5	9	12	4	2
8	29.6 \pm 6.8	9.6	0.9	2.5	5.7	0.6	0.8
9	1.26 \pm 0.41	1.0	<-0.5	0.2	0.6	0.0	0.3
10	21.4 \pm 3.2	6.6	1.0	1.2	4.6	0.6	0.1
11	4.48 \pm 1.72	2.7	0.6	0.5	1.9	0.3	0.0
12	0.0262 \pm 0.0095	0.3	<-0.5	0.0	0.15	0.15	0.0
13	1.06 \pm 0.19	1.1	0.6	0.0	0.4	0.6	0.0
14	0.89 \pm 0.263	0.3	<-0.5	0.0	0.3	0.0	0.0
15	0.0137 \pm 0.0048	0	<-0.5	0	0	0	0
$75\text{GeV} < m_{\tilde{g}} < 105\text{GeV}$							
1	111000 \pm 16000	97	<-0.5	79	11	5	3
2	45900 \pm 7700	170	<-0.5	140	20	10	10
3	7490 \pm 1390	210	<-0.5	140	50	10	10
4	4640 \pm 490	26	<-0.5	12	10	2	1
5	994 \pm 278	31	<-0.5	13	13	4	1
6	55.4 \pm 40.3	16	<-0.5	2	10	2	2
7	444 \pm 75	30	<-0.5	9	17	2	1
8	26.2 \pm 5.0	26	2.9	6	16	4	2
9	1.91 \pm 0.47	2.0	1.0	0.2	1.5	0.0	0.4
10	16.4 \pm 1.4	6.7	1.4	0.9	4.7	0.8	0.4
11	5.01 \pm 0.93	3.0	1.0	0.6	1.6	0.0	0.8
12	0.058 \pm 0.0176	0.12	<-0.5	0.0	0.0	0.0	0.12
13	1.77 \pm 0.2	0.15	<-0.5	0.0	0.15	0.0	0.0
14	2.32 \pm 0.35	0.31	<-0.5	0.0	0.31	0.0	0.0
15	0.113 \pm 0.034	0	<-0.5	0	0	0	0
$m_{\tilde{g}} > 105\text{GeV}$							
1	2380 \pm 320	22	<-0.5	11	7	3	1
2	1720 \pm 240	34	<-0.5	18	10	4	2
3	614 \pm 157	61	<-0.5	21	24	10	6
4	217 \pm 47	10	<-0.5	3	6	1	1
5	57.6 \pm 11.5	11	0.7	2	6	3	1
6	13.7 \pm 3.5	8.0	1.2	1.4	3.9	1.8	0.8
7	70.2 \pm 6.0	15	1.4	1	13	2	0
8	12.6 \pm 2.7	15	2.6	2	10	2	1
9	0.812 \pm 0.174	1.4	1.0	0.2	0.7	0.2	0.4
10	11.7 \pm 1.4	3.5	0.8	0.2	2.3	1.1	0.0
11	1.94 \pm 0.41	2.8	1.4	0.3	1.5	0.3	0.7
12	0.0391 \pm 0.0114	0	<-0.5	0	0	0	0
13	1.36 \pm 0.14	0	<-0.5	0	0	0	0
14	0.674 \pm 0.176	0.78	<-0.5	0.0	0.15	0.15	0.47
15	0.0137 \pm 0.0055	0.24	<-0.5	0.0	0.0	0.0	0.24

STCx @ LHC14: Search for bosinos

- *Dominating* SUSY channel at LHC is

$\tilde{\chi}_2^0 \tilde{\chi}_1^\pm$ production: > 1 pb.

- \Rightarrow search for direct *un-coloured* bosino production (“whiteinos”) !
- Golden channel: \geq three leptons.
- Plethora of channels, and of backgrounds (WZ, $t\bar{t}$, Drell-Yan)
- \Rightarrow 45 different searches binned in
 - $m_{\ell\ell}$, m_T , and E_T^{miss}

- No single bin highly significant, but together, $> 5\sigma$ @ < 200 fb $^{-1}$.

- Signal is a mainly $\tilde{\chi}_2^0 \tilde{\chi}_1^\pm$, followed by $\tilde{\chi}_2^\pm \chi_x$.

Ch	Total SM \pm unc.	STCs	Z_β	$\sqrt{s_{\tilde{\chi}_2^0 \tilde{\chi}_1^\pm}}$	$\sqrt{s_{\tilde{\chi}_2^\pm \chi_x}}$	Other EWK	No EWK
$m_{\ell\ell} < 75\text{GeV}$							
1	10900 \pm 3300	88	<-0.5	76	7	3	2
2	5900 \pm 960	130	<-0.5	110	10	10	0
3	1390 \pm 340	140	<-0.5	110	20	10	0
4	1290 \pm 210	26	<-0.5	16	7	2	2
5	348 \pm 121	19	<-0.5	10	6	2	1
6	45.1 \pm 32.4	8.5	<-0.5	2.3	4.5	0.5	1.2
7	469 \pm 125	26	<-0.5	9	12	4	2
8	29.6 \pm 6.8	9.6	0.9	2.5	5.7	0.6	0.8
9	1.26 \pm 0.41	1.0	<-0.5	0.2	0.6	0.0	0.3
10	21.4 \pm 3.2	6.6	1.0	1.2	4.6	0.6	0.1
11	4.48 \pm 1.72	2.7	0.6	0.5	1.9	0.3	0.0
12	0.0262 \pm 0.0095	0.3	<-0.5	0.0	0.15	0.15	0.0
13	1.06 \pm 0.19	1.1	0.6	0.0	0.4	0.6	0.0
14	0.89 \pm 0.263	0.3	<-0.5	0.0	0.3	0.0	0.0
15	0.0137 \pm 0.0048	0	<-0.5	0	0	0	0
$75\text{GeV} < m_{\ell\ell} < 105\text{GeV}$							
1	111000 \pm 16000	97	<-0.5	79	11	5	3
2	45900 \pm 7700	170	<-0.5	140	20	10	10
3	7490 \pm 1390	210	<-0.5	140	50	10	10
4	4640 \pm 490	26	<-0.5	12	10	2	1
5	994 \pm 278	31	<-0.5	13	13	4	1
6	55.4 \pm 40.3	16	<-0.5	2	10	2	2
7	444 \pm 75	30	<-0.5	9	17	2	1
8	26.2 \pm 5.0	26	2.9	6	16	4	2
9	1.91 \pm 0.47	2.0	1.0	0.2	1.5	0.0	0.4
10	16.4 \pm 1.4	6.7	1.4	0.9	0.8	0.8	0.4
11	5.01 \pm 0.93	3.0	1.0	0.6	1.6	0.0	0.8
12	0.058 \pm 0.0176	0.12	<-0.5	0.0	0.0	0.0	0.12
13	1.77 \pm 0.2	0.15	<-0.5	0.0	0.15	0.0	0.0
14	2.32 \pm 0.35	0.31	<-0.5	0.0	0.31	0.0	0.0
15	0.113 \pm 0.034	0	<-0.5	0	0	0	0
$m_{\ell\ell} > 105\text{GeV}$							
1	2380 \pm 320	22	<-0.5	11	7	3	1
2	1720 \pm 240	34	<-0.5	18	10	4	2
3	614 \pm 157	61	<-0.5	21	24	10	6
4	217 \pm 47	10	<-0.5	3	6	1	1
5	57.6 \pm 11.5	11	0.7	2	6	3	1
6	13.7 \pm 3.5	8.0	1.2	1.4	3.9	1.8	0.8
7	70.2 \pm 6.0	15	1.4	1	13	2	0
8	12.6 \pm 2.7	15	2.6	2	10	2	1
9	0.812 \pm 0.174	1.4	1.0	0.2	0.7	0.2	0.4
10	11.7 \pm 1.4	3.5	0.8	0.2	2.3	1.1	0.0
11	1.94 \pm 0.41	2.8	1.4	0.3	1.5	0.3	0.7
12	0.0391 \pm 0.0114	0	<-0.5	0	0	0	0
13	1.36 \pm 0.14	0	<-0.5	0	0	0	0
14	0.674 \pm 0.176	0.78	<-0.5	0.0	0.15	0.15	0.47
15	0.0137 \pm 0.0055	0.24	<-0.5	0.0	0.0	0.0	0.24

STCx @ LHC14: Search for bosinos

- *Dominating* SUSY channel at LHC is $\tilde{\chi}_2^0 \tilde{\chi}_1^\pm$ production: > 1 pb.
- \Rightarrow search for direct *un-coloured* bosino production (“whiteinos”) !
- Golden channel: \geq **three leptons**.
- **Plethora** of channels, and of backgrounds (WZ, $t\bar{t}$, Drell-Yan)
- \Rightarrow 45 different searches binned in
 - $m_{\ell\ell}$, m_T , and E_T^{miss}
- No single bin highly significant, but together, $> 5\sigma$ @ < 200 fb $^{-1}$.
- Signal is a mainly $\tilde{\chi}_2^0 \tilde{\chi}_1^\pm$, followed by $\tilde{\chi}_2^\pm \chi_x$.

Ch	Total SM \pm unc.	STCs	Z_μ	$\sqrt{s_{\text{eff}}}$	$\sqrt{s_{\text{eff}}}$	Other EWK	No EWK
$m_{\ell\ell} < 75\text{GeV}$							
1	10900 \pm 3300	88	-0.5	76	7	3	2
2	5900 \pm 960	130	-0.5	110	10	10	0
3	1390 \pm 340	140	-0.5	110	20	10	0
4	1290 \pm 210	26	-0.5	16	7	2	2
5	348 \pm 121	19	-0.5	10	6	2	1
6	45.1 \pm 32.4	8.5	-0.5	2.3	4.5	0.5	1.2
7	469 \pm 125	26	-0.5	9	12	4	2
8	29.6 \pm 6.8	9.6	0.9	2.5	5.7	0.6	0.8
9	1.26 \pm 0.41	1.0	-0.5	0.2	0.6	0.0	0.3
10	21.4 \pm 3.2	6.6	1.0	1.2	4.6	0.6	0.1
11	4.48 \pm 1.72	2.7	0.6	0.5	1.9	0.3	0.0
12	0.0262 \pm 0.0095	0.3	-0.5	0.0	0.15	0.15	0.0
13	1.06 \pm 0.19	1.1	0.6	0.0	0.4	0.6	0.0
14	0.89 \pm 0.263	0.3	-0.5	0.0	0.3	0.0	0.0
15	0.0137 \pm 0.0048	0	-0.5	0	0	0	0
$75\text{GeV} < m_{\ell\ell} < 105\text{GeV}$							
1	111000 \pm 16000	97	-0.5	79	11	5	3
2	45900 \pm 7700	170	-0.5	140	20	10	10
3	7490 \pm 1390	210	-0.5	140	50	10	10
4	4640 \pm 490	26	-0.5	12	10	2	1
5	994 \pm 278	31	-0.5	13	13	4	1
6	55.4 \pm 40.3	16	-0.5	2	10	2	2
7	444 \pm 75	30	-0.5	9	17	2	1
8	26.2 \pm 5.0	26	2.9	6	16	4	2
9	1.91 \pm 0.47	2.0	1.0	0.2	1.5	0.0	0.4
10	16.4 \pm 1.4	6.7	1.4	0.9	4.7	0.8	0.4
11	5.01 \pm 0.93	3.0	1.0	0.6	1.6	0.0	0.8
12	0.058 \pm 0.0176	0.12	-0.5	0.0	0.0	0.0	0.12
13	1.77 \pm 0.2	0.15	-0.5	0.0	0.15	0.0	0.0
14	2.32 \pm 0.35	0.31	-0.5	0.0	0.31	0.0	0.0
15	0.113 \pm 0.034	0	-0.5	0	0	0	0
$m_{\ell\ell} > 105\text{GeV}$							
1	2380 \pm 320	22	-0.5	11	7	3	1
2	1720 \pm 240	34	-0.5	18	10	4	2
3	614 \pm 157	61	-0.5	21	24	10	6
4	217 \pm 47	10	-0.5	3	6	1	1
5	57.6 \pm 11.5	11	0.7	2	6	3	1
6	13.7 \pm 3.5	8.0	1.2	1.4	3.9	1.8	0.8
7	70.2 \pm 6.0	15	1.4	1	13	2	0
8	12.6 \pm 2.7	15	2.6	2	10	2	1
9	0.812 \pm 0.174	1.4	1.0	0.2	0.7	0.2	0.4
10	11.7 \pm 1.4	3.5	0.8	0.2	2.3	1.1	0.0
11	1.94 \pm 0.41	2.8	1.4	0.3	1.5	0.3	0.7
12	0.0391 \pm 0.0114	0	-0.5	0	0	0	0
13	1.36 \pm 0.14	0	-0.5	0	0	0	0
14	0.674 \pm 0.176	0.78	-0.5	0.0	0.15	0.15	0.47
15	0.0137 \pm 0.0055	0.24	-0.5	0.0	0.0	0.0	0.24

STCx @ LHC14: Search for bosinos

- *Dominating* SUSY channel at LHC is

$\tilde{\chi}_2^0 \tilde{\chi}_1^\pm$ production: > 1 pb.

- \Rightarrow search for direct *un-coloured* bosino production (“whiteinos”) !
- Golden channel: $>$ three leptons

- Plethora of background
 - Good discovery reach, directly related to ILC reach.

- $\Rightarrow 45$
 - ILC-LHC synergy: precise masses from ILC into LHC long decay-chains \Rightarrow determine

- No sir

properties of heavier bosinos .
together, $> 5\sigma$ @ $\sqrt{s} < 200$ GeV .

- Signal is a mainly $\tilde{\chi}_2^0 \tilde{\chi}_1^\pm$, followed by $\tilde{\chi}_2^\pm \chi_x$.

Ch	Total SM \pm unc.	STCs	Z_μ	\sqrt{s}_{11}	\sqrt{s}_{12}	Other EWK	No EWK
$m_{\tilde{t}^{\pm}} < 75 \text{ GeV}$							
1	10900 \pm 3300	88	<-0.5	76	7	3	2
2	5900 \pm 960	130	<-0.5	110	10	10	0
3	1390 \pm 340	140	<-0.5	110	20	10	10
4	1290 \pm 210	26	<-0.5	16	7	2	2
5	348 \pm 121	19	<-0.5	10	6	2	1
6	45.1 \pm 32.4	8.5	<-0.5	2.3	4.5	0.5	1.2
7	469 \pm 125	26	<-0.5	9	12	4	2
8	29.6 \pm 6.8	9.6	0.9	2.5	5.7	0.6	0.8
9	1.26 \pm 0.41	1.0	<-0.5	0.2	0.6	0.0	0.3
10	21.4 \pm 3.2	6.6	1.0	1.2	4.6	0.6	0.1
11	4.48 \pm 1.72	2.7	0.6	0.5	1.9	0.3	0.0
12	0.0262 \pm 0.0095	0.3	<-0.5	0.0	0.15	0.15	0.0
13	1.06 \pm 0.19	1.1	0.6	0.0	0.4	0.6	0.0
14	0.89 \pm 0.263	0.3	<-0.5	0.0	0.3	0.0	0.0
15	0.0137 \pm 0.0048	0	<-0.5	0	0	0	0
$75 \text{ GeV} < m_{\tilde{t}^{\pm}} < 105 \text{ GeV}$							
						5	3
						10	10
						10	10
						2	1
						4	1
						2	2
						2	1
						4	2
						0.0	0.4
						0.8	0.4
						0.0	0.8
						0.0	0.12
						0.0	0.0
						0.0	0.0
						0	0
						3	1
						4	2
						10	6
						1	1
						3	1
						1.8	0.8
						2	0
8	12.6 \pm 2.7	15	2.6	2	10	2	1
9	0.912 \pm 0.174	1.4	1.0	0.2	0.7	0.2	0.4
10	11.7 \pm 1.4	3.5	0.8	0.2	2.3	1.1	0.0
11	1.94 \pm 0.41	2.8	1.4	0.3	1.5	0.3	0.7
12	0.0391 \pm 0.0114	0	<-0.5	0	0	0	0
13	1.36 \pm 0.14	0	<-0.5	0	0	0	0
14	0.674 \pm 0.176	0.78	<-0.5	0.0	0.15	0.15	0.47
15	0.0137 \pm 0.0055	0.24	<-0.5	0.0	0.0	0.0	0.24

Summary of STCx@LHC

If STCx is realised in nature

- LHC **will** discover something non-SM:
 - Central $t\bar{t} \rightarrow \text{hadrons}$ events with lots of missing ET and one isolated lepton.
 - Central $b\bar{b}$ events with lots of missing ET, and a tell-tale contraverse mass distribution.
 - A diffuse, highly significant, excess of three-lepton, transversly unbalanced events.
 - Insignificant (3σ) excess in central, fully hadronic events with high missing ET.
 - Largely systematics limited.
- So, there is BSM, but:

Summary of STCx@LHC

If STCx is realised in nature

- LHC **will** discover something non-SM:
 - Central $t\bar{t} \rightarrow \text{hadrons}$ events with lots of missing ET and one **isolated lepton**.
 - Central $b\bar{b}$ events with lots of missing ET, and a tell-tale **contraverse mass** distribution.
 - A diffuse, highly significant, excess of **three-lepton**, transversely unbalanced events.
 - Insignificant (3σ) excess in central, fully hadronic events with high missing ET.
 - Largely **systematics limited**.
- So, there is BSM, but:

Summary of STCx@LHC

If STCx is realised in nature

- LHC **will** discover something non-SM:
 - Central $t\bar{t} \rightarrow \text{hadrons}$ events with lots of missing ET and one **isolated lepton**.
 - Central $b\bar{b}$ events with lots of missing ET, and a tell-tale **contraverse mass** distribution.
 - A diffuse, highly significant, excess of **three-lepton**, transversely unbalanced events.
 - Insignificant (3σ) excess in central, fully hadronic events with high missing ET.
 - Largely **systematics limited**.
- So, there **is** BSM, but:


Summary of STCx@LHC

If STCx is realised in nature

- LHC **will** discover something non-SM:
 - Central $t\bar{t} \rightarrow \text{hadrons}$ events with lots of missing ET and one **isolated lepton**.
 - Central $b\bar{b}$ events with lots of missing ET and a tell-tale **contraverse mass** distribution.
 - A diffuse, high energy **lepton**, transversely unbalanced e .
 - Insignificant (3σ) excess in central, fully hadronic events with high missing ET.
 - Largely **systematics limited**.
- So, there **is** BSM, but:

What is it ?
Is it SUSY?

The ILC



- A linear e^+e^- collider.
- E_{CMS} tunable between **250 and 500 GeV**, upgradable to 1 TeV.
- Total length 31 km
- $\int \mathcal{L} \sim 250 \text{ fb}^{-1}/\text{year}$
- **Polarisation** e^- : 80% (e^+ : $\geq 30\%$)
- 2 experiments, but (possibly) only one interaction region.
- Concurrent running with the LHC

The ILC is not LHC


- Lepton-collider: Initial state is known.
- Production is EW \Rightarrow
 - Small theoretical uncertainties.
 - No "underlying event".
 - Low cross sections wrt. LHC, also for background.
 - Trigger-less operation.
- Extremely small beam-spot: $5 \text{ nm} \times 100 \text{ nm} \times 150 \mu\text{m}$.
- Low background \Rightarrow detectors can be:
 - Thin: few % X_0 in front of calorimeters
 - Very close to IP: first layer of VXD at 1.5 cm.
 - Close to 4 π : holes for beam-pipe only few cm = 0.2 msr un-covered
= Area of Schleswig-Holstein relative to earth.


The ILC is not LHC


- Lepton-collider: Initial state is **known**.
- Production is **EW** \Rightarrow
 - Small **theoretical uncertainties**.
 - No “underlying event”.
 - **Low cross-sections** wrt. LHC, also for background.
 - **Trigger-less** operation.
- **Extremely small beam-spot**: $5 \text{ nm} \times 100 \text{ nm} \times 150 \mu\text{m}$.
- **Low background** \Rightarrow **detectors can be**:
 - **Thin**: few % X_0 in front of calorimeters
 - **Very close to IP**: first layer of VXD at 1.5 cm.
 - **Close to 4 π** : holes for beam-pipe only few cm = 0.2 msr un-covered
= Area of Schleswig-Holstein relative to earth.


The ILC is not LHC

- Lepton-collider: Initial state is **known**.
- Production is **EW** \Rightarrow
 - Small **theoretical uncertainties**.
 - No “underlying event”.
 - **Low cross-sections** wrt. LHC, also for background.
 - **Trigger-less** operation.
- Extremely **small beam-spot**: $5 \text{ nm} \times 100 \text{ nm} \times 150 \mu\text{m}$.
- **Low background** \Rightarrow detectors can be:
 - **Thin** : few % X_0 in front of calorimeters
 - **Very close to IP**: first layer of VXD at 1.5 cm.
 - **Close to 4π** : holes for beam-pipe only few cm = 0.2 msr un-covered = Area of Schleswig-Holstein relative to earth.

Doing physics at ILC: the $Zh \rightarrow \mu\mu\tau^+\tau^-$ example

Doing physics at ILC: the $Zh \rightarrow \mu\mu\tau^+\tau^-$ example

Doing physics at ILC: the $Zh \rightarrow \mu\mu\tau^+\tau^-$ example


Doing physics at ILC: the $Zh \rightarrow \mu\mu\tau^+\tau^-$ example

Doing physics at ILC: the $Zh \rightarrow \mu\mu\tau^+\tau^-$ example

At the ILC:


When You're looking at the event-display, You're looking at the Feynman-diagram ! \Rightarrow

- $\eta, E_T, M_T, \dots \Rightarrow \theta, (E, \vec{p}), M \dots$
- Can measure full (polarised) differential quantities



The STCx benchmark @ ILC

Zoomed STCx mass-spectrum



The STCx benchmark @ ILC

Zoomed STCx mass-spectrum

At the ILC@500 GeV:

Signal:

- Typically : a few leptons + LSP:s \Rightarrow
 - Low multiplicity events.
 - Central, much missing energy.
- Cross-sections up to 1 pb+.
- Often cascades over $\tilde{\tau}_1$.
- $\Delta(M) \sim 10$ GeV $\Rightarrow E_\tau \in [2.3, 45.5]$ GeV.

Background:

- Real missing energy = ZZ, WW \rightarrow $ll\nu\nu$
- Fake missing energy = $\gamma\gamma$ processes, ISR, single IVB.

The STCx benchmark @ ILC

Zoomed STCx mass-spectrum

At the ILC@500 GeV:

Signal:

- Typically : a few leptons + LSP:s \Rightarrow
 - Low multiplicity events.
 - Central, much missing energy.
- Cross-sections up to 1 pb+.
- Often cascades over $\tilde{\tau}_1$.
- $\Delta(M) \sim 10 \text{ GeV} \Rightarrow E_\tau \in [2.3, 45.5] \text{ GeV}$.

Background:

- Real missing energy = $ZZ, WW \rightarrow ll\nu\nu$
- Fake missing energy = $\gamma\gamma$ processes, ISR, single IVB.

The STCx benchmark @ ILC

Zoomed STCx mass-spectrum

At the ILC@500 GeV:

Signal:

- Typically : a few leptons + LSP:s \Rightarrow
 - Low multiplicity events.
 - Central, much missing energy.
- Cross-sections up to 1 pb+.
- Often cascades over $\tilde{\tau}_1$.
- $\Delta(M) \sim 10 \text{ GeV} \Rightarrow E_\tau \in [2.3, 45.5] \text{ GeV}$.


Background:

- Real missing energy = $ZZ, WW \rightarrow ll\nu\nu$
- Fake missing energy = $\gamma\gamma$ processes, ISR, single IVB.

STC4 global

After a few very general cuts:

- Missing energy > 100
- Less than 10 charged tracks
- $|\cos \theta_{P_{tot}}| < 0.95$
- Exactly two τ -jets
- Visible mass < 300 GeV
- θ_{acop} between 0.15 and 3.1





Magenta: $\gamma\gamma$, Blue: 3f,
Red: Rest of SM, Green: SUSY.

STC4 sleptons @ 500 GeV: \tilde{e} , $\tilde{\mu}$

- **Selections** for $\tilde{\mu}$ and \tilde{e} :
 - Correct charge.
 - P_T wrt. beam and one ℓ wrt the other.
 - Tag and probe, ie. accept one jet if the other is “in the box”.
- **Further selections** for R:
 - Cuts on polar angle and angle between leptons.
- E_{jet} , beam-pol 80%, -30%...
- Determine masses from edge-positions.

STC4 sleptons @ 500 GeV: \tilde{e} , $\tilde{\mu}$


- **Selections** for $\tilde{\mu}$ and \tilde{e} :
 - Correct charge.
 - P_T wrt. beam and one ℓ wrt the other.
 - Tag and probe, ie. accept one jet if the other is “in the box”.
- **Further selections** for R:
 - Cuts on polar angle and angle between leptons.
- E_{jet} , beam-pol 80%, -30%...
- Determine masses from edge-positions.



STC4 sleptons @ 500 GeV: \tilde{e} , $\tilde{\mu}$

- **Selections** for $\tilde{\mu}$ and \tilde{e} :

Results from edges ($E_{CMS}=500$, 500 fb^{-1} @ $[+0.8, -0.3]$)



selectrons:

$$M_{\tilde{e}_R} = 126.20 \pm 0.21 \text{ GeV}/c^2$$

$$M_{\tilde{\chi}_1^0} = 95.47 \pm 0.16 \text{ GeV}/c^2$$

smuons:


$$M_{\tilde{\mu}_R} = 126.01 \pm 0.51 \text{ GeV}/c^2$$

$$M_{\tilde{\chi}_1^0} = 95.47 \pm 0.38 \text{ GeV}/c^2$$

combined:

$$\sigma M_{\tilde{\chi}_1^0} = 147 \text{ MeV}/c^2$$

$$\sigma M_{\tilde{\ell}_R} = 194 \text{ MeV}/c^2$$



\tilde{e}_R and $\tilde{\mu}_R$ threshold scans

From these spectra, we can estimate $M_{\tilde{e}_R}$, and $M_{\tilde{\chi}_1^0}$ to < 0.2 GeV, and $M_{\tilde{\mu}_R}$ to < 0.5 GeV.


$\tilde{\epsilon}_R$ and $\tilde{\mu}_R$ threshold scans

From these spectra, we can estimate $M_{\tilde{\epsilon}_R}$, and $M_{\tilde{\chi}_1^0}$ to < 0.2 GeV, and $M_{\tilde{\mu}_R}$ to < 0.5 GeV.

So: Next step is $M_{\tilde{\ell}}$ from threshold:

- 10 points, $10 \text{ fb}^{-1}/\text{point}$.
- Fit either P- or S-wave excitation curves (ie. scalar or fermion pairs produced)

- Error on $M_{\tilde{\mu}_R}$ and $M_{\tilde{\epsilon}_R} = 200$ MeV.
- Fermion hypothesis excluded with fit-probabilities $< 10^{-5}$




$\tilde{\epsilon}_R$ and $\tilde{\mu}_R$ threshold scans

From these spectra, we can estimate $M_{\tilde{\epsilon}_R}$, and $M_{\tilde{\chi}_1^0}$ to < 0.2 GeV, and $M_{\tilde{\mu}_R}$ to < 0.5 GeV.

So: Next step is $M_{\tilde{\ell}}$ from threshold:

- 10 points, $10 \text{ fb}^{-1}/\text{point}$.
- Fit either P- or S-wave excitation curves (ie. scalar or fermion pairs produced)

- Error on $M_{\tilde{\mu}_R}$ and $M_{\tilde{\epsilon}_R} = 200$ MeV.
- Fermion hypothesis excluded with fit-probabilities $< 10^{-5}$




$\tilde{\epsilon}_R$ and $\tilde{\mu}_R$ threshold scans

From these spectra, we can estimate $M_{\tilde{\epsilon}_R}$, and $M_{\tilde{\chi}_1^0}$ to < 0.2 GeV, and $M_{\tilde{\mu}_R}$ to < 0.5 GeV.

So: Next step is $M_{\tilde{\ell}}$ from threshold:


- 10 points, $10 \text{ fb}^{-1}/\text{point}$.
 - Fit either P- or S-wave excitation curves (ie. scalar or fermion pairs produced)
- Error on $M_{\tilde{\mu}_R}$ and $M_{\tilde{\epsilon}_R} = 200$ MeV.
 - **Fermion hypothesis excluded** with fit-probabilities $< 10^{-5}$



STC4 sleptons @ 500 GeV: $\tilde{\tau}_1$

Selections for $\tilde{\tau}_1$:


- Correct charge.
- P_T wrt. beam and one τ wrt the other.
- $M_{jet} < M_\tau$
- $E_{vis} < 120 \text{ GeV}, M_{vis} \in [20, 87] \text{ GeV}$.
- At least one τ -jet should be hadronic.
- Anti- $\gamma\gamma$ likelihood.
- Fit background
- Fit upper end-point of data – background fit



STC4 sleptons @ 500 GeV: $\tilde{\tau}_1$

Selections for $\tilde{\tau}_1$:

- Correct **charge**.
- P_T wrt. beam and one τ wrt the other.
- $M_{jet} < M_\tau$
- $E_{vis} < 120$ GeV, $M_{vis} \in [20, 87]$ GeV.
- At least one τ -jet should be **hadronic**.
- **Anti- $\gamma\gamma$ likelihood**.
- Fit background
- Fit upper end-point of data – background fit



STC4 sleptons @ 500 GeV: $\tilde{\tau}_1$

Selections for $\tilde{\tau}_1$:

- Correct charge

Results for $\tilde{\tau}_1$

$$E_{max, \tilde{\tau}_1} = 44.49^{+0.11}_{-0.09} \text{ GeV}$$

Translates to an error on the mass of $0.27 \text{ GeV}/c^2$, dominated by the error from $M_{\tilde{\chi}_1^0}$.

- At least one τ -jet should be

Results for $\tilde{\tau}_2$

$$E_{max, \tilde{\tau}_2} = 145.4^{+5.9}_{-4.4} \text{ GeV}$$

Translates to an error on the mass of $5 \text{ GeV}/c^2$, dominated by the error from the end-point.

background fit



Reminder: SPS1a' results (Phys.Rev.D82:055016,2010)

The previous $\tilde{\tau}$ study in the very similar model SPS1a' gave:

Results for $\tilde{\tau}_1$

$$M_{\tilde{\tau}_1} = 107.73^{+0.03}_{-0.05} \text{ GeV}/c^2 \oplus 1.3\Delta(M_{\tilde{\chi}_1^0})$$

The error from $M_{\tilde{\chi}_1^0}$ largely dominates

Cross-section for $\tilde{\tau}_1$ and $\tilde{\tau}_2$

$$\Delta(N_{\text{signal}})/N_{\text{signal}} = 3.1 \%$$

$$\Delta(N_{\text{signal}})/N_{\text{signal}} = 4.2 \%$$

\Rightarrow Can determine if **sleptons couple like leptons**.

Results for $\tilde{\tau}_2$

$$M_{\tilde{\tau}_2} = 183^{+11}_{-5} \text{ GeV}/c^2 \oplus 18\Delta(M_{\tilde{\chi}_1^0})$$

The error from the endpoint largely dominates

Also: τ polarisation in $\tilde{\tau}_1$ decays

$$\Delta(\mathcal{P}_\tau)/\mathcal{P}_\tau = 9 \%$$

Reminder: SPS1a' results (Phys.Rev.D82:055016,2010)

The previous $\tilde{\tau}$ study in the very similar model SPS1a' gave:

Results for $\tilde{\tau}_1$

$$M_{\tilde{\tau}_1} = 107.73_{-0.05}^{+0.03} \text{ GeV}/c^2 \oplus 1.3\Delta(M_{\tilde{\chi}_1^0})$$

The error from $M_{\tilde{\chi}_1^0}$ largely dominates

Cross-section for $\tilde{\tau}_1$ and $\tilde{\tau}_2$

$$\Delta(N_{\text{signal}})/N_{\text{signal}} = 3.1 \%$$

$$\Delta(N_{\text{signal}})/N_{\text{signal}} = 4.2 \%$$

\Rightarrow Can determine if **sleptons couple like leptons**.

Results for $\tilde{\tau}_2$

$$M_{\tilde{\tau}_2} = 183_{-5}^{+11} \text{ GeV}/c^2 \oplus 18\Delta(M_{\tilde{\chi}_1^0})$$

The error from the endpoint largely dominates

Also: τ polarisation in $\tilde{\tau}_1$ decays

$$\Delta(\mathcal{P}_\tau)/\mathcal{P}_\tau = 9 \%$$

For **implications on DM** from the numbers presented here, listen to Suvi-Leena Lehtinen's talk !

Summary

We have done a combined ILC-LHC study of a SUSY model that fulfils all current constraints. The model has good visibility at *both* machines.

- At LHC: study
 - Methods for discovery
 - Control of systematics essential.
 - Insight in coloured sector, in synergy with ILC.
- The observations will discover BSM physics, both in the coloured and EW sectors.
- But hard to say what physics it is.
- At ILC: Study best method to analyse spectra, eg
 - Optimal statistic for clean signals.
 - Specific reconstruction methods for e , μ , and τ .
- At ILC, will be able to un-ambiguously identify the new physics as SUSY, and to constrain the model.

Summary

We have done a combined ILC-LHC study of a SUSY model that fulfils all current constraints. The model has good visibility at *both* machines.

- At **LHC**: study
 - Methods for **discovery**
 - Control of **systematics** essential.
 - Insight in **coloured sector**, in synergy with ILC.
- The observations **will discover BSM physics**, both in the coloured and EW sectors.
- But **hard to say** what physics it is.
- At **ILC**: Study best method to analyse spectra, eg
 - Optimal statistic for clean signals.
 - Specific reconstruction methods for e , μ , and τ .
- At ILC, will be able to un-ambiguously **identify the new physics as SUSY**, and to constrain the model.

Summary

We have done a combined ILC-LHC study of a SUSY model that fulfils all current constraints. The model has good visibility at *both* machines.

- At **LHC**: study
 - Methods for **discovery**
 - Control of **systematics** essential.
 - Insight in **coloured sector**, in synergy with ILC.
- The observations **will discover BSM physics**, both in the coloured and EW sectors.
- But **hard to say** what physics it is.
- At **ILC**: Study best method to analyse spectra, eg
 - Optimal statistic for clean signals.
 - Specific reconstruction methods for e , μ , and τ .
- At ILC, will be able to un-ambiguously **identify the new physics as SUSY**, and to constrain the model.

Summary

We have done a combined ILC-LHC study of a SUSY model that fulfils all current constraints. The model has good visibility at *both* machines.

- At **LHC**: study
 - Meth **ILC and LHC**
 - Cont **Combining ILC precision with LHC reach**
 - Insig **the will allow to extend the model study to the coloured**
- The obser **the colored sector and to the heavier**
and EW **bosinos, ie. to the full spectrum.**
- But hard **to study the colored sector**
- At **ILC**: Study best method to analyse spectra, eg
 - Optimal statistic for clean signals.
 - Specific reconstruction methods for e , μ , and τ .
- At ILC, will be able to un-ambiguously **identify the new physics as SUSY**, and to constrain the model.

Thank You !

BACKUP

BACKUP SLIDES

Observables: Pair-production, two-body decay (less text)

- So, there are two SUSY parameters, and two independent observables in the spectrum.
- Any pair of observables can be chosen, edges, average, standard deviation, width, ...
- Which choice is the best depends on the situation.
- Just a bit of algebra to extract the two SUSY masses.
- Note that if $E_{beam} \gg M_X$, there is just one observable (low edge becomes 0, width becomes average/2), so one should not operate too far above threshold !
- Note that there are two decays in each event: two measurements per event.
- Also note that there are not enough measurements to make a constrained fit, even assuming that the two SUSY particles in the two decays are the same: $(2 \times 4 \text{ unknown components of 4-momentum } (=8)) - (\text{total E and p conservation } (=4) + 2$

Observables: Pair-production, two-body decay (less text)

- So, there are **two SUSY parameters**, and **two independent observables** in the spectrum.
- **Any pair** of observables can be chosen, edges, average, standard deviation, width, ...
- Which choice is the best **depends on the situation**.
- Just a bit of algebra to extract the two SUSY masses.
- Note that if $E_{beam} \gg M_X$, there is **just one observable** (low edge becomes 0, width becomes average/2), so one should not operate **too far above threshold** !
- Note that there are **two decays in each event**: two measurements per event.
- Also note that there are **not enough measurements** to make a constrained fit, even assuming that the two SUSY particles in the two decays are the same: $(2 \times 4 \text{ unknown components of 4-momentum } (=8)) - (\text{total E and p conservation } (=4) + 2$

Observables: Pair-production, two-body decay (less text)

- So, there are **two SUSY parameters**, and **two independent observables** in the spectrum.
- **Any pair** of observables can be chosen, edges, average, standard deviation, width, ...
- Which choice is the best **depends on the situation**.
- Just a bit of algebra to extract the two SUSY masses.
- Note that if $E_{beam} \gg M_X$, there is **just one observable** (low edge becomes 0, width becomes average/2), so one should not operate **too far above threshold** !
- Note that there are **two decays** in each event: two measurements per event.
- Also note that there are not enough measurements to make a constrained fit, even assuming that the two SUSY particles in the two decays are the same: $(2 \times 4 \text{ unknown components of 4-momentum } (=8)) - (\text{total E and p conservation } (=4) + 2$

Observables: Pair-production, two-body decay (less text)

- So, there are **two SUSY parameters**, and **two independent observables** in the spectrum.
- **Any pair** of observables can be chosen, edges, average, standard deviation, width, ...
- Which choice is the best **depends on the situation**.
- Just a bit of algebra to extract the two SUSY masses.
- Note that if $E_{beam} \gg M_X$, there is **just one observable** (low edge becomes 0, width becomes average/2), so one should not operate **too far above threshold** !
- Note that there are **two decays** in each event: two measurements per event.
- Also note that there are **not enough measurements to make a constrained fit**, even assuming that the two SUSY particles in the two decays are the same: $(2 \times 4 \text{ unknown components of 4-momentum (=8)}) - (\text{total E and p conservation (=4)}) + 2$

Observables: Pair-production, two-body decay

However:

- If the masses are known from **other measurements**, there are **enough constraints**.
- Then the events can be **completely reconstructed** ...
- ... and the **angular distributions** both in production and decay can be measured.
- From this the **spins** can be determined, which is **essential** to determine that what we are seeing is **SUSY**.

Furthermore:

- Looking at more **complicated decays**, such as cascade decays, there are enough constraints if some (but not all) masses are known.
- Allows to reconstruct eg. the **stopion mass** in $\tilde{\chi}_2^0 \rightarrow \tilde{\ell} \rightarrow \ell \tilde{\chi}_1^0$ if chargino and LSP masses are known.
- Order-of-magnitude better mass resolution.

Observables: Pair-production, two-body decay

However:

- If the masses are known from **other measurements**, there are **enough constraints**.
- Then the events can be **completely reconstructed** ...
- ... and the **angular distributions** both in production and decay can be measured.
- From this the **spins** can be determined, which is **essential** to determine that what we are seeing is **SUSY**.



Furthermore:

- Looking at more **complicated decays**, such as cascade decays, there are enough constraints if some (but not all) masses are known.
- Allows to reconstruct eg. **the slepton mass** in $\tilde{\chi}_2^0 \rightarrow \tilde{\ell}\ell \rightarrow \ell\ell\tilde{\chi}_1^0$ if chargino and LSP masses are known.
- Order-of-magnitude better mass resolution.

Observables: Pair-production, two-body decay

However:

- If the masses are known from **other measurements**, there are



- Order-of-magnitude better mass resolution.

Observables

But this is not all !

- The **cross-section** in $e^+e^- \rightarrow XX$ close to threshold depends both on coupling, spin and kinematics ($= \beta$).
- The **distribution of the angle** between the two SM-particles depends on β , in a complicated, but calculable way.
- The **cross-section is different** for L and R SUSY particles.
- So checking how much the cross-section changes when switching beam-polarisations measures mixing.
- Measure the helicity of the SM particle \rightarrow properties of the particles in the decay, ie. in addition to the produced X, also the invisible U. In one case this is possible: In $\tilde{\tau} \rightarrow \tau \tilde{\chi}_1^0 \rightarrow X \nu_\tau \tilde{\chi}_1^0$.

Observables

But this is not all !

- The **cross-section** in $e^+e^- \rightarrow XX$ close to threshold depends both on coupling, spin and kinematics ($= \beta$).
- The distribution of the **angle between the two SM-particles** depends on β , in a complicated, but calculable way.
- The cross-section is different for L and R SUSY particles.
- So checking how much the cross-section changes when switching beam-polarisations measures mixing.
- Measure the helicity of the SM particle \rightarrow properties of the particles in the decay, ie. in addition to the produced X, also the invisible U. In one case this is possible: In $\tilde{\tau} \rightarrow \tau \tilde{\chi}_1^0 \rightarrow X \nu_\tau \tilde{\chi}_1^0$.

Observables

But this is not all !

- The **cross-section** in $e^+e^- \rightarrow XX$ close to threshold depends both on coupling, spin and kinematics ($= \beta$).
- The distribution of the **angle between the two SM-particles** depends on β , in a complicated, but calculable way.
- The cross-section is **different for L and R SUSY particles**.
- So checking how much the cross-section changes when **switching beam-polarisations** measures mixing.
- **Measure the helicity of the SM particle \rightarrow properties of the particles in the decay, ie. in addition to the produced X, also the invisible U. In one case this is possible: In $\tilde{\tau} \rightarrow \tau \tilde{\chi}_1^0 \rightarrow X \nu_\tau \tilde{\chi}_1^0$.**

Observables

But this is not all !

- The **cross-section** in $e^+e^- \rightarrow XX$ close to threshold depends both on coupling, spin and kinematics ($= \beta$).
- The distribution of the **angle between the two SM-particles** depends on β , in a complicated, but calculable way.
- The cross-section is **different for L and R SUSY particles**.
- So checking how much the cross-section changes when **switching beam-polarisations** measures mixing.
- Measure the **helicity of the SM particle** \rightarrow properties of the particles in the decay, ie. in addition to the produced X, **also the invisible U**. In one case this is possible: In $\tilde{\tau} \rightarrow \tau \tilde{\chi}_1^0 \rightarrow X \nu_\tau \tilde{\chi}_1^0$.

Extracting the $\tilde{\tau}$ properties

See Phys.Rev.D82:055016,2010

Use polarisation (0.8,-0.22) to reduce bosino background.

From decay kinematics:

- $M_{\tilde{\tau}}$ from $M_{\tilde{\chi}_1^0}$ and end-point of spectrum = $E_{\tau,max}$.
- Other end-point hidden in $\gamma\gamma$ background: **Must get $M_{\tilde{\chi}_1^0}$ from other sources.** ($\tilde{\mu}$, \tilde{e} , ...)

From cross-section:

- $\sigma_{\tilde{\tau}} = A(\theta_{\tilde{\tau}}, \mathcal{P}_{beam}) \times \beta^3 / s$, so
- $M_{\tilde{\tau}} = E_{beam} \sqrt{1 - (\sigma s / A)^{2/3}}$: **no $M_{\tilde{\chi}_1^0}$!**

From decay spectra:

- \mathcal{P}_{τ} from exclusive decay-mode(s): handle on mixing angles $\theta_{\tilde{\tau}}$ and $\theta_{\tilde{\chi}_1^0}$

Topology selection

Take over SPS1a' $\tilde{\tau}$ analysis principle

$\tilde{\ell}$ properties:

- Only two particles (possibly τ :s:s) in the final state.
- Large missing energy and momentum.
- High Acolinearity, with little correlation to the energy of the τ decay-products.
- Central production.
- No forward-backward asymmetry.

+ anti $\gamma\gamma$ cuts.

Select this by:

- Exactly two jets.
- $N_{ch} < 10$
- Vanishing total charge.
- Charge of each jet = ± 1 ,
- $M_{jet} < 2.5 \text{ GeV}/c^2$,
- E_{vis} significantly less than E_{CMS} .
- M_{miss} significantly less than M_{CMS} .
- No particle with momentum close to E_{beam} .

Topology selection

Take over SPS1a' $\tilde{\tau}$ analysis principle

$\tilde{\ell}$ properties:

- Only two particles (possibly τ :s:s) in the final state.
- Large missing energy and momentum.
- High Acolinearity, with little correlation to the energy of the τ decay-products.
- Central production.
- No forward-backward asymmetry.

+ anti $\gamma\gamma$ cuts.

Select this by:

- Exactly two jets.
- $N_{ch} < 10$
- Vanishing total charge.
- Charge of each jet = ± 1 ,
- $M_{jet} < 2.5 \text{ GeV}/c^2$,
- E_{vis} significantly less than E_{CMS} .
- M_{miss} significantly less than M_{CMS} .
- No particle with momentum close to E_{beam} .

Topology selection

Take over SPS1a' $\tilde{\tau}$ analysis principle

$\tilde{\ell}$ properties:

- Only two particles (possibly τ :s:s) in the final state.
- Large missing energy and momentum.
- High Acolinearity, with little correlation to the energy of the τ decay-products.
- Central production.
- No forward-backward asymmetry.

+ anti $\gamma\gamma$ cuts.


Select this by:

- Exactly two jets.
- $N_{ch} < 10$
- Vanishing total charge.
- Charge of each jet = ± 1 ,
- $M_{jet} < 2.5 \text{ GeV}/c^2$,
- E_{vis} significantly less than E_{CMS} .
- M_{miss} significantly less than M_{CMS} .
- No particle with momentum close to E_{beam} .

$\tilde{\tau}_1$ and $\tilde{\tau}_2$ further selections

- $\tilde{\tau}_1$:
 - $(E_{jet1} + E_{jet2}) \sin \theta_{acop} < 30$ GeV.
- $\tilde{\tau}_2$:
 - Other side jet not e or μ
 - Most energetic jet not e or μ
 - Cut on Signal-SM LR of $f(q_{jet1} \cos \theta_{jet1}, q_{jet2} \cos \theta_{jet2})$


Efficiency 15 (22) %



$\tilde{\tau}_1$ and $\tilde{\tau}_2$ further selections

- $\tilde{\tau}_1$:
 - $(E_{jet1} + E_{jet2}) \sin \theta_{acop} < 30$ GeV.
- $\tilde{\tau}_2$:
 - Other side jet not e or μ
 - Most energetic jet not e or μ
 - Cut on Signal-SM LR of $f(q_{jet1} \cos \theta_{jet1}, q_{jet2} \cos \theta_{jet2})$


Efficiency 15 (22) %



$\tilde{\tau}_1$ and $\tilde{\tau}_2$ further selections

- $\tilde{\tau}_1$:
 - $(E_{jet1} + E_{jet2}) \sin \theta_{acop} < 30$ GeV.
- $\tilde{\tau}_2$:
 - Other side jet not e or μ
 - Most energetic jet not e or μ
 - Cut on Signal-SM LR of $f(q_{jet1} \cos \theta_{jet1}, q_{jet2} \cos \theta_{jet2})$


Efficiency 15 (22) %



$\tilde{\tau}_1$ and $\tilde{\tau}_2$ further selections

- $\tilde{\tau}_1$:
 - $(E_{jet1} + E_{jet2}) \sin \theta_{acop} < 30$ GeV.
- $\tilde{\tau}_2$:
 - Other side jet not e or μ
 - Most energetic jet not e or μ
 - Cut on Signal-SM LR of $f(q_{jet1} \cos \theta_{jet1}, q_{jet2} \cos \theta_{jet2})$


Efficiency 15 (22) %



$\tilde{\tau}_1$ and $\tilde{\tau}_2$ further selections

- $\tilde{\tau}_1$:
 - $(E_{jet1} + E_{jet2}) \sin \theta_{acop} < 30$ GeV.
- $\tilde{\tau}_2$:
 - Other side jet not e or μ
 - Most energetic jet not e or μ
 - Cut on Signal-SM LR of $f(q_{jet1} \cos \theta_{jet1}, q_{jet2} \cos \theta_{jet2})$


Efficiency 15 (22) %



$\tilde{\tau}_1$ and $\tilde{\tau}_2$ further selections

- $\tilde{\tau}_1$:
 - $(E_{jet1} + E_{jet2}) \sin \theta_{acop} < 30$ GeV.
- $\tilde{\tau}_2$:
 - Other side jet not e or μ
 - Most energetic jet not e or μ
 - Cut on Signal-SM LR of $f(q_{jet1} \cos \theta_{jet1}, q_{jet2} \cos \theta_{jet2})$

Efficiency 15 (22) %



$\tilde{\tau}_1$ and $\tilde{\tau}_2$ further selections

- $\tilde{\tau}_1$:
 - $(E_{jet1} + E_{jet2}) \sin \theta_{acop} < 30$ GeV.
- $\tilde{\tau}_2$:
 - Other side jet not e or μ
 - Most energetic jet not e or μ
 - Cut on Signal-SM LR of $f(q_{jet1} \cos \theta_{jet1}, q_{jet2} \cos \theta_{jet2})$


Efficiency 15 (22) %

$\tilde{\mu}$ channels

Use “normal” polarisation (-0.8,0.22).

- $\tilde{\mu}_L \tilde{\mu}_L \rightarrow \mu\mu\tilde{\chi}_1^0\tilde{\chi}_1^0$
- $\tilde{\chi}_1^0\tilde{\chi}_2^0 \rightarrow \mu\tilde{\mu}_R\tilde{\chi}_1^0 \rightarrow \mu\mu\tilde{\chi}_1^0\tilde{\chi}_1^0$

- Momentum of μ :s
- E_{miss}
- $M_{\mu\mu}$




$\tilde{\mu}$ channels

Use “normal” polarisation (-0.8,0.22).

- $\tilde{\mu}_L \tilde{\mu}_L \rightarrow \mu\mu\tilde{\chi}_1^0\tilde{\chi}_1^0$
- $\tilde{\chi}_1^0\tilde{\chi}_2^0 \rightarrow \mu\tilde{\mu}_R\tilde{\chi}_1^0 \rightarrow \mu\mu\tilde{\chi}_1^0\tilde{\chi}_1^0$

- Momentum of μ :s
- E_{miss}
- $M_{\mu\mu}$




$\tilde{\mu}$ channels

Use “normal” polarisation (-0.8,0.22).

- $\tilde{\mu}_L \tilde{\mu}_L \rightarrow \mu\mu\tilde{\chi}_1^0\tilde{\chi}_1^0$
- $\tilde{\chi}_1^0\tilde{\chi}_2^0 \rightarrow \mu\tilde{\mu}_R\tilde{\chi}_1^0 \rightarrow \mu\mu\tilde{\chi}_1^0\tilde{\chi}_1^0$

- Momentum of μ :s
- E_{miss}
- $M_{\mu\mu}$



$\tilde{\mu}_L \tilde{\mu}_L$



Selections

- $\theta_{\text{missing}p} \in [0.1\pi; 0.9\pi]$
- $E_{\text{miss}} \in [200, 430]\text{GeV}$
- $M_{\mu\mu} \notin [80, 100]\text{GeV}$ and $> 30 \text{ GeV}/c^2$

Masses from edges. Beam-energy spread dominates error.

$$\Delta(M_{\tilde{\chi}_1^0}) = 920 \text{ MeV}/c^2$$

$$\Delta(M_{\tilde{\mu}_L}) = 100 \text{ MeV}/c^2$$



$\tilde{\mu}_L \tilde{\mu}_L$



Selections

- $\theta_{missing p} \in [0.1\pi; 0.9\pi]$
- $E_{miss} \in [200, 430]\text{GeV}$
- $M_{\mu\mu} \notin [80, 100]\text{GeV}$ and $> 30\text{ GeV}/c^2$

Masses from edges. Beam-energy spread dominates error.

$$\Delta(M_{\tilde{\chi}_1^0}) = 920\text{MeV}/c^2$$

$$\Delta(M_{\tilde{\mu}_L}) = 100\text{MeV}/c^2$$





$$\tilde{\chi}_1^0 \tilde{\chi}_2^0$$

Selections

- $\theta_{\text{missing } p} \in [0.2\pi; 0.8\pi]$
- $p_{T\text{miss}} > 40\text{GeV}/c$
- β of μ system > 0.6 .
- $E_{\text{miss}} \in [355, 395]\text{GeV}$

Masses from edges. Beam-energy spread dominates error.

$$\Delta(M_{\tilde{\chi}_2^0}) = 1.38\text{GeV}/c^2$$





$$\tilde{\chi}_1^0 \tilde{\chi}_2^0$$

Selections

- $\theta_{\text{missing } p} \in [0.2\pi; 0.8\pi]$
- $p_{T\text{miss}} > 40 \text{ GeV}/c$
- β of μ system > 0.6 .
- $E_{\text{miss}} \in [355, 395] \text{ GeV}$

Masses from edges. Beam-energy spread dominates error.

$$\Delta(M_{\tilde{\chi}_2^0}) = 1.38 \text{ GeV}/c^2$$



Observables and background:

Observable	Gives	If
Edges (or average and width)	Masses	... not too far from threshold
Shape of spectrum	Spin	
Angular distributions	Mass, Spin	
Invariant mass distributions from full reconstruction	Mass	... cascade decays
Angular distributions from full reconstruction	Spin, CP,	... masses known
Un-polarised Cross-section in continuum	Mass, coupling	
Polarised Cross-section in continuum	Mass, coupling, mixing	
Decay product polarisation	Mixing	... $\tilde{\tau}$ decays
Threshold-scan	Mass(es), Spin	

Channels and observables at 250, 350 and 500 GeV

Channel	Threshold	Available at	Can give
$\tilde{\tau}_1 \tilde{\tau}_1$	212	250	$M_{\tilde{\tau}_1}$, $\tilde{\tau}_1$ nature, τ polarisation
$\tilde{\mu}_R \tilde{\mu}_R$	252	250+	+ $M_{\tilde{\mu}_R}$, $M_{\tilde{\chi}_1^0}$, $\tilde{\mu}_R$ nature
$\tilde{e}_R \tilde{e}_R$	252	250+	+ $M_{\tilde{e}_R}$, $M_{\tilde{\chi}_1^0}$, \tilde{e}_R nature
$\tilde{\chi}_1^0 \tilde{\chi}_2^0$ *)	302	350	+ $M_{\tilde{\chi}_2^0}$, $M_{\tilde{\chi}_1^0}$, nature of $\tilde{\chi}_1^0$, $\tilde{\chi}_2^0$
$\tilde{\tau}_1 \tilde{\tau}_2$ *)	325	350	+ $M_{\tilde{\tau}_2} \theta_{mix} \tilde{\tau}$
$\tilde{e}_R \tilde{e}_L$ *)	339	350	+ $M_{\tilde{e}_L}$, $\tilde{\chi}_1^0$ mixing, \tilde{e}_L nature
$\tilde{\nu}_\tau \tilde{\nu}_\tau$	392	500	7 % visible BR ($\rightarrow \tilde{\tau}_1 W$)
$\tilde{\chi}_1^\pm \tilde{\chi}_1^\pm$ *)	412	500	+ $M_{\tilde{\chi}_1^\pm}$, nature of $\tilde{\chi}_1^\pm$
$\tilde{e}_L \tilde{e}_L$ *)	416	500	+ $M_{\tilde{e}_L}$, $M_{\tilde{\chi}_1^0}$, \tilde{e}_L nature
$\tilde{\mu}_L \tilde{\mu}_L$ *)	416	500	+ $M_{\tilde{\mu}_R}$, $M_{\tilde{\chi}_1^0}$, $\tilde{\mu}_R$ nature
$\tilde{\tau}_2 \tilde{\tau}_2$ *)	438	500	+ $M_{\tilde{\tau}_2}$, $M_{\tilde{\chi}_1^0}$, $\tilde{\tau}_2$ nature, $\theta_{mix} \tilde{\tau}$
$\tilde{\chi}_1^0 \tilde{\chi}_3^0$ *)	503	500+	+ $M_{\tilde{\chi}_3^0}$, $M_{\tilde{\chi}_1^0}$, nature of $\tilde{\chi}_1^0$, $\tilde{\chi}_3^0$

*) : Cascade decays.

+ invisible $\tilde{\chi}_1^0 \tilde{\chi}_1^0$, $\tilde{\nu}_{\tilde{e}}, \tilde{\mu} \tilde{\nu}_{\tilde{e}}, \tilde{\mu}$.

SUSY backgrounds at the ILC

The generic STCx signal at ILC is a few SM-particles, usually leptons, and lots of missing energy and momentum.

So: **Background from SM:**

- **Real missing energy** + pair of SM-particles = di-boson production, with neutrinos:
 - $WW \rightarrow \ell\nu\ell\nu$
 - $ZZ \rightarrow f\bar{f}\nu\nu$
- **Fake missing energy** + pair of SM-particles = $\gamma\gamma$ processes, ISR, single IVB.
 - $e^+e^- \rightarrow e^+e^-\gamma\gamma \rightarrow e^+e^-\bar{f}f$, with both e^+e^- un-detected.
 - $e^+e^- \rightarrow \bar{f}f\gamma$, with γ un-detected.

SUSY backgrounds at the ILC

The generic STCx signal at ILC is a few SM-particles, usually leptons, and lots of missing energy and momentum.

So: **Background from SM:**

- **Real missing energy** + pair of SM-particles = di-boson production, with neutrinos:
 - $WW \rightarrow \ell\nu\ell\nu$
 - $ZZ \rightarrow f\bar{f}\nu\nu$
- **Fake missing energy** + pair of SM-particles = $\gamma\gamma$ processes, ISR, single IVB.
 - $e^+e^- \rightarrow e^+e^-\gamma\gamma \rightarrow e^+e^-\bar{f}f$, with both e^+e^- un-detected.
 - $e^+e^- \rightarrow \bar{f}f\gamma$, with γ un-detected.

SUSY backgrounds at the ILC

The generic STCx signal at ILC is a few SM-particles, usually leptons, and lots of missing energy and momentum.

So: **Background from SM:**

- **Real missing energy** + pair of SM-particles = di-boson production, with neutrinos:
 - $WW \rightarrow \ell\nu\ell\nu$
 - $ZZ \rightarrow f\bar{f}\nu\nu$
- **Fake missing energy** + pair of SM-particles = $\gamma\gamma$ processes, ISR, single IVB.
 - $e^+e^- \rightarrow e^+e^-\gamma\gamma \rightarrow e^+e^-\bar{f}f$, with both e^+e^- un-detected.
 - $e^+e^- \rightarrow \bar{f}f\gamma$, with γ un-detected.

SUSY backgrounds at the ILC

The generic STCx signal at ILC is a few SM-particles, usually leptons, and lots of missing energy and momentum.

So: **Background from SM:**

- **Real missing energy** + pair of SM-particles = di-boson production, with neutrinos:
 - $WW \rightarrow \ell\nu\ell\nu$
 - $ZZ \rightarrow f\bar{f}\nu\nu$
- **Fake missing energy** + pair of SM-particles = $\gamma\gamma$ processes, ISR, single IVB.
 - $e^+e^- \rightarrow e^+e^-\gamma\gamma \rightarrow e^+e^-f\bar{f}$, with both e^+e^- un-detected.
 - $e^+e^- \rightarrow f\bar{f}\gamma$, with γ un-detected.

Features of STC4 @ 500 GeV

- The $\tilde{\tau}_1$ is the NLSP.
- For $\tilde{\tau}_1$: $E_{\tau,min} = 2.3$ GeV, $E_{\tau,max} = 45.5$ GeV:
 $\gamma\gamma - \text{background} \Leftrightarrow \text{pairs} - \text{background}$.
- For $\tilde{\tau}_2$: $E_{\tau,min} = 52.4$ GeV, $E_{\tau,max} = 150.0$ GeV:
 $WW \rightarrow l\nu l\nu - \text{background} \Leftrightarrow \text{Polarisation}$.
- For \tilde{e}_R or $\tilde{\mu}_R$: $E_{l,min} = 7.3$ GeV, $E_{l,max} = 99.2$ GeV: Neither $\gamma\gamma$ nor $WW \rightarrow l\nu l\nu$ background severe.
- For $\text{pol}=(1,-1)$: $\sigma(\tilde{e}_R\tilde{e}_R) = 1.3$ pb !
- $\tilde{\tau}$ NLSP $\rightarrow \tau$:s in most SUSY decays \rightarrow SUSY is background to SUSY.
- For $\text{pol}=(-1,1)$: $\sigma(\tilde{\chi}_2^0\tilde{\chi}_2^0)$ and $\sigma(\tilde{\chi}_1^+\tilde{\chi}_1^-) = \text{several hundred fb}$ and $\text{BR}(X \rightarrow \tilde{\tau}) > 70\%$. For $\text{pol}=(1,-1)$: $\sigma(\tilde{\chi}_2^0\tilde{\chi}_2^0)$ and $\sigma(\tilde{\chi}_1^+\tilde{\chi}_1^-) \approx 0$.

Features of STC4 @ 500 GeV

- The $\tilde{\tau}_1$ is the NLSP.
- For $\tilde{\tau}_1$: $E_{\tau,min} = 2.3$ GeV, $E_{\tau,max} = 45.5$ GeV:
 $\gamma\gamma - \text{background} \Leftrightarrow \text{pairs} - \text{background}$.
- For $\tilde{\tau}_2$: $E_{\tau,min} = 52.4$ GeV, $E_{\tau,max} = 150.0$ GeV:
 $WW \rightarrow l\nu l\nu - \text{background} \Leftrightarrow \text{Polarisation}$.
- For \tilde{e}_R or $\tilde{\mu}_R$: $E_{l,min} = 7.3$ GeV, $E_{l,max} = 99.2$ GeV: Neither $\gamma\gamma$ nor $WW \rightarrow l\nu l\nu$ background severe.
- For $\text{pol}=(1,-1)$: $\sigma(\tilde{e}_R\tilde{e}_R) = 1.3$ pb !
- $\tilde{\tau}$ NLSP $\rightarrow \tau$:s in most SUSY decays \rightarrow SUSY is background to SUSY.
- For $\text{pol}=(-1,1)$: $\sigma(\tilde{\chi}_2^0\tilde{\chi}_2^0)$ and $\sigma(\tilde{\chi}_1^+\tilde{\chi}_1^-) = \text{several hundred fb}$ and $\text{BR}(X \rightarrow \tilde{\tau}) > 70\%$. For $\text{pol}=(1,-1)$: $\sigma(\tilde{\chi}_2^0\tilde{\chi}_2^0)$ and $\sigma(\tilde{\chi}_1^+\tilde{\chi}_1^-) \approx 0$.

Features of STC4 @ 500 GeV

- The $\tilde{\tau}_1$ is the NLSP.
- For $\tilde{\tau}_1$: $E_{\tau,min} = 2.3$ GeV, $E_{\tau,max} = 45.5$ GeV:
 $\gamma\gamma - \text{background} \Leftrightarrow \text{pairs} - \text{background}$.
- For $\tilde{\tau}_2$: $E_{\tau,min} = 52.4$ GeV, $E_{\tau,max} = 150.0$ GeV:
 $WW \rightarrow l\nu l\nu - \text{background} \Leftrightarrow \text{Polarisation}$.
- For \tilde{e}_R or $\tilde{\mu}_R$: $E_{l,min} = 7.3$ GeV, $E_{l,max} = 99.2$ GeV: Neither $\gamma\gamma$ nor $WW \rightarrow l\nu l\nu$ background severe.
- For $\text{pol}=(1,-1)$: $\sigma(\tilde{e}_R\tilde{e}_R) = 1.3$ pb !
- $\tilde{\tau}$ NLSP $\rightarrow \tau$:s in most SUSY decays \rightarrow SUSY is background to SUSY.
- For $\text{pol}=(-1,1)$: $\sigma(\tilde{\chi}_2^0\tilde{\chi}_2^0)$ and $\sigma(\tilde{\chi}_1^+\tilde{\chi}_1^-) =$ several hundred fb and $\text{BR}(X \rightarrow \tilde{\tau}) > 70\%$. For $\text{pol}=(1,-1)$: $\sigma(\tilde{\chi}_2^0\tilde{\chi}_2^0)$ and $\sigma(\tilde{\chi}_1^+\tilde{\chi}_1^-) \approx 0$.

Features of STC4 @ 500 GeV

- The $\tilde{\tau}_1$ is the NLSP.
- For $\tilde{\tau}_1$: $E_{\tau,min} = 2.3$ GeV, $E_{\tau,max} = 45.5$ GeV:
 $\gamma\gamma - \text{background} \Leftrightarrow \text{pairs} - \text{background}$.
- For $\tilde{\tau}_2$: $E_{\tau,min} = 52.4$ GeV, $E_{\tau,max} = 150.0$ GeV:
 $WW \rightarrow l\nu l\nu - \text{background} \Leftrightarrow \text{Polarisation}$.
- For \tilde{e}_R or $\tilde{\mu}_R$: $E_{l,min} = 7.3$ GeV, $E_{l,max} = 99.2$ GeV: Neither $\gamma\gamma$ nor $WW \rightarrow l\nu l\nu$ background severe.
- For $\text{pol}=(1,-1)$: $\sigma(\tilde{e}_R\tilde{e}_R) = 1.3$ pb !
- $\tilde{\tau}$ NLSP $\rightarrow \tau$:s in most SUSY decays \rightarrow SUSY is background to SUSY.
- For $\text{pol}=(1,-1)$: $\sigma(\tilde{\chi}_2^0\tilde{\chi}_2^0)$ and $\sigma(\tilde{\chi}_1^+\tilde{\chi}_1^-) = \text{several hundred fb}$ and $\text{BR}(X \rightarrow \tilde{\tau}) > 70\%$. For $\text{pol}=(1,-1)$: $\sigma(\tilde{\chi}_2^0\tilde{\chi}_2^0)$ and $\sigma(\tilde{\chi}_1^+\tilde{\chi}_1^-) \approx 0$.

STC4 @ 500 GeV

Strategy:

- Global preselection to reduce SM, while efficiency for **all** signals stays above $\sim 90\%$.
- The further select for **all sleptons** ($\tilde{e}_R, \tilde{e}_L, \tilde{\mu}_R, \tilde{\mu}_L, \tilde{\tau}_1$).
- Next step: **specific** selections for \tilde{e}_R and $\tilde{\mu}_R$, for \tilde{e}_L and $\tilde{\mu}_L$, and for $\tilde{\tau}_1$.
- Last step: add **particle id** to separate \tilde{e} and $\tilde{\mu}$, special cuts for $\tilde{\tau}_1$.
- Check results both for **RL and LR** beam-polarisation.

Features of STC4 @ 500 GeV

- The $\tilde{\tau}_1$ is the NLSP.
- For $\tilde{\tau}_1$: $E_{\tau,min} = 2.3$ GeV, $E_{\tau,max} = 45.5$ GeV:
 $\gamma\gamma - \text{background} \Leftrightarrow \text{pairs} - \text{background}$.
- For $\tilde{\tau}_2$: $E_{\tau,min} = 52.4$ GeV, $E_{\tau,max} = 150.0$ GeV:
 $WW \rightarrow l\nu l\nu - \text{background} \Leftrightarrow \text{Polarisation}$.
- For \tilde{e}_R or $\tilde{\mu}_R$: $E_{l,min} = 7.3$ GeV, $E_{l,max} = 99.2$ GeV: Neither $\gamma\gamma$ nor $WW \rightarrow l\nu l\nu$ background severe.
- For $\text{pol}=(1,-1)$: $\sigma(\tilde{e}_R\tilde{e}_R) = 1.3$ pb !
- $\tilde{\tau}$ NLSP $\rightarrow \tau$:s in most SUSY decays \rightarrow SUSY is background to SUSY.
- For $\text{pol}=(1,-1)$: $\sigma(\tilde{\chi}_2^0\tilde{\chi}_2^0)$ and $\sigma(\tilde{\chi}_1^+\tilde{\chi}_1^-) = \text{several hundred fb}$ and $\text{BR}(X \rightarrow \tilde{\tau}) > 70\%$. For $\text{pol}=(1,-1)$: $\sigma(\tilde{\chi}_2^0\tilde{\chi}_2^0)$ and $\sigma(\tilde{\chi}_1^+\tilde{\chi}_1^-) \approx 0$.

Features of STC4 @ 500 GeV

- The $\tilde{\tau}_1$ is the NLSP.
- For $\tilde{\tau}_1$: $E_{\tau,min} = 2.3$ GeV, $E_{\tau,max} = 45.5$ GeV:
 $\gamma\gamma - \text{background} \Leftrightarrow \text{pairs} - \text{background}$.
- For $\tilde{\tau}_2$: $E_{\tau,min} = 52.4$ GeV, $E_{\tau,max} = 150.0$ GeV:
 $WW \rightarrow l\nu l\nu - \text{background} \Leftrightarrow \text{Polarisation}$.
- For \tilde{e}_R or $\tilde{\mu}_R$: $E_{l,min} = 7.3$ GeV, $E_{l,max} = 99.2$ GeV: Neither $\gamma\gamma$ nor $WW \rightarrow l\nu l\nu$ background severe.
- For $\text{pol}=(1,-1)$: $\sigma(\tilde{e}_R\tilde{e}_R) = 1.3$ pb !
- $\tilde{\tau}$ NLSP $\rightarrow \tau$:s in most SUSY decays \rightarrow SUSY is background to SUSY.
- For $\text{pol}=(1,-1)$: $\sigma(\tilde{\chi}_2^0\tilde{\chi}_2^0)$ and $\sigma(\tilde{\chi}_1^+\tilde{\chi}_1^-) = \text{several hundred fb}$ and $\text{BR}(X \rightarrow \tilde{\tau}) > 70\%$. For $\text{pol}=(1,-1)$: $\sigma(\tilde{\chi}_2^0\tilde{\chi}_2^0)$ and $\sigma(\tilde{\chi}_1^+\tilde{\chi}_1^-) \approx 0$.

Features of STC4 @ 500 GeV

- The $\tilde{\tau}_1$ is the NLSP.
- For $\tilde{\tau}_1$: $E_{\tau,min} = 2.3$ GeV, $E_{\tau,max} = 45.5$ GeV:
 $\gamma\gamma - \text{background} \Leftrightarrow \text{pairs} - \text{background}$.
- For $\tilde{\tau}_2$: $E_{\tau,min} = 52.4$ GeV, $E_{\tau,max} = 150.0$ GeV:
 $WW \rightarrow l\nu l\nu - \text{background} \Leftrightarrow \text{Polarisation}$.
- For \tilde{e}_R or $\tilde{\mu}_R$: $E_{l,min} = 7.3$ GeV, $E_{l,max} = 99.2$ GeV: Neither $\gamma\gamma$ nor $WW \rightarrow l\nu l\nu$ background severe.
- For $\text{pol}=(1,-1)$: $\sigma(\tilde{e}_R\tilde{e}_R) = 1.3$ pb !
- $\tilde{\tau}$ NLSP $\rightarrow \tau$:s in most SUSY decays \rightarrow SUSY is background to SUSY.
- For $\text{pol}=(-1,1)$: $\sigma(\tilde{\chi}_2^0\tilde{\chi}_2^0)$ and $\sigma(\tilde{\chi}_1^+\tilde{\chi}_1^-) =$ several hundred fb and $\text{BR}(X \rightarrow \tilde{\tau}) > 70$ %. For $\text{pol}=(1,-1)$: $\sigma(\tilde{\chi}_2^0\tilde{\chi}_2^0)$ and $\sigma(\tilde{\chi}_1^+\tilde{\chi}_1^-) \approx 0$.

Features of STC4 @ 500 GeV

- The $\tilde{\tau}_1$ is the NLSP.
- For $\tilde{\tau}_1$: $E_{\tau,min} = 2.3$ GeV, $E_{\tau,max} = 45.5$ GeV:
 $\gamma\gamma - \text{background} \Leftrightarrow \text{pairs} - \text{background}$.
- For $\tilde{\tau}_2$: $E_{\tau,min} = 52.4$ GeV, $E_{\tau,max} = 150.0$ GeV:
 $WW \rightarrow l\nu l\nu - \text{background} \Leftrightarrow \text{Polarisation}$.
- For \tilde{e}_R or $\tilde{\mu}_R$: $E_{l,min} = 7.3$ GeV, $E_{l,max} = 99.2$ GeV: Neither $\gamma\gamma$ nor $WW \rightarrow l\nu l\nu$ background severe.
- For $\text{pol}=(1,-1)$: $\sigma(\tilde{e}_R\tilde{e}_R) = 1.3$ pb !
- $\tilde{\tau}$ NLSP $\rightarrow \tau$:s in most SUSY decays \rightarrow SUSY is background to SUSY.
- For $\text{pol}=(-1,1)$: $\sigma(\tilde{\chi}_2^0\tilde{\chi}_2^0)$ and $\sigma(\tilde{\chi}_1^+\tilde{\chi}_1^-) = \text{several hundred fb}$ and $\text{BR}(X \rightarrow \tilde{\tau}) > 70\%$. For $\text{pol}=(1,-1)$: $\sigma(\tilde{\chi}_2^0\tilde{\chi}_2^0)$ and $\sigma(\tilde{\chi}_1^+\tilde{\chi}_1^-) \approx 0$.

Example: The STCx models

STC4-10

- 11 parameters.
- Separate gluino
- Higgs, un-coloured, and coloured scalar parameters separate

Parameters chosen to deliver all constraints (LHC, LEP, cosmology, low energy). In particular, the $\tilde{\tau}_1$ is the NLSP, with a mass-difference to the LSP ~ 10 GeV \Rightarrow **Co-anhilitaion**.


At $E_{CMS} = 500$ GeV:

- All sleptons available.
- No squarks.
- Lighter bosinos, up to $\tilde{\chi}_3^0$ (in $e^+e^- \rightarrow \tilde{\chi}_1^0\tilde{\chi}_3^0$)

(See H. Baer, J. List, arXiv:1307:0782.)


Fitting the $\tilde{\tau}$ end-points

- Only the **upper end-point** is relevant.
- Background subtraction:
 - $\tilde{\tau}_1$: Important SUSY background, but region above 45 GeV is **signal free**. Fit exponential and extrapolate.
 - $\tilde{\tau}_2$: \sim no SUSY background above 45 GeV. Take background from SM-only simulation and fit exponential.
- Fit line to (data-background fit).




Fitting the $\tilde{\tau}$ end-points

- Only the **upper end-point** is relevant.
- Background subtraction:
 - $\tilde{\tau}_1$: Important SUSY background, but region above 45 GeV is **signal free**. Fit exponential and extrapolate.
 - $\tilde{\tau}_2$: \sim no SUSY background above 45 GeV. Take background from SM-only simulation and fit exponential.
- Fit line to (data-background fit).




Fitting the $\tilde{\tau}$ end-points

- Only the **upper end-point** is relevant.
- Background subtraction:
 - $\tilde{\tau}_1$: Important SUSY background, but region above 45 GeV is **signal free**. Fit exponential and extrapolate.
 - $\tilde{\tau}_2$: \sim no SUSY background above 45 GeV. Take background from SM-only simulation and fit exponential.
- Fit line to (data-background fit).



Fitting the $\tilde{\tau}$ end-points

- Only the **upper end-point** is relevant.
- Background subtraction:
 - $\tilde{\tau}$: Important SUSY



Results for $\tilde{\tau}_1$

$$E_{max, \tilde{\tau}_1} = 44.49^{+0.11}_{-0.09} \text{ GeV}$$

Translates to an error on the mass of $0.27 \text{ GeV}/c^2$, dominated by the error from $M_{\tilde{\chi}_1^0}$.

Results for $\tilde{\tau}_2$

$$E_{max, \tilde{\tau}_2} = 145.4^{+5.9}_{-4.4} \text{ GeV}$$

Translates to an error on the mass of $5 \text{ GeV}/c^2$, dominated by the error from the end-point.

Reminder: SPS1a' results (Phys.Rev.D82:055016,2010)

The previous $\tilde{\tau}$ study in the very similar model SPS1a' gave:

Results for $\tilde{\tau}_1$

$$M_{\tilde{\tau}_1} = 107.73_{-0.05}^{+0.03} \text{ GeV}/c^2 \oplus 1.3\Delta(M_{\tilde{\chi}_1^0})$$

The error from $M_{\tilde{\chi}_1^0}$ largely dominates

Results for $\tilde{\tau}_2$

$$M_{\tilde{\tau}_2} = 183_{-5}^{+11} \text{ GeV}/c^2 \oplus 18\Delta(M_{\tilde{\chi}_1^0})$$

The error from the endpoint largely dominates

Results from cross-section for $\tilde{\tau}_1$

$$\Delta(N_{\text{signal}})/N_{\text{signal}} = 3.1\% \rightarrow \Delta(M_{\tilde{\tau}_1}) = 3.2 \text{ GeV}/c^2$$

Results from cross-section for $\tilde{\tau}_2$

$$\Delta(N_{\text{signal}})/N_{\text{signal}} = 4.2\% \rightarrow \Delta(M_{\tilde{\tau}_2}) = 3.6 \text{ GeV}/c^2$$

End-point + Cross-section
 $\rightarrow \Delta(M_{\tilde{\chi}_1^0}) = 1.7 \text{ GeV}/c^2$

Also: τ polarisation in $\tilde{\tau}_1$ decays

$$\Delta(\mathcal{P}_\tau)/\mathcal{P}_\tau = 9\%.$$