


# The OLYMPUS Experiment at DESY

Michael Kohl for the OLYMPUS Collaboration

Physics Department, Hampton University, Hampton, VA 23668  
Jefferson Laboratory, Newport News, VA 23606




## Abstract


The proton is not an elementary particle but has a substructure governed by quarks and gluons. The spatial distributions of the electric charge and magnetization determine the response to electromagnetic interaction. Recently, it has been demonstrated that the observed difference in the elastic electric-to-magnetic proton form factor ratio with unpolarized and polarized methods can be explained by hard two-photon exchange as a radiative correction previously unaccounted for. Calculations of this correction are model-dependent and need to be validated with precision measurements. The most direct verification is a comparison of positron-proton and electron-proton elastic scattering. The OLYMPUS experiment at DESY was designed to make a definitive, high precision measurement of the elastic  $e^+p/e^-p$  cross section ratio, in order to provide an end to the controversy. The status of the OLYMPUS experiment and analysis will be reported.

## Physics case

- Proton form factor ratio discrepancy between unpolarized and polarized measurements



## Textbook physics




- Interference term depends on lepton charge sign (**C-odd**)

$$\sigma_{e^\pm p} = |\mathcal{M}_{1\gamma}|^2 \pm 2\Re\{\mathcal{M}_{1\gamma}^\dagger \mathcal{M}_{2\gamma}\} + \dots$$


- $e^+/e^-$  ratio deviates from unity by two-photon contribution

$$\frac{\sigma_{e^+p}}{\sigma_{e^-p}} \approx 1 + 4 \frac{\Re\{\mathcal{M}_{1\gamma}^\dagger \mathcal{M}_{2\gamma}\}}{|\mathcal{M}_{1\gamma}|^2}$$


## Projection



## Experimental setup




## Data taking: DORIS at DESY




- 2007 Letter of Intent
- 2008 Proposal
- Spring 2011 Target test run
- Summer 2011 Detector installed
- Fall 2011 Commissioning
- First run Jan 30 – Feb 27, 2012 ... acquired < 0.3 fb^-1
- Summer 2012 Repairs and upgrades
- Second run Oct 24, 2012 – Jan 2, 2013 ... acquired > 4.0 fb^-1
- Smooth performance of machine, target, detector
- Spring 2013 Survey & field mapping
- Analysis progressing – framework, calibrations, tracking, simulations

## Kinematics




## Analysis

- Monte Carlo geometry, acceptance, efficiencies fully integrated in analysis
- Redundant luminosity monitoring:  $e^+p$  elastic at 12° and 1.2° (“pile-up”); sym. Möller/Bhabha; slow control
- Tracking, efficiencies → normalized yields → extraction of  $e^+p/e^-p$  ratio




Slow control luminosity induced asymmetry: 0.60% Analysis by Axel Schmidt, MIT




## Radiative effects

Baseline: Dipole F. F., Maximon Tjon rad. corrections



Simulations by Rebecca Russell, MIT  
Figure by Jan Bernauer, MIT

## Toward final results



## Publications

The OLYMPUS Experiment,  
R. Milner et al., NIMA 741, 1 (2014)

The OLYMPUS Internal Hydrogen Target,  
J.C. Bernauer et al., NIMA 755, 20 (2014)

Measurement and tricubic interpolation of the magnetic field  
for the OLYMPUS experiment,  
J.C. Bernauer et al., NIMA 823, 9 (2016)

## Acknowledgment

This work has been supported by the U.S. National Science Foundation and Department of Energy.