

Search for a dark vector gauge boson decaying to $\pi^+\pi^-$ using $\eta \rightarrow \pi^+\pi^-\gamma$ decays

E. Won,^{29,*} I. Adachi,^{13,10} H. Aihara,⁶⁸ S. Al Said,^{61,27} D. M. Asner,⁵² T. Aushev,⁴¹ R. Ayad,⁶¹ I. Badhrees,^{61,26} A. M. Bakich,⁶⁰ V. Bansal,⁵² E. Barberio,³⁸ P. Behera,¹⁸ B. Bhuyan,¹⁷ J. Biswal,²² A. Bobrov,^{3,50} A. Bozek,⁴⁸ M. Bračko,^{36,22} D. Červenkov,⁴ V. Chekelian,³⁷ A. Chen,⁴⁵ B. G. Cheon,¹¹ K. Chilikin,^{32,40} R. Chistov,^{32,40} K. Cho,²⁸ V. Chobanova,³⁷ Y. Choi,⁵⁹ D. Cinabro,⁷³ N. Dash,¹⁶ S. Di Carlo,⁷³ Z. Doležal,⁴ Z. Drásal,⁴ D. Dutta,⁶² S. Eidelman,^{3,50} D. Epifanov,^{3,50} H. Farhat,⁷³ J. E. Fast,⁵² T. Ferber,⁷ B. G. Fulsom,⁵² V. Gaur,⁶² N. Gabyshev,^{3,50} A. Garmash,^{3,50} R. Gillard,⁷³ P. Goldenzweig,²⁴ D. Greenwald,⁶⁴ J. Haba,^{13,10} K. Hayasaka,⁴⁹ H. Hayashii,⁴⁴ W.-S. Hou,⁴⁷ T. Iijima,^{43,42} K. Inami,⁴² G. Inguglia,⁷ A. Ishikawa,⁶⁶ R. Itoh,^{13,10} Y. Iwasaki,¹³ I. Jaegle,⁸ H. B. Jeon,³⁰ D. Joffe,²⁵ K. K. Joo,⁵ T. Julius,³⁸ K. H. Kang,³⁰ T. Kawasaki,⁴⁹ D. Y. Kim,⁵⁷ J. B. Kim,²⁹ K. T. Kim,²⁹ M. J. Kim,³⁰ S. H. Kim,¹¹ Y. J. Kim,²⁸ K. Kinoshita,⁶ P. Kodyš,⁴ P. Križan,^{33,22} P. Krokovny,^{3,50} T. Kuhr,³⁴ R. Kulasiri,²⁵ Y.-J. Kwon,⁷⁵ J. S. Lange,⁹ I. S. Lee,¹¹ C. H. Li,³⁸ L. Li,⁵⁵ Y. Li,⁷² L. Li Gioi,³⁷ J. Libby,¹⁸ D. Liventsev,^{72,13} T. Luo,⁵³ M. Masuda,⁶⁷ T. Matsuda,³⁹ D. Matvienko,^{3,50} K. Miyabayashi,⁴⁴ H. Miyata,⁴⁹ R. Mizuk,^{32,40,41} G. B. Mohanty,⁶² E. Nakano,⁵¹ M. Nakao,^{13,10} H. Nakazawa,⁴⁷ T. Nanut,²² K. J. Nath,¹⁷ Z. Natkaniec,⁴⁸ M. Nayak,^{73,13} S. Nishida,^{13,10} S. Ogawa,⁶⁵ S. Okuno,²³ P. Pakhlov,^{32,40} B. Pal,⁶ C.-S. Park,⁷⁵ S. Paul,⁶⁴ T. K. Pedlar,³⁵ L. E. Piilonen,⁷² C. Pulvermacher,¹³ J. Rauch,⁶⁴ M. Ritter,³⁴ H. Sahoo,¹² Y. Sakai,^{13,10} S. Sandilya,⁶ L. Santelj,¹³ T. Sanuki,⁶⁶ Y. Sato,⁴² V. Savinov,⁵³ T. Schlüter,³⁴ O. Schneider,^{1,15} G. Schnell,^{1,15} C. Schwanda,²⁰ Y. Seino,⁴⁹ D. Semmler,⁹ K. Senyo,⁷⁴ O. Seon,⁴² I. S. Seong,¹² V. Shebalin,^{3,50} C. P. Shen,² T.-A. Shibata,⁶⁹ J.-G. Shiu,⁴⁷ F. Simon,^{37,63} M. Starič,²² T. Sumiyoshi,⁷⁰ M. Takizawa,^{56,14,54} U. Tamponi,^{21,71} F. Tenchini,³⁸ K. Trabelsi,^{13,10} M. Uchida,⁶⁹ S. Uehara,^{13,10} T. Uglov,^{32,41} Y. Unno,¹¹ S. Uno,^{13,10} P. Urquijo,³⁸ Y. Usov,^{3,50} C. Van Hulse,¹ G. Varner,¹² K. E. Varvell,⁶⁰ V. Vorobyev,^{3,50} C. H. Wang,⁴⁶ M.-Z. Wang,⁴⁷ M. Watanabe,⁴⁹ Y. Watanabe,²³ E. Widmann,⁵⁸ J. Yamaoka,⁵² H. Ye,⁷ Y. Yook,⁷⁵ C. Z. Yuan,¹⁹ Y. Yusa,⁴⁹ Z. P. Zhang,⁵⁵ V. Zhilich,^{3,50} V. Zhukova,⁴⁰ V. Zhulanov,^{3,50} and A. Zupanc^{33,22}

(Belle Collaboration)

¹University of the Basque Country UPV/EHU, 48080 Bilbao, Spain

²Beihang University, Beijing 100191, China

³Budker Institute of Nuclear Physics SB RAS, Novosibirsk 630090, Russia

⁴Faculty of Mathematics and Physics, Charles University, 121 16 Prague, Czech Republic

⁵Chonnam National University, Kwangju 660-701, Korea

⁶University of Cincinnati, Cincinnati, Ohio 45221, USA

⁷Deutsches Elektronen-Synchrotron, 22607 Hamburg, Germany

⁸University of Florida, Gainesville, Florida 32611, USA

⁹Justus-Liebig-Universität Gießen, 35392 Gießen, Germany

¹⁰SOKENDAI (The Graduate University for Advanced Studies), Hayama 240-0193, Japan

¹¹Hanyang University, Seoul 133-791, Korea

¹²University of Hawaii, Honolulu, Hawaii 96822, USA

¹³High Energy Accelerator Research Organization (KEK), Tsukuba 305-0801, Japan

¹⁴J-PARC Branch, KEK Theory Center, High Energy Accelerator Research Organization (KEK), Tsukuba 305-0801, Japan

¹⁵IKERBASQUE, Basque Foundation for Science, 48013 Bilbao, Spain

¹⁶Indian Institute of Technology Bhubaneswar, Satya Nagar 751007, India

¹⁷Indian Institute of Technology Guwahati, Assam 781039, India

¹⁸Indian Institute of Technology Madras, Chennai 600036, India

¹⁹Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049, China

²⁰Institute of High Energy Physics, Vienna 1050, Austria

²¹INFN—Sezione di Torino, 10125 Torino, Italy

²²J. Stefan Institute, 1000 Ljubljana, Slovenia

²³Kanagawa University, Yokohama 221-8686, Japan

²⁴Institut für Experimentelle Kernphysik, Karlsruher Institut für Technologie, 76131 Karlsruhe, Germany

²⁵Kennesaw State University, Kennesaw, Georgia 30144, USA

²⁶King Abdulaziz City for Science and Technology, Riyadh 11442, Saudi Arabia

²⁷Department of Physics, Faculty of Science, King Abdulaziz University, Jeddah 21589, Saudi Arabia

²⁸Korea Institute of Science and Technology Information, Daejeon 305-806, Korea

²⁹Korea University, Seoul 02841, Korea

³⁰Kyungpook National University, Daegu 702-701, Korea

³¹École Polytechnique Fédérale de Lausanne (EPFL), Lausanne 1015, France

- ³²*P.N. Lebedev Physical Institute of the Russian Academy of Sciences, Moscow 119991, Russia*
³³*Faculty of Mathematics and Physics, University of Ljubljana, 1000 Ljubljana, Slovenia*
³⁴*Ludwig Maximilians University, 80539 Munich, Germany*
³⁵*Luther College, Decorah, Iowa 52101, USA*
³⁶*University of Maribor, 2000 Maribor, Slovenia*
³⁷*Max-Planck-Institut für Physik, 80805 München, Germany*
³⁸*School of Physics, University of Melbourne, Victoria 3010, Australia*
³⁹*University of Miyazaki, Miyazaki 889-2192, Japan*
⁴⁰*Moscow Physical Engineering Institute, Moscow 115409, Russia*
⁴¹*Moscow Institute of Physics and Technology, Moscow Region 141700, Russia*
⁴²*Graduate School of Science, Nagoya University, Nagoya 464-8602, Japan*
⁴³*Kobayashi-Maskawa Institute, Nagoya University, Nagoya 464-8602, Japan*
⁴⁴*Nara Women's University, Nara 630-8506, Japan*
⁴⁵*National Central University, Chung-li 32054, Taiwan*
⁴⁶*National United University, Miao Li 36003, Taiwan*
⁴⁷*Department of Physics, National Taiwan University, Taipei 10617, Taiwan*
⁴⁸*H. Niewodniczanski Institute of Nuclear Physics, Krakow 31-342, Poland*
⁴⁹*Niigata University, Niigata 950-2181, Japan*
⁵⁰*Novosibirsk State University, Novosibirsk 630090, Russia*
⁵¹*Osaka City University, Osaka 558-8585, Japan*
⁵²*Pacific Northwest National Laboratory, Richland, Washington 99352, USA*
⁵³*University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA*
⁵⁴*Theoretical Research Division, Nishina Center, RIKEN, Saitama 351-0198, Japan*
⁵⁵*University of Science and Technology of China, Hefei 230026, China*
⁵⁶*Showa Pharmaceutical University, Tokyo 194-8543, Japan*
⁵⁷*Soongsil University, Seoul 156-743, Korea*
⁵⁸*Stefan Meyer Institute for Subatomic Physics, Vienna 1090, Austria*
⁵⁹*Sungkyunkwan University, Suwon 440-746, Korea*
⁶⁰*School of Physics, University of Sydney, New South Wales 2006, Australia*
⁶¹*Department of Physics, Faculty of Science, University of Tabuk, Tabuk 71451, Saudi Arabia*
⁶²*Tata Institute of Fundamental Research, Mumbai 400005, India*
⁶³*Excellence Cluster Universe, Technische Universität München, 85748 Garching, Germany*
⁶⁴*Department of Physics, Technische Universität München, 85748 Garching, Germany*
⁶⁵*Toho University, Funabashi 274-8510, Japan*
⁶⁶*Department of Physics, Tohoku University, Sendai 980-8578, Japan*
⁶⁷*Earthquake Research Institute, University of Tokyo, Tokyo 113-0032, Japan*
⁶⁸*Department of Physics, University of Tokyo, Tokyo 113-0033, Japan*
⁶⁹*Tokyo Institute of Technology, Tokyo 152-8550, Japan*
⁷⁰*Tokyo Metropolitan University, Tokyo 192-0397, Japan*
⁷¹*University of Torino, 10124 Torino, Italy*
⁷²*Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061, USA*
⁷³*Wayne State University, Detroit, Michigan 48202, USA*
⁷⁴*Yamagata University, Yamagata 990-8560, Japan*
⁷⁵*Yonsei University, Seoul 120-749, Korea*

(Received 19 September 2016; published 29 November 2016)

We report a search for a dark vector gauge boson U' that couples to quarks in the decay chain $D^{*+} \rightarrow D^0 \pi^+$, $D^0 \rightarrow K_S^0 \eta$, $\eta \rightarrow U' \gamma$, $U' \rightarrow \pi^+ \pi^-$. No signal is found and we set a mass-dependent limit on the baryonic fine structure constant of $10^{-3} - 10^{-2}$ in the U' mass range of 290 to 520 MeV/ c^2 . This analysis is based on a data sample of 976 fb $^{-1}$ collected by the Belle experiment at the KEKB asymmetric-energy e^+e^- collider.

DOI: [10.1103/PhysRevD.94.092006](https://doi.org/10.1103/PhysRevD.94.092006)

*Corresponding author.
eunil@hep.korea.ac.kr

The standard model (SM) of particle physics cannot explain the nature of dark matter that is understood to have mostly gravitational effects on visible matter, radiation, and the large-scale structure of the Universe [1–4]. The dark matter can be naturally explained by the introduction of a

weakly interacting particle predicted in the supersymmetric extension of the SM [5]. The absence of observation of any supersymmetric particles in hadron collider experiments [6] motivates studies of new classes of models, commonly referred to as dark models, which introduce new gauge symmetries [7] and predict the existence of new particles that couple weakly to SM particles. Most accelerator-based experiments have focused on the dark photon or dark particles coupling to the SM photons [8], though many dark models suggest a new gauge boson that could couple predominantly to quarks [9,10]. This new dark boson (hereafter referred to as the U' boson, instead of B as is originally proposed in Ref. [9], to avoid confusion with the SM B meson) can be produced from light SM meson decays through $P \rightarrow U'\gamma$ or $V \rightarrow U'P$, where P refers to a pseudoscalar meson (e.g., π^0, η, η') and V to a vector meson (e.g., ω, ϕ). Two recent experimental limits on searches for a dark photon A' via $\pi^0 \rightarrow A'\gamma, A' \rightarrow e^+e^-$ [11] and $\phi \rightarrow A'\gamma, A' \rightarrow e^+e^-$ [12] can be applied to the U' boson search in a model-dependent way to constrain the baryonic fine structure constant $\alpha_{U'} \equiv g_{U'}^2/(4\pi)$, where $g_{U'}$ is the universal gauge coupling between the U' boson and the quarks [10]. There are also limits from $\eta \rightarrow \pi^0\gamma\gamma$ and $\phi \rightarrow \eta\pi^0\gamma$ decays based on their total rate, as well as from the analysis of hadronic $\Upsilon(1S)$ decays [10].

We search for U' bosons decaying to $\pi^+\pi^-$ pairs using $\eta \rightarrow \pi^+\pi^-\gamma$ decays, where η is produced in the decay chain $D^{*+} \rightarrow D^0\pi^+, D^0 \rightarrow K_S^0\eta$ [13]. The kinematics here allows us to suppress the combinatorial background significantly. The decay $U' \rightarrow \pi^+\pi^-$ is expected to have a relatively small branching fraction of 2%–4% [10] but nevertheless provides a very clean signature for a possible dark vector gauge boson. The dominant decay modes are $\pi^0\gamma$ at low U' mass and $\pi^+\pi^-\pi^0$ at higher U' mass; however they suffer from higher combinatorial background and therefore are not used in the analysis. We use the decay $\eta \rightarrow \pi^+\pi^-\pi^0$ to validate our event reconstruction by measuring the branching fraction of $\eta \rightarrow \pi^+\pi^-\gamma$ relative to that of $\eta \rightarrow \pi^+\pi^-\pi^0$.

The data used in this analysis were recorded at the $\Upsilon(nS)$ resonances ($n = 1, \dots, 5$) and 60 MeV below the $\Upsilon(4S)$ resonance with the Belle detector [14] at the e^+e^- asymmetric-energy collider KEKB [15]. The sample corresponds to an integrated luminosity of 976 fb^{-1} . We generated two million Monte Carlo (MC) events [16] each for $\eta \rightarrow \pi^+\pi^-\gamma$, $\eta \rightarrow \pi^+\pi^-\pi^0$, and $\eta \rightarrow U'\gamma \rightarrow \pi^+\pi^-\gamma$ at a particular U' mass selected in the range from 280 to 540 MeV/c^2 in steps of 10 MeV/c^2 (i.e., 58 million events in all). The lifetime of the U' is assumed to be negligible. The U' samples are used to determine the $M(\pi^+\pi^-)$ resolution. The U' signal shape parameters for intermediate U' mass values are determined using spline interpolation.

Except for tracks from K_S^0 decays, we require that the charged tracks originate from the vicinity of the interaction

point (IP) with impact parameters along the beam direction (z axis) and perpendicular to it of less than 4 and 2 cm, respectively. All such charged tracks are required to have at least two associated hits in the silicon vertex detector (SVD), both in the z and perpendicular directions. Such charged tracks are identified as pions or kaons by requiring that the ratio of particle identification likelihoods, $\mathcal{L}_K/(\mathcal{L}_K + \mathcal{L}_\pi)$, constructed using information from the central drift chamber (CDC), time-of-flight scintillation counters, and aerogel threshold Cherenkov counters, be larger or smaller than 0.6, respectively. For both kaons and pions, the efficiencies and misidentification probabilities are 86% and 14%, respectively.

For photon selection, we require the energy of the candidate photon to be greater than 60 MeV (100 MeV) when the candidate photon is reconstructed in the barrel (end cap) calorimeter that covers $32^\circ < \theta < 130^\circ$ ($12^\circ < \theta < 32^\circ$ or $130^\circ < \theta < 157^\circ$) in the polar angle θ with respect to the $+z$ axis. To reject neutral hadrons, the ratio of the energy deposited by a photon candidate in the 3×3 and 5×5 calorimeter arrays centered on the crystal with the largest signal is required to exceed 0.85.

Candidate π^0 mesons are reconstructed from pairs of γ candidates; we require $M_{\gamma\gamma} \in [120, 150] \text{ MeV}/c^2$ and refit γ momenta with the π^0 mass constraint.

Candidate $K_S^0 \rightarrow \pi^+\pi^-$ mesons are reconstructed from two tracks, assumed to be pions, using a neural network technique [17] that uses the following information: the K_S^0 momentum in the laboratory frame; the distance along z between the two track helices at their closest approach; the K_S^0 flight length in the transverse plane; the angle between the K_S^0 momentum and the vector joining the K_S^0 decay vertex to the IP; the angles between the pion momenta and the laboratory-frame direction in the K_S^0 rest frame; the distances of closest approach in the transverse plane between the IP and the two pion helices; and the pion hit information in the SVD and CDC. We also require that the $\pi^+\pi^-$ invariant mass be within $\pm 9 \text{ MeV}/c^2$ (about 3σ in resolution [18]) of the nominal K_S^0 mass [19].

For the $\eta \rightarrow \pi^+\pi^-\gamma$ candidates, we require that the photon not be associated with a π^0 candidate and its transverse momentum be greater than 200 MeV/c to remove $D^{*+} \rightarrow D^+(\rightarrow K_S^0\pi^+\pi^-\pi^+)\gamma$ background. For both $\eta \rightarrow \pi^+\pi^-\gamma$ and $\eta \rightarrow \pi^+\pi^-\pi^0$ candidates, we perform a vertex fit with the two charged pions and require the reduced χ^2 to be less than 10. The efficiency of this requirement is 94%. We require the reconstructed mass of each η candidate to be in the range $[500, 600] \text{ MeV}/c^2$ and refit momenta of its daughters with the constraint of the nominal η mass.

Combinations of a K_S^0 candidate and η candidate are fit to a common vertex and their invariant mass is required to be within $\pm 40 \text{ MeV}/c^2$ of the nominal D^0 mass. The D^0 and π^+ combinations are fitted to the IP, and the mass difference




FIG. 1. Invariant mass of the $K_S^0\eta$ combinations (left) and the $D^* - D^0$ mass difference (right) for $\eta \rightarrow \pi^+\pi^-\gamma$ decays.

$\Delta M_{D^*} = M(K_S^0\eta\pi^+) - M(K_S^0\eta)$ is required to satisfy $\Delta M_{D^*} \in [143, 148] \text{ MeV}/c^2$. To remove the combinatorial background, the momentum of the D^{*+} candidates, measured in the center-of-mass system, is required to be greater than 2.5, 2.6, and 3.0 GeV/c for the data taken below, at, and above the $\Upsilon(4S)$ resonance, respectively. Figure 1 shows the invariant mass of the $K_S^0\eta$ combinations (left) and the mass difference (right) for $\eta \rightarrow \pi^+\pi^-\gamma$ decays after applying all selection criteria described above, except the mass requirements themselves. Figure 2 shows the invariant mass of the $\pi^+\pi^-\gamma$ combinations after all requirements. There are clear peaks of signal events in all distributions; the increase of the background at low masses in the $M(\pi^+\pi^-\gamma)$ distribution is due to the feed-down from the $\eta \rightarrow \pi^+\pi^-\pi^0$ decays when a photon from π^0 is not reconstructed.

To extract the signal yield, we perform a binned maximum likelihood fit to the $M(\pi^+\pi^-\gamma)$ distribution. The fit function is the sum of the signal, the combinatorial background, and the feed-down background components. The signal probability density function (PDF) is the sum of a Gaussian and a bifurcated Gaussian with the ratios of widths fixed from the MC simulation. A linear function is




FIG. 2. Invariant mass distribution of the $\pi^+\pi^-\gamma$ combinations (points with error bars), fit result (solid curve), and combinatorial background component (dashed line) of the fit function. Arrows with lines indicate boundaries of the signal and sideband regions.

used for the combinatorial background PDF. The feed-down contribution is described by a Gaussian with shape parameters fixed from the MC simulation. The confidence level (p -value) of the fit is 12% and the $\eta \rightarrow \pi^+\pi^-\gamma$ signal yield is $N_\eta = 2974 \pm 90$ events. The feed-down yield agrees well with the expectation.

As a cross-check, we measure the ratio of branching fractions $\mathcal{B}(\eta \rightarrow \pi^+\pi^-\gamma)/\mathcal{B}(\eta \rightarrow \pi^+\pi^-\pi^0)$. The fit to the $\pi^+\pi^-\pi^0$ invariant mass distribution is similar to the one described above, except that the combinatorial background is described by a second-order polynomial and there is no feed-down background. The reconstruction efficiencies, determined from the MC simulation, are $\varepsilon(\pi^+\pi^-\gamma) = 5.1\%$ and $\varepsilon(\pi^+\pi^-\pi^0) = 4.8\%$. The measured ratio of branching fractions, 0.185 ± 0.007 , where the uncertainty is statistical only, is in good agreement with the world-average value of 0.184 ± 0.004 [19].

We define the η signal region as $M(\pi^+\pi^-\gamma) \in [535.5, 560.5] \text{ MeV}/c^2$, and the sideband regions used for background subtraction as $M(\pi^+\pi^-\gamma) \in [520.0, 532.5]$ or $[563.5, 576.0] \text{ MeV}/c^2$. The $M(\pi^+\pi^-)$ distribution for the background-subtracted η signal is shown in Fig. 3.

To describe the $M(\pi^+\pi^-)$ distribution, we use an expression of the differential decay rate based on low-energy quantum chromodynamics phenomenology [20,21] using a combination of chiral perturbation theory and dispersive analysis,

$$\frac{d\Gamma}{ds} \propto |P(s)F_V(s)|^2(m_\eta^2 - s)^3 s(1 - 4m_\pi^2/s)^{3/2}, \quad (1)$$

where $s \equiv M(\pi^+\pi^-)^2$, $P(s)$ is a reaction-specific perturbative part, and $F_V(s)$ is the pion vector form factor. We use $|P(s)| = 1 + (1.89 \pm 0.64)s$ [20] and $|F_V(s)| = 1 + (2.12 \pm 0.01)s + (2.13 \pm 0.01)s^2 + (13.80 \pm 0.14)s^3$




FIG. 3. $\pi^+\pi^-$ invariant mass distribution from the $\eta \rightarrow \pi^+\pi^-\gamma$ signal (points with error bars), the fitted differential decay rate described in Eq. (1) (solid curve), and an example U' signal at a mass of $400 \text{ MeV}/c^2$ from $\eta \rightarrow U'\gamma$, $U' \rightarrow \pi^+\pi^-$ (histogram with arbitrary normalization).

[21] (s in GeV^2/c^4). The numerical values and the uncertainties of the expansion coefficients of $|P(s)|$ and $|F_V(s)|$ are taken from fits to data of $\eta^{(\prime)} \rightarrow \pi^+\pi^-\gamma$ decays. We multiply the $d\Gamma/ds$ expression from Eq. (1) by the reconstruction efficiency. The efficiency as a function of $M(\pi^+\pi^-)$ is approximately flat but drops to 0 at the kinematic limit of m_η . The fit results are presented in Fig. 3. Equation (1) describes the $M(\pi^+\pi^-)$ distribution well, and the confidence level of the fit is 95%.

We add the U' signal to the above fit function and perform fits while fixing the U' mass at a value between 290 and 520 MeV/c^2 in steps of 1 MeV/c^2 . The U' signal is described by the sum of two Gaussians. The signal resolution of the core Gaussian is about 1 MeV/c^2 near the $2m_\pi$ threshold and 2 MeV/c^2 at the m_η kinematic limit. An example of the U' signal with the mass of 400 MeV/c^2 and arbitrary normalization is shown in Fig. 3. We do not find a significant U' signal at any mass value. The typical uncertainty in the U' yield $N_{U'}$ is $\mathcal{O}(1-10)$ events.

We express the baryonic fine structure constant $\alpha_{U'}$ using the equation for the partial width ratio $\Gamma(\eta \rightarrow U'\gamma)/\Gamma(\eta \rightarrow \gamma\gamma)$ from Ref. [10] as

$$\alpha_{U'} = \left[\frac{\alpha}{2} \left(1 - \frac{m_{U'}^2}{m_\eta^2} \right)^{-3} \left| \mathcal{F}(m_{U'}^2) \right|^{-2} \frac{1}{\mathcal{B}(U' \rightarrow \pi^+\pi^-)} \right] \times \left[\frac{\Gamma(\eta \rightarrow \pi^+\pi^-\gamma)}{\Gamma(\eta \rightarrow \gamma\gamma)} \right] \left[\frac{\Gamma(\eta \rightarrow U'\gamma \rightarrow \pi^+\pi^-\gamma)}{\Gamma(\eta \rightarrow \pi^+\pi^-\gamma)} \right], \quad (2)$$

where α is the electromagnetic fine structure constant. The first factor in Eq. (2), which is purely theoretical, contains the phase space, the form factor $\mathcal{F}(m_{U'}^2)$, and the branching fraction of $U' \rightarrow \pi^+\pi^-$ decay. The branching fraction is about 2%–4%, as computed from formulas provided in Ref. [10] and references therein. The second factor is obtained from the latest measurements [19]. The third factor is determined from the η and U' yields and reconstruction efficiencies ($N_{U'}/\varepsilon(\eta \rightarrow U'\gamma \rightarrow \pi^+\pi^-\gamma)/N_\eta/\varepsilon(\eta \rightarrow \pi^+\pi^-\gamma)$).

To estimate the systematic uncertainties in the $\eta \rightarrow \pi^+\pi^-\gamma$ and $\eta \rightarrow U'\gamma \rightarrow \pi^+\pi^-\gamma$ yields, we change the parametrization of the combinatorial background in the $M(\pi^+\pi^-\gamma)$ fit from a first- to a second-order polynomial and account for the background nonlinearity while subtracting the sidebands. The change in the η yield is at the 1% level, while the change in the U' yield is negligible. The systematic effect due to the uncertainties of the expansion coefficients in $|P(s)|$ and $|F_V(s)|$ is negligible in the U' yield. The systematic uncertainty in the ratio of the reconstruction efficiencies $\varepsilon(\eta \rightarrow U'\gamma \rightarrow \pi^+\pi^-\gamma)/\varepsilon(\eta \rightarrow \pi^+\pi^-\gamma)$ is conservatively estimated to be 4% (1% per track and 3% per photon). The total systematic uncertainties are estimated by adding the above contributions in quadrature.




FIG. 4. Computed 95% upper limit on the baryonic fine structure constant $\alpha_{U'}$ as a function of the unknown U' mass (solid curve).

Using Eq. (2), we set a 95% confidence level upper limit on $\alpha_{U'}$ using the Feldman-Cousins approach [22], adding the statistical and systematic uncertainties in quadrature. The upper limit as a function of the U' boson mass is shown in Fig. 4. Considering other results in this mass region, we find that our limit is stronger than that from a model-dependent analysis [10] of the $\phi \rightarrow e^+e^-\gamma$ decays [12] for $m_{U'} > 450 \text{ MeV}/c^2$, but weaker than the limit based on the $\eta \rightarrow \pi^0\gamma\gamma$ total rate [10]. Recently, we learned that the data set in Ref. [23] contains many more $\eta \rightarrow \pi^+\pi^-\gamma$ decays and can provide a more stringent limit on $\alpha_{U'}$ in future.

To conclude, we perform a search for a dark vector gauge boson U' that couples to quarks [10], using the decay chain $D^{*+} \rightarrow D^0\pi^+$, $D^0 \rightarrow K_S^0\eta$, $\eta \rightarrow U'\gamma$, $U' \rightarrow \pi^+\pi^-$. Our results limit the baryonic fine structure constant $\alpha_{U'}$ to below 10^{-3} – 10^{-2} at 95% confidence level over the U' mass range 290 to 520 MeV/c^2 . This is the first search for U' in the $\pi^+\pi^-$ mode. We find that our limit is stronger than that from a model-dependent analysis [10] of the $\phi \rightarrow e^+e^-\gamma$ decays [12] for $m_{U'} > 450 \text{ MeV}/c^2$, but weaker than the limit based on the $\eta \rightarrow \pi^0\gamma\gamma$ total rate [10].

We thank the KEKB group for the excellent operation of the accelerator; the KEK cryogenics group for the efficient operation of the solenoid; and the KEK computer group, the National Institute of Informatics, and the PNNL/EMSL computing group for valuable computing and SINET4 network support. We acknowledge support from the Ministry of Education, Culture, Sports, Science, and Technology (MEXT) of Japan, the Japan Society for the Promotion of Science (JSPS), and the Tau-Lepton Physics Research Center of Nagoya University; the Australian Research Council; Austrian Science Fund under Grants No. P 22742-N16 and No. P 26794-N20; the National Natural Science Foundation of China under Contracts No. 10575109, No. 10775142, No. 10875115, No. 11175187, No. 11475187 and No. 11575017; the

Chinese Academy of Science Center for Excellence in Particle Physics; the Ministry of Education, Youth and Sports of the Czech Republic under Contract No. LG14034; the Carl Zeiss Foundation, the Deutsche Forschungsgemeinschaft, the Excellence Cluster Universe, and the VolkswagenStiftung; the Department of Science and Technology of India; the Istituto Nazionale di Fisica Nucleare of Italy; the World Class University program of the Ministry of Education, National Research Foundation (NRF) of Korea Grants No. 2011-0029457, No. 2012-0008143, No. 2012R1A1A2008330, No. 2013R1A1A3007772, No. 2014R1A2A2A01005286, No. 2014R1A2A2A01002734, No. 2015R1A2A2A01003280 and No. 2015H1A2A1033649; the Basic Research Lab program under NRF Grant No. KRF-2011-0020333, Center for Korean J-PARC Users, Grant No. NRF-2013K1A3A7A06056592; the Brain Korea 21-Plus program

and Radiation Science Research Institute; the Polish Ministry of Science and Higher Education and the National Science Center; the Ministry of Education and Science of the Russian Federation and the Russian Foundation for Basic Research; the Slovenian Research Agency; Ikerbasque, Basque Foundation for Science and the Euskal Herriko Unibertsitatea (UPV/EHU) under program UFI 11/55 (Spain); the Swiss National Science Foundation; the Ministry of Education and the Ministry of Science and Technology of Taiwan; and the U.S. Department of Energy and the National Science Foundation. This work is supported by a Grant-in-Aid from MEXT for Science Research in a Priority Area (“New Development of Flavor Physics”) and from JSPS for Creative Scientific Research (“Evolution of Tau-lepton Physics”). E. W. acknowledges a partial NRF grant of Korea Grant No. NRF-2011-0030865 and Korea University Future Research grant, and thanks B. R. Ko for his suggestion on an inclusive version of this analysis.

-
- [1] F. Zwicky, *Astrophys. J.* **86**, 217 (1937).
 [2] D. Clowe, A. Gonzalez, and M. Markevitch, *Astrophys. J.* **604**, 596 (2004).
 [3] E. Komatsu *et al.*, *Astrophys. J. Suppl. Ser.* **180**, 330 (2009).
 [4] G. R. Blumenthal, S. M. Faber, J. R. Primack, and M. J. Rees, *Nature (London)* **311**, 517 (1984).
 [5] B. Kane and M. Shifman, *The Supersymmetric World* (World Scientific, Singapore, 2000).
 [6] G. Aad *et al.* (ATLAS Collaboration), *J. High Energy Phys.* **10** (2015) 134; S. Chatrchyan *et al.* (CMS Collaboration), *Phys. Rev. D* **88**, 052017 (2013).
 [7] B. Holdom, *Phys. Lett.* **166B**, 196 (1986); P. Fayet, *Phys. Rev. D* **75**, 115017 (2007).
 [8] G. Agakishiev *et al.* (HADES Collaboration), *Phys. Lett. B* **731**, 265 (2014); I. Jaegle *et al.* (Belle Collaboration), *Phys. Rev. Lett.* **114**, 211801 (2015); J. P. Lees *et al.* (BABAR Collaboration), *Phys. Rev. Lett.* **108**, 211801 (2012); S. Giovannella *et al.* (KLOE Collaboration), *J. Phys. Conf. Ser.* **335**, 012067 (2011).
 [9] A. E. Nelson and N. Tetradis, *Phys. Lett. B* **221**, 80 (1989).
 [10] S. Tulin, *Phys. Rev. D* **89**, 114008 (2014).
 [11] P. Adlarson *et al.* (WASA-at-COSY Collaboration), *Phys. Lett. B* **726**, 187 (2013).
 [12] D. Babusci *et al.* (KLOE-2 Collaboration), *Phys. Lett. B* **720**, 111 (2013).
 [13] Throughout this paper, inclusion of the charge-conjugate decay is implied unless stated otherwise.
 [14] A. Abashian *et al.* (Belle Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **479**, 117 (2002); also see the detector section in J. Brodzicka *et al.*, *Prog. Theor. Exp. Phys.* **2012**, 4D001 (2012).
 [15] S. Kurokawa and E. Kikutani, *Nucl. Instrum. Methods Phys. Res., Sect. A* **499**, 1 (2003), and other papers included in this volume; T. Abe *et al.*, *Prog. Theor. Exp. Phys.* **2013**, 03A001 (2013) and the following articles up to 03A011.
 [16] $e^+e^- \rightarrow c\bar{c}$ events are generated with PYTHIA [T. Sjöstrand, *Comput. Phys. Commun.* **135**, 238 (2001)]; particle decay is simulated with EVTGEN [D. J. Lange, *Nucl. Instrum. Methods Phys. Res., Sect. A* **462**, 152 (2001)]; the detector response is simulated with GEANT 3.21 (R. Brun *et al.*, CERN Report No. DD/EE/84-1, 1984).
 [17] M. Feindt and U. Kerzel, *Nucl. Instrum. Methods Phys. Res., Sect. A* **559**, 190 (2006).
 [18] E. Won *et al.* (Belle Collaboration), *Phys. Rev. D* **80**, 111101(R) (2009).
 [19] K. A. Olive *et al.* (Particle Data Group), *Chin. Phys. C* **38**, 090001 (2014).
 [20] P. Adlarson *et al.* (WASA-at-COSY Collaboration), *Phys. Lett. B* **707**, 243 (2012).
 [21] F. Stollenwerk, C. Hanhart, A. Kupsc, U.-G. Meißner, and A. Wirzba, *Phys. Lett. B* **707**, 184 (2012).
 [22] G. J. Feldman and R. D. Cousins, *Phys. Rev. D* **57**, 3873 (1998).
 [23] D. Babusci *et al.* (KLOE/KLOE-2 Collaboration), *Phys. Lett. B* **718**, 910 (2013).