


DESY 76/35  
July 1976



## Dilations and Interaction

by

D. Buchholz and K. Fredenhagen

*in: Institut für Theoretische Physik der Universität Hamburg*

### Abstract

As a consequence of the geometrical features of dilations, massless particles do not interact in a local, dilationally invariant quantum theory. This result also holds in models in which dilations are only an asymptotically visible symmetry of the S-matrix.

2 HAMBURG 52 . NOTKESTIEG 1

To be sure that your preprints are promptly included in the  
HIGH ENERGY PHYSICS INDEX ,  
send them to the following address ( if possible by air mail ) :

DESY  
Bibliothek  
2 Hamburg 52  
Notkestieg 1  
Germany

# Dilations and Interaction

by

Detlev Buchholz and Klaus Fredenhagen

II. Institut für Theoretische Physik der Universität Hamburg

## Abstract:

As a consequence of the geometrical features of dilations massless particles do not interact in a local, dilationally invariant quantum theory. This result also holds in models in which dilations are only an asymptotically visible symmetry of the S-matrix.

## 1. Introduction and Main Results

The conventional argument showing that massless particles do not interact in a local, dilationally invariant quantum theory is in the simplest case the following one (see e.g. [1]): suppose  $\phi$  is a scalar Wightman field transforming under dilations according to

$$D(\lambda) \phi(\lambda^{-1}x) D(\lambda)^{-1} = \lambda^d \cdot \phi(x). \quad (1)$$

If  $\phi$  has a non-vanishing matrix element between the vacuum and a massless one-particle state,  $d$  can only be one. Then  $\phi$  has canonical dimension and this implies that it is a free field. This reasoning is quite correct. However, since the argument depends upon the existence of a field  $\phi$  with the special properties mentioned above the conclusion appears to us to be rather premature. First, there is no physical reason to rule out ab initio all models in which the basic fields do not transform like a finite dimensional representation under dilations. And secondly, even if the fields transform in this way, it could happen that they do not interpolate between the vacuum and the massless one-particle states. In general one should only expect that suitable polynomials in the fields have this property. It is the aim of the present note to close these apparent loopholes. Using only the geometrical features of dilations and the basic properties of local field theory, we give a fairly general argument confirming the above no-go theorem.

The setting used for the analysis may be sketched as follows: we deal with an irreducible field algebra  $\mathcal{F}$  of bounded operators acting on a Hilbert space  $\mathcal{H}$ .  $\mathcal{F}$  is generated by a net  $\mathcal{O} \rightarrow \mathcal{F}(\mathcal{O})$  of local algebras attached to the regions  $\mathcal{O}$  of Minkowski space. We may forego here a formal specification of the usual structural assumptions on the theory like locality, covariance, spectrum condition and uniqueness of the vacuum. (For a detailed discussion see for example [2]). In addition to these familiar properties we require that there is a continuous, unitary representation  $\lambda \rightarrow D(\lambda)$  of the multiplicative group of the positive reals in  $\mathcal{H}$ . The operators  $D(\lambda)$ , the dilations, satisfy

$$D(\lambda) \mathcal{U}(x) = \mathcal{U}(\lambda x) D(\lambda) \quad \text{and} \quad D(\lambda) \mathcal{U}(\Lambda) = \mathcal{U}(\Lambda) D(\lambda) \quad (2a)$$

where  $x = (x, \underline{x}) \rightarrow \mathcal{U}(x)$  are the translations and  $\Lambda \rightarrow \mathcal{U}(\Lambda)$  the Lorentz

transformations. Moreover, the dilations  $D(\lambda)$  induce automorphisms of the field algebra  $\mathcal{F}$  with appropriate geometric properties:

$$D(\lambda) \mathcal{F}(\phi) D(\lambda)^{-1} = \mathcal{F}(\lambda \cdot \phi). \quad (2b)$$

These rather general assumptions suffice to prove the following statement:

If there exist massless particles in the model, (i.e. a family of subspaces  $\mathcal{H}_i^{(0)} \subset \mathcal{H}$  on which the unitaries  $U(\infty)$ ,  $U(1)$  act like an irreducible representation of the Poincaré group with mass zero and helicity  $S_i$ ), then the S-matrix for these particles is trivial.

Our interest in this problem arose in discussions with R. Haag on supersymmetric field theories. In a recent article Haag, Lopuszanski and Sohnius have analyzed the structure of all possible supersymmetries of the S-matrix [3]. They found out that in a pure S-matrix formalism there is essentially only one way of a complete fusion between internal and geometrical symmetries, including dilations. Since such a structure looks very promising from the point of view of physics, one may ask whether it can be embedded into a conventional field theoretical setting. As a consequence of our analysis the answer to this question is negative: if the theory is to describe collisions of massless particles and if dilations are to be a proper, unbroken symmetry one has to abandon some of the usual field theoretical assumptions. At present it is unclear how the assumptions have to be modified and we refrain from speculations. However, we want to emphasize that even in a modified scheme the local observables (the currents etc.) should have a structure similar to that of  $\mathcal{F}$  given above. What may then be learned from our analysis is that massless particles in the vacuum sector of the observable algebra do not interact. It is therefore unlikely that particles like the photon and the  $\eta'$ -meson (both of which carry the charge quantum numbers of the vacuum) can be incorporated into such a scheme. This apparently restricts the possible range of application of these models to weak interaction physics.

## 2. The Proof

The central idea of the proof is very simple: we derive an asymptotic expansion for the function  $\lambda \rightarrow D(\lambda) A D(\lambda)^{-1}$  at  $\lambda=0$ , where  $A$  is a suitable local operator taken from  $\mathcal{F}$ . It turns out that

$$D(\lambda) A D(\lambda)^{-1} = (\Omega, A \Omega) \cdot 1 + \lambda \cdot \phi + o(\lambda) \quad (3)$$

where this expansion is understood in the sense of operator valued distributions;  $\Omega$  denotes the vector representing the vacuum and  $\phi$  is some local field. Now the crucial point is that if  $\phi$  is not zero it creates a massless particle from the vacuum. It then follows from Huyghens' principle (i.e. the timelike commutation relations between local and asymptotic fields given in [4]) that the S-matrix of this particle can only be trivial.

Unfortunately, there are models in which, for kinematical reasons, all local operators  $A$  give rise to a vanishing  $\phi$ . However, this defect can be cured by a slight modification of the above expansion: dilating and boosting the operator  $A$  simultaneously one arrives at an expression similar to (3), but with a non-trivial  $\phi$ . To abbreviate the argument we confine our attention to models involving only one type of massless particles with helicity  $s=0$ . But we shall give a brief outline of how to proceed in more complicated situations.

Now let  $A$  be any operator from  $\mathcal{F}$  which is localized in a bounded region  $\mathcal{O} \subset \mathbb{R}^4$ . We regularize  $A$  according to

$$A_\varphi = \int dt \varphi(t) U(t) A U(t)^{-1} \quad (4)$$

where  $t \rightarrow U(t)$  are the time translations.  $\varphi(t)$  is a test function with compact support which has a Fourier transform  $\tilde{\varphi}(\omega)$  with a twofold zero at  $\omega=0$ . The smoothed operator  $A_\varphi$  is still local and we get the following bound on its two-point function:

Lemma 1:

Let  $\Delta \rightarrow E(\Delta)$  be the spectral projections of the mass operator  $M = (p^2)^{1/2}$  where  $\Delta \subset \mathbb{R}^+$  is any Borel set of mass values. Then

$$|(A_\varphi \Omega, E(\Delta) U(x) A_\varphi \Omega)| \leq c \cdot (1+|x|)^{-1} \cdot \{ \|E(\Delta) A \Omega\|^2 + \|E(\Delta) A^* \Omega\|^2 \}$$

where the constant  $c$  depends neither on  $x$  nor on  $\Delta$ .

Proof:

Using the methods of the Jost-Lehmann-Dyson representation one can show

that the function

$$h_\Delta(x) = (A_\varphi \Omega, E(\Delta) U(x) A_\Omega) - (A^* \Omega, E(\Delta) U(-x) A^* \Omega)$$

vanishes in the spacelike complement of some bounded region  $\mathcal{O}_1$  which depends only on the localisation region  $\mathcal{O}$  of  $A$  (see e.g. [5, Lemma 6.2]). Now if one puts  $\psi(t) = \int ds \tilde{\varphi}(s) \varphi(s+t)$  one gets owing to the spectrum condition

$$\begin{aligned} & (A_\varphi \Omega, E(\Delta) U(x) A_\varphi \Omega) \\ &= \int dt \psi(t) (A_\Omega, E(\Delta) U(t, x) A_\Omega) = \int dt \psi^*(t) (A_\Omega, E(\Delta) U(t, x) A_\Omega) \\ &= \int dt \psi^*(t) \{ (A_\Omega, E(\Delta) U(t, x) A_\Omega) - (A^* \Omega, E(\Delta) U(-t, x) A^* \Omega) \} = \int dt \psi^*(t) h_\Delta(t, x) \end{aligned}$$

where

$$\psi^*(t) = (2\pi)^{-1/2} \int_0^\infty d\omega \tilde{\varphi}(\omega) e^{-i\omega t} = \int_0^\infty d\omega |\tilde{\varphi}(\omega)|^2 e^{-i\omega t}.$$

Since  $|\tilde{\varphi}(\omega)|^2$  is a test function with a fourfold zero at  $\omega = 0$  it is easy to verify that  $\psi^*(t)$  is continuous and  $|\psi^*(t)| \leq c \cdot (1+|t|)^{-4}$ .

Taking the support properties of  $h_\Delta(x)$  into account, one arrives at

$$\begin{aligned} |(A_\varphi \Omega, E(\Delta) U(x) A_\varphi \Omega)| &\leq \int_{|t| \geq |x| - R} dt |\psi^*(t)| \cdot |h_\Delta(t, x)| \\ &\leq c \cdot \int_{|t| \geq |x| - R} dt (1+|t|)^{-4} \cdot \{ \|E(\Delta) A_\Omega\|^2 + \|E(\Delta) A^* \Omega\|^2 \} \end{aligned}$$

where  $R$  is some length which depends only on  $A$ . From this inequality the statement of the lemma follows at once. ■

We take now the operator  $A_\varphi$  and carry out the following manipulations: first we dilate it, then we boost it in a fixed direction and finally we smear it in the two remaining spatial directions. For the boosts we take those in the  $x_1$ -direction:

$$K_\lambda = \begin{pmatrix} \frac{1}{2}(\lambda + \lambda^{-1}) & \frac{1}{2}(\lambda - \lambda^{-1}) & & \\ \frac{1}{2}(\lambda - \lambda^{-1}) & \frac{1}{2}(\lambda + \lambda^{-1}) & & \\ & & 1 & \\ & & & 1 \end{pmatrix}, \quad \lambda > 0. \quad (5)$$

Then if  $\frac{x^\perp}{2} = (0, x_2, x_3)$  denotes the projection of  $x$  onto the  $x_2, x_3$ -plane and if  $\frac{dx^\perp}{2} = dx_2 dx_3$  we set

$$B_\lambda = \lambda^{-1} \cdot \int d^2 x^\perp f(x^\perp) U(x^\perp) U(K_\lambda) D\omega A_\varphi D\omega^{-1} U(K_\lambda)^{-1} U(x^\perp)^{-1} \quad (6)$$

where  $f(x^\perp)$  is any test function with compact support. To begin with we examine the localisation properties of  $B_\lambda$ : since  $A_\varphi$  is localized in some bounded region  $\mathcal{O}$  it follows from (6) that  $B_\lambda$  is localized in  $\{\lambda \cdot K_\lambda \mathcal{O} + \text{supp } f\}$ . Now  $\lim_{\lambda \rightarrow 0} \lambda \cdot K_\lambda = P$  where  $P$  is the projection onto the ray  $(a, -a, 0, 0)$ ,  $a \in \mathbb{R}$ . Therefore the operators  $B_\lambda$  are, for sufficiently small  $\lambda$ , localized in a fixed bounded region  $\mathcal{O}_1$ . The next step is to show that the sequence  $B_\lambda \Omega$  converges to a (possibly zero) one-particle state in the limit of small  $\lambda$ :

Proposition 2:

Let  $B_\lambda$  be the operator defined in relation (6). Then the weak limit  $\omega - \lim_{\lambda \rightarrow 0} B_\lambda \Omega$  exists and is an element of  $\mathcal{H}_1$ .

Proof:

The proof of this assertion is based on Lemma 1. Since  $U(K_\lambda)$  commutes with  $U(x^\perp)$  and  $E(\Delta)$  we may write

$$\|E(\Delta) B_\lambda \Omega\|^2 = \lambda^{-2} \cdot \int d^2 x^\perp \int d^2 y^\perp \bar{f}(x^\perp) f(y^\perp) (A_\varphi \Omega, E(\Delta) U(x^\perp) U(y^\perp) A_\varphi \Omega)$$

where we have made use of relation (2a). If we set  $g(x^\perp) = \int d^2 y^\perp \bar{f}(y^\perp) f(x^\perp + y^\perp)$  we get, using Lemma 1,

$$\begin{aligned} \|E(\Delta) B_\lambda \Omega\|^2 &= \lambda^{-2} \cdot \int d^2 x^\perp g(x^\perp) (A_\varphi \Omega, E(\Delta) U(x^\perp) A_\varphi \Omega) \\ &\leq \sup_{y^\perp} |g(y^\perp)| \cdot c \int d^2 x^\perp (1+|x^\perp|)^{-4} \cdot \{ \|E(\Delta) A_\Omega\|^2 + \|E(\Delta) A^* \Omega\|^2 \}. \end{aligned}$$

Putting  $\Delta = \mathbb{R}^+$  it follows that the sequence  $B_\lambda \Omega$  is uniformly bounded in  $\lambda$ . Putting  $\Delta_c = [a, b]$  where  $0 < a \leq b < \infty$  it follows that

$\lim_{\lambda \rightarrow 0} \|E(\Delta_c) B_\lambda \Omega\| = 0$  because the continuity properties of the spectral resolution imply  $\lim_{\lambda \rightarrow 0} \|E(\Delta_c) \Phi\| = 0$  for every vector  $\Phi \in \mathcal{H}$ .

Thus the sequence  $B_\lambda \Omega$  converges weakly to zero on the orthogonal complement of the one-particle space  $\mathcal{H}_1$ . It remains to establish its convergence on  $\mathcal{H}_1$ .

Since we are dealing with only one type of massless particles with helicity  $s = 0$  we may identify the one particle states  $\Psi \in \mathcal{H}_1$  with their momentum space wave functions  $\Psi(p)$  in  $L^2(\mathbb{R}^3, \frac{d^3 p}{2|p|})$ .

The dilations and Poincaré transformations act on these functions as follows:

$$(D(\lambda) \Psi)(p) = \lambda \Psi(\lambda p), \quad (U(t, x) \Psi)(p) = e^{i(t|p| - x \cdot p)} \Psi(p) \quad (7a)$$

and

$$(U(\Lambda) \Psi)(p) = \Psi(\Lambda^{-1} \circ p) \quad (7b)$$

where  $\Lambda^{-1} \circ \underline{p}$  denotes the spatial components of the 4-vector  $\Lambda^{-1}(|\underline{p}|, \underline{p})$ . What is crucial now is that the wave function  $(A_q \Omega)(\underline{p})$  of the one-particle state  $E(\{0\}) A_q \Omega$  is continuous in  $\underline{p}$  if  $A_q$  is the operator defined in relation (4). To verify this we fix a set  $\mathcal{L}$  of Lorentz transformations  $\Lambda$  which are close to the identity  $I$ , e.g.  $\mathcal{L} = \{\Lambda: \|\Lambda - I\| \leq \frac{1}{2}\}$ . Since  $A$  is local it is obvious that all operators  $(U(\Lambda)^{-1} A U(\Lambda) - A)$ ,  $\Lambda \in \mathcal{L}$  are localized in a bounded region  $\mathcal{O}$  of configuration space. Therefore we get the estimate, using relation 7 and Lemma 1,

$$\begin{aligned} & \frac{\pi}{|\underline{p}|} |\tilde{\varphi}(\underline{p})|^2 \cdot |(A\Omega)(\Lambda \circ \underline{p}) - (A\Omega)(\underline{p})|^2 = \frac{1}{2|\underline{p}|} \cdot |((U(\Lambda)^{-1} A U(\Lambda) - A)_q \Omega)(\underline{p})|^2 \\ & = (2\pi)^{-3} \int d\underline{x} e^{i\underline{x} \cdot \underline{p}} \cdot ((U(\Lambda)^{-1} A U(\Lambda) - A)_q \Omega, E(\{0\}) U(\underline{x}) (U(\Lambda)^{-1} A U(\Lambda) - A)_q \Omega) \\ & \leq c \cdot \{ \| (U(\Lambda) - 1) E(\{0\}) A \Omega \|^2 + \| (U(\Lambda) - 1) E(\{0\}) A^* \Omega \|^2 \}, \end{aligned}$$

and this inequality holds for all  $\Lambda \in \mathcal{L}$  and  $\underline{p} \in \mathbb{R}^3$ . Since we may take for  $\tilde{\varphi}$  a test function which has a zero only at the origin, it is evident that  $\lim_{\Lambda \rightarrow I} (A\Omega)(\Lambda \circ \underline{p}) = (A\Omega)(\underline{p})$  for  $\underline{p} \neq 0$ . But this shows that  $(A\Omega)(\underline{p})$  and therefore also  $(A_q \Omega)(\underline{p}) = (2\pi)^{1/2} \tilde{\varphi}(|\underline{p}|) (A\Omega)(\underline{p})$  are continuous at  $\underline{p} \neq 0$  because for every sequence  $\underline{p}_n$  converging to  $\underline{p}$  we can specify a sequence of Lorentz transformations  $\Lambda_n$  such that for sufficiently large  $n$ :  $\Lambda_n \circ \underline{p} = \underline{p}_n$  and  $\lim_{n \rightarrow \infty} \Lambda_n = I$ . In order to establish the continuity of  $(A_q \Omega)(\underline{p})$  at  $\underline{p} = 0$  we estimate

$$\frac{1}{2|\underline{p}|} |(A_q \Omega)(\underline{p})|^2 = (2\pi)^{-3} \int d\underline{x} e^{i\underline{x} \cdot \underline{p}} \cdot (A_q \Omega, E(\{0\}) U(\underline{x}) A_q \Omega) \leq c.$$

This bound holds uniformly for all  $\underline{p} \in \mathbb{R}^3$  and implies  $\lim_{\underline{p} \rightarrow 0} (A_q \Omega)(\underline{p}) = 0$ . Now we are almost finished: using relation (7) we get for the wavefunction  $(B_\lambda \Omega)(\underline{p})$  of the one-particle state  $E(\{0\}) B_\lambda \Omega$

$$(B_\lambda \Omega)(\underline{p}) = 2\pi \cdot \tilde{f}(\underline{p}^\perp) \cdot (A_q \Omega)(\lambda K_\lambda^{-1} \underline{p})$$

where

$$\tilde{f}(\underline{p}^\perp) = (2\pi)^{-1} \int d\underline{x}^\perp e^{-i\underline{x}^\perp \cdot \underline{p}^\perp} f(\underline{x}^\perp) \text{ with } \underline{p}^\perp = (0, p_2, p_3).$$

An easy calculation shows that  $\lim_{\lambda \rightarrow 0} \lambda \cdot K_\lambda^{-1} \circ \underline{p} = \frac{1}{2} (|\underline{p}| + p_1) \underline{e}_1$  where  $\underline{e}_1 = (1, 0, 0)$ . Taking into account the continuity of  $\underline{p} \rightarrow (A_q \Omega)(\underline{p})$  we get

$$\lim_{\lambda \rightarrow 0} (B_\lambda \Omega)(\underline{p}) = 2\pi \cdot \tilde{f}(\underline{p}^\perp) \cdot (A_q \Omega)(\frac{1}{2} [|\underline{p}| + p_1] \underline{e}_1). \quad (8)$$

It then follows from the bounded convergence theorem that the limit  $\lim_{\lambda \rightarrow 0} \int \frac{d\underline{p}}{2|\underline{p}|} \Psi(\underline{p}) (B_\lambda \Omega)(\underline{p})$  exists for all test functions  $\Psi(\underline{p})$  with compact support. These functions are dense in  $L^2(\mathbb{R}^3, \frac{d\underline{p}}{2|\underline{p}|})$  and since the vectors  $B_\lambda \Omega$  are uniformly bounded in  $\lambda$  we conclude that the weak limit  $\omega - \lim_{\lambda \rightarrow 0} E(\{0\}) B_\lambda \Omega$  exists. This finishes the proof of the statement. ■

#### Remark:

Using the above proposition and the localisation properties of the operators  $B_\lambda$ , one can show that  $\lim_{\lambda \rightarrow 0} B_\lambda$  also exists on a dense set of vectors in  $\mathcal{H}$ .

The wave function of the one-particle state  $\omega - \lim_{\lambda \rightarrow 0} B_\lambda \Omega$  is given by the right hand side of equation (8). It is therefore easy to specify a local operator  $A$  for which this vector is non-trivial: pick for example a one-particle state  $\Phi \in \mathcal{H}_1$  which is invariant under spatial rotations  $R \rightarrow U(R)$ . Since  $\mathcal{F}$  is irreducible there exists a local operator  $A_1 \in \mathcal{F}$  such that the matrix element  $(\Phi, A_1 \Omega)$  is not zero. The operator  $A = \int d\mu(R) U(R) A_1 U(R)^{-1}$ , where  $d\mu(R)$  is the Haar measure on the group of rotations, then has the desired property. If one takes  $A_1$  hermitian and the functions  $\varphi, f$  real one can even arrange for the approximating operators  $B_\lambda$  to be hermitian.

In the remainder of this section we shall show that the existence of an operator sequence  $B_\lambda$  with properties mentioned above implies that the massless particles do not scatter. The argument is based on results recently derived in [4] in the context of collision theory for massless particles. We recapitulate the main facts briefly: as in the massive case, there are collision states  $\Phi_1^{in} \dots \Phi_n^{in}$  and  $\Phi_1^{out} \dots \Phi_n^{out}$  in  $\mathcal{H}$  corresponding to incoming and outgoing configurations  $\Phi_1, \dots, \Phi_n \in \mathcal{H}_1$  of massless particles. These vectors have the familiar Fock structure known from a free theory. They can be generated from the vacuum  $\Omega$  with the aid of asymptotic fields  $A^{in}$  and  $A^{out}$ . The bounded functions of the fields which are localized in a region  $\mathcal{O}$  constitute the local asymptotic field algebras  $\mathcal{F}^{in}(\mathcal{O})$  and  $\mathcal{F}^{out}(\mathcal{O})$  respectively. They have commutation relations with the basic fields which may be interpreted as the field theoretical version of Huyghens' principle: if  $\mathcal{O}$  is any bounded region and if  $\mathcal{O}_+, \mathcal{O}_-$  are two regions which have a positive and negative timelike distance from  $\mathcal{O}$  then

$$[F, F_+^{in}] = 0 \quad \text{and} \quad [F, F_-^{out}] = 0 \quad (9)$$

for arbitrary  $F \in \mathcal{F}(0)$ ,  $F_+^{\text{in}} \in \mathcal{F}^{\text{in}}(0_+)$  and  $F_-^{\text{out}} \in \mathcal{F}^{\text{out}}(0_-)$ . This relation is the key to the proof of the following statement.

**Proposition 3:**

If there exists a bounded region  $\mathcal{O} \subset \mathbb{R}^4$  and a sequence of hermitian operators  $B_\lambda \in \mathcal{F}(\mathcal{O})$  which converges weakly on the vacuum to some non-zero vector in  $\mathcal{H}_1$ , then the collision states  $\Phi_1^{\text{in}} \dots \Phi_m^{\text{in}}$  and  $\Phi_1^{\text{out}} \dots \Phi_n^{\text{out}}$  coincide for arbitrary configurations  $\Phi_1, \dots, \Phi_n \in \mathcal{H}_1$ . Consequently the S-matrix is trivial.

**Proof:**

We define  $B_\lambda^L = U(L) B_\lambda U(L)^{-1}$  where  $L = (1, x)$  is an arbitrary Poincaré transformation and  $U(L) = U(x)U(1)$  is the corresponding unitary in  $\mathcal{H}$ . Since  $B_\lambda$  converges weakly on the vacuum to some non-trivial one-particle state  $\Phi \in \mathcal{H}_1$  we get  $\omega\text{-}\lim_{\lambda \rightarrow 0} B_\lambda^L \Omega = U(L)\Phi = \Phi_L$ . These vectors form a total set in  $\mathcal{H}_1$  because the Poincaré transformations are irreducibly represented in  $\mathcal{H}_1$ . Now the operators  $B_\lambda^L$  are localized in the region  $L\mathcal{O}$ . Using relation (9) we get therefore

$$(F_+^{\text{in}} \Phi_L, F_-^{\text{out}} \Omega) = \lim_{\lambda \rightarrow 0} (F_+^{\text{in}} B_\lambda^L \Omega, F_-^{\text{out}} \Omega) = \lim_{\lambda \rightarrow 0} (F_+^{\text{in}} \Omega, F_-^{\text{out}} B_\lambda^L \Omega) = (F_+^{\text{in}} \Omega, F_-^{\text{out}} \Phi_L), \quad (10)$$

provided  $F_+^{\text{in}} \in \mathcal{F}^{\text{in}}(L\mathcal{O}_+)$  and  $F_-^{\text{out}} \in \mathcal{F}^{\text{out}}(L\mathcal{O}_-)$ . Since the operators  $A^{\text{in}}$  and  $A^{\text{out}}$  are free fields it is straightforward to verify that the bounded operators  $F_+^{\text{in}}$  and  $F_-^{\text{out}}$  in this relation may be replaced by products of smeared field operators  $A_1^{\text{in}}, \dots, A_m^{\text{in}}$  and  $A_{m+1}^{\text{out}}, \dots, A_n^{\text{out}}$  which are localized in  $L\mathcal{O}_+$  and  $L\mathcal{O}_-$  respectively. Thus we arrive at

$$(A_1^{\text{in}} \dots A_m^{\text{in}} \Phi_L, A_{m+1}^{\text{out}} \dots A_n^{\text{out}} \Omega) = (A_1^{\text{in}} \dots A_m^{\text{in}} \Omega, A_{m+1}^{\text{out}} \dots A_n^{\text{out}} \Phi_L). \quad (11)$$

Now we can prove the proposition by induction. For a one-particle state there is nothing to show, so let us assume that  $\Phi_1^{\text{in}} \dots \Phi_m^{\text{in}} = \Phi_1^{\text{out}} \dots \Phi_m^{\text{out}}$  for arbitrary configurations  $\Phi_1, \dots, \Phi_m \in \mathcal{H}_1$ . This implies in particular that  $A_1^{\text{in}} \dots A_m^{\text{in}} \Omega = A_1^{\text{out}} \dots A_m^{\text{out}} \Omega$  and using relation (11) we get

$$(A_1^{\text{in}} \dots A_m^{\text{in}} \Phi_L, A_{m+1}^{\text{out}} \dots A_n^{\text{out}} \Omega) = (A_1^{\text{in}} \dots A_m^{\text{in}} \Omega, A_{m+1}^{\text{out}} \dots A_n^{\text{out}} \Phi_L) = (A_1^{\text{out}} \dots A_m^{\text{out}} \Omega, A_{m+1}^{\text{out}} \dots A_n^{\text{out}} \Phi_L) = (A_1^{\text{out}} \dots A_m^{\text{out}} \Phi_L, A_{m+1}^{\text{out}} \dots A_n^{\text{out}} \Omega),$$

where the last equality sign follows from an explicit calculation of the scalar products. If we set  $\Phi_1 = A_1^{\text{in}} \Omega, \dots, \Phi_n = A_n^{\text{out}} \Omega$  we can reexpress this equation in terms of the collision states,

$$(\Phi_1^{\text{in}} \dots \Phi_m^{\text{in}} \Phi_L, \Phi_{m+1}^{\text{out}} \dots \Phi_n^{\text{out}}) = (\Phi_1^{\text{out}} \dots \Phi_m^{\text{out}} \Phi_L, \Phi_{m+1}^{\text{out}} \dots \Phi_n^{\text{out}}),$$

provided  $A_1^{\text{in}}, \dots, A_n^{\text{out}}$  are operators with the special localisation properties mentioned above. However, keeping in mind that the vectors  $\Phi_L$  form a total set in  $\mathcal{H}_1$  one can extend this equation by continuity to arbitrary configurations  $\Phi_1, \dots, \Phi_n, \Phi_L \in \mathcal{H}_1$  [4] and it is then obvious that  $\Phi_1^{\text{in}} \dots \Phi_m^{\text{in}} = \Phi_1^{\text{out}} \dots \Phi_m^{\text{out}}$ . ■

Combining the two propositions it follows that the massless particles in  $\mathcal{H}_1$  do not interact if the dilations are a true symmetry. We have established this result only for one type of massless particles with helicity  $s = 0$ . In the presence of a family of one-particle spaces  $\mathcal{H}_1^{(i)} \subset \mathcal{H}$  on which the unitaries  $U(x), U(1)$  act like an irreducible representation of the Poincaré group with mass zero and helicity  $s_i$  the main modifications are in the second part of the proof of Proposition 2: for vectors  $\Psi \in \mathcal{H}_1^{(k)}$  relation (7b) changes according to

$$(U(\Lambda)\Psi)_k(p) = e^{is_k \alpha(\Lambda, p)} (\Psi)_k(\Lambda^{-1} p)$$

where the index  $k$  refers to the space  $\mathcal{H}_1^{(k)}$ . The functions  $\alpha(\Lambda, p)$  are the Wigner phases [6]. They are not completely fixed by the structural relations imposed by the Lorentz group. As a matter of fact we may choose a convention such that the functions  $\alpha(\Lambda, p)$  are simultaneously continuous in  $\Lambda$  and  $p$  except at  $p = 0$ ; moreover we may require that  $\alpha(K_\lambda, p) = 0$  where  $K_\lambda$  are the boosts in the  $x_1$ -direction introduced in relation (5). It is then easy to verify that the functions  $(B_\lambda \Omega)_k(p)$  are continuous and that the analogue of relation (8) holds. The proof of Proposition 3 carries over almost literally and we may therefore omit the details.

Finally we want to point out a further generalisation of our main result.

In an asymptotically complete theory of massless particles there always exist two representations  $D^{\text{in}}(\lambda)$  and  $D^{\text{out}}(\lambda)$  of the group of dilations which act on the asymptotic fields  $A^{\text{in}}$  and  $A^{\text{out}}$ , respectively, as in a free field theory. Their commutation relations with the translations  $U(x)$  and Lorentz transformations  $U(\Lambda)$  are again given by (2a). However, they do not, in general, act on the basic fields according to relation (2b). In order that the dilations are an asymptotically visible symmetry it would be sufficient to require

$$D^{\text{in}}(\lambda) = D^{\text{out}}(\lambda) = D(\lambda) \quad (12)$$

and relation (2b) could be dropped. But this assumption still implies that the S-matrix is trivial! To verify this one has only to realize that Propositions 2 and 3 still hold in this case. The proof of Proposition 2 depends on the clustering properties of the vacuum and relation (2a) and therefore applies. Of course the operators  $B_\lambda$  are in general not local. However, relation (10) which was crucial for the proof of Proposition 3 can still be established. This follows simply from the fact that the asymptotic nets  $\mathcal{O} \rightarrow \mathcal{F}^{\text{in}}(\mathcal{O})$  and  $\mathcal{O} \rightarrow \mathcal{F}^{\text{out}}(\mathcal{O})$  transform under the dilations  $D(\lambda) = D^{\text{in}}(\lambda) = D^{\text{out}}(\lambda)$  according to relation (2b). Hence if for example  $A \in \mathcal{F}(\mathcal{O})$  where  $\mathcal{O}$  is any bounded region which contains the origin and if  $F_+^{\text{in}} \in \mathcal{F}^{\text{in}}(\mathcal{O}_+)$  where  $\mathcal{O}_+$  has a positive timelike separation from  $\mathcal{O}$  one gets for  $\lambda < 1$

$$[D(\lambda) A D(\lambda)^{-1}, F_+^{\text{in}}] = D(\lambda) [A, D(\lambda^{-1}) F_+^{\text{in}} D(\lambda^{-1})^{-1}] D(\lambda)^{-1} = 0$$

by Huyghens' principle. A similar relation holds for  $F_-^{\text{out}} \in \mathcal{F}^{\text{out}}(\mathcal{O}_-)$ . It is then easy to verify that the operators  $B_\lambda$  commute for small  $\lambda$  with the operators in  $\mathcal{F}^{\text{in}}(\mathcal{O}_+)$  and  $\mathcal{F}^{\text{out}}(\mathcal{O}_-)$  where the regions  $\mathcal{O}_+, \mathcal{O}_-$  depend only on the localisation properties of  $\mathcal{F}$  and  $A_\varphi$ . The rest of the argument can then be carried over.

#### Acknowledgements

We are indebted to R. Haag for stimulating discussions and to G. Mack and G. Roepstorff for useful remarks on helicity representations.

#### References

- [1] G. F. Dell-Antonio; Dilation Invariance and the Wilson Expansion; *Il Nuovo Cimento* 12A, 756 (1972)
- [2] S. Doplicher, R. Haag and J. E. Roberts; Fields, Observables and Gauge-Transformations I; *Commun. math. Phys.* 13, 1 (1969)
- [3] R. Haag, J. T. Lopuszanski and M. Sohnius; All Possible Generators of Supersymmetries of the S-matrix; *Nucl. Phys.* B88, 257 (1975)
- [4] D. Buchholz; Collision Theory for Massless Fermions; *Commun. math. Phys.* 42, 269 (1975)  
D. Buchholz; Collision Theory for Massless Bosons; DESY Report 76/22 (1976)
- [5] S. Doplicher, R. Haag and J. E. Roberts; Local Observables and Particle Statistics II; *Commun. math. Phys.* 35, 49 (1974)
- [6] E. P. Wigner; Unitary Representations of the Inhomogeneous Lorentz Group; *Ann. Math.* 40, 149 (1939)  
A. S. Wightman; l' invariance dans la mécanique quantique relativiste in: *Dispersion Relations and Elementary Particles*, Paris: Hermann 1960.