

From NANOGrav to LIGO with metastable cosmic strings

Wilfried Buchmüller,^{1,*} Valerie Domcke,^{2,3,†} and Kai Schmitz^{2,‡}

¹*Deutsches Elektronen Synchrotron DESY, 22607 Hamburg, Germany*

²*Theoretical Physics Department, CERN, 1211 Geneva 23, Switzerland*

³*Institute of Physics, Laboratory for Particle Physics and Cosmology, EPFL, CH-1015, Lausanne, Switzerland*
 (Dated: September 23, 2020)

We interpret the recent NANOGrav results in terms of a stochastic gravitational wave background from metastable cosmic strings. The observed amplitude of a stochastic signal can be translated into a range for the cosmic string tension and the mass of magnetic monopoles arising in theories of grand unification. In a sizable part of the parameter space, this interpretation predicts a large stochastic gravitational wave signal in the frequency band of ground-based interferometers, which can be probed in the very near future. We confront these results with predictions from successful inflation, leptogenesis and dark matter from the spontaneous breaking of a gauged $B-L$ symmetry.

Introduction

The direct observation of gravitational waves (GWs) generated by merging black holes [1–3] has led to an increasing interest in further explorations of the GW spectrum. Astrophysical sources can lead to a stochastic gravitational background (SGWB) over a wide range of frequencies, and the ultimate hope is the detection of a SGWB of cosmological origin. So far, transient merger events have been observed at frequencies around 100 Hz. Moreover, stringent upper bounds on a SGWB have been obtained by pulsar timing array (PTA) experiments which are sensitive to frequencies around $f_{\text{yr}} = 1/\text{yr}$. Over the past years the European Timing Array (EPTA) [4], the Parkes Pulsar Timing Array (PPTA) [5] and the North American Nanohertz Observatory for Gravitational Waves (NANOGrav) [6] have reached upper bounds on the amplitude $h^2\Omega_{\text{gw}}(1/\text{yr})$ of order 10^{-9} .

Searching for an isotropic SGWB, the NANOGrav Collaboration has recently reported strong evidence of a stochastic process in their lowest frequency bins, which can be modeled as a power-law with common amplitude and slope across all pulsars [7]. The amplitude of the signal is of the order of the previously obtained upper bounds. The current data is not conclusive with respect to a quadrupolar spatial correlation and therefore the discovery of a SGWB cannot be claimed. Nevertheless, the result of the analysis is very intriguing, and the NANOGrav Collaboration finds that the signal is consistent, within 2σ of a Bayesian analysis, with a SGWB from supermassive black hole binaries, the expected dominant astrophysical source at frequencies around $1/\text{yr}$ [8, 9].

There are also cosmological interpretations of the NANOGrav results. Examples are the formation of primordial black holes from high-amplitude curvature perturbations during inflation [10, 11] or dark sector phase transitions [12]. Another prominent possibility is cosmic strings formed in a $U(1)$ symmetry-breaking phase transition in the early universe [13, 14]. Indeed, it has been demonstrated that GWs from a network of stable strings with an amplitude $h^2\Omega_{\text{gw}}(1/\text{yr}) \sim 10^{-9}$ can account for the NANOGrav stochastic background [15, 16]. This sig-

nal is too small to be observed by Virgo [17], LIGO [18] and KAGRA [19] but will be probed by LISA [20] and other planned GW observatories.

In this Letter we study a further possibility, metastable cosmic strings. Recently, it has been shown that GWs emitted from a metastable cosmic string network can probe the seesaw mechanism of neutrino physics and high-scale leptogenesis [21] as well as the energy scale of grand unification [22, 23]. Such metastable cosmic strings arise when connecting hybrid inflation, high-scale leptogenesis and dark matter with gravitational waves through $U(1)_{B-L}$ breaking in a cosmological phase transition [24, 25]. Here $B-L$ denotes the difference of baryon number and lepton number, and the product of $U(1)_{B-L}$ and the Standard Model gauge group is embedded into the GUT group $SO(10)$. If the $U(1)_{B-L}$ cosmic strings are not protected by an additional unbroken discrete symmetry, this embedding leads to the existence of magnetic monopoles, allowing the cosmic strings to decay via the Schwinger production of monopole-antimonopole pairs with a rate per string unit length of [26–28]

$$\Gamma_d = \frac{\mu}{2\pi} \exp(-\pi\kappa) \ , \quad \kappa = \frac{m^2}{\mu} \ , \quad (1)$$

where $m \sim v_{\text{GUT}}$ is the monopole mass and $\mu \sim v_{B-L}^2$ is the string tension. Here v_{GUT} and v_{B-L} are the scales of $SO(10)$ and $U(1)_{B-L}$ symmetry breaking, respectively.

At frequencies around 100 Hz the model of [24] predicts a GW amplitude close to the present upper bound found by the LIGO/Virgo collaboration, and upper bounds on a SGWB by PTA experiments lead to an upper bound on the ratio κ and therefore on the monopole mass [22]. With the new NANOGrav data [7], κ and hence the scale of grand unification v_{GUT} can now be determined.

GWs from metastable cosmic strings

We briefly review the calculation of the stochastic gravitational wave background arising from metastable cosmic strings [22]. The present-day GW spectrum can be

expressed as [20]

$$\Omega_{\text{gw}}(f) = \frac{\partial \rho_{\text{gw}}(f)}{\rho_c \partial \ln f} = \frac{8\pi f (G\mu)^2}{3H_0^2} \sum_{n=1}^{\infty} C_n(f) P_n, \quad (2)$$

where ρ_{gw} denotes the GW energy density, ρ_c is the critical energy density of the universe, $G\mu$ denotes the dimensionless string tension with the gravitational constant $G = 6.7 \cdot 10^{-39} \text{ GeV}^{-2}$, $H_0 = 100 h \text{ km/s/Mpc}$ is today's Hubble parameter, $P_n \simeq 50/\zeta[4/3] n^{-4/3}$ is the power spectrum of GWs emitted by the n^{th} harmonic of a cosmic string loop¹, and $C_n(f)$ indicates the number of loops emitting GWs that are observed at a given frequency f ,

$$C_n(f) = \frac{2n}{f^2} \int_{z_{\min}}^{z_{\max}} dz \frac{\mathcal{N}(\ell(z), t(z))}{H(z) (1+z)^6}, \quad (3)$$

which is a function of the number density of cosmic string loops $\mathcal{N}(\ell, t)$, with $\ell = 2n/((1+z)f)$, selecting the loops that contribute to the spectrum at frequency f today. Modeling the evolution and GW emission of a cosmic string network is a challenging task, resulting in several competing models for the loop number density in the literature (see [20] for an overview). For concreteness, we will base our analysis on the BOS model [29] and fix the cosmic string loop size to $\alpha = \ell/H = 0.1$ at formation, as found in the largest and most recent simulations [20].² For loops generated and decaying during the radiation-dominated era, this in particular yields [20, 29]

$$\mathcal{N}_r(\ell, t) = \frac{0.18}{t^{3/2} (\ell + \Gamma G\mu t)^{5/2}}, \quad (4)$$

where $\Gamma \simeq 50$ parametrizes the cosmic string decay rate into GWs, $\dot{\ell} = -\Gamma G\mu$. This yields the dominant contribution to the GW spectrum in most of the parameter range of interest, but in our numerical computation of the spectrum we also include the loops created and/or decaying in the matter dominated era. The integration range in Eq. (3) accounts for the lifetime of the cosmic string network, from the formation at $z_{\max} \simeq T_{\text{rh}}/(2.7 \text{ K})$ until their decay at $z_{\min}$ when the decay rate of a string loop with average length equals the Hubble rate [26],

$$z_{\min} = \left(\frac{70}{H_0} \right)^{1/2} (\Gamma \Gamma_d G\mu)^{1/4}. \quad (5)$$

For cosmic string loops formed and emitting GWs in the radiation dominated era, this results in an approximately scale invariant GW spectrum. The finite lifetime of the

cosmic strings leads to a fall-off $\propto f^{3/2}$ of this spectrum at small frequencies $f < f_*$ with [22]

$$f_* \simeq 4.4 \times 10^{-8} \text{ Hz} \frac{e^{-\pi\kappa/4}}{e^{-16\pi}} \left(\frac{10^{-7}}{G\mu} \right)^{1/2}, \quad (6)$$

see Fig. 2 for some examples of GW spectra for different values of the two dimensionless model parameters $G\mu$ and κ .

For the numerical evaluation of Eq. (2), we refine the analysis of Ref. [22] by resumming the first 20,000 modes and taking into account the changes in the number of effective degrees of freedom in the thermal bath (see also [30]). Our final results prove rather insensitive to both these refinements. Approximating $\mathcal{N} \simeq \mathcal{N}_r$, we can extract the n -dependence of $C_n P_n$ analytically if ℓ is much smaller or larger than $\Gamma G\mu t$. As discussed in Ref. [22], this distinction corresponds to the $f^{3/2}$ slope and the plateau regime. For the former, we find $C_n P_n \propto n^{-17/3}$, such that the resummation yields $\Omega_{\text{gw}} = \zeta(17/3) \Omega_{\text{gw}}^{(1)} \simeq 1.02 \Omega_{\text{gw}}^{(1)}$, with $\Omega_{\text{gw}}^{(1)}$ denoting the result for $n = 1$. For the plateau value, we instead obtain a factor $\zeta(4/3) \simeq 3.6$, which implies an $\mathcal{O}(1)$ correction.

For the evolution of the degrees of freedom we use the results of [31] for the SM degrees of freedom and moreover include supersymmetric degrees of freedom at a threshold value of 2 TeV. This does not impact the predictions in the NANOGrav frequency range.

Explaining the NANOGrav results

We now proceed to comparing the GW signal predicted by metastable cosmic strings to the recent NANOGrav results [7], which constrain the amplitude and slope of a stochastic process. Expressing the dimensionless characteristic strain as $h_c = A(f/f_{\text{yr}})^\alpha$ with the reference frequency $f_{\text{yr}} = 32 \text{ nHz}$, the amplitude of the SGWB is obtained as

$$\Omega_{\text{gw}}(f) = \frac{2\pi^2 f_{\text{yr}}^2 A^2}{3H_0^2} \left(\frac{f}{f_{\text{yr}}} \right)^{2\alpha+2} \equiv \Omega_{\text{gw}}^{\text{yr}} \left(\frac{f}{f_{\text{yr}}} \right)^{n_t}. \quad (7)$$

To compare with the results of Ref. [7], we determine the spectral index n_t by fitting a power law to our numerically determined spectrum $\Omega_{\text{gw}} h^2(f)$ from Eq. (2) in the range of maximal sensitivity of NANOGrav, i.e. $f = 2.4..12 \text{ nHz}$ (this corresponds to the first 5 bins of the NANOGrav data set). The amplitude $\Omega_{\text{gw}}^{\text{yr}}$ is the amplitude of this power law at the reference frequency $f = f_{\text{yr}}$.

In Fig. 1, we compare these predictions from metastable cosmic strings (mesh of solid and dotted curves) with the constraints on the amplitude and tilt from [7] (orange shaded region). We vary $G\mu$ from the lowest value capable of explaining the NANOGrav results at 2 sigma, $G\mu \simeq 10^{-10}$ to the largest value compatible with the constraints from LIGO/Virgo [32], $G\mu \simeq 10^{-6}$. Note that the CMB constraint $G\mu < 1.3 \times 10^{-7}$ [33] only

¹ Here we focus on cusps as the main source of GW emission, kinks and kink-kink collisions yield a different $\mathcal{O}(1)$ factor in both the argument of the ζ function and the power of n in P_n .

² For an interpretation of the NANOGrav results for smaller values of the parameter α , see [16].




FIG. 1: Gravitational wave signals from metastable cosmic strings compared to the NANOGrav observations for different values of the string tension $G\mu$ and the hierarchy between the GUT and U(1) breaking scale κ . The solid coloured lines indicate fixed values of $G\mu = 10^{-10}, \dots, 10^{-6}$, the dotted lines indicate contours of $\sqrt{\kappa} = 7.8, 7.9, \dots, 9$. The orange region with the solid (dashed) contours show the 68% and 95% regions reported by NANOGrav when performing a fit to the first 5 frequency bins (performing a fit with a broken power law).

applies to cosmic strings with a life-time longer than CMB decoupling, corresponding to $\sqrt{\kappa} \gtrsim 8.6$ (indicated by gray points in the upper left corner). For each value of $G\mu$, we consider the range $\sqrt{\kappa} = 7.8..9.0$; smaller values lead to an unobservably small spectrum at nHz frequencies, while all values $\sqrt{\kappa} \gtrsim 9$ quickly converge towards the result for stable cosmic strings, see [15, 16]. Contours of fixed $G\mu$ (κ) are indicated by solid (dotted) lines in Fig. 1.

The cyan shaded band in Fig. 1 indicates the prediction from B - L breaking in the early Universe [22, 24]. Remarkably the predicted GW signal at nHz frequencies is compatible with the NANOGrav results at 2 sigma.

It is intriguing that the values of the cosmic string tension $G\mu$ found in the context of metastable cosmic strings can be significantly larger than the values found for stable cosmic strings [15, 16], implying the possibility of observing this signal with the existing ground-based detectors Virgo, LIGO and KAGRA. The reason for this is twofold. Firstly, the finite lifetime of the cosmic strings leads to a suppression of the low-frequency spectrum, implying a blue tilt of the GW spectrum between the PTA and the frequency band of ground-based interferometers. In particular, the production of GWs after matter-radiation equality is suppressed, which for stable cosmic strings leads to an enhancement at low frequencies, see e.g. the dotted red curve in Fig. 2. Secondly, the NANOGrav data exhibit a sizable correlation between the amplitude and tilt of the spectrum, allowing for larger amplitudes for positive values of n_t .




FIG. 2: Gravitational wave spectra from metastable cosmic strings explaining the NANOGrav excess (at 2σ CL). The colored blue (green) region accounts for successful inflation, baryogenesis and dark matter [22, 24] for the maximal (minimal) allowed value of $G\mu$ with $\sqrt{\kappa} = 8.0..8.1$ ($8.1..8.3$), corresponding to the $\star$ ($\diamond$) markers in Fig. 1. For reference, the dotted red line shows the spectrum for stable cosmic strings for the best-fit value $G\mu = 10^{-10}$ [15, 16]; the dashed red line shows the corresponding curve for the $\sqrt{\kappa} = 9$, the largest value considered here. The (lighter) gray-shaded areas indicate the sensitivities of the (planned) experiments SKA [34], LISA [35], LIGO/Virgo [32] and ET [36]. The orange band indicates the frequency range of NANOGrav.

Discussion

In the model of cosmological U(1) $_{B-L}$ breaking [22, 24], successful inflation, leptogenesis and dark matter restrict the allowed values of $G\mu$ to a narrow band around $G\mu \sim 3 \times 10^{-7}$, depicted by the cyan region in Fig. 1. Interpreting the NANOGrav results as originating from a metastable cosmic string network determines the ratio between the GUT and the B - L breaking scales to lie around $\sqrt{\kappa} \simeq 8$, excluding stable cosmic strings. More precisely, the predictions of [22, 24] are consistent with the recent NANOGrav results in the range from $G\mu = 1.0 \times 10^{-7}$, with $\sqrt{\kappa} = 8.1..8.3$, to $G\mu = 5.6 \times 10^{-7}$, with $\sqrt{\kappa} = 8.0..8.1$. The corresponding values of B - L breaking scales and monopoles masses are $v_{B-L} = 3.0 \times 10^{15}$ GeV, with $m = (3.3..3.4) \times 10^{16}$ GeV and $v_{B-L} = 5.8 \times 10^{15}$ GeV, with $m = (7.8..7.9) \times 10^{16}$ GeV, respectively. The precise connection between GUT symmetry breaking, inflation and U(1) $_{B-L}$ is a challenging theoretical question.³

A second important outcome of our analysis are the

³ Determining the $\mathcal{O}(1)$ factors between the ratio of monopole mass and string tension (parametrized by $\sqrt{\kappa}$) and the ratio of the underlying scales v_{GUT}/v_{B-L} requires a careful and consistent treatment of both types of topological defects under consideration of the gauge coupling and the symmetry breaking potentials.

expectations for ground-based GW interferometers. In Fig. 2 the GW spectrum is shown for the upper and the lower boundary of the range in $G\mu$ that is predicted by the considered $U(1)_{B-L}$ model [24]. The prediction of this model will be probed by Advanced LIGO [32].⁴ The observation of a SGWB with PTA experiments as well as with LIGO would give stunning support for grand unified theories, with important implications for inflation, baryogenesis and dark matter [22].

An improved determination of the tilt of the spectrum at PTA frequencies together with upcoming results on SGWBs at LIGO frequencies will soon rule out or further support the model presented here. This encourages further refinements of the analysis, e.g. going beyond the instantaneous decay approximation for the cosmic string network and taking into account the dynamics of cosmic string decay induced by monopole formation, which may lead to an additional GW contribution [26, 37]. One may also consider relaxing some of the model-building assumptions within the model of cosmological $U(1)_{B-L}$ breaking [24]. However, the core of the model — inflation ending in a GUT-scale phase transition in combination with leptogenesis and dark matter in a supersymmetric extension of the SM — is intrinsically tied to the GW signals discussed here.

Acknowledgements. We thank Daniel Figueroa for helpful discussions on the modeling of cosmic strings. This project has received funding from the European's Union Horizon 2020 Research and Innovation Programme under grant agreement number 796961, 'Ax-iBAU' (K.S.).

* Electronic address: wilfried.buchmueller@desy.de

† Electronic address: valerie.domcke@cern.ch

‡ Electronic address: kai.schmitz@cern.ch

- [1] **LIGO Scientific, Virgo** Collaboration, B. Abbott et al., *Observation of Gravitational Waves from a Binary Black Hole Merger*, *Phys. Rev. Lett.* **116** (2016), no. 6 061102, [[arXiv:1602.03837](#)].
- [2] **LIGO Scientific, Virgo** Collaboration, B. P. Abbott et al., *GW151226: Observation of Gravitational Waves from a 22-Solar-Mass Binary Black Hole Coalescence*, *Phys. Rev. Lett.* **116** (2016), no. 24 241103, [[arXiv:1606.04855](#)].
- [3] **LIGO Scientific, VIRGO** Collaboration, B. P. Abbott et al., *GW170104: Observation of a 50-Solar-Mass Binary Black Hole Coalescence at Redshift 0.2*, *Phys. Rev. Lett.* **118** (2017), no. 22

⁴ On the contrary, interpreting the NANOGrav signal as originating from stable cosmic strings forces $G\mu$ to values too low to be observed by current ground-based GW interferometers [15].

- 221101, [[arXiv:1706.01812](#)]. [Erratum: *Phys. Rev. Lett.* **121**, no. 12, 129901 (2018)].
- [4] R. M. Shannon et al., *Gravitational waves from binary supermassive black holes missing in pulsar observations*, *Science* **349** (2015), no. 6255 1522–1525, [[arXiv:1509.07320](#)].
- [5] M. Kerr et al., *The Parkes Pulsar Timing Array Project: Second data release*, *Publ. Astron. Soc. Austral.* **37** (2020) e020, [[arXiv:2003.09780](#)].
- [6] **NANOGrav** Collaboration, Z. Arzoumanian et al., *The NANOGrav 11-year Data Set: Pulsar-timing Constraints On The Stochastic Gravitational-wave Background*, *Astrophys. J.* **859** (2018), no. 1 47, [[arXiv:1801.02617](#)].
- [7] **NANOGrav** Collaboration, Z. Arzoumanian et al., *The NANOGrav 12.5-year Data Set: Search For An Isotropic Stochastic Gravitational-Wave Background*, [arXiv:2009.04496](#).
- [8] M. Rajagopal and R. W. Romani, *Ultralow frequency gravitational radiation from massive black hole binaries*, *Astrophys. J.* **446** (1995) 543–549, [[astro-ph/9412038](#)].
- [9] E. Phinney, *A Practical theorem on gravitational wave backgrounds*, [astro-ph/0108028](#).
- [10] V. Vaskonen and H. Veerme, *Did NANOGrav see a signal from primordial black hole formation?*, [arXiv:2009.07832](#).
- [11] V. De Luca, G. Franciolini, and A. Riotto, *NANOGrav Hints to Primordial Black Holes as Dark Matter*, [arXiv:2009.08268](#).
- [12] Y. Nakai, M. Suzuki, F. Takahashi, and M. Yamada, *Gravitational Waves and Dark Radiation from Dark Phase Transition: Connecting NANOGrav Pulsar Timing Data and Hubble Tension*, [arXiv:2009.09754](#).
- [13] T. Kibble, *Topology of Cosmic Domains and Strings*, *J. Phys. A* **9** (1976) 1387–1398.
- [14] R. Jeannerot, J. Rocher, and M. Sakellariadou, *How generic is cosmic string formation in SUSY GUTs*, *Phys. Rev. D* **68** (2003) 103514, [[hep-ph/0308134](#)].
- [15] J. Ellis and M. Lewicki, *Cosmic String Interpretation of NANOGrav Pulsar Timing Data*, [arXiv:2009.06555](#).
- [16] S. Blasi, V. Brdar, and K. Schmitz, *Has NANOGrav found first evidence for cosmic strings?*, [arXiv:2009.06607](#).
- [17] F. Acernese et al., *Advanced virgo: a second-generation interferometric gravitational wave detector*, *Classical and Quantum Gravity* **32** (2015), no. 2 024001.
- [18] **LIGO Scientific Collaboration and Virgo Collaboration** Collaboration, B. P. Abbott, R. Abbott, T. D. Abbott, M. R. Abernathy, F. Acernese, K. Ackley, C. Adams, T. Adams, P. Addesso, R. X. Adhikari, et al., *Gw150914: The advanced LIGO detectors in the era of first discoveries*, *Phys. Rev. Lett.* **116** (Mar, 2016) 131103.
- [19] **KAGRA** Collaboration, T. Akutsu et al., *KAGRA: 2.5 Generation Interferometric Gravitational Wave Detector*, *Nature Astron.* **3** (2019), no. 1 35–40, [[arXiv:1811.08079](#)].
- [20] P. Auclair et al., *Probing the gravitational wave background from cosmic strings with LISA*, *JCAP* **04** (2020) 034, [[arXiv:1909.00819](#)].
- [21] J. A. Dror, T. Hiramatsu, K. Kohri, H. Murayama, and G. White, *Testing Seesaw and Leptogenesis with Gravitational Waves*, [arXiv:1908.03227](#).
- [22] W. Buchmuller, V. Domcke, H. Murayama, and

- K. Schmitz, *Probing the scale of grand unification with gravitational waves*, *Phys. Lett. B* (2020) 135764, [[arXiv:1912.03695](#)].
- [23] S. F. King, S. Pascoli, J. Turner, and Y.-L. Zhou, *Gravitational waves and proton decay: complementary windows into GUTs*, [[arXiv:2005.13549](#)].
- [24] W. Buchmuller, V. Domcke, and K. Schmitz, *Spontaneous $B-L$ Breaking as the Origin of the Hot Early Universe*, *Nucl. Phys. B* **862** (2012) 587–632, [[arXiv:1202.6679](#)].
- [25] W. Buchmuller, V. Domcke, K. Kamada, and K. Schmitz, *The Gravitational Wave Spectrum from Cosmological $B-L$ Breaking*, *JCAP* **1310** (2013) 003, [[arXiv:1305.3392](#)].
- [26] L. Leblond, B. Shlaer, and X. Siemens, *Gravitational Waves from Broken Cosmic Strings: The Bursts and the Beads*, *Phys. Rev. D* **79** (2009) 123519, [[arXiv:0903.4686](#)].
- [27] A. Monin and M. B. Voloshin, *The Spontaneous breaking of a metastable string*, *Phys. Rev. D* **78** (2008) 065048, [[arXiv:0808.1693](#)].
- [28] A. Monin and M. B. Voloshin, *Destruction of a metastable string by particle collisions*, *Phys. Atom. Nucl.* **73** (2010) 703–710, [[arXiv:0902.0407](#)].
- [29] J. J. Blanco-Pillado, K. D. Olum, and B. Shlaer, *The number of cosmic string loops*, *Phys. Rev. D* **89** (2014), no. 2 023512, [[arXiv:1309.6637](#)].
- [30] Y. Gouttenoire, G. Servant, and P. Simakachorn, *Beyond the Standard Models with Cosmic Strings*, *JCAP* **07** (2020) 032, [[arXiv:1912.02569](#)].
- [31] L. Husdal, *On Effective Degrees of Freedom in the Early Universe*, *Galaxies* **4** (2016), no. 4 78, [[arXiv:1609.04979](#)].
- [32] **LIGO Scientific, Virgo** Collaboration, B. P. Abbott et al., *Search for the isotropic stochastic background using data from Advanced LIGOs second observing run*, *Phys. Rev. D* **100** (2019), no. 6 061101, [[arXiv:1903.02886](#)].
- [33] **Planck** Collaboration, P. A. R. Ade et al., *Planck 2015 results. XIII. Cosmological parameters*, *Astron. Astrophys.* **594** (2016) A13, [[arXiv:1502.01589](#)].
- [34] R. Smits, M. Kramer, B. Stappers, D. R. Lorimer, J. Cordes, and A. Faulkner, *Pulsar searches and timing with the square kilometre array*, *Astron. Astrophys.* **493** (2009) 1161–1170, [[arXiv:0811.0211](#)].
- [35] **LISA** Collaboration, P. Amaro-Seoane et al., *Laser Interferometer Space Antenna*, [[arXiv:1702.00786](#)].
- [36] **Einstein Telescope** Collaboration <http://www.et-gw.eu/index.php/etsensitivities>.
- [37] A. Vilenkin, *Cosmological evolution of monopoles connected by strings*, *Nucl. Phys. B* **196** (1982) 240–258.