

PAPER • OPEN ACCESS

Top quark pair production at complete NLO accuracy with NNLO+NNLL' corrections in QCD^{*}

To cite this article: Micha Czakon *et al* 2020 *Chinese Phys. C* **44** 083104

View the [article online](#) for updates and enhancements.

Top quark pair production at complete NLO accuracy with NNLO+NNLL' corrections in QCD*

Michał Czakon¹ Andrea Ferroglia^{2,3} Alexander Mitov⁴ Davide Pagani^{5,6} Andrew S. Papanastasiou⁴
 Benjamin D. Pecjak⁷ Darren J. Scott^{8,9} Ioannis Tsinikos¹⁰ Xing Wang¹¹
 Li Lin Yang(杨李林)^{12,1)} Marco Zaro¹³

¹Institut für Theoretische Teilchenphysik und Kosmologie, RWTH Aachen University, D-52056 Aachen, Germany

²New York City College of Technology, Brooklyn, NY 11201, USA

³The Graduate School and University Center, The City University of New York, New York, NY 10016 USA

⁴Cavendish Laboratory, University of Cambridge, CB3 0HE Cambridge, UK

⁵Technische Universität München, James-Frank-Str. 1, D-85748 Garching, Germany

⁶Theory Group, Deutsches Elektronen-Synchrotron (DESY), D-22607 Hamburg, Germany

⁷Institute for Particle Physics Phenomenology, University of Durham, DH1 3LE Durham, UK

⁸Institute for Theoretical Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands

⁹Nikhef, Theory Group, Science Park 105, 1098 XG, Amsterdam, The Netherlands

¹⁰Theoretical Particle Physics, Department of Astronomy and Theoretical Physics, Lund University, Sölvegatan 14A, SE-223 62 Lund, Sweden

¹¹PRISMA+ Cluster of Excellence, Johannes Gutenberg University, D-55128 Mainz, Germany

¹²Zhejiang Institute of Modern Physics, Department of Physics, Zhejiang University, Hangzhou 310027, China

¹³INFN, Sezione di Milano, Via Celoria 16, 20133 Milano, Italy

Abstract: We describe predictions for top quark pair differential distributions at hadron colliders, by combining the next-to-next-to-leading order quantum chromodynamics calculations and next-to-leading order electroweak corrections with double resummation at the next-to-next-to-leading logarithmic accuracy of threshold logarithms and small-mass logarithms. To the best of our knowledge, this is the first study to present such a combination, which incorporates all known perturbative information. Numerical results are presented for the invariant-mass distribution, transverse-momentum distribution, and rapidity distributions.

Keywords: top quark production, QCD, electroweak

DOI: 10.1088/1674-1137/44/8/083104

1 Introduction

Top quark pair production is one of the most important processes at the Large Hadron Collider (LHC). It allows one to precisely study the properties of the top quark, which are related to many important questions in particle physics, such as the hierarchy problem, the stability of the electroweak (EW) vacuum, and the origin of

fermion masses. Further, top quark pair production is a major background in searches for many rare processes in the Standard Model (SM) and in new physics models beyond the SM.

Currently, the most precise fixed-order calculation in quantum chromodynamics (QCD) for top quark pair production reaches the next-to-next-to-leading order (NNLO) [1–9]. The high precision of the NNLO result is sufficient for the total cross section and generic phase-

Received 11 March 2020, Published online 17 June 2020

* L. L. Yang and X. Wang are supported in part by the National Natural Science Foundation of China (11975030, 11635001, 11575004). X. Wang is also supported in part by the Cluster of Excellence PRISMA⁺ (project ID 39083149). D. J. S. is supported under the ERC grant ERC-STG2015-677323. The work of D. P. and I. T. is supported by the Alexander von Humboldt Foundation, in the framework of the Sofja Kovalevskaja Award Project "Event Simulation for the Large Hadron Collider at High Precision". D. P. is also supported by the Deutsche Forschungsgemeinschaft (DFG) under Germany's Excellence Strategy – EXC 2121 "Quantum Univers" – 390833306. I. T. is also supported by the Swedish Research Council under contract number 2016-05996. The research of A. M. and A. P. has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (683211) as well as from UK STFC grants ST/L002760/1 and ST/K004883/1. The work of M. C. was supported in part by a grant of the BMBF and by the Deutsche Forschungsgemeinschaft under grant 396021762 - TRR 257

1) E-mail: yanglilin@zju.edu.cn



Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Article funded by SCOAP³ and published under licence by Chinese Physical Society and the Institute of High Energy Physics of the Chinese Academy of Sciences and the Institute of Modern Physics of the Chinese Academy of Sciences and IOP Publishing Ltd

space regions of differential distributions. However, in certain extreme kinematic regions, it is desirable to include even high-order corrections, for instance through resummation. In particular, top quarks produced at the LHC may have energies that are considerably larger than their rest mass m_t , because of the large collider energy. Such boosted top quarks require dedicated studies beyond the fixed-order calculations because the NNLO results in this regime are sensitive to the choice of factorization and renormalization scales [7]. This scale dependence can be significantly reduced by resumming two kinds of large logarithms to all orders in the strong coupling [10]. The first are threshold logarithms, which are important when the partonic center-of-mass energy is close to the $t\bar{t}$ invariant mass $M_{t\bar{t}}$, and the second are the small-mass logarithms of the form $\ln^n(m_t^2/M_{t\bar{t}}^2)$, which are enhanced in the boosted region $M_{t\bar{t}} \gg m_t$.

Besides QCD corrections, at high energies, EW corrections also become important [11–28]. In [26], the complete next-to-leading order (NLO) corrections of QCD and EW origin were combined with the NNLO QCD results using the multiplicative approach (denoted as QCD×EW in this paper). Their results show that at high transverse momentum, the EW effects can significantly reduce the differential cross section, and they should be considered for the accurate modeling of the spectrum.

In this study, we combined four calculations for the differential cross sections in $t\bar{t}$ production:

1. The NNLO QCD calculation of [1–9];
2. the soft gluon resummation of [29] at next-to-next-to-leading logarithmic (NNLL) accuracy;
3. the boosted soft gluon resummation of [30–32] at NNLL' accuracy; and
4. the complete-NLO predictions of QCD and EW origin [26,28,33].

In this work, all the currently-available perturbative contributions to these observables are combined. Therefore, the results presented here are the state-of-the-art theoretical predictions from the perturbation theory within the SM. To the best of our knowledge, this is the first study to perform such a complicated combination of radiative corrections.

2 Matching of various corrections

In this section, we briefly introduce the four sets of corrections used in the combination and then discuss the matching procedure employed to combine them. The matching procedure is necessary to remove the overlap of various calculations.

To illustrate the idea, we discuss the matching procedure for the particular case of the invariant-mass distribution; the combination procedure for other distributions follows exactly the same pattern. We start with the factor-

ization formula

$$\frac{d\sigma(\tau)}{dM_{t\bar{t}}} = \frac{8\pi\beta_t}{3sM_{t\bar{t}}} \sum_{ij} \int d\Theta \int_{\tau}^1 \frac{dz}{z} \mathcal{L}_{ij}(\tau/z, \mu_f) \times C_{ij}(z, M_{t\bar{t}}, m_t, \Theta, \mu_f), \quad (1)$$

where $M_{t\bar{t}}$ is the invariant mass of the $t\bar{t}$ pair; $\tau \equiv M_{t\bar{t}}^2/s$ and $z \equiv M_{t\bar{t}}^2/\hat{s}$, with \sqrt{s} and $\sqrt{\hat{s}}$ being the hadronic and partonic center-of-mass energies, respectively; $\beta_t = \sqrt{1 - 4m_t^2/M_{t\bar{t}}^2}$; $\mathcal{L}_{ij}(x, \mu_f)$ is the parton luminosity function, where μ_f is the factorization scale; C_{ij} is the partonic hard-scattering kernel where Θ is used to collectively denote additional kinematic variables. The sum in the above formula runs over the initial-state partons $i, j = q, \bar{q}, g$, and the prefactor is introduced by convention.

It is convenient to perform a Mellin transform of Eq. (1) with respect to τ . After the transform, the differential cross section becomes

$$\frac{d\tilde{\sigma}(N)}{dM_{t\bar{t}}} = \frac{8\pi\beta_t}{3sM_{t\bar{t}}} \sum_{ij} \int d\Theta \tilde{\mathcal{L}}_{ij}(N, \mu_f) \tilde{C}_{ij}(N, M_{t\bar{t}}, m_t, \Theta, \mu_f), \quad (2)$$

where N is the Mellin moment, and the functions with a tilde are the Mellin transform of the corresponding functions in Eq. (1). In the following, we deal with the perturbative contributions to the hard-scattering kernel \tilde{C}_{ij} within the SM.

In the fixed-order perturbation theory, \tilde{C}_{ij} can be expanded as a double series in the strong coupling constant α_s and the fine-structure constant α . The NNLO QCD result contains the α_s^2 , α_s^3 , and α_s^4 terms in the expansion; while the complete-NLO result includes the α_s^2 , $\alpha_s\alpha$, α^2 terms at the leading order (LO), and the α_s^3 , $\alpha_s^2\alpha$, $\alpha_s\alpha^2$, α^3 terms at NLO. Such a fixed-order expansion is formally correct in the generic phase-space regions. However, in certain kinematic limits, the fixed-order expansion breaks down due to the appearance of large logarithms at each order in the perturbation theory. In such cases, especially for pure QCD, the resummation of these logarithms is necessary to avoid bad convergence and/or the large scale dependence of the fixed-order results.

One of the limits in which potentially large logarithms arise is the threshold limit $z \rightarrow 1$ in the momentum space that corresponds to the $N \rightarrow \infty$ limit in the Mellin space. In this limit, the Mellin-space hard-scattering kernel \tilde{C}_{ij} develops large logarithms of the form $\alpha_s^n L^k$, where $L \sim \ln N$. The all-order resummation of these logarithms was studied in [29,34,35]. In [29], the resummation was carried out at the NNLL accuracy with the soft scale chosen in the momentum space. The NNLL threshold resummation was re-evaluated in [10,31] with the soft scale chosen in the Mellin space, to match the settings used in the boosted-soft resummation. By ignoring the technical

subtleties such as matrix-formed renormalization group (RG) evolution, the resummed hard-scattering kernel in the Mellin space can be schematically written as

$$\tilde{c}_{ij} \sim \alpha_s^2 h_{ij}(\alpha_s) \exp[g_{ij}(\alpha_s, \alpha_s L)] + O(1/N), \quad (3)$$

for $ij = q\bar{q}, \bar{q}q, gg$. All other partonic channels are power-suppressed in the $N \rightarrow \infty$ limit. The coefficient function $h_{ij}(\alpha_s)$ comes from the fixed-order calculation of the hard and soft functions [29], while the exponent $g_{ij}(\alpha_s, \alpha_s L)$ comes from the RG evolution. At NNLL accuracy, $h_{ij}(\alpha_s)$ needs to be evaluated up to NLO, namely, order α_s^1 . For the exponent g_{ij} , $\alpha_s L \sim 1$ or $L \sim 1/\alpha_s$, and the orders α_s^{-1} , α_s^0 and α_s^1 are maintained. By doing this, all terms of the form $\alpha_s^n L^m$ resummed into the exponent, with $n-1 \leq m \leq n+1$. Upon expansion, this generates the terms $\alpha_s^n L^k$ in the fixed-order perturbation theory up to $k = 2n$. The various elements required for achieving this level of logarithmic accuracy are given in [29,36,37]. In this work, this result is denoted as NNLL_m, where the subscript "m" means "massive", to indicate that full dependence on the top quark mass is retained.

In the threshold resummation framework discussed above, additional large logarithms of the form $\alpha_s^n \ln^l(m_t^2/M_{H}^2)$ ($l \leq 2n$) might arise in the boosted limit $M_{H} \gg m_t$ or $\beta_t \rightarrow 1$. In this limit both the top and anti-top quarks are highly boosted in the $t\bar{t}$ rest frame. In [30], a framework was developed to simultaneously resum the two kinds of logarithms $\ln N$ and $\ln(m_t^2/M_{H}^2)$. The form of the resummed result is very similar to Eq. (3), with much more complicated functions h_{ij} and g_{ij} . In addition, $O(m_t^2/M_{H}^2)$ power corrections are neglected in this boosted-soft resummation. With the elements evaluated in [30,38-40], the boosted-soft resummation was carried out at the NNLL' accuracy in [31], where the prime indicates that the coefficient function h_{ij} has to be evaluated to one order higher, namely, to NNLO or order α_s^2 . The net effect of computing the h_{ij} functions to one higher order is that the resummation captures the effect of one additional logarithm at each order in α_s . In the following, we will denote this result as NNLL'_b, where "b" implies "boosted."

Table 1 lists the contributions included (and not in-

cluded) in the four types of corrections used in the final combination. One immediately sees that there are overlaps among them, particularly among the three QCD-based calculations. These need to be carefully removed to avoid double-counting or triple-counting of certain sets of corrections. This matching was done in [10,31] for the three purely QCD contributions. We first combine the NNLL'_b and the NNLL_m results to obtain an NNLL'_{b+m} result. For this, we need to remove the overlap between the NNLL'_b and NNLL_m results to all orders in α_s . This can be done by exploiting the fact that the boosted-soft resummation formula is the small-mass limit of the soft-gluon resummation formula at any given order in α_s . Therefore,

$$d\sigma^{\text{NNLL}'_{b+m}} = d\sigma^{\text{NNLL}'_b} + (d\sigma^{\text{NNLL}_m} - d\sigma^{\text{NNLL}_m}|_{m_t \rightarrow 0}), \quad (4)$$

where the terms in the parenthesis account for the contributions suppressed by $\alpha_s^n m_t^2/M_{H}^2$ for $n > 2$.

Subsequently, the matching with the NNLO QCD calculation proceeds by subtracting the NNLO expansion of the resummed formula

$$d\sigma^{\text{NNLO}+\text{NNLL}'} = d\sigma^{\text{NNLL}'_{b+m}} + \left(d\sigma^{\text{NNLO}} - d\sigma^{\text{NNLL}'_{b+m}} \Big|_{\text{NNLO expansion}} \right), \quad (5)$$

where the terms in the parentheses account for contributions which are suppressed by α_s^n/N for $n = 3, 4$.

Finally, the complete-NLO contributions can be incorporated by first combining them with the NNLO QCD contributions in the multiplicative approach,¹⁾ arriving at the QCD×EW result, and then matching them against the resummation results as in Eq. (5). This gives us the final matching formula:

$$d\sigma^{\text{QCD} \times \text{EW} + \text{NNLL}'} = d\sigma^{\text{NNLL}'_{b+m}} + \left(d\sigma^{\text{QCD} \times \text{EW}} - d\sigma^{\text{NNLL}'_{b+m}} \Big|_{\text{NNLO expansion}} \right). \quad (6)$$

Table 1. Contributions included in the four types of corrections in the combination. See the text for detailed explanation.

	Included	Not included
NNLO QCD	$\alpha_s^n (n = 2, 3, 4)$	$\alpha_s^n (n > 4); \alpha_s^n \alpha^m (m > 0)$
NNLL _m	$\alpha_s^n \ln^k N (n \geq 2)$	$\alpha_s^n O(1/N) (n > 2); \alpha_s^n \alpha^m (m > 0)$
NNLL' _b	$\alpha_s^n \ln^k N \ln^l(m_t^2/M_{H}^2) (n \geq 2)$	$\alpha_s^n O(1/N, m_t^2/M_{H}^2) (n > 2); \alpha_s^n \alpha^m (m > 0)$
Complete NLO	$\alpha_s^n \alpha^m (n + m = 2, 3)$	$\alpha_s^n \alpha^m (n + m > 3)$

1) Orders $\alpha_s^n \alpha^m (n + m \leq 3)$ and α_s^4 are summed and, bin-by-bin in any distributions, the order $\alpha_s^3 \alpha$ contribution is approximated via rescaling the order $\alpha_s^2 \alpha$ contribution by the NLO QCD K -factor.

3 Numerical results

In this section, we present the numerical results based on the matching formula Eq. (6) and compare them with the predictions of previous studies. For all the results, we take the top quark mass as $m_t=172.5$ GeV. The results for other top quark masses can be obtained from the authors

$$\mu_r = \mu_f = \begin{cases} \frac{m_{T,t}}{2} \equiv \frac{1}{2} \sqrt{p_{T,t}^2 + m_t^2} & \text{for } p_{T,t} \text{ distribution} \\ \frac{H_T}{4} \equiv \frac{1}{4} \left(\sqrt{p_{T,t}^2 + m_t^2} + \sqrt{p_{T,\bar{t}}^2 + m_t^2} \right) & \text{for all other distributions} \end{cases},$$

$$\mu_h = \frac{H_T}{2}, \quad \mu_s = \frac{H_T}{\bar{N}} \equiv \frac{H_T}{N e^{\gamma_E}}, \quad \mu_{dh} = m_t, \quad \mu_{ds} = \frac{m_t}{\bar{N}}. \quad (7)$$

For the meanings of these scales, refer to [10,31]. The variations in the unphysical scales around the default values listed above are used to estimate the impact of the higher order corrections that are not included in the calcu-

upon request. For purely QCD-based predictions, we used the NNPDF3.1 NNLO PDF sets with $\alpha_s(m_Z)=0.118$ [41]. When EW corrections are included, we used the NNPDF3.1 NNLO LUXQED PDF sets [42] with the same $\alpha_s(m_Z)$. There are a few unphysical scales in the fixed-order and resummed calculations. Their default choices in the results shown below are

lations. The scales are varied individually up and down by a factor of 2, and the resulting variations in the cross section are added in the quadrature to give the combined uncertainty [10,31]. For the settings on the EW paramet-




Fig. 1. (color online) Theoretical predictions compared with CMS data in the di-lepton channel [43].

ers, refer to [26]. The NLO EW calculation was performed by using the latest public version of MADGRAPH5_AMC@NLO [33].

Fig. 1 shows the predictions for the distributions differential with respect to

- i) the $t\bar{t}$ invariant mass $M_{t\bar{t}}$,
- ii) the transverse momentum $p_{T,t}$ of the top quark,
- iii) the rapidity $Y_{t\bar{t}}$ of the $t\bar{t}$ pair,
- iv) and the rapidity y_t of the top quark.

The four kinds of vertical bars correspond to the four kinds of theoretical predictions discussed in the last section: NNLO QCD, NNLO+NNLL', QCD×EW, and QCD×EW+NNLL'. The bands in red correspond to the CMS measurement in the di-lepton channel at the 13 TeV LHC using 35.9 fb^{-1} of data [43].

From the plots, one can conclude that the predictions are generically stable against the inclusion of various sets of corrections. This indicates that the convergence of the perturbative series and the estimate of the residual theoretical uncertainty affecting the predictions are well under-control. The effects of including QCD resummation and EW corrections are more evident in the large $M_{t\bar{t}}$ region and in the high $p_{T,t}$ tail (see inset in the first and second panel in Fig. 1). In these cases, QCD resummation and EW corrections both tend to reduce the differential cross sections, which appear to be more compatible with the experimental data than when the corrections are not included. In addition, the resummation effects enlarge the scale uncertainty in the first $M_{t\bar{t}}$ bin near the $2m_t$ threshold, where a small discrepancy is present between the theoretical predictions and the experimental measurement, thus slightly reducing the discrepancy. A recent study [44] showed that a Coulomb resummation can sig-

nificantly enhance the differential cross section in this region and can partly resolve the discrepancy.¹⁾ In principle, Coulomb resummation can be combined with the results of this work to provide a good description of the $M_{t\bar{t}}$ spectrum in the whole phase space. For the rapidity distributions, we can see that all the theoretical predictions almost completely lie within the uncertainty bands associated with the experimental measurements, indicating excellent agreement for this observable. While the effect of the resummation on the uncertainty bands for these observables is minimal, we can see that the effect of the higher order terms captured by the resummation slightly soften the rapidity spectrum compared with the corresponding fixed order predictions.

4 Summary

In this paper, we described a combination of four calculations for the differential cross sections in $t\bar{t}$ production: the NNLO QCD calculations, NNLL QCD threshold resummation, NNLL' QCD resummation for boosted top quarks, and complete-NLO predictions of QCD and EW origin. To the best of our knowledge, this is the first work to present such a complicated combination. The result represents the state-of-the-art prediction in the perturbation theory for $t\bar{t}$ differential distributions within the SM, which includes all sets of corrections available at the moment. Numerical results are presented for the invariant-mass distribution, transverse-momentum distribution, and rapidity distributions. We compared our predictions with the CMS measurements in the di-lepton channel at the 13 TeV LHC with an integrated luminosity of 35.9 fb^{-1} , and we found good agreement overall.

References

- 1 P. Bärnreuther, M. Czakon, and A. Mitov, *Phys. Rev. Lett.*, **109**: 132001 (2012), arXiv:1204.5201[hep-ph]
- 2 M. Czakon and A. Mitov, *JHEP*, **1212**: 054 (2012), arXiv:1207.0236[hep-ph]
- 3 M. Czakon and A. Mitov, *JHEP*, **1301**: 080 (2013), arXiv:1210.6832[hep-ph]
- 4 M. Czakon, P. Fiedler, and A. Mitov, *Phys. Rev. Lett.*, **110**: 252004 (2013), arXiv:1303.6254 [hep-ph]
- 5 M. Czakon, P. Fiedler, and A. Mitov, *Phys. Rev. Lett.*, **115**(5): 052001 (2015), arXiv:1411.3007[hep-ph]
- 6 M. Czakon, D. Heymes, and A. Mitov, *Phys. Rev. Lett.*, **116**(8): 082003 (2016), arXiv:1511.00549[hep-ph]
- 7 M. Czakon, D. Heymes, and A. Mitov, *JHEP*, **1704**: 071 (2017), arXiv:1606.03350[hep-ph]
- 8 S. Catani, S. Devoto, M. Grazzini *et al.*, *Phys. Rev. D*, **99**(5): 051501 (2019), arXiv:1901.04005[hep-ph]
- 9 S. Catani, S. Devoto, M. Grazzini *et al.*, *JHEP*, **1907**: 100 (2019), arXiv:1906.06535[hep-ph]
- 10 M. Czakon, A. Ferroglia, D. Heymes *et al.*, *JHEP*, **1805**: 149 (2018), arXiv:1803.07623[hep-ph]
- 11 W. Beenakker, A. Denner, W. Hollik *et al.*, *Nucl. Phys. B*, **411**: 343 (1994)
- 12 W. Bernreuther, M. Fückner, and Z. G. Si, *Phys. Lett. B*, **633**: 54(2006) Erratum: [*Phys. Lett. B*, **644**: 386(2007)] [hep-ph/0508091]
- 13 J. H. Kuhn, A. Scharf, and P. Uwer, *Eur. Phys. J. C*, **45**: 139 (2006), arXiv:hep-ph/0508092
- 14 W. Bernreuther, M. Fückner, and Z. G. Si, *Phys. Rev. D*, **74**: 113005 (2006), arXiv:hep-ph/0610334
- 15 J. H. Kuhn, A. Scharf, and P. Uwer, *Eur. Phys. J. C*, **51**: 37 (2007), arXiv:hep-ph/0610335
- 16 W. Hollik and M. Kollar, *Phys. Rev. D*, **77**: 014008 (2008), arXiv:0708.1697[hep-ph]
- 17 W. Bernreuther, M. Fückner, and Z. G. Si, *Phys. Rev. D*, **78**: 017503 (2008), arXiv:0804.1237 [hep-ph]
- 18 W. Bernreuther and Z. G. Si, *Nucl. Phys. B*, **837**: 90 (2010), arXiv:1003.3926[hep-ph]
- 19 W. Hollik and D. Pagani, *Phys. Rev. D*, **84**: 093003 (2011),

1) Note that the differential cross section in the first bin is very sensitive to m_t . The discrepancy here decreases for smaller values of m_t .

- arXiv:1107.2606[hep-ph]
- 20 J. H. Kuhn and G. Rodrigo, JHEP, **1201**: 063 (2012), arXiv:1109.6830[hep-ph]
 - 21 A. V. Manohar and M. Trott, Phys. Lett. B, **711**: 313 (2012), arXiv:1201.3926[hep-ph]
 - 22 W. Bernreuther and Z. G. Si, Phys. Rev. D, **86**: 034026 (2012), arXiv:1205.6580[hep-ph]
 - 23 J. H. Kühn, A. Scharf, and P. Uwer, Phys. Rev. D, **91**(1): 014020 (2015), arXiv:1305.5773 [hep-ph]
 - 24 J. M. Campbell, D. Wackeroth, and J. Zhou, PoS DIS, **2015**: 130 (2015), arXiv:1508.06247 [hep-ph]
 - 25 D. Pagani, I. Tsinikos, and M. Zaro, Eur. Phys. J. C, **76**(9): 479 (2016), arXiv:1606.01915 [hep-ph]
 - 26 M. Czakon, D. Heymes, A. Mitov *et al.*, JHEP, **1710**: 186 (2017), arXiv:1705.04105[hep-ph]
 - 27 M. Czakon, D. Heymes, A. Mitov *et al.*, Phys. Rev. D, **98**(1): 014003 (2018), arXiv:1711.03945[hep-ph]
 - 28 C. Gütschow, J. M. Lindert, and M. Schönherr, Eur. Phys. J. C, **78**(4): 317 (2018), arXiv:1803.00950[hep-ph]
 - 29 V. Ahrens, A. Ferroglia, M. Neubert *et al.*, JHEP, **1009**: 097 (2010), arXiv:1003.5827[hep-ph]
 - 30 A. Ferroglia, B. D. Pecjak, and L. L. Yang, Phys. Rev. D, **86**: 034010 (2012), arXiv:1205.3662 [hep-ph]
 - 31 B. D. Pecjak, D. J. Scott, X. Wang *et al.*, Phys. Rev. Lett., **116**(20): 202001 (2016), arXiv:1601.07020[hep-ph]
 - 32 B. D. Pecjak, D. J. Scott, X. Wang *et al.*, JHEP, **1903**: 060 (2019), arXiv:1811.10527[hep-ph]
 - 33 R. Frederix, S. Frixione, V. Hirschi *et al.*, JHEP, **1807**: 185 (2018), arXiv:1804.10017[hep-ph]
 - 34 N. Kidonakis and G. F. Sterman, Phys. Lett. B, **387**: 867 (1996)
 - 35 N. Kidonakis and G. F. Sterman, Nucl. Phys. B, **505**: 321 (1997), arXiv:hep-ph/9705234
 - 36 A. Ferroglia, M. Neubert, B. D. Pecjak *et al.*, Phys. Rev. Lett., **103**: 201601 (2009), arXiv:0907.4791[hep-ph]
 - 37 A. Ferroglia, M. Neubert, B. D. Pecjak *et al.*, JHEP, **0911**: 062 (2009), arXiv:0908.3676[hep-ph]
 - 38 A. Ferroglia, B. D. Pecjak, L. L. Yang *et al.*, JHEP, **1210**: 180 (2012), arXiv:1207.4798[hep-ph]
 - 39 A. Ferroglia, B. D. Pecjak, and L. L. Yang, JHEP, **1309**: 032 (2013), arXiv:1306.1537[hep-ph]
 - 40 A. Broggio, A. Ferroglia, B. D. Pecjak *et al.*, JHEP, **1412**: 005 (2014), arXiv:1409.5294[hep-ph]
 - 41 R. D. Ball *et al.* (NNPDF Collaboration), Eur. Phys. J. C, **77**(10): 663 (2017), arXiv:1706.00428[hep-ph]
 - 42 V. Bertone *et al.* (NNPDF Collaboration), SciPost Phys., **5**(1): 008 (2018), arXiv:1712.07053[hep-ph]
 - 43 A. M. Sirunyan *et al.* (CMS Collaboration), JHEP, **1902**: 149 (2019), arXiv:1811.06625 [hep-ex]
 - 44 W. L. Ju, G. Wang, X. Wang *et al.*, accepted by Chinese Physics C, arXiv: 1908.02179[hep-ph]; accepted by JHEP, arXiv:2004.03088[hep-ph]