

Fornax globular cluster orbits in different models of dark matter

Nitsan Bar,¹ Diego Blas,² Kfir Blum,¹ and Hyungjin Kim^{3,1}

¹*Department of Particle Physics and Astrophysics,
Weizmann Institute of Science, Rehovot 7610001, Israel*

²*Theoretical Particle Physics and Cosmology Group, Department of Physics,
King's College London, Strand, London WC2R 2LS, UK*

³*DESY, Notkestrasse 85, 22607 Hamburg, Germany*

(Dated: February 15, 2021)

We investigate what the orbits of globular clusters (GCs) in the Fornax dwarf spheroidal (dSph) galaxy can teach us about dark matter (DM), both with respect to DM microphysics and to its density profile. This problem was addressed recently for ultralight dark matter (ULDM). We consider two additional models: (i) fermionic degenerate dark matter (DDM), where Pauli blocking must be taken into account in the dynamical friction computation; and (ii) self-interacting dark matter (SIDM). We give a simple and direct Fokker-Planck derivation of dynamical friction, new in the case of DDM and reproducing previous results in the literature for ULDM and cold DM. ULDM, DDM and SIDM were considered in the past as leading to cores in dSphs, a feature that acts to suppress dynamical friction and prolong GC orbits. For DDM we derive a robust version of the cosmological free streaming limit that is independent of the assumed DM production mechanism, finding that DDM cannot produce an appreciable core in Fornax without violating Ly- α limits. If the Ly- α limit is discounted for some reason, then stellar kinematics data does allow a DDM core which could prolong GC orbits. For SIDM we find that significant prolongation of GC orbits, consistent with stellar kinematics, could be obtained for values of the self-interaction cross section considered in previous works. In addition to reassessing the inspiral time using updated observational data, we give a new perspective on the so-called GC timing problem, demonstrating that for a cuspy cold DM profile dynamical friction predicts a zero redshift (current) radial cumulative distribution function for GCs that is independent of the poorly known initial conditions. The observed projected orbits of Fornax GCs is consistent with this expectation with a mild apparent fine-tuning at the level of $\sim 25\%$, associated with the innermost GCs. We also assess observational uncertainties, notably projection and phase-space effects, that enter the estimation of GC orbits and inspiral times.

CONTENTS

I. Introduction	1	A. Dynamical friction in exotic media: derivation from the Boltzman equation	14
II. Dynamical friction: microphysics	3	1. A classical gas medium	15
A. Classical gas	3	2. Degenerate fermionic dark matter	16
B. Degenerate Fermi gas	3	3. Ultralight dark matter	16
C. Bose gas	4	B. Maximum entropy DDM halos	17
III. Dynamical friction in a CDM halo: core vs. cusp	4	C. Jeans modeling	18
IV. Degenerate dark matter (DDM)	6	D. CDM velocity dispersion in a cored profile	19
A. Stellar LOSVD constraints on DDM in Fornax	7	E. Orbits under dynamical friction	19
B. Structure formation constraints on DDM	8	F. The radial and projected CDF of GCs	20
C. Orbital decay time in DDM	9	1. CDF of GCs in a cuspy halo	20
V. Self-interacting dark matter (SIDM)	9	2. CDF of GCs in a cored halo	21
VI. Discussion	11	3. Accounting for distribution of GC masses	21
A. Comments regarding GC data used in this work	11	4. Projected radius distribution	21
B. Statistical discussion	13	G. Exploration of initial conditions	22
VII. Summary	14	References	24
Acknowledgments	14		

I. INTRODUCTION

The Milky Way dwarf spheroidal (dSph) satellite galaxies are broadly believed to be dominated by dark

matter (DM) [1, 2]¹, and this fact combined with their small sizes and near-by locations makes them interesting test beds of the small-scale behaviour of DM [4–10]. In fact, some of the basic predictions of the most commonly considered paradigm of DM — collisionless cold dark matter (CDM) — may be in tension with observations (see, e.g., [10, 11]). Conclusive kinematic data for a decisive test of CDM in dSphs is difficult to obtain, but upcoming observatories may supply it [12].

One intriguing puzzle about the dSph galaxies concerns the globular clusters (GCs) of the Fornax dSph [13]: some of Fornax’s six known GCs [2, 14] have orbital decay times due to dynamical friction (DF) which seem to fall significantly short of their age [13]. If estimated naïvely based on the Chandrasekhar formula [15], assuming the usual CDM cusp density profile (see, e.g. [16]), one obtains an instantaneous DF time of less than 1 Gyr for the most troublesome GC4 [17]. On the other hand, the stellar content of the GCs is old, > 10 Gyr [18, 19], as is much of the stellar content of Fornax itself [20, 21]. It may seem unlikely then, that we observe some of the GCs just a short time before they fall to the center of the galaxy. We show a visualization of this problem in Fig. 1.




FIG. 1. Orbital radius vs. time, calculated for the Fornax GC4 assuming a slightly eccentric orbit. $t = 0$ represents today. The orbit calculation assumes the CDM Navarro-Frenk-White (NFW) [22] profile of [16].

Many explanations were suggested for the Fornax GC timing puzzle [17, 23–28], of which an exciting class of ideas entails a modification to the nature of dark matter, going beyond CDM [17, 29, 30]. In particular, Refs. [17, 31, 32] studied ultralight dark matter (ULDM) and showed that in the particle mass window $m \lesssim 10^{-21}$ eV, ULDM would suppress DF enough to solve the timing puzzle. However, most of this mass range for ULDM has been scrutinised in the last few years, resulting with disfavoured evidence [33–37].

Motivated by the fact that the combination of GC age and orbit measurements probes the details of the DM halo and microphysics, we extend the DF analysis to additional DM models. The first model is degenerate dark matter (DDM), in which the phase space distribution of DM in dSph cores is affected by Pauli blocking [38, 39]. The second model is self-interacting DM (SIDM), in which self interactions between DM particles produce a cored isothermal distribution.

In Sec. II we focus on the microphysics and calculate DF for CDM, DDM, and ULDM. Our results for DDM are new; for ULDM, we make contact with a different derivation in the literature; while for SIDM the microphysics of the DF calculation is argued to be similar to that in CDM.

All three DM models can, in principle, naturally produce cored isothermal halos. As we show in Sec. III, a cored isothermal distribution of DM suppresses DF in part due to a phase-space effect (associated with the “core stalling” [40] identified in past numerical work), as the velocity of the inspiraling GC can become parametrically lower than the DM velocity dispersion.

For ULDM, DF and the Fornax GC timing puzzle were studied in recent works [17, 31, 32] and we do not review them again. As noted above, constraints from galaxy dynamics and from cosmological Ly- α analyses suggest a similar behaviour to CDM.

For DDM (Sec. IV), we formulate a robust version of the Ly- α bound that is insensitive to DM model building and cosmological history, finding that it disfavors an appreciable core. If one chooses to discount the Ly- α bound (see, e.g. [17] for a qualitative discussion of concerns regarding systematic uncertainties), then stellar kinematics does allow a considerable DDM core which could lead to significant suppression of DF and prolong the settling time of the innermost GCs.

For SIDM (Sec. V), stellar kinematics allows a considerable core. If the SIDM cross section is as large as that considered in [41], then the DF settling time for the innermost GCs can be significantly longer than in the cuspy halo CDM model.

Part of the scope of this work is to give an analytical perspective on DF. We start this analysis in Sec. III, in which we track GC orbits in cuspy and cored halos, and continue in App. F and in the Discussion (Sec. VI) with a derivation of the time evolution of a population of GCs. When the dust settles we obtain reasonably robust predictions for the late-time distribution of GCs. Our conclusion (summarised in the second half of Sec. VI) is that even for a cuspy CDM halo, when one takes into account that Fornax hosts not just one, but a collection of GCs, then the timing puzzle boils down to a rather mild (but quite persistent) fine-tuning in the ballpark of 25% or so.

Our approach is mostly analytical. Of course, this has limitations and one may be justified in expecting that conclusive results would require numerical simulations. In most current simulations, however, both GC formation

¹ See, however, a contrary claim in [3].

and DF correspond to sub-grid phenomena. A state-of-the-art example of numerical simulation work targeting the GC timing puzzle in the context of CDM was recently published by Ref. [42]. According to Ref. [42], reasonable initial conditions for the GCs (derived from the simulations) can naturally lead to the observed present configuration in a standard cuspy CDM halo. The timing puzzle may thus be even less significant than the mild 25% that we find with analytical tools. However, the fact remains that both GC formation and DF were treated in [42] at the sub-grid and post-processing level, and it is not clear (to us) if the resolution of the simulation was high enough to resolve the small-scale region of Fornax, that is detrimental for the dynamics of the innermost GCs. Moreover, we also feel that theoretical insight about the role that initial conditions play in shaping the present distribution of GCs is important. For these reasons we believe that our analytical approach remains useful.

We summarize in Sec. VII. Many details of the calculations are deferred to the Appendices.

II. DYNAMICAL FRICTION: MICROPHYSICS

Dynamical friction can be described in terms of the Fokker-Planck theory for the motion of a probe particle (a GC in our case) traveling through a gas of spectator particles (DM particles in our case). In App. A we derive the Fokker-Planck equation as the small-momentum-exchange limit of the Boltzmann equation, governing the motion of a probe object in different background media, accounting for the gravitational interaction between the probe and the medium particles. Our calculation is direct, in the sense that it simply amounts to computing the collision integral while taking care to account for the quantum statistics of spectator gas particles. Here we bypass the details of the calculation, while utilising the main results.

The Fokker-Planck equation is characterised by a set of momentum space diffusion coefficients, calculated in App. A 1, A 2, and A 3 for the case of a medium composed of a classical gas, degenerate Fermi gas, and Bose gas, respectively. Of particular importance for our analysis is the diffusion coefficient $D_{||}$, corresponding to the diffusion in momentum parallel to the probe object's instantaneous velocity. The DF deceleration acting on a probe with mass m_* moving with instantaneous velocity \mathbf{V} w.r.t. the medium is computed as [43]

$$\begin{aligned} \frac{d\mathbf{V}}{dt} &= \frac{D_{||}}{m_*} \hat{\mathbf{V}} \\ &= -\frac{4\pi G^2 m_* \rho}{V^3} C \mathbf{V}. \end{aligned} \quad (1)$$

In the second line, to compare the DF arising in different types of medium we define the dimensionless coefficient C as follows [17],

$$C = -\frac{V^2 D_{||}}{4\pi G^2 m_*^2 \rho}, \quad (2)$$

where ρ is the mass density of the medium.

Different microphysics properties of the medium (in our case, the DM galactic halo) predict different results for C . In the next subsection we discuss three scenarios.

A. Classical gas

This is the appropriate limit for a halo composed of a gas of classical particles. We will adopt this limit to describe DF in the ordinary CDM model, as well as for the SIDM model². For a homogeneous classical gas with an isotropic distribution function $f_v(v)$, DF is described by the Chandrasekhar formula [15] (see also App. A 1),

$$C_{\text{class}} = 4\pi \ln \Lambda \int_0^V dv_m v_m^2 f_v(v_m), \quad (3)$$

where $\ln \Lambda$ is the Coulomb logarithm. If the gas distribution function is a Maxwellian with velocity dispersion σ , $f_v(v) = (2\pi\sigma^2)^{-3/2} \exp(-v^2/(2\sigma^2))$, we have

$$\begin{aligned} C_{\text{Max}} &= \ln \Lambda \left(\text{erf}(X) - \frac{2X}{\sqrt{\pi}} e^{-X^2} \right) \\ &\rightarrow \ln \Lambda \begin{cases} 1 & V \gg \sigma \\ \frac{\sqrt{2}}{3\sqrt{\pi}} \frac{V^3}{\sigma^3} & V \ll \sigma \end{cases}, \end{aligned} \quad (4)$$

where $X \equiv V/(\sqrt{2}\sigma)$ and where in the second line we show the asymptotic scaling of C at large and small X .

B. Degenerate Fermi gas

This is the relevant limit for DF at the core of a halo supported by the degeneracy pressure of light fermionic DM (DDM model). In the high-degeneracy limit we have $f_v(v) = 3/(4\pi v_F^3) \theta(v_F - v)$, where $\theta(x)$ is the Heaviside function, the Fermi velocity v_F is related to the medium density via

$$\rho = \frac{gm^4 v_F^3}{6\pi^2}, \quad (5)$$

m is the mass of the particles and g the number of degrees of freedom (e.g. $g = 2$ for Weyl fermions). The calculation in App. A 2 gives the following limiting behaviour,

$$C_{\text{DDM}} \rightarrow \ln \Lambda \begin{cases} 1 & V \gg v_F \\ \frac{V^3}{v_F^3} & V \ll v_F \end{cases}. \quad (6)$$

Thus, in both limits $V \gg v_F$ and $v \ll v_F$, we find that DF in a degenerate medium is equivalent to DF in a

² This is a good approximation for the SIDM cross-sections of interest, which are small enough such that SIDM particles travel across distances larger than the size of the system without colliding with each other. See Sec. V.

classical medium with the replacement $\sigma \rightarrow \left(\frac{2}{9\pi}\right)^{\frac{2}{3}} v_F \approx 0.17v_F$. Note that the 3-dimensional velocity dispersion associated with the classical isotropic Maxwellian distribution is $\langle v_x^2 + v_y^2 + v_z^2 \rangle = \langle v^2 \rangle = 3\sigma^2$, while the dispersion for the degenerate distribution is $\langle v^2 \rangle = (3/5)v_F^2$. Therefore, the pressure in the different types of medium matches when $v_F \approx 2.2\sigma$. Similarly, Eqs. (6) and (4) tell us that DF in these media match when $v_F \approx 5.8\sigma$.

As an aside, it is interesting to note that if one inserts the zero temperature velocity distribution of degenerate fermions into Eq. (3), one obtains exactly Eq. (6). This is somewhat surprising, because the Fokker-Planck calculation took into account Pauli exclusion in the medium whereas Eq. (3) does not. Moreover, according to Eq. (3), only particles with velocities smaller than the probe object's contribute to the DF. For the case of degenerate matter, one would have expected that the opposite should happen: only particles close to the Fermi surface contribute to DF. We refer the reader again to App. A 2 for the detailed computation that leads us to Eq. (6).

Finally, note that above we evaluated DF in the zero-temperature limit and not in the finite-temperature limit. In Sec. IV we consider a finite-temperature density profile, so we should keep this caveat in mind. We have not explored DF of degenerate matter within the more sophisticated treatment of Refs. [44, 45].

C. Bose gas

This is the relevant limit for the case where halo particles follow the Bose-Einstein statistics, as in the ULDM model. The diffusion coefficients can be obtained either by solving a Langevin equation with stochastic fluctuations of the gravitational potential [31] or, as we do in App. A 3, by using a kinetic equation³. Both approaches provide identical results.

Up to a slight modification of Coulomb logarithm, DF for the bosonic gas includes a contribution to the C term that is identical to that of the classical gas in Eq. (4). In addition to this, ULDM large-scale density fluctuations (manifested by Bose-enhancement terms in the kinetic theory computation) cause additional velocity drift that can be characterised by an extra term to $C \rightarrow C + \Delta C$, with⁴

$$\Delta C = \ln \Lambda \left(\frac{m_{\text{eff}}}{m_\star} \right) \left(\text{erf}(X_{\text{eff}}) - \frac{2X_{\text{eff}}}{\sqrt{\pi}} e^{-X_{\text{eff}}^2} \right), \quad (7)$$

where $m_{\text{eff}} = \pi^{3/2} \rho / (m\sigma)^3$ is the ULDM mass enclosed in an effective de Broglie volume and

$X_{\text{eff}} \equiv v/\sqrt{2}\sigma_{\text{eff}}$ with $\sigma_{\text{eff}} = \sigma/\sqrt{2}$. Numerically, $m_{\text{eff}} \approx 1.2 \times 10^6 (10^{-21} \text{ eV}/m)^3 [\rho/(3 \times 10^7 M_\odot/\text{kpc}^3)] [(10 \text{ km/s})/\sigma]^3 M_\odot$. With these numbers and keeping in mind a typical GC mass $m_\star \sim 10^5 M_\odot$, the ΔC effect becomes quantitatively important in Fornax for $m \lesssim 3 \times 10^{-20} \text{ eV}$.

The kinetic theory result summarised above assumed that the scale size of the system – e.g., the radius r of a GC orbit – is much larger than the effective de Broglie wavelength of the ULDM particles,

$$r_{\text{dB}} \approx \frac{2\pi}{m\sigma} \approx 300 \left(\frac{10 \text{ km/s}}{\sigma} \right) \left(\frac{10^{-21} \text{ eV}}{m} \right) \text{ pc}, \quad (8)$$

and thus much larger than ULDM quasi-particle excitations or than the soliton core that is ubiquitously found in ULDM simulations (see [17] for a review). For $r < r_{\text{dB}}$, the treatment above breaks down and must be modified by taking into account large-scale coherence effects of the ULDM. This can be done via solving the Schroedinger equation, as shown in [17, 32], which indeed found that DF becomes suppressed at $r \lesssim r_{\text{dB}}$. We refer the reader to [17, 32] for more details on DF and the Fornax GC puzzle in the context of ULDM. Here we only note that for $m \gtrsim 10^{-20} \text{ eV}$, where $r \gg r_{\text{dB}}$ and $m_{\text{eff}} \ll m_\star$ for the Fornax GCs, DF in the ULDM medium becomes quantitatively similar to DF in a classical medium.

III. DYNAMICAL FRICTION IN A CDM HALO: CORE VS. CUSP

It is natural to define an instantaneous DF time, τ , via

$$\tau = \frac{V^3}{4\pi G^2 m_\star \rho C}, \quad (9)$$

such that (including here only the DF effect)

$$\dot{\mathbf{V}} = -\frac{1}{\tau} \mathbf{V}. \quad (10)$$

A crude estimate of the time scale it would take a GC to settle down to the dynamical center of a halo can be obtained by computing τ , using the current instantaneous position and velocity of the GC. Assuming a CDM NFW distribution, and plugging an estimate of the dark matter density and velocity dispersion corresponding to the present observed position of each GC into Eqs. (4) and (9), the result we find is summarised in the column marked τ_{CDM} (highlighted in blue) in Tab. I. For GC3 and GC4 the DF time estimated in this way is 2.6 and 0.9 Gyr, respectively: much shorter than the age of the system.⁵

³ While this paper was being prepared for publication, Ref. [46] appeared which also presents a kinetic theory derivation of the ULDM diffusion coefficients.

⁴ Formally, the ΔC term is there also for standard CDM but is negligible unless the individual DM particles are extremely massive.

⁵ Our estimates are larger than those previously obtained in [17]; we discuss the differences in Sec. VI.

TABLE I. Some details of Fornax GCs. For the galactic center of Fornax we use an updated measurement [21], based on surface brightness modelling. This estimate is ≈ 160 pc off relative to the center defined by previous works [2, 16, 17, 19, 28, 42], leading to different projected radii of GCs. We set the distance to Fornax as 147 ± 4 kpc [18]. We estimate the error on r_\perp by propagating the distance error, added in quadrature with a 13 pc [21] uncertainty on the center. For relative radial velocities Δv_r , we use the galactic radial velocity $RV_{\text{Fornax}} = 55.46 \pm 0.63$ km/s [47] and set $\Delta v_r = RV_{\text{GC}} - RV_{\text{Fornax}}$, adding errors in quadrature. For GC6, the values correspond to a small sample of stars, likely contaminated by background [14]. $r_{c/h}$ refers to King radius for GC1-GC5 and half-light radius for GC6. The CDM DF time estimates are based on the NFW profile of [16], with the semi-analytic procedure, as explained in the text. The DF time of DDM and SIDM are based on the semi-analytic procedure and Secs. IV and V.

	m_\star [$10^5 M_\odot$]	r_\perp [kpc]	Δv_r [km/s]	$r_{c/h}$ [pc]	Refs.	τ_{CDM} [Gyr]	$\tau_{\text{DDM}}^{(135)}$ [Gyr]	τ_{SIDM} [Gyr]
GC1	0.42 ± 0.10	1.73 ± 0.05	3.54 ± 1.18	10.8 ± 0.3	[18, 19, 47–49]	119	122	79.3
GC2	1.54 ± 0.28	0.98 ± 0.03	3.9 ± 0.7	6.2 ± 0.2	[18, 19, 49, 50]	14.7	7.12	8.82
GC3	4.98 ± 0.84	0.64 ± 0.02	4.94 ± 0.66	1.7 ± 0.1	[18, 19, 51, 52]	2.63	1.48	2.21
GC4	0.76 ± 0.15	0.154 ± 0.014	-8.26 ± 0.64	1.9 ± 0.2	[18, 19, 51, 52]	0.91	10.7	14.8
GC5	1.86 ± 0.24	1.68 ± 0.05	3.93 ± 0.77	1.5 ± 0.1	[18, 19, 47, 51, 52]	32.2	30.1	20
GC6	~ 0.29	0.254 ± 0.015	-1.56 ± 1.36	12.0 ± 1.4	[14, 42]	5.45	16.1	22

However, estimating an orbital decay time from the instantaneous value of τ can be misleading. In a realistic galaxy, the DM phase space distribution and with it the instantaneous value of τ could change along the orbit of the GC. To obtain a better estimate of the actual settling time one could track the orbit of the GC semi-analytically, using the phase space-dependent value of τ along the orbit [26, 40, 53]. Some details of this calculation are given in App. E.

The semi-analytic integration reproduces results from N-body simulations [16]. To demonstrate this, we use Eqs. (4) and (9) while reading the CDM density and velocity dispersion from the N-body simulations of [16] to integrate the orbit of a GC. In Fig. 2 we compare our results to two different scenarios from [16].

The first scenario, denoted NFW, contains a cuspy NFW-like halo (the density profile of this model is shown in Fig. 11). The orbit of a GC in this halo is shown by the blue dashed line for the simulation of [16] and by a blue solid line for the semi-analytic tracking. The second scenario, denoted ISO, contains an isothermal core halo (also shown in Fig. 11). The GC orbits are shown by the red lines. Again, the semi-analytic method (solid) compares reasonably well with the simulation (dotted).

The results have a mild dependence on the choice of the Coulomb log, and we make slightly different choices for the different scenarios. For NFW we follow [25] in setting

$$\ln \Lambda_{\text{NFW}} = \ln \frac{b_{\text{max}} \sigma^2}{G m_\star}. \quad (11)$$

However, instead of the $b_{\text{max}} = 0.25$ kpc used in [25], we adopt $b_{\text{max}} = 0.5$ kpc. For ISO we follow [17, 40]

$$\ln \Lambda_{\text{ISO}} = \ln \frac{2V^2 r}{G m_\star}. \quad (12)$$

We have checked that changing the definition of the Coulomb log according to different prescriptions in the literature changes the predicted infall time of GCs at the level of a few tens of percent. This would not be crucial for our main results.




FIG. 2. Radius of an infalling GC with mass $m_\star = 3 \times 10^5 M_\odot$, based on the simulations of [16]. In dotted blue (thick dotted red) we plot the simulation result (Fig. 3 in [16]) for the NFW (ISO) halo. In solid lines, we plot our semi-analytic integration. Horizontal dot-dashed lines show the radii r_f where $M_{\text{halo}}(r_f) = m_\star$, in which the semi-analytic treatment should break down.

We can gain some insight on the difference between the DF settling time in the cusp vs. the core profiles. In the central part of a cuspy NFW halo, the density scales as $\rho \propto 1/r$ and the circular velocity scales as $V_{\text{circ}} \propto r^{1/2}$. Let us simplify matters by assuming (as was often done in previous works) that the GC moves on an approximately circular orbit, $V = V_{\text{circ}}$. With this, considering the NFW halo of [16] we find⁶ $C_{\text{Max}} \approx 0.3(r/\text{kpc})^{0.5} \ln \Lambda$. The DF time τ defined in Eq. (9) then scales as $\tau \propto r^2$. This is a rough estimate: if we use the simulation data of [16] for σ and ρ we find a similar but slightly different scaling, $\tau \propto r^{1.85}$, plotted in solid blue in Fig. 4. The important point is the approximately power law decline

⁶ Ref. [17] assumed that $V = V_{\text{circ}}$ and also took V_{circ} equal to the velocity dispersion σ , which would lead to a constant C_{Max} . While this is roughly correct, for the NFW halo of [16] we find mild radial dependence of $V_{\text{circ}}/\sigma \propto r^{0.23}$, as shown in Fig. 3.

of τ towards small r . This is the cause of the fast orbital decay of the GC in the cuspy halo model.

The situation is different in a cored halo. In a core, the density $\rho \approx \rho_0 = \text{const.}$, the circular velocity $V_{\text{circ}} \propto r$, while a Jeans analysis shows that for an isotropic velocity distribution the velocity dispersion is constant $\sigma \approx \sqrt{G\rho_0 r_c}$ [40], where r_c is the core radius (see App. D). This implies $V_{\text{circ}}/(\sqrt{2}\sigma) \sim r/r_c$, as corroborated in Fig. 3 by comparing to the simulation data from [16]. At $r < r_c$ the low-velocity approximation in Eq. (4) gives $C \propto (r/r_c)^3 \ln \Lambda$. The $(r/r_c)^3$ factor can be thought of as a phase-space suppression of DF: it arises from the factor $\int_0^V dv_m v_m^2 f_v(v_m)$ in Eq. (3), because the velocity dispersion inside an isotropic core is greater than the circular velocity (which we assumed to match the instantaneous GC velocity).

Altogether, referring to Eq. (9), an isotropic core predicts an approximately constant τ . Inside the core we can estimate,

$$\begin{aligned} \tau &\approx \frac{3\sqrt{\pi}}{\sqrt{2}} \frac{\sigma^3}{4\pi G^2 m_\star \rho_0 \ln \Lambda} \\ &\approx \frac{\pi}{2\sqrt{3}} \frac{r_c^3 \sqrt{\rho_0}}{\sqrt{G m_\star \ln \Lambda}} \\ &\approx 1.95 \frac{4}{\ln \Lambda} \left(\frac{r_c}{1 \text{ kpc}} \right)^3 \frac{3 \times 10^5 M_\odot}{m_\star} \left(\frac{\rho_0}{3 \times 10^7 \frac{M_\odot}{\text{kpc}^3}} \right)^{\frac{1}{2}} \text{ Gyr}. \end{aligned} \quad (13)$$

In the second line we used Eq. (D4) and in the third line we used values relevant for Fornax GCs. Again, we can compare this estimate to numerical simulations. The dashed red line in Fig. 4 shows τ as calculated by using the velocity dispersion and density read off the cored ISO model of [16]. τ is approximately constant in the core, exceeding the value of τ found for the NFW halo.

The result that DF is suppressed in a cored halo, in comparison to a cusp, is consistent with the finding of Refs. [2, 25, 54], further confirmed in [16, 26, 40, 53].⁷

IV. DEGENERATE DARK MATTER (DDM)

Ref. [38] (see also [39]) made the interesting observation that light fermionic DM would produce a core in dSphs, if the DM particle mass m is light enough to place the inner part of the dSph halo in the degenerate regime. We call this model degenerate DM (DDM). The DDM




FIG. 3. The ratio of circular velocity V_{circ} to the radial velocity dispersion σ_r , reproduced from [16] for NFW (thick red) and isothermal (ISO, blue) halos. The values correspond to simulations that have been run for $t = 4$ Gyr to allow the halos to reach equilibrium. We also plot power law approximations. For the NFW case, $r^{0.23}$ works fairly well for the entire region, whereas for the isothermal case, r^1 may approximate the inner part of the core, which extends to $r \sim 1$ kpc.




FIG. 4. Orbital decay time calculated using ρ and σ from Fig. 1 of [16] and assuming a test object on a circular orbit. NFW denotes a cuspy profile, and ISO denotes an isothermal cored profile. We used here GC mass $m_\star = 3 \times 10^5 M_\odot$.

core scale radius r_c can be estimated via

$$\begin{aligned} r_c &= \frac{A}{G^{\frac{1}{2}} \rho_0^{\frac{1}{6}} (g m^4)^{\frac{1}{3}}} \\ &\approx 681 \left(\frac{\rho_0}{10^7 M_\odot/\text{kpc}^3} \right)^{-\frac{1}{6}} \left(\frac{g m^4}{2 \times (120 \text{ eV})^4} \right)^{-\frac{1}{3}} \text{ pc}, \end{aligned} \quad (14)$$

where ρ_0 is the core central density and where $A = (9\pi/2^7)^{1/6} \approx 0.78$. In App. B we give a derivation of Eq. (14), modelling the dSph halo by a maximum entropy configuration (at fixed total mass and energy, similarly

⁷ The core stalling observed in N-body simulations was initially ascribed in [54] to a failure of the Chandrasekhar formula. However, as we explained here (see also [40]), semi-analytic tracking using the Chandrasekhar formula along the orbit reproduces this result.

to [55])⁸. The maximum entropy halo is isothermal, scaling as $\rho \propto 1/r^2$ at large r . Between the degenerate core and the $1/r^2$ regime there are intermediate features that depend on the temperature.

Inside $r \lesssim r_c$ DDM particles are described by a degenerate distribution function with Fermi velocity v_F related to their mass density via Eq. (5). DF for this system is characterised by Eq. (6), so inside the DDM core, where ρ and v_F are constant, Eq. (9) yields a constant DF time,

$$\begin{aligned} \tau_{\text{DDM}} &\approx \frac{3\pi}{2G^2 g m^4 m_\star \ln \Lambda} \\ &= 4.8 \frac{4}{\ln \Lambda} \frac{10^5 M_\odot}{m_\star} \frac{2 \times (150 \text{ eV})^4}{g m^4} \text{ Gyr} . \end{aligned} \quad (15)$$

Note that if one inserts the DDM halo core radius Eq. (14) into Eq. (13), one obtains the same parametric dependence as in Eq. (15). This is a result of the similarity between Eqs. (4) and (6). The interesting feature of the DDM model is that it produces the core due to Pauli blocking.

Naively, Eq. (15) suggests that DF in a DDM core could be arbitrarily suppressed by decreasing m . This happens because decreasing m at fixed ρ is tied to increasing v_F . However, m cannot be arbitrarily small, because it is constrained by observational limits.

First, the Fermi velocity cannot be arbitrarily high in a gravitationally bound halo [56]. In Sec. IV A we make this analysis more precise by fitting stellar line-of-sight velocity distribution (LOSVD) data to the DDM halo model; we find that while the Fornax LOSVD data indeed constrains $m \gtrsim 100$ eV or so, this constraint by itself would still allow a significant modification of DF compared to the CDM prediction.

A second and much tighter constraint comes from cosmological structure formation as observed through Ly- α forest statistics. We show in Sec. IV B that this constraint directly affects the same combination, $g m^4$, that appears in Eqs. (15) and (14). Imposing the Ly- α constraint excludes DDM from making an appreciable core in Fornax on scales $r \gtrsim 100$ pc, meaning that DDM could not significantly affect the orbits of GCs. While earlier work on DDM argued that a non-thermal production mechanism for DDM could avoid the cosmological constraint, we formulate a rather robust version of the bound which appears difficult to evade.

A. Stellar LOSVD constraints on DDM in Fornax

In this section we summarise the results of a Jeans analysis for the DDM model in Fornax⁹. The DDM pro-

file is described in App. B and the details of the Jeans analysis are given in App. C.

In Fig. 5 we plot LOSVD data of Fornax [10] alongside fits of the density profile presented in App. B. Our fitting procedure is based on a simple χ^2 minimization, where $\chi^2 \equiv \sum_{i=1}^{N_{\text{data}}} (\sigma_{\text{LOS},i} - \sigma_{\text{LOS}}(r_i))^2 / \sigma_i^2$ and σ_i is the reported uncertainty for radial bin i . At a given particle mass m , our fit has three free parameters: the degeneracy parameter μ_0/T , the central core density ρ_0 , and the stellar velocity anisotropy parameter β , taken to be constant in r .




FIG. 5. LOSVD compared with data [10] for DDM with $g = 2$. The best-fit parameters of the profile are $\mu_0/T = 3$ and $\rho_0 = 3.9 \times 10^7 M_\odot/\text{kpc}^3$.

In Fig. 6 we plot the circular velocity compared to the Fermi velocity, which we define using $v_F(r) \equiv \sqrt{2\mu(r)/m}$, for the $m = 135$ eV fit. As can be expected, the circular velocity rises $V \propto r$, whereas v_F remains constant at small radii. This leads to a suppression of DF, as explained in Sec. III.




FIG. 6. The circular velocity V_{circ} and the Fermi velocity v_F for the DDM halo with $m = 135$ eV, $\rho_0 = 3.9 \times 10^7 M_\odot/\text{kpc}^3$, $\mu_0/T = 3$ and $g = 2$. The vertical lines show the estimated orbital radii ($r = r_\perp \times 2/\sqrt{3}$) of the three GCs closest to the dynamical center of Fornax, c.f. Table I.

⁸ Our approach in App. B is similar to that of [38], but differs in that we include also non-zero temperature solutions. Such solutions were noted but not implemented in [38]. We find that these solutions could expand the range of applicability of the DDM model in dSphs.

⁹ For previous analyses, see [38, 39, 57–60].

B. Structure formation constraints on DDM

The free streaming of light DM suppresses the matter power spectrum [61], notably constrained via Ly- α forest statistics [62–64], with details depending on the cosmological DM production mechanism. Refs. [38, 39] considered non-thermal mechanisms for cosmological production of DDM, aiming to bypass the Ly- α bounds. With such mechanisms in mind, values of m in the ballpark of 100 eV were considered in these works. We now revisit the cosmological bound and formulate a conservative limit that is insensitive to the cosmological production mechanism of DM. Our results suggest that mechanisms of the kind proposed in [38, 39] should not be able to produce $m < 1.4$ keV without tension with the nominal Ly- α bound.

The instantaneous DM free streaming wavelength k_{FS} depends on the DM velocity dispersion¹⁰,

$$k_{\text{FS}}(z) \equiv \sqrt{\frac{3}{2}} \frac{\mathcal{H}(z)}{c_s(z)} \simeq \sqrt{\frac{3}{2}} \frac{\mathcal{H}(z)}{\sigma(z)}. \quad (16)$$

Here z is the redshift, $\mathcal{H}(z)$ is the Hubble rate, we have set $\sigma(z) = \sqrt{\langle v^2 \rangle}$ and

$$\langle v^2 \rangle = \frac{\int d^3p \frac{p^2}{m^2 + p^2} f(p)}{\int d^3p f(p)}. \quad (17)$$

Bounds on warm DM (WDM) [64] effectively constrain $k_{\text{FS}}(z)$. Specifically, they apply to $z \lesssim 10^6$, where density perturbations on comoving scales of the order of $\lambda \approx (1+z)/\mathcal{H}(z) \approx 0.5$ Mpc enter the horizon and begin to evolve under their own gravitational potential.

We can convert the WDM limit of [64] into a bound on DDM by the following prescription. At a given energy density, the coldest possible distribution function of DDM is the fully degenerate distribution $f(p) = \theta(p_F - p)$, where the Fermi momentum p_F is related to the energy density via

$$\begin{aligned} \rho &= \frac{g}{(2\pi)^3} \int d^3p \sqrt{m^2 + p^2} f \\ &= \frac{g}{16\pi^2} \left[p_F \sqrt{m^2 + p_F^2} (m^2 + 2p_F^2) - m^2 \sinh^{-1} \left(\frac{p_F}{m} \right) \right] \\ &\approx \begin{cases} \frac{gm p_F^3}{6\pi^2} & p_F \ll m \\ \frac{g p_F^4}{8\pi^2} & p_F \gg m \end{cases}. \end{aligned} \quad (18)$$

The p_F parameter redshifts as $p_F \propto (1+z)$. The velocity dispersion for this distribution is

$$\begin{aligned} \langle v^2 \rangle &= 1 - 3 \left(\frac{m}{p_F} \right)^2 + 3 \left(\frac{m}{p_F} \right)^3 \arctan \left(\frac{p_F}{m} \right) \\ &\approx \begin{cases} \frac{3p_F^2}{5m^2} & p_F \ll m \\ 1 & p_F \gg m \end{cases}. \end{aligned} \quad (19)$$

Using Eq. (19) we can calculate $k_{\text{FS}}(z)$, compare this to the $k_{\text{FS}}(z)$ of WDM, and cast the bounds of [64] into the most conservative, maximally-cold DDM model by matching the $k_{\text{FS}}(z)$ curves of the two models. To recall, WDM was defined [64] by the distribution function $f_{\text{WDM}} = (\exp(p/T) + 1)^{-1}$, where $T \propto (1+z)$. Comparing the DM mass density for DDM and WDM in the nonrelativistic regime, we have

$$\rho = \begin{cases} \frac{gm p_F^3}{6\pi^2} & \text{DDM} \\ \frac{3\zeta(3)gmT^3}{4\pi^2} & \text{WDM} \end{cases}. \quad (20)$$

Matching the density in the two models implies:

$$\frac{(T/m)_{\text{WDM}}}{(p_F/m)_{\text{DDM}}} = \left[\frac{2}{9\zeta(3)} \frac{(gm^4)_{\text{DDM}}}{(gm^4)_{\text{WDM}}} \right]^{\frac{1}{3}}. \quad (21)$$

On the other hand, still in the nonrelativistic regime we can compare the velocity dispersion in the two models,

$$\begin{aligned} \frac{\langle v^2 \rangle_{\text{DDM}}}{\langle v^2 \rangle_{\text{WDM}}} &\approx \frac{3\zeta(3)}{75\zeta(5)} \left[\frac{(p_F/m)_{\text{DDM}}}{(T/m)_{\text{WDM}}} \right]^2 \\ &= \frac{3\zeta(3)}{75\zeta(5)} \left(\frac{9\zeta(3)}{2} \right)^{\frac{2}{3}} \left[\frac{(gm^4)_{\text{WDM}}}{(gm^4)_{\text{DDM}}} \right]^{\frac{2}{3}}, \end{aligned} \quad (22)$$

where in the second line we used Eq. (21). We illustrate the comparison in Fig. 7. For model parameters of interest to this discussion, the ratio of velocity dispersions is redshift-independent at $z \lesssim 10^6$. We can therefore conclude that the WDM constraint of [64], $m_{\text{WDM}} > 2.96$ keV at the 95%CL for $g = 2$, implies the constraint:

$$gm_{\text{DDM}}^4 > 2 \times (1.4 \text{ keV})^4. \quad (23)$$




FIG. 7. A comparison of the velocity dispersions (in natural units) of WDM and DDM for various particle masses. Including CMB data, Ref. [64] puts a bound of $m > 2.96$ keV on WDM (95% CL), for which we plot the velocity dispersion in green solid line.

We expect that model-building around the bound of Eq. (23) would be quite difficult. No production mecha-

¹⁰ This formula is correct up to an order unity factor relating the speed of sound c_s and the velocity dispersion σ [65]. This factor is unimportant for our results.

nism should be able to create a colder distribution function for DDM. In particular, the skewed momentum distribution scenarios of [39] and the scalar decay models of [38, 66] should all satisfy this bound. Using Eq. (14), we find that Eq. (23) constrains the DDM core in Fornax to $r_c \lesssim 20$ pc, irrelevant for the orbits of GCs.

The Ly- α analyses may be affected by systematic uncertainties related, among other things, to the thermal history of the inter-cluster medium (IGM) and other baryonic effects, and that are still not fully implemented from first-principles in the numerical simulations used to compare theoretical predictions to the data in producing the bound [64]. Keeping this caveat in mind, it seems sensible to take Fig. (7) with a grain of salt. If we allow k_{FS} of the (coldest possible) DDM model to exceed the nominal bound of [64] by, say, a factor of ~ 4.5 , we could relax Eq. (7) to $gm^4 > 2 \times (500 \text{ eV})^4$. With such (rather ad-hoc) relaxed bound we could allow a DDM core $r_c \lesssim 80$ pc; still irrelevant for GC orbits in Fornax. Going down to $m = 150$ eV (still at $g = 2$), which would allow a DDM core radius of $r_c \approx 385$ pc in the ballpark of some GC orbits, would amount to k_{FS} larger by a factor of 20 than the nominal WDM bound.

C. Orbital decay time in DDM

As we have seen, Ly- α analyses exclude DDM from producing a core extending all the way to the observed orbital positions of GCs in Fornax. This constraint then implies that DDM would not change the standard CDM predictions for the DF settling time of the GCs. Nevertheless, given that the Ly- α bound is subject to some debate, it is interesting to see what DDM could do to DF subject only to the LOSVD constraints of Sec. IV A.

In order to estimate the instantaneous DF time scale τ , we use Eq. (9), with a modified Eq. (6). In order to interpolate between a quasi-degenerate core and a classical gas in the outskirts of the halo, we adopt

$$C_{\text{DDM}} = \frac{1}{\frac{1}{0.5} + \frac{v_F^3}{V^3}} \ln \Lambda, \quad (24)$$

such that in the regime $V \gg v_F$, we retrieve CDM-like behavior, c.f. Sec. III. For $\ln \Lambda$, we adopt the choice for $\ln \Lambda_{\text{ISO}}$, c.f. Eq. (12).

We use the GC masses and projected radii collected in Table I, combined with the density profile derived in the LOSVD fits. We correct for the projection effect by relating the assumed true orbital radius to the observed projected radius of the GC via $r_{\text{true}}/r_{\perp} = 2/\sqrt{3}$. We also assume that the GCs are on circular orbits, setting $V_{\text{true}}/V_{\text{circ}}(r_{\text{true}}) = 1$. (This de-projection procedure is, of course, simplistic: we will shortly report a more comprehensive treatment.) The results are summarised in Table I. For $m = 135$ eV, we find naïve orbital decay time-scales of 1.48 Gyr and 10.7 Gyr for GC3 and GC4, respectively. For comparison, using the approach of Ref. [17]

for cuspy CDM we find 2.63 Gyr and 0.99 Gyr. Therefore, while the naïve DF time of GC4 in DDM is much longer than in cuspy CDM, for GC3 the naïve time in cuspy CDM actually exceeds that of DDM. However, as discussed in Sec. III, the instantaneous τ can be misleading when comparing different halo morphologies: a realistic estimate of the GC settling time requires orbit integration. We turn to this analysis next, finding that the real orbit settling times of both GC4 and GC3 are in fact longer in DDM compared with cuspy CDM.

To obtain a more comprehensive estimate of the DF settling time and the impact of different line-of-sight projection angles, we use the orbit integration explained in App. E with initial conditions that we vary as follows. For each GC, we scan the range $r_{\text{true}} \in [1, 2]r_{\perp}$ (the logic behind this range is explained in App. G). For each r_{true} we scan over $V_{\text{true}} \in [0.5, 1.5]V_{\text{circ}}(r_{\text{true}})$. For each V_{true} we test positive and negative $\cos \theta$. Finally, we test the two cases, $\Delta v_y = \sqrt{v_{\text{true}}^2 - \Delta v_r^2}, \Delta v_z = 0$ and $\Delta v_y = 0, \Delta v_z = \sqrt{v_{\text{true}}^2 - \Delta v_r^2}$.

For each starting point in phase-space, we integrate the equations of motion, stopping the integration when $(r_{\text{apoenter}} + r_{\text{pericenter}})/2 \lesssim 0.3r_{\text{initial}} \equiv f_r r_{\text{initial}}$, or after 10 Gyr (the earlier of the two). We then report the integration time as τ_{inspiral} .

In Fig. 8 we plot the result of this procedure for GC3, comparing the DDM halo for $m = 135$ eV (top panel) and the cuspy CDM halo from [16] (bottom panel). For the representative phase-space point $r_{\text{true}}/r_{\perp} = 2/\sqrt{3}$, $V_{\text{true}}/V_{\text{circ}}(r_{\text{true}}) = 1$, highlighted in Fig. 8 by a red dot, we find that the inspiral time in the DDM halo is in fact longer (~ 4 Gyr) than in the cuspy CDM one (~ 1.5 Gyr). This result is in reversed order to the naïve estimate in Table I, demonstrating that the naïve DF time can be misleading in judging the future of GC orbits.

We note that the inspiral time in the NFW case is not very sensitive to the stopping radius fraction f_r (set as 0.3), whereas the DDM case is, and so are other cored halo models. As explained in III, a cuspy profile predicts approximately $\tau(r) \propto r^2$, therefore the inspiral time is mostly sensitive to the initial radius. In a cored model, $\tau(r)$ is a weak function of r , potentially even falling and then rising again (see bottom right panel of Fig. 11). Therefore, the definition of the inspiral time in the cored model becomes sensitive to the radius at which our orbit integration is stopped.

V. SELF-INTERACTING DARK MATTER (SIDM)

Self-interacting DM (SIDM) is a simple modification of CDM, that could arise in many models [41, 67–74]. The self interactions can be expressed in terms of the cross section per unit mass, $\sigma/m \sim 0.1 \div 10 \text{ cm}^2/\text{gr}$, which could be velocity-dependent [72]. The scattering mean




FIG. 8. Contours of the inspiral time of GC3, defined in App. G, for DDM (**top**) and cuspy CDM model (**bottom**). The “naïve” estimates written on top are those given in Table I based on an evaluation of the instantaneous DF time at $r_{\text{true}}/r_{\perp} = 2/\sqrt{3}$ and $V_{\text{true}}/V_{\text{circ}}(r_{\text{true}}) = 1$. The different line types correspond to different discrete choices in our scan of the initial conditions in phase-space, explained in more detail in App. G.

free path is

$$l = \frac{m}{\rho\sigma} = 48 \frac{10^8 M_{\odot}/\text{kpc}^3}{\rho} \frac{1 \text{ cm}^2/\text{gr}}{\sigma/m} \text{ kpc}, \quad (25)$$

and the time between scatterings l/v is

$$t_{\text{scat}} = 2.35 \frac{20 \text{ km/s}}{v} \frac{10^8 M_{\odot}/\text{kpc}^3}{\rho} \frac{1 \text{ cm}^2/\text{gr}}{\sigma/m} \text{ Gyr} \quad (26)$$

When l is larger than the distance across the halo, we expect that the microphysics of DF in the SIDM model will be similar to that of CDM. On the other hand, the morphology of an SIDM halo could be different to that in CDM as long as t_{scat} is smaller than the age of the system.

Given a large enough cross section, SIDM produces cored halos which affect the orbital settling time of GCs as discussed in Sec. III.

We follow Ref. [41] in modelling the SIDM halo profile. Inside some radius r_1 , we assume an hydrostatic profile with central density ρ_c and pressure $P = \sigma_0^2 \rho$. For a self-gravitating spherical halo, the density profile obeys

$$\frac{1}{r^2} \partial_r (r^2 \partial_r \ln \rho_{\text{iso}}) = -\frac{1}{r_{\text{ic}}^2} \frac{\rho_{\text{iso}}}{\rho_c}, \quad (27)$$

where $r_{\text{ic}} \equiv \sigma_0 / \sqrt{4\pi G \rho_c}$ (similar to the King radius [43]). Beyond r_1 we match the density to the NFW profile, $\rho_{\text{NFW}} = \rho_s (r/r_s)^{-1} (1 + r/r_s)^{-2}$, fixing ρ_s and r_s by imposing continuity of $\rho_{\text{iso}}(r_1) = \rho_{\text{NFW}}(r_1)$ and of the enclosed mass $M_{\text{iso}}(r_1) = M_{\text{NFW}}(r_1)$. This procedure is consistent with an initially NFW-like cusp profile that was deformed into a cored isothermal profile due to the SIDM scatterings. Altogether, the halo model has three free parameters, ρ_c , r_1 and σ_0 , that we can constrain with LOSVD data.

In Fig. 9 we plot a LOSVD fit, following the same χ^2 procedure as in Sec. IV. The model is taken to illustrate a large core solution that would include the orbit of GC3.




FIG. 9. **Top:** LOSVD data of Fornax dSph modeled by different SIDM profiles. The central density is $\rho_c = 2.6 \times 10^7 M_{\odot}/\text{kpc}^3$, the velocity dispersion is $\sigma_0 = 17 \text{ km/sec}$ and $r_1/r_{\text{ic}} = 6$.

In Fig. 10 we show the inspiral time of GC3, using the same procedure as in Fig. 8. We use Eq. (4) with σ_0 from the LOSVD fit and adopt $\ln \Lambda = \ln \Lambda_{\text{ISO}}$ as in Eq. (12). We find that the large core SIDM model significantly increases the inspiral time of GC3.

Ref. [41] pointed out that in the SIDM core region, DM particles have undergone about a single collision during the age of the system, i.e.

$$\frac{\langle \sigma v \rangle}{m} \rho_c t_{\text{age}} = \frac{t_{\text{age}}}{t_{\text{scat}}} \approx 1. \quad (28)$$

With this assumption,¹¹ we can estimate the cross section

¹¹ It has been pointed out [70, 75] that this assumption may be simplistic.

implied by the LOSVD fit

$$\frac{\langle \sigma v \rangle}{m} \approx 18 \frac{10 \text{ Gyr}}{t_{\text{age}}} \frac{2.6 \times 10^7 M_{\odot} / \text{kpc}^3}{\rho_c} \frac{\text{cm}^2 \text{ km}}{\text{g s}}. \quad (29)$$

This result is compatible with the baseline model of [41], which predicted $\sim 25 \text{ cm}^2 \text{ g}^{-1} \text{ km s}^{-1}$. It would significantly increase the DF settling time of the innermost Fornax GCs.




FIG. 10. DF time for SIDM halo, analogously to Fig. 8.

VI. DISCUSSION

In Fig. 11 we summarise the key features of different models of DM discussed in this work, including both cuspy and core halo models. In the spirit of [2], we also add the hybrid **coreNFW** model of [76] with density¹²

$$\rho_{\text{coreNFW}} = \tilde{f}^n \rho_{\text{NFW}} + \frac{n \tilde{f}^{n-1} (1 - \tilde{f}^2)}{4\pi r^2 r_c} M_{\text{NFW}}, \quad (30)$$

where $\tilde{f} = \tanh(r/r_c)$ and $M_{\text{NFW}} = \int_0^r d^3 r' \rho_{\text{NFW}}$.

The LOSVD data (top left of Fig. 11) is described reasonably well in all models, with the fit of the ISO model of [16] being slightly worse.

The instantaneous DF time for a GC with $m_* = 3 \times 10^5 M_{\odot}$ is shown in the bottom right of Fig. 11. It illustrates the fact that the main impact of the microphysics of DM (as in DDM and SIDM) on DF comes from their prediction of a cored halo morphology, and not from the exotic microphysics per-se. The formation

of a core due to baryonic feedback in CDM [25, 54] would therefore have similar consequences.

The density profiles (top right of Fig. 11) demonstrate the cusp for NFW, large-cores and the intermediate **coreNFW**. We also plot an estimate of the stellar density. This may become important for large-core models, whose density is only larger by a factor of two or so than the stellar density at small r . In these cases, accounting for the stellar-induced potential could slightly change our results numerically, but not qualitatively: as far as the DF microphysics is concerned, background stars would contribute to the DF of a GC just like CDM particles, and since the total mass density is constrained by the LOSVD fit, the separation into DM and stellar components is not essential for the DF computation.

The circular velocity profiles of different models (bottom left of Fig. 11) can place constraints on the phase-space parameters of GCs. We discuss this point in the next subsection, where we consider in turn each of the six GCs.

A. Comments regarding GC data used in this work

Let us re-evaluate the role of each GC in the timing puzzle, in light of the observational data used in this work (see Tab. I), noting that the projected radii are different than those of earlier analyses [2, 17, 19, 28] and that the LOS velocities, that were often ignored in past analyses, are in fact known fairly well from observations.

1. GC1 and GC5 – these GCs do not seem to pose a timing problem as they are located at fairly large projected radii, $r_{\perp} \approx 1.7 \text{ kpc}$, not far below the (somewhat model-dependent) tidal radius of Fornax at $1.8 \div 2.8 \text{ kpc}$ [2, 24, 77]. This said, it is interesting to note that while the circular velocity at the GC radii is $20 \div 30 \text{ km/s}$, the measured GC LOS velocities are smaller than V_{circ} by a factor of 5 or so. This could hint that GC1 and GC5 are close to the apocenter of a fairly eccentric orbit. If this is indeed the case, then the naïve instantaneous DF time overestimates the true orbital settling time because the GCs typically experience stronger DF when they venture into smaller radii, as expected if the orbit is eccentric.
2. GC2 – does not seem to present a timing problem.
3. GC3 – with the orbit integration of App. G, we find that both a simple cusp profile and the intermediate **coreNFW** profile predict an inspiral time $\approx 1.5 \text{ Gyr}$, whereas models with a large core predict $4 \div 5 \text{ Gyr}$. For a cuspy profile, our “naïve” instantaneous DF time of 2.6 Gyr comes in some disagreement with the $\tau \approx 0.6 \text{ Gyr}$ quoted in [17]. The main reasons for the difference are the new estimate of r_{\perp} and the updated LOSVD data [10] compared to the older data [1].

¹² For the **coreNFW** model we adopt the best-fit of [10], with $\rho_0 \approx 10^{7.1} M_{\odot} / \text{kpc}^3$, $r_s \approx 2.1 \text{ kpc}$, $n \approx 0.8$ and $r_c \approx 0.52 \text{ kpc}$. We derive the velocity-dispersion of the halo by solving the Jeans equation, Eq. (D1), assuming isotropy.




FIG. 11. Comparison of models. **Top left:** LOSVD data and fits. The M19 NFW and M19 ISO models refer to the halos of [16], for which we only fit the velocity anisotropy. The DDM and SIDM models are based on Sec. IV and Sec. V. The $\chi^2/\text{d.o.f}$ is ≈ 1.9 for NFW, DDM and SIDM, and ≈ 2.5 for ISO. The velocity anisotropy is taken to be constant in each fit. We find $\beta_{\text{NFW}} = -0.4$, $\beta_{\text{ISO}} = 0.2$, $\beta_{\text{DDM}} = -0.1$, $\beta_{\text{SIDM}} = 0.1$ and $\beta_{\text{coreNFW}} = -0.1$. **Top right:** Density profiles. In addition to DM, we also plot an estimate of the stellar density, assuming a Plummer profile with scale $r_p = 851$ pc [21] and mass $4 \times 10^7 M_\odot$ [18]. **Bottom left:** Circular velocity. (Note how cored models require some tuning to explain the large radial velocity of GC4, $|\Delta v_r| = 8.26 \pm 0.64$ km/s, at its small projected radius $r_\perp \approx 0.154$ pc). **Bottom right:** Instantaneous DF time, evaluated for $m_\star = 3 \times 10^5 M_\odot$.

- GC4 – the cuspy CDM orbital decay time is short ≈ 1 Gyr, but not as short as previously estimated [17]. Beyond the reasons listed for GC3, we find that the approximation $C \approx 0.5 \ln \Lambda$, adopted in [17], is not accurate for small radii, c.f. Sec. III.

A large core would stabilise the orbit of this GC to the 10 Gyr time scale, and even the intermediate coreNFW profile predicts inspiral time of ≈ 5 Gyr.

GC4 is younger and more metal-rich compared to the other GCs [18, 19]. It has been debated in the literature whether GC4 is the nuclear star cluster of Fornax dSph [18, 78–80]. The large observed LOS velocity of this GC, $\gtrsim 8$ km/s, appears potentially at odds with this interpretation [81].

The large LOS velocity is also somewhat difficult to accommodate within a large core halo model. In all of our cored halo models (see bottom left of Fig. 11), GC4 needs to be on a circular orbit with $r_{\text{true}}/r_\perp \gtrsim 2$ or close to the pericenter of an eccentric orbit with $r_{\text{true}}/r_\perp \gtrsim 1.6$ (see Fig. ??). These possibilities are somewhat tuned, either w.r.t. to the projection angle or w.r.t. to the orbital phase. In comparison, an NFW profile can comfortably

accommodate the radial velocity of GC4. For the coreNFW model, it is marginally possible to have GC4 on a circular orbit without tuning in radius.

- GC6 – the newly rediscovered GC [14] probably has a smaller mass ($\approx 0.29 \times 10^5 M_\odot$ [42]) than the other five GCs. It does not seem to reinforce the timing puzzle.

Ref. [14] notes that GC6 has an elongated shape and may be undergoing tidal disruption. This may comprise some evidence in favour of a cuspy halo.

To summarise, the usual suspects for a GC timing puzzle, GC3 and GC4, are found here to have somewhat longer settling times than previously thought [17]. Within a cuspy profile, we find that GC3 and GC4 have an inspiral time of order 1 Gyr. Within the framework of SIDM, substantially longer DF times are possible. For DDM, Ly- α constraints suggest the same DF time as in cuspy CDM, while if one ignores the Ly- α limit, long DF time is again possible at $m \lesssim 150$ eV. The intermediate coreNFW profile would also prolong the inspiral times of GC3 and GC4 to a few Gyr.

B. Statistical discussion

We finally have the tools to provide an analytical perspective on the timing puzzle. Consider the cuspy NFW profile, illustrated in Fig. 11. Using the results developed in App. F, we can map a distribution of GC initial radii into the cumulative distribution function (CDF) of GC projected radii today. An example of such calculation is shown in Fig. 12. As shown in App. F, before projection




FIG. 12. Example of a calculated CDF of projected radii using $\tau(r)$ from the NFW fit in Fig. 11, compared to the observed distribution of Fornax GCs. Solid curve shows the CDF after projection effects are taken into account. Dashed curve shows the result before projection.

effects are taken into account the CDF of GC radii at $r \ll r_{\text{cr}} \approx 1$ kpc takes the form

$$F_{\Delta t}(r) \approx A \frac{\tau(r)}{\Delta t}, \quad (31)$$

independent on initial conditions, with A an $\mathcal{O}(1)$ coefficient. The dashed line in Fig. 12 shows the unprojected GC CDF and is consistent with this expectation. Projection has a significant effect where GCs at projected radius r_{\perp} are in fact located at $r > r_{\perp}$. The solid line shows the projected radius CDF: for $r_{\perp} \lesssim 0.5$ kpc, projection roughly doubles the GC count inside a given r_{\perp} .

The GC timing puzzle is reflected in Fig. 12 in two ways, one quite obvious (but apparently mild) and the other more subtle. First, while Fig. 12 shows the CDF resulting from just one example of GC initial conditions, we could not find a reasonable distribution of initial conditions which fits the innermost GC4 and GC6, while not overshooting GC3 and GC2 further out; nor could we find initial conditions which fit GC3, and do not undershoot GC4 and GC6. The reason for this (mild) inconsistency is the growth in r of $F_{\Delta t}$ in Eq. (31). This inconsistency, however, appears quite mild: in the example we show in Fig. 12, GC4 and GC6 make 2 GCs within a projected radius where the CDF predicts 1. The probability to see

2 GCs or more where only 1 GC is expected is about 25%. This does not seem like severe fine-tuning.

The second and more subtle issue is again related to Eq. (31). As we explain in App. F, Eq. (31) arises as all of the GCs which started their life inside true radius $r < r_{\text{cr}}$ at time $-\Delta t$ in the past, have already settled down to $r \approx 0$ and were tidally disrupted. These GCs are lost to the GC count today; as a result, the form of the initial distribution of GCs at $r < r_{\text{cr}}$ is unconstrained by the GC distribution today. The issue regarding this disappearance of GCs is that unless the initial distribution of GCs was strongly suppressed at small r , then a few Gyr ago Fornax must have had many more than 6 GCs. For example, the initial distribution we used to make Fig. 12 was $f_0(r) \propto r^2 e^{-r^2}$ (with r measured in kpc). This function is suppressed at small r , but nevertheless it predicts that Fornax started out with around 13 GCs, of which only about half survived until today. If we drop the r^2 small- r suppression factor, taking $f_0(r) \propto e^{-r^2}$ for example, then we have a very similar-looking $F_{\Delta t}$ but we find that Fornax must have started off with more than 50 GCs.

It seems to us difficult to accept that Fornax initially hosted many more than 6 GCs, but lost most of them by now. If GC formation in dSphs is so prolific, why don't we find GCs around many other dSphs? This said, without a first-principle understanding of GC formation in the early Universe we do not know how (if at all) this potential difficulty may turn into a real constraint. A possible resolution could come if the initial distribution of GCs is suppressed at small radii.

More insight could come from numerical simulations, even though both DF and GC formation involve sub-grid physics in most existing simulations. Recently, Ref. [42] explored the survival of GCs in cuspy Fornax-like halos using hydrodynamical cosmological simulations [82, 83], to which sub-grid formation of GCs was added and their orbital evolution under DF was tracked in post-processing. Ref. [42] calculated the projected radius CDF of GCs at $z = 0$, finding that their simulated CDF is consistent with the observed positions of the Fornax GCs. According to [42], only around 33% of GCs are tidally disrupted in the simulations; a somewhat smaller number than the 50% in our example in Fig. 12. Fornax is found to be special, in that only about 3% of Fornax-like galaxies in the simulation ended up with 5 or more surviving GCs today.¹³

¹³ Note that [42] considered GC6 as a candidate for a tidally disrupted GC, assuming that it's projected distance from the center of Fornax is just 30 pc. In comparison, using the Fornax center of [21], we find a projected distance of 254 pc for GC6.

VII. SUMMARY

We revisit the calculation of globular cluster (GC) orbits under dynamical friction (DF), considering different microscopic models of dark matter (DM) and different halo morphologies that they predict. We focus on the Fornax dwarf spheroidal (dSph) galaxy, which hosts 6 GCs and which has been noted in previous literature to pose a GC timing problem, that is, the future orbits of some of its GCs are much shorter than their current age.

We presented analytical and semi-analytical computations of DF and of GC orbits under DF. For a cuspy DM halo, we showed that the current cumulative distribution function (CDF) of GC radii takes an approximately power law form that can be deduced from stellar kinematics and age measurements. Including projection effects, we demonstrated that the GC timing problem does not appear very severe: the existence of the innermost GCs could be accounted for at the cost of moderate small-numbers fluctuation with a probability of about 25%. A cuspy halo, in conjunction with GC orbits, does place interesting constraints on the initial distribution of GCs in dSphs. Testing these constraints would likely require high resolution numerical simulations including baryonic effects, where GC formation can be modelled from first principles.

The recent numerical simulations of Ref. [42] made an interesting step in this direction, although they did not contain a direct simulation of GC formation or of DF. According to Ref. [42] a reasonable distribution of initial conditions for the GCs can naturally lead to the observed present configuration.

The fact remains that the combination of GC age and orbit measurements could probe the Fornax DM halo and microphysics. At the level of the microphysics, in Sec. II we calculated DF for three models of DM: fermionic degenerate DM (DDM), where Pauli blocking affects the DF derivation; bosonic ultralight DM (ULDM), where an astronomical de-Broglie scale comes into play; and self-interacting DM (SIDM), where – in the limit we were mostly interested in – the microphysics of DF should mostly follow that of CDM, but the halo morphology is different.

For ULDM, DF and specifically the Fornax GC problem were studied in a number of works. Constraints from galaxy dynamics and, independently, cosmological Ly- α analyses exclude a soliton core reaching out to the orbits of Fornax GCs, and lead to a similar behaviour as in CDM.

For DDM, we gave a new derivation of DF. We then formulated a robust (in terms of DM model building) version of the Ly- α bound, showing that it excludes an appreciable core, leading again to CDM-like behaviour at the scale of GC orbits. At the same time, uncertainties and degeneracies in the stellar kinematics in Fornax could still allow a considerable DDM core. If the Ly- α bound is discounted, for some reason, then DDM could lead to significant suppression of DF and prolongation of the

settling time of GC3 and GC4.

For SIDM, stellar kinematics allows a considerable core. If the SIDM cross section is as large as that considered in [41], then the DF settling time for GC3 and GC4 can be significantly longer than in the cuspy halo CDM model.

ACKNOWLEDGMENTS

We thank Justin Read for clarifications on the kinematic data. We thank Joshua Eby and Daniel Kaplan for useful discussions. NB is grateful for the support of the Clore scholarship of the Clore Israel Foundation. KB is incumbent of the Dewey David Stone and Harry Levine career development chair at the Weizmann Institute of Science, and was supported by grant 1784/20 from the Israel Science Foundation.

Appendix A: Dynamical friction in exotic media: derivation from the Boltzman equation

In this section we provide an economical derivation of gravitational DF acting on a nonrelativistic probe object moving in a medium, with different medium microphysics including a classical gas as well as quantum Fermi and Bose gases. We neglect interactions apart from minimal gravity. We start with a quick recap of the derivation of the Fokker-Planck equation, governing the phase space distribution functions of the probe and medium particles.

We consider the following elastic scattering process of two particle species,

$$1(p) + 2(k) \rightarrow 1(p') + 2(k').$$

The phase space distribution function for the particle species 1 evolves according to the Boltzmann equation,

$$\frac{df_1}{dt} = C[f_1]. \quad (\text{A1})$$

The collision integral $C[f_1]$ contains information about the elastic scattering processes, and is written as

$$\begin{aligned} C[f_1] = & \frac{(2\pi)^4}{2E_p} \int d\Pi_k d\Pi_{p'} d\Pi_{k'} \delta^{(4)}(p + k - p' - k') |\overline{\mathcal{M}}|^2 \\ & \times \left[f_1(p') f_2(k') (1 \pm f_1(p)) (1 \pm f_2(k)) \right. \\ & \left. - f_1(p) f_2(k) (1 \pm f_1(p')) (1 \pm f_2(k')) \right], \quad (\text{A2}) \end{aligned}$$

where $|\overline{\mathcal{M}}|^2$ is a squared matrix element averaged over initial and final spins, and $d\Pi_k = \frac{g}{2E_k} \frac{d^3k}{(2\pi)^3}$ is the Lorentz invariant phase element with the number of internal degrees of freedom g . The sign in $1 \pm f_i$ refers to bosons (+) or fermions (−), respectively. It is convenient to write the

above Boltzmann equation in the following form,

$$\frac{df_1}{dt} = \int \frac{d^3 p'}{(2\pi)^3} \left[S(\mathbf{p}', \mathbf{p}) f_1(p') (1 \pm f_1(p)) - S(\mathbf{p}, \mathbf{p}') f_1(p) (1 \pm f_1(p')) \right], \quad (\text{A3})$$

where the function S encodes the response of the medium, and is defined as

$$S(\mathbf{p}, \mathbf{p}') \equiv \frac{(2\pi)^4}{2E_p 2E_{p'}} \int d\Pi_k d\Pi_{k'} \delta^{(4)}(p + k - p' - k') \times |\overline{\mathcal{M}}|^2 f_2(k) (1 \pm f_2(k')). \quad (\text{A4})$$

The function $S(\mathbf{p}, \mathbf{p}')$ can be interpreted as a differential rate at which a particle of momentum \mathbf{p} is converted into a particle with momentum \mathbf{p}' .

The Boltzmann equation can be greatly simplified if the momentum exchange

$$q = p' - p \quad (\text{A5})$$

is smaller than the typical momentum given by the distribution function f_1 . In such cases, the Boltzmann equation is reduced to the nonlinear Fokker-Planck equation,

$$\frac{df_1}{dt} = - \frac{\partial}{\partial p^i} [f_1 (1 \pm f_1) D_i] + \frac{1}{2} \frac{\partial}{\partial p^i} \left[\frac{\partial}{\partial p^j} (D_{ij} f_1) \pm f_1^2 \frac{\partial}{\partial p^j} D_{ij} \right], \quad (\text{A6})$$

where the diffusion coefficients are defined as

$$D_i(\mathbf{p}) = \int \frac{d^3 q}{(2\pi)^3} q^i S(\mathbf{p}, \mathbf{p} + \mathbf{q}), \quad (\text{A7})$$

$$D_{ij}(\mathbf{p}) = \int \frac{d^3 q}{(2\pi)^3} q^i q^j S(\mathbf{p}, \mathbf{p} + \mathbf{q}). \quad (\text{A8})$$

are the diffusion coefficients.

The gravitational scattering of a probe particle of mass M and a particle in the medium with mass m is described by the spin-averaged matrix element

$$|\overline{\mathcal{M}}|^2 = \frac{1}{2s+1} \frac{(16\pi G)^2 m^4 M^4}{[(q^0)^2 - \mathbf{q}^2]^2}, \quad (\text{A9})$$

entering Eq. (A4). In the nonrelativistic limit, we can neglect q^0 and maintain only \mathbf{q} in Eq. (A9).

The problem of calculating the diffusion coefficients for different types of medium amounts to evaluating Eqs. (A7) and (A8), where in the response function Eq. (A4) we can select the appropriate sign in $1 \pm f_2$ corresponding to the medium particle's spin-statistics (or setting $1 \pm f_2 \rightarrow 1$ if we wish to compute the classical gas limit).

For the calculation of DF we are particularly interested in $D_{||}$, the first diffusion coefficient corresponding to motion parallel to the probe object's instantaneous velocity. $D_{||}$ is simply given by Eq. (A7) when we select q^i to align with the direction of \mathbf{p} .

1. A classical gas medium

We first re-derive the relaxation of massive classical objects, such as supermassive black holes or GCs, in a background medium consisting of other classical objects such as stars or CDM particles. In the nonrelativistic limit, the function $S(\mathbf{p}, \mathbf{p}')$ is simplified as

$$S(\mathbf{p}, \mathbf{p}') \simeq g_\chi \frac{(4\pi G m M)^2}{q^4} \int \frac{d^3 k}{(2\pi)^3} \frac{d^3 k'}{(2\pi)^3} \times (2\pi)^4 \delta^{(4)}(p + k - p' - k') f_2(k) \quad (\text{A10})$$

where g_χ is the number of internal degrees of freedom of dark matter. Here, M and m are the mass of the particle species 1 and 2, respectively. In the small momentum exchange limit, the δ -function for the energy conservation can be expanded as

$$\delta(E_p + E_k - E_{p'} - E_{k'}) \simeq \frac{1}{q} \left(1 + \frac{M}{2\mu_r} \mathbf{q} \cdot \frac{\partial}{\partial \mathbf{p}} \right) \delta \left[\hat{\mathbf{q}} \cdot \left(\frac{\mathbf{k}}{m} - \frac{\mathbf{p}}{M} \right) \right], \quad (\text{A11})$$

where $\mu_r = mM/(m+M)$ is the reduced mass. Using these approximate expressions in nonrelativistic and small momentum exchange limit, we obtain the diffusion coefficients as

$$D_i(\mathbf{p}) = \int \frac{d^3 q}{(2\pi)^3} q^i S(\mathbf{p}, \mathbf{p} + \mathbf{q}) = 4\pi G^2 m^2 M^2 \left(1 + \frac{M}{m} \right) \ln \Lambda \frac{\partial}{\partial p^i} h(\mathbf{p}; f_2) \quad (\text{A12})$$

and

$$D_{ij}(\mathbf{p}) = \int \frac{d^3 q}{(2\pi)^3} q^i q^j S(\mathbf{p}, \mathbf{p} + \mathbf{q}) = 4\pi G^2 m^2 M^4 \ln \Lambda \frac{\partial^2}{\partial p^i \partial p^j} g(\mathbf{p}; f_2) \quad (\text{A13})$$

where $\ln \Lambda = \int_{q_{\min}}^{q_{\max}} dq/q$ is the Coulomb logarithm, and we have used identities Eqs. (26)–(27) of [31] to perform the angular integration at the second step in each equation. The Rosenbluth potentials $h(\mathbf{p})$ and $g(\mathbf{p})$ are defined as [84]

$$h(\mathbf{p}; f) = g_\chi \int \frac{d^3 k}{(2\pi)^3} \frac{f(k)}{\left| \frac{\mathbf{k}}{m} - \frac{\mathbf{p}}{M} \right|}, \quad (\text{A14})$$

$$g(\mathbf{p}; f) = g_\chi \int \frac{d^3 k}{(2\pi)^3} \left| \frac{\mathbf{k}}{m} - \frac{\mathbf{p}}{M} \right| f(k) \quad (\text{A15})$$

This reproduces the well known diffusion coefficients in a classical system, see Eq. (7.83) in Binney & Tremaine [85]. For the Maxwell-Boltzmann distribution $f_2(k) = (2\pi)^{3/2} n_2 / [g_\chi (m\sigma)^3] e^{-v_k^2/2\sigma^2}$, it is straightfor-

ward to find

$$\begin{aligned}\frac{\partial h}{\partial p^i} &= -\frac{v^i}{v} \frac{n_2}{M\sigma^2} \frac{1}{2X^2} \left[\text{erf}(X) - \frac{2X}{\sqrt{\pi}} e^{-X^2} \right] \\ &\equiv -\frac{v^i}{v} \frac{n_2}{M\sigma^2} G(X),\end{aligned}\quad (\text{A16})$$

$$\begin{aligned}\frac{\partial g}{\partial p^i \partial p^j} &= \frac{\sqrt{2}\sigma^2}{M^2} \left[\frac{3}{2} \frac{X^i X^j}{X^3} \left(G(X) - \frac{1}{3} \text{erf}(X) \right) \right. \\ &\quad \left. + \frac{\delta^{ij}}{2} \frac{\text{erf}(X) - G(X)}{X} \right],\end{aligned}\quad (\text{A17})$$

where $\mathbf{v} = \mathbf{p}/M$, $v = |\mathbf{v}|$ and

$$X = \frac{v}{\sqrt{2}\sigma}. \quad (\text{A18})$$

2. Degenerate fermionic dark matter

We now consider the diffusion of astrophysical objects such as GCs in a halo of fermionic dark matter. In this case, the response function S becomes

$$\begin{aligned}S(\mathbf{p}, \mathbf{p}') &\simeq g_\chi \frac{(4\pi G m M)^2}{q^4} \int \frac{d^3 k}{(2\pi)^3} \frac{d^3 k'}{(2\pi)^3} \times \\ &\quad (2\pi)^4 \delta^{(4)}(p + k - p' - k') f_2(k) (1 - f_2(k'))\end{aligned}\quad (\text{A19})$$

Expanding $f_2(k')$ around k , we find additional contribution to the function S due to quantum statistics as

$$\begin{aligned}\Delta S(\mathbf{p}, \mathbf{p}') &\simeq -2\pi g_\chi \frac{(4\pi G m M)^2}{q^5} \times \\ &\quad \int \frac{d^3 k}{(2\pi)^3} \left(1 + \frac{\mathbf{q}}{2} \cdot \frac{\partial}{\partial \mathbf{p}} \right) \delta \left[\hat{\mathbf{q}} \cdot \left(\frac{\mathbf{k}}{m} - \frac{\mathbf{p}}{M} \right) \right] f_2^2(k),\end{aligned}\quad (\text{A20})$$

which is the same as Eq. (A10) upon substituting $M/\mu \rightarrow 1$ and $f_2 \rightarrow f_2^2$. We find

$$D_i(\mathbf{p}) = \frac{4\pi G^2 m^2 M^3 \ln \Lambda}{\mu_r} \frac{\partial}{\partial p^i} \left[h(\mathbf{p}; f_2) - \frac{\mu_r}{M} h(\mathbf{p}; f_2^2) \right] \quad (\text{A21})$$

$$D_{ij}(\mathbf{p}) = 4\pi G^2 m^2 M^4 \ln \Lambda \frac{\partial^2}{\partial p^i \partial p^j} \left[g(\mathbf{p}; f_2) - g(\mathbf{p}; f_2^2) \right] \quad (\text{A22})$$

Again, this result is obtained by expanding $f_2(k')$ around k assuming small momentum transfer.

For degenerate fermions with momentum $k < k_F$, we have $1 - f_2(k) \simeq e^{-\mu/T} = e^{-k_F^2/2mT}$ and thus the second diffusion coefficient is exponentially suppressed as a function of the chemical potential. In such case, the above formula for D_{ij} may not be the leading order term for the diffusion and a more careful treatment is required.

For the degenerate case, one can perform the \mathbf{k} and \mathbf{k}' integrations without expanding $f_2(k')$. This computation was already done in the context of neutrino transport in

a hot and dense medium [86] and dark matter thermalization in neutron stars [87]. We find:

$$S(\mathbf{p}, \mathbf{p}') = g_\chi \frac{(4\pi G m M)^2}{q^4} \frac{m^2 T}{2\pi q} \frac{z}{1 - e^{-z}} \left(1 + \frac{\xi_-}{z} \right) \quad (\text{A23})$$

where $z = -q^0/T$, $E_-^2 = m^2 + k_-^2$, $k_-^2 = (m^2/q^2)(q^0 + q^2/2m)^2$, and

$$\xi_- = \ln \left[\frac{1 + e^{(E_- - \mu)/T}}{1 + e^{(E_- - \mu)/T} e^z} \right]. \quad (\text{A24})$$

Integrating this response function with respect to \mathbf{q} , one obtains the diffusion coefficients for a degenerate medium.

3. Ultralight dark matter

It is discussed in [88] that the dynamical relaxation of stars in ultralight dark matter halo proceeds as stars scatter off a quasi-particle of ultralight dark matter whose size is order of its de Broglie wavelength, $\lambda_{\text{dB}} \sim 1/mv$. This observation is confirmed by Bar-Or et al [31], where the dynamical relaxation time scale as well as diffusion coefficients are computed in more rigorous way by using Fokker-Planck equation and stochastic gravitational potential.

The Boltzmann equation approach can also reproduce the dynamical relaxation time scale and diffusion coefficients. The gravitational scattering between ultralight dark matter and star can be described by the same matrix element, Eq. (A9), where M and m are the mass of star and ultralight dark matter, respectively. We treat the star as a pointlike particle, and this can be justified since the maximum momentum exchange $q \sim mv$ is much smaller than $1/r$ with a typical radius of star r . The function $S(\mathbf{p}, \mathbf{p}')$ is

$$\begin{aligned}S(\mathbf{p}, \mathbf{p}') &\simeq g_\chi \frac{(4\pi G m M)^2}{q^4} \int \frac{d^3 k}{(2\pi)^3} \frac{d^3 k'}{(2\pi)^3} \times \\ &\quad (2\pi)^4 \delta^{(4)}(p + k - p' - k') f_2(k) (1 + f_2(k')).\end{aligned}\quad (\text{A25})$$

The quantum correction is the same as Eq. (A20) with an opposite sign. Therefore, the diffusion coefficients are

$$D_i(\mathbf{p}) = \frac{4\pi G^2 m^2 M^3 \ln \Lambda}{\mu_r} \frac{\partial}{\partial p^i} \left[h(\mathbf{p}; f_2) + \frac{\mu_r}{M} h(\mathbf{p}; f_2^2) \right] \quad (\text{A26})$$

$$D_{ij}(\mathbf{p}) = 4\pi G^2 m^2 M^4 \ln \Lambda \frac{\partial^2}{\partial p^i \partial p^j} \left[g(\mathbf{p}; f_2) + g(\mathbf{p}; f_2^2) \right] \quad (\text{A27})$$

This reproduces the results of [31].

Appendix B: Maximum entropy DDM halos

In the derivation of the quasi-degenerate density profile, we adopt the assumption that a galactic structure may be described as a statistical ensemble close to equilibrium, in the sense of a maximal Boltzmann-Gibbs entropy. A similar approach to ours can be found in a number of earlier works [38, 55, 89–92].

The phase space distribution function $f(\mathbf{r}, \mathbf{p})$ and differential particle number density dN are related via

$$(2\pi)^3 \frac{dN}{d^3x d^3p} = g f(\mathbf{r}, \mathbf{p}). \quad (\text{B1})$$

The entropy of the gas is then given by the functional

$$S = -g \int \frac{d^3p d^3r}{(2\pi)^3} [f \ln f + (1-f) \ln(1-f)]. \quad (\text{B2})$$

Supplemented with Lagrange multipliers for the total energy and the total number of particles, the variation problem can be carried out along the lines of [55]. The maximum entropy result is

$$f(\mathbf{r}, \mathbf{p}) = \frac{1}{1 + \exp[z(\mathbf{r}, \mathbf{p})]}, \quad (\text{B3})$$

where

$$z(\mathbf{r}, \mathbf{p}) = \frac{\beta \mathbf{p}^2}{2m} + \beta m \Phi(\mathbf{r}) + \alpha, \quad (\text{B4})$$

with β and α the energy and particle number Lagrange multipliers. The gravitational potential is given by

$$\Phi(\mathbf{r}) = -Gmg \int \frac{d^3p' d^3r'}{(2\pi)^3} \frac{f(\mathbf{r}', \mathbf{p}')}{|\mathbf{r} - \mathbf{r}'|}. \quad (\text{B5})$$

By construction, the gravitational potential solves the Poisson equation, $\nabla^2 \Phi = 4\pi G\rho$, where the density is

$$\rho(\mathbf{r}) = mg \int \frac{d^3p}{(2\pi)^3} f(\mathbf{r}, \mathbf{p}). \quad (\text{B6})$$

We can make progress by evaluating the density,

$$\rho(\mathbf{r}) = -mg \left(\frac{m}{2\pi\beta} \right)^{3/2} \text{PolyLog} \left[\frac{3}{2}, -e^\varphi \right], \quad (\text{B7})$$

where $\varphi \equiv -\beta m \Phi(\mathbf{r}) - \alpha \equiv \beta \mu = \mu/T$, defining also the “chemical potential” $\mu(\mathbf{r}) = -m\Phi(\mathbf{r}) - \tilde{\alpha}$, with $\tilde{\alpha} \equiv \alpha/\beta$.

Using the Poisson equation and the definition of φ , we have

$$\nabla^2 \Phi = -\frac{1}{\beta m} \nabla^2 \varphi = 4\pi G\rho. \quad (\text{B8})$$

In the degenerate limit, $\mu/T = \varphi \gg 1$, The PolyLog function asymptotes to

$$-\text{PolyLog} \left[\frac{3}{2}, -e^\varphi \right] \rightarrow \frac{4\varphi^{3/2}}{3\sqrt{\pi}} = \frac{4(\mu/T)^{3/2}}{3\sqrt{\pi}}. \quad (\text{B9})$$

It’s therefore useful to rewrite Eq. (B8) as

$$\begin{aligned} \nabla^2 \mu(\mathbf{r}) & \\ &= \frac{4\sqrt{2}}{3\pi} g G m^{7/2} \mu_0^{1/2} \frac{\text{PolyLog} \left[\frac{3}{2}, -e^{\beta \mu_0 f} \right]}{\frac{4}{3\sqrt{\pi}} (\beta \mu_0)^{3/2}}, \end{aligned} \quad (\text{B10})$$

where we defined $\mu(0) \equiv \mu_0$ and $\mu(\mathbf{r}) = \mu_0 f(\mathbf{r})$. Let us also define a dimensionless radius x via $r = r_0 x$, with r_0 given by

$$r_0 = \sqrt{\frac{3\pi}{4\sqrt{2} g G m^{7/2} \mu_0^{1/2}}}. \quad (\text{B11})$$

Then, one finds the equation

$$\partial_x (x^2 \partial_x f) = x^2 \frac{\text{PolyLog} \left[\frac{3}{2}, -e^{(\mu_0/T)f} \right]}{\frac{4}{3\sqrt{\pi}} (\mu_0/T)^{3/2}}. \quad (\text{B12})$$

Given the solution for f , the density is simply

$$\rho = \frac{\sqrt{2}}{3\pi^2} g m^{\frac{5}{2}} \mu_0^{\frac{3}{2}} \frac{-\text{PolyLog} \left[\frac{3}{2}, -e^{\frac{\mu_0}{T} f} \right]}{\frac{4}{3\sqrt{\pi}} (\mu_0/T)^{3/2}}. \quad (\text{B13})$$

In the limit $\mu_0/T \gg 1$, the right-hand side of Eq. (B12) becomes $-x^2 f^{3/2}$, which is a Lane-Emden equation with a scale r_0 . The central density becomes $\rho(0) \equiv \rho_0 \approx (\sqrt{2}/3\pi^2) g m^{5/2} \mu_0^{3/2}$, i.e. $\mu_0 = (3\pi^2/\sqrt{2})^{2/3} (\rho_0/g)^{2/3} / m^{5/3}$, which implies¹⁴

$$r_0 \approx \frac{1}{2} \left(\frac{9\pi}{2G \rho_0 g^2 m^8} \right)^{\frac{1}{6}}. \quad (\text{B14})$$

The solution for ρ is constant near the origin and falls as $\rho \propto 1/r^2$ at $r \gg r_0$, with a “wriggle” feature near $r \sim r_0$. An example with $\mu_0/T = 10$ is shown in Fig. 13.

¹⁴ Our natural unit notation, which includes $\hbar \rightarrow 1$, may mask the fact that the characteristic radius Eq. (B14) is determined by quantum degeneracy pressure. This is easy to unmask by restoring $(2\pi)^3 \rightarrow (2\pi\hbar)^3$ on the left-hand side of Eq. (B1). Tracking \hbar through the computation gives a factor of \hbar on the right-hand side of Eq. (B14).




FIG. 13. The density profile found by solving Eq. (B12) and inserting into Eq. (B13) (blue line), compared with an analytical ansatz (orange) that demonstrates asymptotic behavior of the solution. Evidently, for $x \lesssim 1$ the density profile is constant, whereas the density asymptotes to $1/x^2$ for large x .

It is interesting to compare this class of solutions to the solutions obtained from the prescription of [39]. Rescaling the equations of [39] by r_0 from Eq. (B14), we plot the solution we find (at constant ρ_0) in Fig. 14 (named RSU), along with different solutions corresponding different μ_0/T . Evidently, the profile used by [39] bears a strong resemblance to the $\mu_0/T \sim 1$ case. Also, Fig. 14 shows that in the limit $\mu_0/T \rightarrow \infty$, the solution is a core with finite radius. We show it both by solving the density profile for $\mu_0/T \gg 1$ and by solving the Lane-Emden (LE) approximation that appears above.

In Fig. 15 we plot the circular velocities induced by the density profiles in Fig. 14.




FIG. 14. The density profile found by solving Eq. (B12) and inserting into Eq. (B13) for different values of μ_0/T , keeping ρ_0 constant. The dashed blue line (RSU) is based on the treatment of [39]. The dotted red line (LE) is based on solving the Lane-Emden equation, which is the $\mu_0/T \rightarrow \infty$ limit of the equations, as described in the text.




FIG. 15. The circular velocities $\sqrt{GM(r)/r}$ for the density profiles that appear in Fig. 14.

Appendix C: Jeans modeling

Following [43], the equation for the second velocity moments of a static spherical system of particles under the influence of a gravitational potential with enclosed mass $M(r)$ is

$$\frac{1}{\nu} \frac{d}{dr} (\nu \bar{v}_r^2) + 2 \frac{\beta \bar{v}_r^2}{r} = - \frac{GM}{r^2}, \quad (C1)$$

where $\nu(r)$ is the particles' density, $\bar{v}_r^2(r)$ is the radial second velocity moment, \bar{v}_θ^2 is the angular second velocity moment and $\beta \equiv 1 - \bar{v}_\theta^2/\bar{v}_r^2$ is the velocity anisotropy. For constant β , Eq. (C1) is solved by

$$\nu \bar{v}_r^2(r) = \frac{G}{r^{2\beta}} \int_r^\infty r'^{2\beta-2} \nu(r') M(r') dr'. \quad (C2)$$

This can be related to the line-of-sight (LOS) velocity,

$$\sigma_{\text{LOS}}^2(r) = \frac{2}{I(r)} \int_r^\infty \left(1 - \beta \frac{r^2}{r'^2}\right) \frac{\nu \bar{v}_r^2(r') r'}{\sqrt{r'^2 - r^2}} dr'. \quad (C3)$$

In modelling stellar kinematics in Fornax, we assume a Plummer profile with density and surface density given by, respectively,

$$\nu(r) = \frac{1}{(1 + r^2/r_p^2)^{5/2}} \frac{3L}{4\pi r_p^3} \quad (C4)$$

$$I(r) = \frac{1}{(1 + r^2/r_p^2)^2} \frac{L}{\pi r_p^2}. \quad (C5)$$

We use the radius parameter $r_p = 710$ pc [10], consistent with the stellar sample on which the kinematics data is based. (This radius parameter is about 20% smaller than the $r_p \approx 851$ pc reported in a new morphological study [21]. The difference is not crucial for our analysis. Moreover, we prefer to consider the photometry and spectroscopy of the same data set.)

Appendix D: CDM velocity dispersion in a cored profile

In this appendix we discuss some features of cored CDM halos, notably DF, partially following [40].

The velocity dispersion of dark matter is important to the discussion, therefore let us write the Jeans equation for the second radial velocity moment with a constant velocity anisotropy β , which has the solution [43]

$$\bar{v}_r^2(r) = \frac{G}{r^{2\beta}\rho(r)} \int_r^\infty r'^{2\beta} \frac{\rho(r')M(r')}{r'^2} dr', \quad (\text{D1})$$

which is the same as Eq. (C2) but in slightly different notation.

Consider a finite-core toy model, where the density is $\rho(0)$ for $r < r_c$ and 0 for $r > r_c$. Then, the solution of Eq. (D1) is [40]

$$\bar{v}_r^2(r) = \frac{2\pi G\rho(0)}{3(\beta+1)} \frac{1}{r^{2\beta}} (r_c^{2\beta+2} - r^{2\beta+2}). \quad (\text{D2})$$

For isotropic velocity dispersion $\beta = 0$, this reduces to

$$\bar{v}_r^2(r; \beta = 0) = \frac{2\pi G\rho(0)}{3} (r_c^2 - r^2). \quad (\text{D3})$$

Thus, for $r \ll r_c$,

$$\begin{aligned} \sigma_r &\equiv \sqrt{\bar{v}_r^2(r \ll r_c; \beta = 0)} \approx \sqrt{\frac{2\pi G\rho(0)}{3}} r_c \\ &\approx 30 \left(\frac{\rho(0)}{10^8 \frac{M_\odot}{\text{kpc}^3}} \right)^{\frac{1}{2}} \frac{r_c}{1 \text{ kpc}} \frac{\text{km}}{\text{s}}. \end{aligned} \quad (\text{D4})$$

We can also note the ratio,

$$X \equiv \frac{V_{\text{circ}}}{\sqrt{2}\sigma_r} \approx \sqrt{\frac{GM(r)}{2r}} / \sqrt{\frac{2\pi G\rho(0)r_c^2}{3}} = \frac{r}{r_c}, \quad (\text{D5})$$

indicating that the low-velocity approximation of the Chandrasekhar deceleration may apply inside a core, see Eq. (4). This implies a “phase-space suppression” to DF, as discussed in the main text (Sec. III).

Appendix E: Orbits under dynamical friction

In this appendix we review the solution of an orbit under the influence of DF. We write the equations of motion (EoM) in circular coordinates,

$$\ddot{\mathbf{r}} = (\ddot{r} - r\dot{\varphi}^2)\hat{r} + (2\dot{r}\dot{\varphi} + r\ddot{\varphi})\hat{\varphi} \quad (\text{E1})$$

$$= -\frac{GM(r)}{r^2}\hat{r} - \left| \frac{d\dot{\mathbf{r}}}{dt} \right|_{\text{DF}} \frac{\dot{\mathbf{r}}}{|\dot{\mathbf{r}}|}. \quad (\text{E2})$$

We express the deceleration $|d\dot{\mathbf{r}}/dt|_{\text{DF}}$ as $|\dot{\mathbf{r}}|/\tau$, where τ appears in Eq. (9).

Defining $r = R_0 x$, $t = T_0 \bar{t}$, $T_0^2 = R_0^3/GM(R_0)$, we find

$$x'' - x\varphi'^2 = -\frac{1}{x^2} \frac{M(R_0 x)}{M(R_0)} - \frac{x'}{\tau/T_0} \quad (\text{E3})$$

$$2x'\varphi' + x\varphi'' = -\frac{x\varphi'}{\tau/T_0} \quad (\text{E4})$$

where \prime is differentiation with respect to \bar{t} . Note, τ can depend on r and $|\dot{\mathbf{r}}|$. For a circular orbit, for example, the initial conditions can be set as $x(0) = 1$, $x'(0) = 0$, $\varphi(0) = 0$ and $\varphi'(0) = 1/x(0) = 1$, which has a revolution time of $\Delta\bar{t} = 2\pi$.

Solving the orbit of a decelerating test object generally requires numerical integration. We can understand some features of the solution analytically, however. Defining $v_\varphi \equiv r\dot{\varphi}$, the $\hat{\varphi}$ part of the EoM has the solution

$$rv_\varphi = (rv_\varphi)_0 \exp\left(-\int_0^{\bar{t}} \frac{dt'}{\tau}\right). \quad (\text{E5})$$

This solution expresses the decay of angular momentum of the test object. Using the circular velocity $v_{\text{circ}}^2 = GM(r)/r$, we can express the \hat{r} part of the EoM as

$$v_\varphi^2 - v_{\text{circ}}^2 = r\left(\ddot{r} + \frac{\dot{r}}{\tau}\right). \quad (\text{E6})$$

We can gain more analytical intuition by considering nearly-circular orbits, assuming that the inspiral rate is much smaller than the circular velocity, $r/\tau \ll v_{\text{circ}}$. Assuming that $\dot{r} \sim r/\tau$, $\ddot{r} \sim r/\tau^2$ and $r/\tau \ll v_{\text{circ}}$, Eq. (E6) implies $v_\varphi \approx v_{\text{circ}}$. We can use this to write

$$-\frac{rv_\varphi}{\tau} = \dot{r}v_\varphi + r\dot{v}_\varphi \approx \dot{r}v_{\text{circ}} + r\dot{v}_{\text{circ}} \quad (\text{E7})$$

$$= \frac{1}{2}v_{\text{circ}}\dot{r}\left(1 + \frac{d\ln M}{d\ln r}\right). \quad (\text{E8})$$

Rearranging, we find

$$\frac{\dot{r}}{r} \approx -\frac{2}{\left(1 + \frac{d\ln M}{d\ln r}\right)\tau}. \quad (\text{E9})$$

Using this, we can estimate the time it takes a test object to fall from r_0 down to $r < r_0$:

$$t(r; r_0) = \int_r^{r_0} \frac{dr}{2r} \left(1 + \frac{d\ln M}{d\ln r}\right) \tau(r, v_{\text{circ}}(r)). \quad (\text{E10})$$

Given the mass profile of the halo, $M(r)$, and a DF model encapsulated by τ , Eq. (E10) is a simple and quick estimate of the inspiral time of a test object.

For eccentric orbits, the approximation above is less justified. Defining eccentricity as $e \equiv (r_{\text{apo}} - r_{\text{peri}})/(r_{\text{apo}} + r_{\text{peri}})$ with apocenter radius r_{apo} and pericenter radius r_{peri} , we numerically tested Eq. (E10) for $e > 0$. In these calculations we defined r_0 and r via $(r_{\text{apo}} + r_{\text{peri}})/2$, where r_{apo} and r_{peri} are obtained per cycle of the orbital phase. With these definition, in numerical experiments representative of Fornax GCs we find that Eq. (E10) holds to better than 30% accuracy for $e \lesssim 0.5$.

Appendix F: The radial and projected CDF of GCs

Consider a population of identical GCs (all with the same mass), that start off their life at some initial time $t = 0$ on approximately circular orbits with a radial probability distribution function (PDF) $f_0(r_0)$ w.r.t. to initial radial coordinate r_0 . The CDF of initial GC positions is $F_0(r_0) = \int_0^{r_0} dy f_0(y)$. We are interested in computing the PDF and CDF of GC radial positions today, at $t = \Delta t$; call these $f_{\Delta t}(r)$ and $F_{\Delta t}(r)$.

DF causes GC orbits to inspiral inwards, and integrating along the orbit we can compute the function $r = r(r_0; \Delta t)$ and invert it to obtain $r_0 = r_0(r; \Delta t)$.¹⁵ Neglecting tidal disruption, we have

$$F_{\Delta t}(r) = F_0(r_0(r; \Delta t)). \quad (\text{F1})$$

Now we can use explicit results for $r_0(r; \Delta t)$ to connect $F_{\Delta t}(r)$ with $F_0(r_0)$ in different halo models. To this end we can use Eq. (E10),

$$\Delta t = \int_r^{r_0} \frac{dr'}{2r'} (1 + \alpha(r)) \tau(r'), \quad (\text{F2})$$

where $\alpha(r) \equiv d \ln M / d \ln r$. Let us consider the general features of $F_{\Delta t}$ for different halo shapes.

1. CDF of GCs in a cuspy halo

We have seen in the main analysis that a cuspy halo (i.e. the inner region of an NFW halo, where $\alpha \approx 2$) exhibits an approximately power law form for the DF time τ . For an approximately constant α and power law $\tau = \bar{\tau}(r/\bar{r})^\beta$, it is useful to define the *critical radius* r_{cr} via

$$\tau(r_{\text{cr}}) = \frac{2\beta}{1+\alpha} \Delta t. \quad (\text{F3})$$

The physical meaning of r_{cr} is that GCs that start their life at $r_0 \leq r_{\text{cr}}$ arrive at the origin within $t \leq \Delta t$. Using our power law form for τ , we have

$$r_{\text{cr}} = \bar{r} \left(\frac{2\beta}{1+\alpha} \frac{\Delta t}{\bar{\tau}} \right)^{1/\beta}. \quad (\text{F4})$$

In terms of r_{cr} , the solution of Eq. (F2) evaluates to

$$r_0(r; \Delta t) = r_{\text{cr}} \left(1 + \left(\frac{r}{r_{\text{cr}}} \right)^\beta \right)^{1/\beta}. \quad (\text{F5})$$

For GCs that satisfy $r \ll r_{\text{cr}}$ today, we can expand their starting point:

$$\begin{aligned} r_0(r; \Delta t) &= r_{\text{cr}} + \frac{r_{\text{cr}}}{\beta} \left(\frac{r}{r_{\text{cr}}} \right)^\beta + \dots \\ &= r_{\text{cr}} \left(1 + \frac{2}{1+\alpha} \frac{\tau(r)}{\Delta t} + \dots \right). \end{aligned} \quad (\text{F6})$$

In other words, for cuspy CDM halos, GCs that are currently seen at $r \ll r_{\text{cr}}$ must have originated near $r_0 \approx r_{\text{cr}}$. This means that for GCs with $r \ll r_{\text{cr}}$ today, the radial distribution today is not very sensitive to the (difficult to predict) initial distribution. We can make this point manifest by expanding Eq. (F1), using Eqs. (F4) and (F6):

$$F_{\Delta t}(r) \approx F_0(r_{\text{cr}}) + \frac{(1+\alpha)}{2\beta^2} f_0(r_{\text{cr}}) r_{\text{cr}} \frac{\tau(r)}{\Delta t} + \dots, \quad (\text{F7})$$

where the ... refer to higher powers of the small ratio $\tau(r)/\Delta t$. Above, the r -independent constant $F_0(r_{\text{cr}})$ counts GCs that have already settled to the center of the halo. These GCs at $r \approx 0$ were likely tidally disrupted, suggesting that in actually counting GCs in the system, we should eliminate the term $F_0(r_{\text{cr}})$ on the RHS of Eq. (F7). We thus have $F_{\Delta t}(r) \approx A(\tau(r)/\Delta t)$, where $A = \frac{(1+\alpha)}{2\beta^2} f_0(r_{\text{cr}}) r_{\text{cr}}$ is an order unity constant (r -independent) coefficient.¹⁶

We can summarise this section with two important conclusions. First, for a cuspy halo, all GCs that are born at $r < r_{\text{cr}}$ have arrived at $r \approx 0$ by today and are plausibly tidally disrupted. This means that observations today are not sensitive to initial conditions, characterised by different $f_0(r_0)$, that differ from each other only at $r < r_{\text{cr}}$; unless stellar age and metallicity measurements can identify the remnants and approximately count tidally disrupted GCs, on time scales of Gyrs after the disruption. Second, the radial CDF of GCs at small radii $r \ll r_{\text{cr}}$ should follow $F_{\Delta t}(r) \approx A(\tau(r)/\Delta t)$, with order unity A , irrespective of initial conditions.

Fig. 16 illustrates both of these two points, by showing two examples of f_0 and the resulting $F_{\Delta t}$. The DF time $\tau(r)$ and the critical radius r_{cr} are measurable given a model of the DM halo, fitted to stellar kinematics, and given GC age measurements that define Δt . This makes the predicted shape of $F_{\Delta t}$ measurable, in principle. In practice, however, projection effects (explained below) complicate the interpretation. In addition, the collection of GCs in Fornax seems too sparse to draw robust conclusions.

¹⁵ The monotonous decrease of r with time, that allowed this inversion, is lost for non-circular orbits. We could accommodate elliptical orbits approximately, by letting r represent the average between peri- and apo-center per cycle.

¹⁶ For the inner region of an NFW profile, we have seen that $\alpha \approx 2$ and $\beta \approx 2$, so $A \approx 0.4 N_{\text{cr}}$, where $N_{\text{cr}} = f_0(r_{\text{cr}}) r_{\text{cr}}$ counts the number of GCs that were located in a region of order r_{cr} around r_{cr} . Predicting the actual value of N_{cr} would require understanding of the initial cosmological formation of GCs, which is still not under full theory control.




FIG. 16. Initial radial PDF f_0 (**left**) and resulting current radial CDF $F_{\Delta t}$ (**right**) for a cuspy halo with $\tau(r) \propto r^{1.85}$ and $r_{\text{cr}} = 1.12$ (units on x-axis are arbitrary). Two different examples for f_0 are shown, leading to nearly identical $F_{\Delta t}$. The two versions of f_0 are normalised to yield $F_{\Delta t}(\infty) = 6$. On both panels, r_{cr} is marked with a vertical black line. On the left, the $r \ll r_{\text{cr}}$ approximation $F_{\Delta t} \approx A(\tau(r)/\Delta t)$ is shown by the red dotted curve.

2. CDF of GCs in a cored halo

Inside a core we expect $\alpha \approx 3$ and an approximately constant τ . Eq. (F2) is evaluated to

$$\Delta t = \frac{1 + \alpha}{2} \tau \ln \frac{r_0}{r} \quad (\text{F8})$$

and the radial CDF today is

$$F_{\Delta t}(r) \approx F_0(re^{\frac{2\Delta t}{(1+\alpha)\tau}}) \approx F_0(re^{\frac{\Delta t}{2\tau}}). \quad (\text{F9})$$

The distribution of GCs inside a core reflects a stretched version of the initial conditions. Because of this sensitivity to initial conditions, the degree of possible fine-tuning in the current positions of GCs may be difficult to assess.

3. Accounting for distribution of GC masses

The masses of GCs in Fornax vary over about an order of magnitude around $10^5 M_\odot$, and the instantaneous DF time satisfies $\tau \propto 1/m_*$ up to logarithmic corrections that we neglect here. It is therefore necessary to revise the prediction of the radial CDF of GCs to account for different GC masses.

In the case of a cuspy profile, where $\tau \propto r^\beta$, the critical radius scales as $r_{\text{cr}} \propto m_*^{1/\beta}$. For example, using the NFW fit of Fornax (which gives $\beta \approx 1.85$), relevant GC masses, and $\Delta t = 12$ Gyr we have:

$$r_{\text{cr}} \approx 0.7 \left(\frac{m_*}{M_{\text{GC4}}} \right)^{0.54} \text{ kpc} \quad (\text{F10})$$

$$\approx 1.6 \left(\frac{m_*}{M_{\text{GC3}}} \right)^{0.54} \text{ kpc}. \quad (\text{F11})$$

Suppose we have a set of GC masses m_{*i} with initial radial distribution functions $f_{0,i}(r_0)$. Summing over all

GC masses we find that as long as $r \ll r_{\text{cr},i}$, Eq. (F7) predicts that the total radial CDF today is (again omitting GCs that have already settled to the center of the halo)

$$\sum_i F_{\Delta t,i}(r) \approx F_{\Delta t,1}(r) \sum_i \frac{f_{0,i}(r_{\text{cr},i})}{f_{0,1}(r_{\text{cr},1})} \left(\frac{m_{*i}}{m_{*1}} \right)^{\frac{1}{\beta}-1}, \quad (\text{F12})$$

where $F_{\Delta t,1}(r)$ is the radial CDF of GCs of mass m_{*1} . We see that Eq. (F12) simply reproduces Eq. (F7) up to a modified over-all multiplicative constant.

4. Projected radius distribution

In reality we only know the projected distance of GCs from the center of Fornax, r_\perp , and not the true radial distance r . To obtain the CDF of projected radii, we can start with the surface density of GCs,

$$\Sigma_{\Delta t}(r_\perp) = \int_{-\infty}^{\infty} dz n(r) = 2 \int_{r_\perp}^{\infty} dr \frac{rn(r)}{\sqrt{r^2 - r_\perp^2}}, \quad (\text{F13})$$

where the 3D number density $n(r)$ is related to the radial PDF via $n = f_{\Delta t}(r)/4\pi r^2$. Using this relation we have

$$\Sigma_{\Delta t}(r_\perp) = \frac{1}{2\pi} \int_{r_\perp}^{\infty} dr \frac{f_{\Delta t}(r)}{r \sqrt{r^2 - r_\perp^2}}. \quad (\text{F14})$$

The CDF in r_\perp , that we define by $F_{\Delta t}^\perp(r_\perp)$, is given by:

$$F_{\Delta t}^\perp(r_\perp) = 2\pi \int_0^{r_\perp} dR R \Sigma_{\Delta t}(R) \quad (\text{F15})$$

$$= F_{\Delta t}(r_\perp) + \int_{r_\perp}^\infty dr f_{\Delta t}(r) \left(1 - \sqrt{1 - \frac{r_\perp^2}{r^2}} \right).$$

The CDF of projected radii contains the CDF of true radii, evaluated at $r = r_\perp$, plus another term that counts GCs at $r > r_\perp$ which projection casts into LOS inside of r_\perp . The added projection term can exceed the unprojected term, meaning that most GCs seen inside $r < r_\perp$ could be physically located at $r > r_\perp$. The effect is illustrated in Fig. 17.




FIG. 17. The effect of projection. Solid blue: un-projected radial CDF computed at the projected radius, $F_{\Delta t}(r_\perp)$. Dashed black: CDF of projected radii $F_{\Delta t}^\perp(r_\perp)$.

Appendix G: Exploration of initial conditions

The goal of this section is to explore the implications of uncertainties due to the line-of-sight projection in the true positions and velocities of GCs. Different projection angles lead to different true positions and velocities of a GC, affecting the orbital settling time under DF.

Consider the orbits of test bodies in a spherically-symmetric gravitational potential $\Phi(r)$. A given orbit lies on an orbital plane. In the coordinate system in Fig. 18, one can parameterize the orbital plane with the unit vector $\hat{n} = (\cos \alpha \sin \beta, \sin \alpha \sin \beta, \cos \beta)$. On the orbital plane, the radius $r(\varphi)$ and phase $\varphi(t)$ completely define the orbit. As in App. E, defining $r = r_{\text{true}}x$, $t = T_0\bar{t}$, $T_0^2 = r_{\text{true}}^3/GM(r_{\text{true}})$, one finds

$$x'' - x\varphi'^2 \approx -\frac{1}{x^2} \frac{M(r_{\text{true}}x)}{M(r_{\text{true}})} \quad (\text{G1})$$

$$(x^2\varphi')' \approx 0 \quad (\text{G2})$$

where \prime is differentiation with respect to \bar{t} . (Here we tentatively neglect DF.) The initial conditions are $x(0) = 1$, $\varphi(0) = 0$, $x'(0) = (-\Delta v_r \cos \theta + \Delta v_y \sin \theta)/(r_{\text{true}}/T_0)$, $\theta(0) = 0$ and $|\varphi'(0)| = \sqrt{\Delta v_z^2 + (\Delta v_r \sin \theta + \Delta v_y \cos \theta)^2}/(x(0)r_{\text{true}}/T_0)$. Evidently, $x'(0)^2 + (x(0)\varphi'(0))^2 = (\Delta v_r^2 + \Delta v_y^2 + \Delta v_z^2)/(r_{\text{true}}/T_0)^2$.

Given measured r_\perp , Δv_r and a model $\Phi(r)$, we explore the remaining orbital parameters which affect the inspiral time. We start with the true radius r_{true} . Given r_{true} and, for simplicity, assuming a circular orbit, the probability of observing $r_\perp < x r_{\text{true}}$ is $P(r_\perp/r_{\text{true}} < x) = (2/\pi) \arcsin x$, because $\sin \theta = r_\perp/r_{\text{true}}$ and θ is distributed uniformly for a circular orbit. Numerically, $P(r_\perp/r_{\text{true}} < 1/2) = 1/3$. We therefore explore orbits with true radius in the range $r_{\text{true}} = r_\perp \times [1, 2]$.

Next, consider the velocity. We explore a total velocity in the range $v_{\text{true}} \in [\max(\Delta v_r, 0.5V_{\text{circ}}(r)), \max(\Delta v_r, 1.5V_{\text{circ}}(r))]$. Defining the eccentricity $e \equiv (r_{\text{apo}} - r_{\text{peri}})/(r_{\text{apo}} + r_{\text{peri}})$, we find a maximal $e \sim 0.2 - 0.5$ for this range of v_{true} in the central $\lesssim 1$ kpc of Fornax. We note that if we decrease the lower bound of v_{true} we expect smaller inspiral times. Increasing the upper bound, however, results in larger inspiral times – but also in more tuning. An test object spends relatively little time near the pericenter. Specifically, $T_{\text{peri}}/T_{\text{apo}} \equiv (r_{\text{peri}}/v_{\text{peri}})/(r_{\text{apo}}/v_{\text{apo}}) \approx (1-e)^2/(1+e)^2$, yielding about $1/9$ for $e = 0.5$.

Considering the velocity components, we can take $\Delta v_z > 0$ without loss of generality. The sign of Δv_y is, however, important: under the transformation $\Delta v_y \rightarrow -\Delta v_y$, $\cos \theta \rightarrow -\cos \theta$, so $|\varphi'(0)|$ remains constant, but $x'(0) \rightarrow -x'(0)$. Since the specific energy is

$$\epsilon \approx \frac{v^2}{2} + \Phi(r) = \frac{\dot{r}^2}{2} + \frac{l^2}{2r^2} + \Phi(r), \quad (\text{G3})$$

where l is the specific angular momentum, this transformation returns the same orbit. The inspiral time is therefore invariant under this transformation. We shall explore then $\Delta v_y > 0$ and $\cos \theta$ positive or negative.

To sum up, for each GC (and a given model of the halo), we scan the range $r_{\text{true}} \in [1, 2]r_{\text{proj}}$. For each r_{true} we scan over $V_{\text{true}} \in [0.5, 1.5]V_{\text{circ}}(r_{\text{true}})$. For each true velocity we scan positive and negative $\cos \theta$. Finally, we test the two cases, $\Delta v_y = \sqrt{v_{\text{true}}^2 - \Delta v_r^2}$, $\Delta v_z = 0$ and $\Delta v_y = 0$, $\Delta v_z = \sqrt{v_{\text{true}}^2 - \Delta v_r^2}$. For each point in phase-space, we integrate the full equations of motion as in App. E. For each integration, we stop when $(r_{\text{apo}} + r_{\text{peri}})/2 \lesssim 0.3r_{\text{initial}}$ or after 10 Gyr (the first of the two). We then denote the integration time as τ_{inspiral} .




FIG. 18. The coordinate system that we adopt to analyse a given GC. The galactic dynamical center is in the origin. The observer is located at a very large X . The GC is located somewhere on the line $Z = 0, Y = r_{\perp}$. The true radius is therefore $r_{\text{true}} = r_{\perp} / \sin \alpha$. We assume Δv_r , the component of velocity in the X direction, can be measured. We assume that the rest of the components cannot be measured for now. The dotted line is the quasi-stable orbit of the GC.

- [1] M. G. Walker, M. Mateo, E. W. Olszewski, J. Penarrubia, N. W. Evans, and G. Gilmore, "A universal mass profile for dwarf spheroidal galaxies?," *The Astrophysical Journal* **704** no. 2, (2009) 1274.
- [2] D. R. Cole, W. Dehnen, J. I. Read, and M. I. Wilkinson, "The mass distribution of the Fornax dSph: constraints from its globular cluster distribution," *Mon. Not. Roy. Astron. Soc.* **426** (2012) 601, [arXiv:1205.6327 \[astro-ph.CO\]](#).
- [3] F. Hammer, Y. Yang, F. Arenou, J. Wang, H. Li, P. Bonifacio, and C. Babusiaux, "Orbital evidences for dark-matter-free Milky Way dwarf spheroidal galaxies," [arXiv:2002.09493 \[astro-ph.GA\]](#).
- [4] R. A. Flores and J. R. Primack, "Observational and theoretical constraints on singular dark matter halos," *Astrophys. J. Lett.* **427** (1994) L1–4, [arXiv:astro-ph/9402004](#).
- [5] B. Moore, "Evidence against dissipation-less dark matter from observations of galaxy haloes," *Nature* **370** no. 6491, (Aug., 1994) 629–631.
- [6] W. De Blok, "The core-cusp problem," *Advances in Astronomy* **2010** (2010) .
- [7] A. Fattahi, J. F. Navarro, T. Sawala, C. S. Frenk, L. V. Sales, K. Oman, M. Schaller, and J. Wang, "The cold dark matter content of Galactic dwarf spheroidals: no cores, no failures, no problem," [arXiv:1607.06479 \[astro-ph.GA\]](#).
- [8] J. I. Read, G. Iorio, O. Agertz, and F. Fraternali, "The stellar mass–halo mass relation of isolated field dwarfs: a critical test of LambdaCDM at the edge of galaxy formation," *Monthly Notices of the Royal Astronomical Society* **467** no. 2, (01, 2017) 2019–2038, <https://academic.oup.com/mnras/article-pdf/467/2/2019/10873983/mnras-stx147.pdf>, <https://doi.org/10.1093/mnras/stx147>.
- [9] L. J. Chang and L. Necib, "Dark Matter Density Profiles in Dwarf Galaxies: Linking Jeans Modeling Systematics and Observation," [arXiv:2009.00613 \[astro-ph.CO\]](#).
- [10] J. Read, M. Walker, and P. Steger, "Dark matter heats up in dwarf galaxies," *Mon. Not. Roy. Astron. Soc.* **484** no. 1, (2019) 1401–1420, [arXiv:1808.06634 \[astro-ph.GA\]](#).
- [11] J. S. Bullock and M. Boylan-Kolchin, "Small-Scale Challenges to the Λ CDM Paradigm," *Ann. Rev. Astron. Astrophys.* **55** (2017) 343–387, [arXiv:1707.04256 \[astro-ph.CO\]](#).
- [12] J. Simon, K. Bechtol, S. Birrer, S. Chakrabarti, F.-Y. Cyr-Racine, I. Dell’Antonio, A. Drlica-Wagner, C. Fassnacht, M. Geha, D. Gilman, Y. Hezaveh, D. Kim, T. Li, L. Strigari, and T. Treu, "Testing the nature of dark matter with extremely large telescopes." Snowmass 2021 - letter of interest, 2020. https://www.snowmass21.org/docs/files/summaries/CF/SNOWMASS21-CF3_CF0_Josh_Simon-038.pdf.
- [13] S. Tremaine, "The formation of the nuclei of galaxies. ii-the local group," *The Astrophysical Journal* **203** (1976) 345–351.
- [14] M.-Y. Wang, S. Koposov, A. Drlica-Wagner, A. Pieres, T. Li, T. de Boer, K. Bechtol, V. Belokurov, A. Pace, D. Bacon, et al., "Rediscovery of the sixth star cluster in the fornax dwarf spheroidal galaxy," *The Astrophysical Journal Letters* **875** no. 2, (2019) L13.
- [15] S. Chandrasekhar, "Dynamical Friction. I. General Considerations: the Coefficient of Dynamical Friction.," *Astrophys. J.* **97** (Mar, 1943) 255.
- [16] N. Meadows, J. F. Navarro, I. Santos-Santos, A. Benítez-Llambay, and C. Frenk, "Cusp or core? Revisiting the globular cluster timing problem in Fornax," *Mon. Not. Roy. Astron. Soc.* **491** no. 3, (Jan, 2020) 3336–3342, [arXiv:1910.11887 \[astro-ph.GA\]](#).
- [17] L. Hui, J. P. Ostriker, S. Tremaine, and E. Witten, "Ultralight scalars as cosmological dark matter," *Phys. Rev. D* **95** no. 4, (2017) 043541, [arXiv:1610.08297 \[astro-ph.CO\]](#).
- [18] T. de Boer and M. Fraser, "Four and one more: The formation history and total mass of globular clusters in the fornax dsph," *Astronomy & Astrophysics* **590** (2016) A35.
- [19] A. D. Mackey and G. F. Gilmore, "Surface brightness profiles and structural parameters for globular clusters in the Fornax and Sagittarius dwarf spheroidal galaxies," *Mon. Not. Roy. Astron. Soc.* **340** (2003) 175, [arXiv:astro-ph/0211396 \[astro-ph\]](#).
- [20] A. del Pino, S. L. Hidalgo, A. Aparicio, C. Gallart, R. Carrera, M. Monelli, R. Buonanno, and G. Marconi, "Spatial dependence of the star formation history in the central regions of the fornax dwarf spheroidal galaxy," *Monthly Notices of the Royal Astronomical Society* **433** no. 2, (2013) 1505–1516.
- [21] M.-Y. Wang, T. de Boer, A. Pieres, T. Li, A. Drlica-Wagner, S. Koposov, A. Vivas, A. Pace, B. Santiago, A. Walker, et al., "The morphology and structure of stellar populations in the fornax dwarf spheroidal galaxy from dark energy survey data," *The Astrophysical Journal* **881** no. 2, (2019) 118.
- [22] J. F. Navarro, C. S. Frenk, and S. D. M. White, "A Universal density profile from hierarchical clustering," *Astrophys. J.* **490** (1997) 493–508, [arXiv:astro-ph/9611107 \[astro-ph\]](#).
- [23] K. S. Oh, D. N. C. Lin, and H. B. Richer, "Globular Clusters in the Fornax Dwarf Spheroidal Galaxy," *Astrophys. J.* **531** no. 2, (Mar., 2000) 727–738.
- [24] G. W. Angus and A. Diaferio, "Resolving the timing problem of the globular clusters orbiting the fornax dwarf galaxy," *Monthly Notices of the Royal Astronomical Society* **396** no. 2, (2009) 887–893.
- [25] T. Goerdt, B. Moore, J. I. Read, J. Stadel, and M. Zemp, "Does the Fornax dwarf spheroidal have a central cusp or core?," *Mon. Not. Roy. Astron. Soc.* **368** no. 3, (May, 2006) 1073–1077, [arXiv:astro-ph/0601404 \[astro-ph\]](#).
- [26] K. Kaur and S. Sridhar, "Stalling of Globular Cluster Orbits in Dwarf Galaxies," *Astrophys. J.* **868** no. 2, (Dec., 2018) 134, [arXiv:1810.00369 \[astro-ph.GA\]](#).
- [27] G. Y. C. Leung, R. Leaman, G. van de Ven, and G. Battaglia, "A dwarf–dwarf merger and dark matter core as a solution to the globular cluster problems in the fornax dsph," *Monthly Notices of the Royal Astronomical Society* **493** no. 1, (Dec, 2019) 320–336. <http://dx.doi.org/10.1093/mnras/stz3017>.
- [28] P. Boldrini, R. Mohayaee, and J. Silk, "Embedding globular clusters in dark matter minihaloes solves the

- cusp-core and timing problems in the fornax dwarf galaxy,” *Monthly Notices of the Royal Astronomical Society* **492** no. 3, (2020) 3169–3178.
- [29] L. Berezhiani, B. Elder, and J. Khoury, “Dynamical Friction in Superfluids,” *JCAP* **1910** no. 10, (2019) 074, [arXiv:1905.09297 \[hep-ph\]](#).
- [30] S. Hartman, H. Winther, and D. Mota, “Dynamical friction in finite temperature superfluids, and the Fornax dwarf spheroidal,” [arXiv:2011.00116 \[astro-ph.CO\]](#).
- [31] B. Bar-Or, J.-B. Fouvry, and S. Tremaine, “Relaxation in a Fuzzy Dark Matter Halo,” *Astrophys. J.* **871** (2019) 28, [arXiv:1809.07673 \[astro-ph.GA\]](#).
- [32] L. Lancaster, C. Giovanetti, P. Mocz, Y. Kahn, M. Lisanti, and D. N. Spergel, “Dynamical Friction in a Fuzzy Dark Matter Universe,” *JCAP* **2001** no. 01, (2020) 001, [arXiv:1909.06381 \[astro-ph.CO\]](#).
- [33] V. Iršič, M. Viel, M. G. Haehnelt, J. S. Bolton, and G. D. Becker, “First constraints on fuzzy dark matter from Lyman- α forest data and hydrodynamical simulations,” *Phys. Rev. Lett.* **119** no. 3, (2017) 031302, [arXiv:1703.04683 \[astro-ph.CO\]](#).
- [34] N. Bar, D. Blas, K. Blum, and S. Sibiryakov, “Galactic rotation curves versus ultralight dark matter: Implications of the soliton-host halo relation,” *Phys. Rev. D* **98** no. 8, (2018) 083027, [arXiv:1805.00122 \[astro-ph.CO\]](#).
- [35] D. J. Marsh and J. C. Niemeyer, “Strong Constraints on Fuzzy Dark Matter from Ultrafaint Dwarf Galaxy Eridanus II,” *Phys. Rev. Lett.* **123** no. 5, (2019) 051103, [arXiv:1810.08543 \[astro-ph.CO\]](#).
- [36] N. Bar, K. Blum, J. Eby, and R. Sato, “Ultralight dark matter in disk galaxies,” *Phys. Rev. D* **99** no. 10, (2019) 103020, [arXiv:1903.03402 \[astro-ph.CO\]](#).
- [37] M. Safarzadeh and D. N. Spergel, “Ultra-light Dark Matter is Incompatible with the Milky Way’s Dwarf Satellites,” [arXiv:1906.11848 \[astro-ph.CO\]](#).
- [38] V. Domcke and A. Urbano, “Dwarf spheroidal galaxies as degenerate gas of free fermions,” *JCAP* **1501** no. 01, (2015) 002, [arXiv:1409.3167 \[hep-ph\]](#).
- [39] L. Randall, J. Scholtz, and J. Unwin, “Cores in dwarf galaxies from fermi repulsion,” *Monthly Notices of the Royal Astronomical Society* (Jan, 2017) stx161. <http://dx.doi.org/10.1093/mnras/stx161>.
- [40] J. A. Petts, A. Gualandris, and J. I. Read, “A semi-analytic dynamical friction model that reproduces core stalling,” *Mon. Not. Roy. Astron. Soc.* **454** no. 4, (Dec., 2015) 3778–3791, [arXiv:1509.07871 \[astro-ph.GA\]](#).
- [41] M. Kaplinghat, S. Tulin, and H.-B. Yu, “Dark Matter Halos as Particle Colliders: Unified Solution to Small-Scale Structure Puzzles from Dwarfs to Clusters,” *Phys. Rev. Lett.* **116** no. 4, (2016) 041302, [arXiv:1508.03339 \[astro-ph.CO\]](#).
- [42] S. Shao, M. Cautun, C. S. Frenk, M. Reina-Campos, A. J. Deason, R. A. Crain, J. D. Kruijssen, and J. Pfeffer, “The survival of globular clusters in a cuspy Fornax,” [arXiv:2012.08058 \[astro-ph.GA\]](#).
- [43] J. Binney and S. Tremaine, *Galactic Dynamics: Second Edition*. 2008.
- [44] S. Tremaine and M. D. Weinberg, “Dynamical friction in spherical systems,” *Mon. Not. Roy. Astron. Soc.* **209** (Aug., 1984) 729–757.
- [45] M. D. Weinberg, “Orbital Decay of Satellite Galaxies in Spherical Systems,” *Astrophys. J.* **300** (Jan., 1986) 93.
- [46] B. Bar-Or, J.-B. Fouvry, and S. Tremaine, “Relaxation in a Fuzzy Dark Matter Halo. II. Self-consistent kinetic equations,” [arXiv:2010.10212 \[astro-ph.GA\]](#).
- [47] B. Hendricks, A. Koch, M. Walker, C. I. Johnson, J. Peñarrubia, and G. Gilmore, “Insights from the outskirts: Chemical and dynamical properties in the outer parts of the fornax dwarf spheroidal galaxy,” *Astronomy & Astrophysics* **572** (Dec, 2014) A82. <http://dx.doi.org/10.1051/0004-6361/201424645>.
- [48] A. Lauberts, *ESO/Uppsala survey of the ESO(B) atlas*. 1982.
- [49] B. Letarte, V. Hill, P. Jablonka, E. Tolstoy, P. François, and G. Meylan, “VLT/UVES spectroscopy of individual stars in three globular clusters in the Fornax dwarf spheroidal galaxy,” *Astron. Astrophys.* **453** no. 2, (July, 2006) 547–554, [arXiv:astro-ph/0603315 \[astro-ph\]](#).
- [50] J. E. Morrison, S. Röser, B. McLean, B. Bucciarelli, and B. Lasker, “The Guide Star Catalog, Version 1.2: An Astrometric Recalibration and Other Refinements,” *Astrophys. J.* **121** no. 3, (Mar., 2001) 1752–1763.
- [51] M. F. Skrutskie, R. M. Cutri, R. Stiening, M. D. Weinberg, S. Schneider, J. M. Carpenter, C. Beichman, R. Capps, T. Chester, J. Elias, J. Huchra, J. Liebert, C. Lonsdale, D. G. Monet, S. Price, P. Seitzer, T. Jarrett, J. D. Kirkpatrick, J. E. Gizis, E. Howard, T. Evans, J. Fowler, L. Fullmer, R. Hurt, R. Light, E. L. Kopan, K. A. Marsh, H. L. McCallon, R. Tam, S. Van Dyk, and S. Wheelock, “The Two Micron All Sky Survey (2MASS),” *Astrophys. J.* **131** no. 2, (Feb., 2006) 1163–1183.
- [52] S. S. Larsen, J. P. Brodie, and J. Strader, “Detailed abundance analysis from integrated high-dispersion spectroscopy: globular clusters in the fornax dwarf spheroidal,” *Astronomy & Astrophysics* **546** (Oct, 2012) A53. <http://dx.doi.org/10.1051/0004-6361/201219895>.
- [53] S. M. Inoue, “The test for suppressed dynamical friction in a constant density core of dwarf galaxies,” *Mon. Not. Roy. Astron. Soc.* **397** (2009) 709, [arXiv:0901.4861 \[astro-ph.CO\]](#).
- [54] J. I. Read, T. Goerdt, B. Moore, A. P. Pontzen, J. Stadel, and G. Lake, “Dynamical friction in constant density cores: A failure of the Chandrasekhar formula,” *Mon. Not. Roy. Astron. Soc.* **373** (2006) 1451–1460, [arXiv:astro-ph/0606636 \[astro-ph\]](#).
- [55] D. Lynden-Bell and R. Wood, “The gravo-thermal catastrophe in isothermal spheres and the onset of red-giant structure for stellar systems,” *Mon. Not. Roy. Astron. Soc.* **138** (Jan., 1968) 495.
- [56] S. Tremaine and J. E. Gunn, “Dynamical Role of Light Neutral Leptons in Cosmology,” *Phys. Rev. Lett.* **42** (1979) 407–410. [66(1979)].
- [57] C. Di Paolo, F. Nesti, and F. L. Villante, “Phase space mass bound for fermionic dark matter from dwarf spheroidal galaxies,” *Mon. Not. Roy. Astron. Soc.* **475** no. 4, (2018) 5385–5397, [arXiv:1704.06644 \[astro-ph.GA\]](#).
- [58] D. Savchenko and A. Rudakovskiy, “New mass bound on fermionic dark matter from a combined analysis of classical dSphs,” *Mon. Not. Roy. Astron. Soc.* **487** no. 4, (2019) 5711–5720, [arXiv:1903.01862 \[astro-ph.CO\]](#).

- [59] A. Boyarsky, O. Ruchayskiy, and D. Iakubovskyi, “A Lower bound on the mass of Dark Matter particles,” *JCAP* **0903** (2009) 005, [arXiv:0808.3902 \[hep-ph\]](#).
- [60] J. Alvey, N. Sabti, V. Tiki, D. Blas, K. Bondarenko, A. Boyarsky, M. Escudero, M. Fairbairn, M. Orkney, and J. I. Read, “New Constraints on the Mass of Fermionic Dark Matter from Dwarf Spheroidal Galaxies,” [arXiv:2010.03572 \[hep-ph\]](#).
- [61] J. Bond, G. Efstathiou, and J. Silk, “Massive Neutrinos and the Large Scale Structure of the Universe,” *Phys. Rev. Lett.* **45** (1980) 1980–1984.
- [62] M. Viel, J. Lesgourgues, M. G. Haehnelt, S. Matarrese, and A. Riotto, “Constraining warm dark matter candidates including sterile neutrinos and light gravitinos with WMAP and the Lyman-alpha forest,” *Phys. Rev. D* **71** (2005) 063534, [arXiv:astro-ph/0501562](#).
- [63] M. Viel, G. D. Becker, J. S. Bolton, and M. G. Haehnelt, “Warm dark matter as a solution to the small scale crisis: New constraints from high redshift Lyman- α forest data,” *Phys. Rev. D* **88** (2013) 043502, [arXiv:1306.2314 \[astro-ph.CO\]](#).
- [64] J. Baur, N. Palanque-Delabrouille, C. Yèche, C. Magneville, and M. Viel, “Lyman-alpha Forests cool Warm Dark Matter,” *JCAP* **08** (2016) 012, [arXiv:1512.01981 \[astro-ph.CO\]](#).
- [65] M. Shoji and E. Komatsu, “Massive Neutrinos in Cosmology: Analytic Solutions and Fluid Approximation,” *Phys. Rev. D* **81** (2010) 123516, [arXiv:1003.0942 \[astro-ph.CO\]](#). [Erratum: *Phys. Rev. D* **82**, 089901 (2010)].
- [66] G. Choi, M. Suzuki, and T. Yanagida, “Degenerate fermion dark matter from a broken $U(1)_{B-L}$ gauge symmetry,” *Phys. Rev. D* **102** no. 3, (2020) 035022, [arXiv:2004.07863 \[hep-ph\]](#).
- [67] D. N. Spergel and P. J. Steinhardt, “Observational evidence for selfinteracting cold dark matter,” *Phys. Rev. Lett.* **84** (2000) 3760–3763, [arXiv:astro-ph/9909386](#).
- [68] M. Kaplinghat, R. E. Keeley, T. Linden, and H.-B. Yu, “Tying Dark Matter to Baryons with Self-interactions,” *Phys. Rev. Lett.* **113** (2014) 021302, [arXiv:1311.6524 \[astro-ph.CO\]](#).
- [69] M. Rocha, A. H. Peter, J. S. Bullock, M. Kaplinghat, S. Garrison-Kimmel, J. Onorbe, and L. A. Moustakas, “Cosmological Simulations with Self-Interacting Dark Matter I: Constant Density Cores and Substructure,” *Mon. Not. Roy. Astron. Soc.* **430** (2013) 81–104, [arXiv:1208.3025 \[astro-ph.CO\]](#).
- [70] A. Sokolenko, K. Bondarenko, T. Brinckmann, J. Zavala, M. Vogelsberger, T. Bringmann, and A. Boyarsky, “Towards an improved model of self-interacting dark matter haloes,” *JCAP* **12** (2018) 038, [arXiv:1806.11539 \[astro-ph.CO\]](#).
- [71] J. Zavala, M. Vogelsberger, and M. G. Walker, “Constraining self-interacting dark matter with the milky way’s dwarf spheroidals,” *Monthly Notices of the Royal Astronomical Society: Letters* **431** no. 1, (Feb, 2013) L20–L24. [http://dx.doi.org/10.1093/mnrasl/sls053](#).
- [72] S. Tulin and H.-B. Yu, “Dark Matter Self-interactions and Small Scale Structure,” *Phys. Rept.* **730** (2018) 1–57, [arXiv:1705.02358 \[hep-ph\]](#).
- [73] A. Fitts et al., “Dwarf Galaxies in CDM, WDM, and SIDM: Disentangling Baryons and Dark Matter Physics,” *Mon. Not. Roy. Astron. Soc.* **490** no. 1, (2019) 962–977, [arXiv:1811.11791 \[astro-ph.GA\]](#).
- [74] A. Robertson, R. Massey, V. Eke, J. Schaye, and T. Theuns, “The surprising accuracy of isothermal Jeans modelling of self-interacting dark matter density profiles,” [arXiv:2009.07844 \[astro-ph.CO\]](#).
- [75] K. Bondarenko, A. Sokolenko, A. Boyarsky, A. Robertson, D. Harvey, and Y. Revaz, “From dwarf galaxies to galaxy clusters: Self-Interacting Dark Matter over 7 orders of magnitude in halo mass,” [arXiv:2006.06623 \[astro-ph.CO\]](#).
- [76] J. I. Read, O. Agertz, and M. L. M. Collins, “Dark matter cores all the way down,” *Mon. Not. Roy. Astron. Soc.* **459** no. 3, (2016) 2573–2590, [arXiv:1508.04143 \[astro-ph.GA\]](#).
- [77] J. I. Read, M. Wilkinson, N. Evans, G. Gilmore, and J. T. Kleyna, “The Tidal stripping of satellites,” *Mon. Not. Roy. Astron. Soc.* **366** (2006) 429–437, [arXiv:astro-ph/0506687](#).
- [78] E. Hardy in *IAU Symposium. 206, Extragalactic Star Clusters*, D. Geisler, E. Grebel, and D. Minniti, eds. (San Francisco: ASP), 62, 2002.
- [79] J. Strader, J. P. Brodie, D. A. Forbes, M. A. Beasley, and J. P. Huchra, “Spectroscopy of globular clusters in the fornax dwarf galaxy,” *The Astronomical Journal* **125** no. 3, (Mar, 2003) 1291–1297. [https://doi.org/10.1086/2F367599](#).
- [80] S. Martocchia, E. Dalessandro, M. Salaris, S. Larsen, and M. Rejkuba, “Is fornax 4 the nuclear star cluster of the fornax dwarf spheroidal galaxy?,” *Monthly Notices of the Royal Astronomical Society* **495** no. 4, (Jun, 2020) 4518–4528. [http://dx.doi.org/10.1093/mnras/staa1482](#).
- [81] B. Hendricks, C. Boeche, C. I. Johnson, M. J. Frank, A. Koch, M. Mateo, and J. I. Bailey, “Evidence for a chemical enrichment coupling of globular clusters and field stars in the fornax dsph,” *Astronomy & Astrophysics* **585** (Dec, 2015) A86. [http://dx.doi.org/10.1051/0004-6361/201526996](#).
- [82] J. Pfeffer, J. M. D. Kruijssen, R. A. Crain, and N. Bastian, “The E-MOSAICS project: simulating the formation and co-evolution of galaxies and their star cluster populations,” *Mon. Not. Roy. Astron. Soc.* **475** no. 4, (Apr., 2018) 4309–4346, [arXiv:1712.00019 \[astro-ph.GA\]](#).
- [83] J. M. D. Kruijssen, J. L. Pfeffer, R. A. Crain, and N. Bastian, “The E-MOSAICS project: tracing galaxy formation and assembly with the age-metallicity distribution of globular clusters,” *Mon. Not. Roy. Astron. Soc.* **486** no. 3, (July, 2019) 3134–3179, [arXiv:1904.04261 \[astro-ph.GA\]](#).
- [84] M. N. Rosenbluth, W. M. MacDonald, and D. L. Judd, “Fokker-Planck Equation for an Inverse-Square Force,” *Phys. Rev.* **107** (1957) 1–6.
- [85] J. Binney and S. Tremaine, *Galactic Dynamics: Second Edition*. 2008.
- [86] S. Reddy, M. Prakash, and J. M. Lattimer, “Neutrino interactions in hot and dense matter,” *Phys. Rev. D* **58** (1998) 013009, [arXiv:astro-ph/9710115](#).
- [87] B. Bertoni, A. E. Nelson, and S. Reddy, “Dark Matter Thermalization in Neutron Stars,” *Phys. Rev. D* **88** (2013) 123505, [arXiv:1309.1721 \[hep-ph\]](#).

- [88] L. Hui, J. P. Ostriker, S. Tremaine, and E. Witten, "Ultralight scalars as cosmological dark matter," Phys. Rev. **D95** no. 4, (2017) 043541, [arXiv:1610.08297](#) [[astro-ph.CO](#)].
- [89] D. Lynden-Bell, "Statistical mechanics of violent relaxation in stellar systems," Mon. Not. Roy. Astron. Soc. **136** (1967) 101.
- [90] P.-H. Chavanis, "Phase transitions in self-gravitating systems: Self-gravitating fermions and hard spheres models," Phys. Rev. E **65** (2002) 056123, [arXiv:cond-mat/0109294](#).
- [91] P. Chavanis, "Statistical mechanics of violent relaxation in stellar systems," [arXiv:astro-ph/0212205](#).
- [92] P.-H. Chavanis, "Statistical mechanics and thermodynamic limit of self-gravitating fermions in dimensions," Physical Review E **69** no. 6, (Jun, 2004) .
<http://dx.doi.org/10.1103/PhysRevE.69.066126>.