



Transverse energy–energy correlations of jets in the electron–proton deep inelastic scattering at HERA

Ahmed Ali^{1,a}, Gang Li^{2,b}, Wei Wang^{3,c}, Zhi-Peng Xing^{3,d}

¹ Deutsches Elektronen-Synchrotron DESY, 22607 Hamburg, Germany

² Amherst Center for Fundamental Interactions, Department of Physics, University of Massachusetts Amherst, Amherst, MA 01003, USA

³ INPAC, Shanghai Key Laboratory for Particle Physics and Cosmology, MOE Key Laboratory for Particle Astrophysics and Cosmology, School of Physics and Astronomy, Shanghai Jiao Tong University, Shanghai 200240, China

Received: 6 August 2020 / Accepted: 29 October 2020 / Published online: 30 November 2020
© The Author(s) 2020

Abstract We study the event shape variables, transverse energy–energy correlation TEEC ($\cos\phi$) and its asymmetry ATEEC ($\cos\phi$) in deep inelastic scattering (DIS) at the electron–proton collider HERA, where ϕ is the angle between two jets defined using a transverse-momentum (k_T) jet algorithm. At HERA, jets are defined in the Breit frame, and the leading nontrivial transverse energy–energy correlations arise from the 3-jet configurations. With the help of the NLOJET++, these functions are calculated in the leading order (LO) and the next-to-leading order (NLO) approximations in QCD at the electron–proton center-of-mass energy $\sqrt{s} = 314$ GeV. We restrict the angular region to $-0.8 \leq \cos\phi \leq 0.8$, as the forward- and backward-angular regions require resummed logarithmic corrections, which we have neglected in this work. Following experimental jet-analysis at HERA, we restrict the DIS-variables x , $y = Q^2/(xs)$, where $Q^2 = -q^2$ is the negative of the momentum transfer squared q^2 , to $0 \leq x \leq 1$, $0.2 \leq y \leq 0.6$, and the pseudo-rapidity variable in the laboratory frame (η^{lab}) to the range $-1 \leq \eta^{\text{lab}} \leq 2.5$. The TEEC and ATEEC functions are worked out for two ranges in Q^2 , defined by $5.5 \text{ GeV}^2 \leq Q^2 \leq 80 \text{ GeV}^2$, called the low- Q^2 -range, and $150 \text{ GeV}^2 \leq Q^2 \leq 1000 \text{ GeV}^2$, called the high- Q^2 -range. We show the sensitivity of these functions on the parton distribution functions (PDFs), the factorization (μ_F) and renormalization (μ_R) scales, and on $\alpha_s(M_Z^2)$. Of these the correlations are stable against varying the scale μ_F and the PDFs, but they do depend on μ_R . For the choice of the scale $\mu_R = \sqrt{\langle E_T \rangle^2 + Q^2}$, advocated in earlier jet analysis at HERA, the shape variables TEEC and ATEEC are found

perturbatively robust. These studies are useful in the analysis of the HERA data, including the determination of $\alpha_s(M_Z^2)$ from the shape variables.

1 Introduction

Event shape variables involving the energy-momentum variables of hadrons and jets have played a crucial role in testing Quantum Chromodynamics (QCD), providing a detailed comparison with the experimentally measured shapes in high energy collisions and in determining the strong interaction coupling constant $\alpha_s(Q^2)$. Of these, the energy–energy correlation (EEC) and its asymmetry (AEEC), introduced by Basham et al. in e^+e^- annihilation [1,2] have received a lot of experimental and theoretical attention. Next-to-leading order (NLO) corrections in $\alpha_s(Q^2)$ were calculated long ago for the EEC in e^+e^- annihilation, using a number of different methods to regulate the soft and collinear divergences [3–10]. Accurate numerical results for the EEC are available from the program Event 2, based on the dipole subtraction technique [11,12]. EEC has also been calculated to NNLO accuracy in perturbative QCD [13,14] and in the next-to-next-to-leading logarithms (NNLL) [15]. Recent advances in theoretical calculational techniques have led to a renaissance of interest in this topic. In particular, an analytic NLO calculation of the EEC in e^+e^- annihilation [16,17], and an all-order factorization formula for the EEC in the back-to-back limit [18–21], are now available. We also mention here the derivation of the EEC function in the maximally supersymmetric $N = 4$ super-Yang-Mills theory in the NLO accuracy [22], which has been recently extended up to NNLO accuracy [23]. Experimental measurements of EEC in e^+e^- annihilation are discussed in [24–28].

^a e-mail: ahmed.ali@desy.de

^b e-mail: ligang@umass.edu

^c e-mail: wei.wang@sjtu.edu.cn (corresponding author)

^d e-mail: zpxing@sjtu.edu.cn

Following EEC in e^+e^- annihilation, transverse energy–energy correlation (TEEC) and the corresponding asymmetry (ATEEC) were introduced in hadronic collisions at the SP \bar{P} S [29], but did not evoke much experimental interest. With the advent of the LHC era, NLO corrections were calculated in pp collisions [30]. They have been used by the ATLAS collaboration for comparison with data and in the determination of $\alpha_s(M_Z^2)$ from these shape functions [31, 32]. Recently, TEEC in the dijet back-to-back limit in hadronic collisions has been derived, achieving an impressive perturbative simplicity [33]. Currently the TEEC-data in pp collisions are restricted in their theoretical interpretation to NLO accuracy.

What concerns deep inelastic scattering (DIS), event shape variables have also received a lot of theoretical attention [34–39]. Prominent among them are the thrust-distribution, 1-jettiness, jet-broadening, and the C parameter, which have been calculated to very high accuracy in fixed order (NNLO) [39], and in the resummed leading logarithms (N^3LL) [38]. Some of these event shape variables have been measured by the H1 [40] and ZEUS [41] collaborations at HERA. The definitions of these shape variables together with some others, such as the jet shape, can be seen in [42], where DIS and photoproduction experiments at HERA are reviewed. However, to the best of our knowledge, the transverse energy–energy correlation between the final state jets in deep inelastic scattering has neither been calculated nor measured so far. Analogous to the TEEC for hadronic collisions [29, 30], TEEC in DIS is introduced in Eq. (1) in the next section. It involves transverse energy correlations in two jets, defined by a jet-definition and jet algorithm, separated by an azimuthal angle ϕ . We calculate TEEC and its asymmetry in DIS at HERA under realistic experimental conditions.

Jets at HERA are defined in the Breit frame, in which the exchanged photon is at rest and the incoming and outgoing quarks are along the z direction. In this frame, the involved hadronic final states have zero total transverse momentum, and thus the leading nontrivial transverse energy–energy correlation comes from the 3-jet configurations. To match the measurements of jets at HERA, we adopt the transverse-momentum (k_T) algorithm to classify the jets [43] and calculate the TEEC and its asymmetry (ATEEC) in the kinematic conditions employed typically in H1 and ZEUS. The calculations are done in the NLO accuracy in the central angular region, $-0.8 \leq \cos \phi \leq 0.8$. This avoids the back-to-back angular configuration, i.e., near $\phi = \pi$, where the leading logs (LL) and the next-to-leading logs (NLL), $\alpha_s^m(\mu) \ln^n \tau$ ($m \leq n$) in the variable $\tau = \ln(1 + \cos \phi)/2$, have to be resummed. For the fixed-order perturbative calculations, we have used the NLOJET++ package [44, 45] and have tested it against the distributions obtained by Madgraph [46]. To achieve numerical stability, we have generated 10^9 DIS events at HERA ($\sqrt{s} = 314$ GeV), allowing us to reach an

statistical accuracy of a few percent over most of the phase space.

Being weighted by the product of transverse energies of jets, both the TEEC and ATEEC are expected to be insensitive to the parton distribution functions (PDFs). To quantify this, we use two PDF sets of relatively recent vintage, the CT18 [47], and MMHT14 [48]. The main theoretical uncertainty in the jet physics comes from the scale-dependence, of these the so-called factorization scale μ_F enters through the PDFs, and the partonic matrix elements depend essentially on the renormalization scale μ_R . Detailed studies done for the inclusive jet and dijet data at HERA show that the μ_F -dependence of the cross sections is small, but the μ_R -dependence is substantial in the NLO accuracy [49, 50]. We study these dependencies in TEEC and ATEEC, following the choice of the nominal scale, $\mu_0 = \sqrt{\langle E_T \rangle^2 + Q^2}$, where $\langle E_T \rangle$ denotes the average of E_T , as advocated in these papers. Varying the scales in the range “ $\mu_F = [0.5, 2]\mu_0$ ”, we find that the μ_F -dependence is small in the TEEC, not exceeding 5% over the $\cos \phi$ range, but the μ_R -dependence is found to be significant. Thus, NNLO improvements are needed to reduce the μ_R -uncertainty. However, fitting the HERA data on TEEC may also effectively reduce the allowed μ_R -range. Finally, we show the sensitivity of the TEEC and ATEEC on the strong coupling constant $\alpha_s(M_Z^2)$, for three representative values $\alpha_s(M_Z) = 0.108, 0.118, 0.128$. With the nominal choice of the scales $\mu_F = \mu_R = \mu_0$, and the current central value of $\alpha_s(M_Z) = 0.118$ [51], we show that the differential distributions TEEC($\cos \phi$) and ATEEC($\cos \phi$) are remarkably stable perturbatively in both the Q^2 -ranges. This remains to be tested in the NNLO accuracy. Our study presented here makes a good case for using the TEEC in DIS-data as a precision test of perturbative QCD, following similar analyses done for the high energy pp data at the LHC.

The rest of this paper is organized as follows. Section 2 collects the definitions of TEEC and its asymmetry. Experimental cuts to calculate these functions are stated in this section together with the jet algorithm used and the jet definitions. In Sect. 3, we present the numerical results calculated at next-to-leading order in α_s and estimate the uncertainty in the shape variables TEEC and ATEEC arising from the different PDFs, and the scale-dependence by varying the scale μ_F and μ_R . Of these, the μ_R -dependence is substantial. Fixing the scale μ_R to the nominal value μ_0 , which provides a good fit of the inclusive-jet and dijet data at HERA [49, 50], we show the sensitivity of the TEEC and ATEEC on $\alpha_s(M_Z^2)$. A comparison of the LO and NLO results is also presented here. We summarise our results in the last section. A check of the NLOJET++ calculation is shown in Appendix-A at the LO, by using the package MadGraph5_aMC@NLO [46] with the MMHT14 PDF set.

2 Transverse energy–energy correlation and its asymmetry

In the Breit frame, the transverse energy–energy correlation in $\gamma(q) + p \rightarrow a + b + X$ involving hadrons or jets is expressed as:

$$\begin{aligned} \frac{1}{\sigma'} \frac{d\Sigma'}{d\cos\phi} &= \frac{\sum_{a,b} \int dE_T d\cos\phi_{ab} \frac{d\sigma_{\gamma p \rightarrow a+b+X}}{dE_T d\cos\phi_{ab}} \frac{2E_{T,a}E_{T,b}}{|\sum_i E_{T,i}|^2} \delta(\cos\phi_{ab} - \cos\phi)}{\int dE_T d\sigma_{\gamma p \rightarrow a+b+X}/dE_T} \\ &= \frac{1}{N} \sum_{A=1}^N \frac{1}{\Delta\cos\phi} \sum_{\text{pairs in } \Delta\cos\phi} \frac{2E_{T,a}^A E_{T,b}^A}{(E_T^A)^2}, \end{aligned} \quad (1)$$

where $E_{T,a}$ and $E_{T,b}$ are transverse energies of two jets or hadrons. The δ -function assures that these hadrons or jets are separated by the azimuthal angle ϕ , and the cross section σ' and Σ' indicate kinematic cuts on the integrals, defined later. The second expression is valid for a sample of N hard-scattering multi-jet events, labelled by the index A . The associated asymmetry (ATEEC) is then defined as the asymmetry between the forward ($\cos\phi > 0$) and backward ($\cos\phi < 0$) parts of the TEEC:

$$\frac{1}{\sigma'} \frac{d\Sigma'^{\text{asym}}}{d\cos\phi} \equiv \frac{1}{\sigma'} \frac{d\Sigma'}{d\cos\phi} \Big|_{\phi} - \frac{1}{\sigma'} \frac{d\Sigma'}{d\cos\phi} \Big|_{\pi-\phi}. \quad (2)$$

Due to the factorization of the amplitudes in QCD, the denominator of the first equation in Eq. (1) $d\sigma_{\gamma p \rightarrow a+b+X}/dE_T$ can be written as a convolution of the parton distribution functions (PDFs) $f_{q/p}(x_1)$, where x_1 is the fractional energy of the proton carried by the parton q , and the parton level cross section. In the leading order, this is given by $\sigma_{\gamma q \rightarrow b_1 b_2}$. As the for numerator, it can also be expressed as the convolution of PDFs with $2 \rightarrow 3$ parton level subprocess, in the leading order, such as $\gamma q \rightarrow qgg$. Thus, TEEC is calculated from the following expression:

$$\frac{1}{\sigma'} \frac{d\Sigma'}{d\cos\phi} = \frac{\sum_{j,a,b} \int dE_T d\cos\phi_{ab} f_{j/p}(x_1) \star d\sigma_{\gamma j \rightarrow b_1 b_2 b_3} / (dE_T d\cos\phi_{ab}) \frac{2E_{T,a}E_{T,b} \delta(\cos\phi - \cos\phi_{ab})}{|\sum_i E_{T,i}|^2}}{\sum_j f_{j/p}(x_1) \star \sigma_{\gamma j \rightarrow b_1 b_2}}, \quad (3)$$

where the symbol \star stands for the convolution and j represents quarks and gluons. Which processes are included in the calculations of the TEEC depends on the theoretical accuracy. In NLO, this involves $2 \rightarrow 2$, $2 \rightarrow 3$ and $2 \rightarrow 4$ partonic subprocesses. Some representative Feynman diagrams of the subprocess are shown in Fig. 1. In the upper row of Fig. 1, we show the leading order (LO) (a), NLO real (b) and NLO virtual diagrams (c) which enter in the calculations of the numerator of Eq. (3). In the lower row of this figure, the Feynman diagrams of the subprocess in the denominator of

Eq. (3) are shown. Of these, (d, e) are LO diagrams, and the NLO virtual corrections are represented by the diagram (f). The NLO real diagram in inclusive two jet cross section are the same as the LO diagrams of three jet cross section of which we have shown a representative diagram (a) in Fig. 1.

As defined in Eq. (3), the TEEC correlation $\frac{1}{\sigma'} \frac{d\Sigma'}{d\cos\phi}$ is a normalized variable. In particular, the dependence of the TEEC on the PDFs is compensated to a large extent. Thus, to a good approximation, a factorized result is expected,

$$\frac{1}{\sigma'} \frac{d\Sigma'}{d\cos\phi} \sim \frac{\alpha_s(\mu)}{\pi} F(\cos\phi), \quad (4)$$

which can be perturbatively improved by including higher orders.

We calculate the TEEC and ATEEC close to experimental conditions used by the HERA experiments H1 and ZEUS, which assume a certain selection criteria based on physical cuts on the kinematic variables. They are defined as follows: The basic DIS kinematic variables x and $y = Q^2/(sx)$ satisfy

$$0 < x < 1, \quad 0.2 < y < 0.6. \quad (5)$$

Besides, we restrict the range of the pseudo-rapidity in the laboratory frame (η^{lab}) as

$$-1 < \eta^{\text{lab}} < 2.5. \quad (6)$$

The pseudorapidity is related to the polar angle θ , defined with respect to the proton beam direction, by $\eta^{\text{lab}} = -\ln \tan(\theta/2)$. We also use the right-handed co-ordinate system of the H1 collaboration, in which the positive z -axis is in the direction of the proton beam, and the nominal interaction point is located at $z = 0$.

We calculate the TEEC and ATEEC in the Breit frame used by experiments at HERA. In this frame, transverse energy E_T is z -axis boost invariant and $\gamma p \rightarrow jjjX$ is the nontrivial process at the leading order. The cuts for the transverse energy

of dijet and trijet events, defined in the Breit frame, are as follows:

$$\begin{aligned} 5 \text{ GeV} &< \langle E_T \rangle_2 < 50 \text{ GeV} \\ 5.5 \text{ GeV} &< \langle E_T \rangle_3 < 40 \text{ GeV}, \end{aligned} \quad (7)$$

where the $\langle E_T \rangle_2$ and $\langle E_T \rangle_3$ denote $\frac{1}{2}(E_T^{\text{jet1}} + E_T^{\text{jet2}})$ and $\frac{1}{3}(E_T^{\text{jet1}} + E_T^{\text{jet2}} + E_T^{\text{jet3}})$, respectively. These cuts are consistent with the measurement at HERA [49]. Following the practice in the HERA experimental analysis, we use the k_T




Fig. 1 Representative Feynman diagrams of the partonic subprocess in $\gamma^* + p$ scattering which are included in the numerator (first line) and denominator (second line) of Eq. (3). Here, the virtual photon is denoted by a wavy line and the gluon by a curled line

jet-algorithm [51], where the distance measure of partons (i, j) is given by

$$d_{ij} = \min(k_{ii}^2, k_{jj}^2) \frac{(\eta_i - \eta_j)^2 + (\phi_i - \phi_j)^2}{R^2},$$

$$d_{iB} = k_{ii}^2. \quad (8)$$

Here B represents the “beam jet” of the proton: particles with small momenta transverse to the beam axis, and R is the cone-size parameter of the jet which we set to $R = 1.0$ in our calculation. We use two different PDF sets, CT18 [47] and MMHT14 [48], and explore the uncertainty on the TEEC ($\cos \phi$) and ATEEC ($\cos \phi$) distributions from these two sets in the next section.

It has become customary to determine the QCD coupling constant at the scale $\mu = M_Z$ [52]. To determine $\alpha_s(M_Z^2)$ from TEEC ($\cos \phi$) and ATEEC ($\cos \phi$) in DIS, the cross section can be expressed as:

$$\sigma = \sum_k \int dx f_k(x, \mu_F) \sigma_k(x, \mu_F, \mu_R), \quad (9)$$

where k denotes a parton (quark or gluon), $f_k(x, \mu_F)$ is the parton density, and $\sigma_k(x, \mu_F, \mu_R)$ is the partonic cross section, which depends on the renormalization scale μ_R and the factorization scale μ_F . The partonic cross section is calculated in perturbative QCD as an expansion in α_s :

$$\sigma_k = \sum_n \alpha_s^n(\mu_R) \sigma_k^{(n)}(x, \mu_R, \mu_F). \quad (10)$$

As the μ_F -dependence is very mild on TEEC, as shown later, the dominant scale-dependence of the cross section enters through the scale μ_R , i.e., from $\alpha_s(\mu_R)$, which we relate to $\alpha_s(M_Z^2)$ on an event-by-event basis in our simulations. The μ_R dependence of α_s is given by renormalization group equation

$$\mu_R \frac{d\alpha_s}{d\mu_R} = \beta(\alpha_s). \quad (11)$$

Table 1 Dijet cross sections at HERA with $\sqrt{s} = 314$ GeV in the two $\langle E_T \rangle_2$ -bins defined in the text and the corresponding HERA data from the Table 7 in H1 collaboration [49]

	Bin1	Bin2
$\sigma_{\text{HERA}}[\text{pb}]$	$299 \pm 9.9 \pm 52.3$	$185 \pm 3.7 \pm 13.9$
$\sigma_{\text{NLOJET++}}[\text{pb}]$	298.03 ± 3.93	199.9 ± 3.04

In the NLO calculation, the two-loop β -function is used for transcribing $\alpha_s(\mu)$ to $\alpha_s(M_Z^2)$ with a certain scale μ which is relevant for the jets defined above. The coupling constant $\alpha_s(\mu)$ is given as

$$\alpha_s(\mu) = \frac{1}{b_0 \log(\mu^2/\Lambda^2)} \left[1 - \frac{b_1 \log(\log(\mu^2/\Lambda^2))}{b_0^2 \log(\mu^2/\Lambda^2)} \right],$$

$$b_0 = \frac{33 - 2n_f}{12\pi}, \quad b_1 = \frac{153 - 19n_f}{24\pi^2}. \quad (12)$$

Here, n_f is the number of quark flavors, which is determined by the scale μ , we have set $n_f = 5$, and Λ is the QCD parameter, which is determined by the value of $\alpha_s(M_Z^2)$. In the LO calculation, we set $b_1 = 0$ in the above expression. In our numerical results, we present TEEC($\cos \phi$) and ATEEC($\cos \phi$) calculated in the LO and NLO for the same value of $\alpha_s(M_Z^2)$, which implies a different value of Λ in the LO and NLO. As already stated, the $\alpha_s(\mu)$ -dependence enters essentially through $\mu = \mu_R$. Since μ_R is not determined uniquely, there will remain a residual scale-dependence in the differential distributions for TEEC ($\cos \phi$) and ATEEC ($\cos \phi$). In the next section, we show the dependence of TEEC ($\cos \phi$) and ATEEC ($\cos \phi$) at HERA on the scales μ_F , μ_R and $\alpha_s(M_Z^2)$.

Before ending this section, we remark that very recently another shape variable involving the azimuthal angle correlation of the lepton and hadron in DIS process has been





Fig. 2 Differential distribution $1/\sigma' d\Sigma'/d(\cos\phi)$ and its asymmetry $1/\sigma' d\Sigma'^{asym}/d(\cos\phi)$, calculated in Next-to-Leading order for the low- Q^2 range $5.5 \text{ GeV}^2 < Q^2 < 80 \text{ GeV}^2$ for the ep center-of-mass



energy $\sqrt{s} = 314 \text{ GeV}$ at HERA. The two input PDFs are indicated on the upper frames. The lower frames show $\Delta[\text{TEEC}(\cos\phi)]_{\text{PDF}}$ and $\Delta[\text{ATEEC}(\cos\phi)]_{\text{PDF}}$, defined in Eq. (15)





Fig. 3 Differential distribution $1/\sigma' d\Sigma'/d(\cos\phi)$ and its asymmetry $1/\sigma' d\Sigma'^{asym}/d(\cos\phi)$ as in Fig. 2, but for the high- Q^2 range $150 \text{ GeV}^2 < Q^2 < 1000 \text{ GeV}^2$ at HERA



proposed and calculated in [53], which is defined as

$$\begin{aligned} \ell HTEC(\cos\phi) &\equiv \sum_a \int d\sigma_{\ell p \rightarrow \ell + a + X} \frac{E_{T,\ell} E_{T,a}}{E_{T,\ell} \sum_i E_{T,i}} \\ &\quad \times \delta(\cos\phi_{\ell a} - \cos\phi) \\ &= \sum_a \int d\sigma_{\ell p \rightarrow \ell + a + X} \frac{E_{T,a}}{\sum_i E_{T,i}} \\ &\quad \times \delta(\cos\phi_{\ell a} - \cos\phi), \end{aligned} \quad (13)$$

where the sum runs over all hadrons and $\cos\phi_{\ell a}$ is the cosine of the azimuthal angle between the lepton and the hadron. As seen in the second of the above equation, transverse energy of the lepton drops out of this variable. As opposed to the shape

variable TEEC, defined here in Eq. (1) for DIS, as well as the EEC/TEEC variables defined earlier in e^+e^- annihilation [1,2] and pp collisions [29], which involve (transverse) energy weighted azimuthal angle correlations between two jets or hadrons, the shape variable defined in [53] is the azimuthal angle correlation between the lepton and a hadron (or a jet) weighted by the transverse energy of a single hadron (or jet). We emphasize that $\ell HTEC(\cos\phi)$, defined in [53] and Eq. (13), while interesting in its own right, is a different variable from TEEC. Lepton-jet correlation in DIS has also been studied in [54], and revisited very recently in [55], where a detailed derivation of the formalism used and a phenomeno-




Fig. 4 Factorization scale dependence of the differential distribution $1/\sigma' d\Sigma'/d(\cos\phi)$ and its asymmetry $1/\sigma' d\Sigma'^{asym}/d(\cos\phi)$ in the leading order (upper frames), and the next to leading order (lower frames), varying μ_F in the range $[0.5, 2] \times \mu_0$, where μ_0 is the nom-

inal scale defined in the text, calculated with the MMHT14 PDFs for the low- Q^2 : $5.5 \text{ GeV}^2 < Q^2 < 80 \text{ GeV}^2$ at HERA. The corresponding μ_F -dependence is also shown in terms of $\mathcal{R}[\text{TEEC}(\cos\phi)]_{\mu_F}$ and $\mathcal{R}[\text{ATEEC}(\cos\phi)]_{\mu_F}$, defined in Eq. (16)

logical study relevant for the jet production at HERA are carried out.

3 Results for TEEC ($\cos\phi$) and its asymmetry ATEEC ($\cos\phi$) in DIS process at HERA

For the numerical results presented here in the LO and NLO accuracy, we have used the program NLOJET++ [44,45]. As a cross check on our calculations, we have also used the program Madgraph to calculate the leading order TEEC and ATEEC functions. The errors shown for the TEEC and ATEEC are of statistical origin, resulting from the Monte Carlo phase space integration. To compare with the results obtained using NLOJET++, parton-level events are gener-

ated in MadGraph5_aMC@NLO [46] with the MMHT14 PDF set. The distributions obtained from the two packages agree well in both the low- Q^2 ($5.5 \text{ GeV}^2 < Q^2 < 80 \text{ GeV}^2$) and high- Q^2 ($150 \text{ GeV}^2 < Q^2 < 1000 \text{ GeV}^2$) ranges. The details are given in Appendix A. From now on, we shall work only with the NLOJET++.

We have generated 10^9 events to obtain the LO and NLO results in each of the two Q^2 ranges. This large statistics is required to obtain an accuracy of a few percent in NLO, which enables us to meaningfully calculate the various parametric dependences intrinsic to the problem at hand. At the very outset, we have calculated the two-jet cross sections at $\sqrt{s} = 314 \text{ GeV}$ for the ranges of the DIS variables given in the preceding section and compared them with the corresponding HERA data [49] in Table 1. The NLOJET++ results are obtained




Fig. 5 Factorization scale dependence of the differential distribution $1/\sigma' d\Sigma'/d(\cos\phi)$ and its asymmetry $1/\sigma' d\Sigma'^{asym}/d(\cos\phi)$ in the leading order (upper frames) and the next to leading order (lower frames) as in Fig. 4, but for the high- Q^2 range $150 \text{ GeV}^2 < Q^2 < 1000 \text{ GeV}^2$ at HERA

from the jet clusters using parton-level cross sections and the HERA data refer to the hadron-level cross section. While the two are not identical, this comparison should hold to a good first approximation. The two-jet events selected for this comparison are defined by the following two bins in $\langle E_T \rangle_2$ and the Q^2 -range given below:

$$\begin{aligned} 5.5 \text{ GeV}^2 < Q^2 < 8 \text{ GeV}^2, \\ \text{bin1} : 5 \text{ GeV} < \langle E_T \rangle_2 < 7 \text{ GeV}, \\ \text{bin2} : 7 \text{ GeV} < \langle E_T \rangle_2 < 11 \text{ GeV}. \end{aligned} \quad (14)$$

Theoretical cross sections are obtained using the CT18 [47] PDFs, the scales set to the values $\mu_R = \mu_F = \sqrt{\langle E_T \rangle^2 + Q^2}$, and $\alpha_s(M_Z) = 0.118$. The NLO cross sections are in excellent agreement with the HERA data.

We start by showing the differential distributions $\frac{1}{\sigma'} \frac{d\Sigma'}{d\cos\phi}$, defining TEEC ($\cos\phi$), and its asymmetry, $\frac{1}{\sigma'} \frac{d\Sigma'^{asym}}{d\cos\phi}$, ATEEC ($\cos\phi$), for the two PDF sets CT18 [47] and

MMHT14 [48]. They are presented for the low- Q^2 range ($5.5 \text{ GeV}^2 \leq Q^2 \leq 80 \text{ GeV}^2$) and the high- Q^2 range ($50 \text{ GeV}^2 \leq Q^2 \leq 1000 \text{ GeV}^2$) in Figs. 2 and 3, respectively. The left frame in these figure shows TEEC ($\cos\phi$) and the right frame ATEEC ($\cos\phi$), calculated in the NLO accuracy.

We restrict $\cos\phi$ in the range $[-0.8, 0.8]$ to avoid the regions $\phi \simeq 0^\circ$ and $\phi \simeq 180^\circ$ which will involve self-correlations ($a = b$) and virtual corrections to $2 \rightarrow 2$ processes. In calculating these functions, we use $\alpha_s(M_Z) = 0.118$ and have set the factorization (μ_F) and the renormalization (μ_R) scales to the following values: $\mu_F = \mu_R = \mu_0 = \sqrt{\langle E_T \rangle^2 + Q^2}$. This scale-setting is discussed in the analysis of the jet-data by the H1 Collaboration [50]. The effect of varying the scale μ_F which enters in the PDFs has little effect in the inclusive- and dijet- cross sections [50], which we also find for the TEEC ($\cos\phi$) and ATEEC($\cos\phi$), shown later in this section. We quantify the uncertainty on the TEEC ($\cos\phi$) and ATEEC ($\cos\phi$) from the two input




Fig. 6 Renormalization scale dependence of the differential distribution $1/\sigma' d\Sigma'/d(\cos\phi)$ and its asymmetry $1/\sigma' d\Sigma'^{asym}/d(\cos\phi)$ in the leading order (upper frames) and the next to leading order (lower frames) varying μ_R in the range $[0.5, 2] \times \mu_0$, where μ_0 is the nom-

inal scale defined in the text, calculated with the MMHT14 PDFs for the low- Q^2 : $5.5 \text{ GeV}^2 < Q^2 < 80 \text{ GeV}^2$ at HERA. The corresponding μ_R -dependence is also shown in terms of $\mathcal{R}[\text{TEEC}(\cos\phi)]_{\mu_R}$ and $\mathcal{R}[\text{ATEEC}(\cos\phi)]_{\mu_R}$, defined in Eq. (17)

PDFs by the following ratios:

$$\begin{aligned} & \Delta[\text{TEEC}(\cos\phi)]_{\text{pdf}} \\ & \equiv \frac{\text{TEEC}(\cos\phi)_{\text{CT18}} - \text{TEEC}(\cos\phi)_{\text{MMHT14}}}{\text{TEEC}(\cos\phi)_{\text{CT18}}} \\ & \Delta[\text{ATEEC}(\cos\phi)]_{\text{pdf}} \\ & \equiv \frac{\text{ATEEC}(\cos\phi)_{\text{CT18}} - \text{ATEEC}(\cos\phi)_{\text{MMHT14}}}{\text{ATEEC}(\cos\phi)_{\text{CT18}}}. \end{aligned} \quad (15)$$

The PDF-related uncertainties $\Delta[\text{TEEC}(\cos\phi)]_{\text{pdf}}$ and $\Delta[\text{ATEEC}(\cos\phi)]_{\text{pdf}}$ are shown in the lower frames in these figures. We note that, within our statistics, the former are well below 5% for most of the range. The asymmetry becomes increasingly small as one approaches $\cos\phi \simeq 0$, and it would require much higher statistics to reduce the numerical error on $\Delta[\text{ATEEC}(\cos\phi)]_{\text{pdf}}$ near the end-point. It should

be stressed that when showing the ratios in the lower frames, the statistical errors have been neglected.

Next, we present the factorization-scale and the renormalization-scale dependence of the TEEC ($\cos\phi$) and ATEEC ($\cos\phi$), by fixing the other parameters to their nominal values, and use the MMHT14 PDF set. Fixing $\mu_R = \mu_0$, we vary μ_F in the range $\mu_F = [0.5, 2]\mu_0$ and show the μ_F -dependence in Figs. 4 and 5 for the low- Q^2 range $5.5 \text{ GeV}^2 < Q^2 < 80 \text{ GeV}^2$ and the high- Q^2 range $150 \text{ GeV}^2 < Q^2 < 1000 \text{ GeV}^2$, respectively, in the LO and the NLO accuracy. The μ_F -uncertainty on TEEC ($\cos\phi$) and ATEEC ($\cos\phi$) are plotted in the lower frames of these Figs. 4 and 5 in terms of the ratios $\mathcal{R}[\text{TEEC}(\cos\phi)]_{\mu_F}$ and $\mathcal{R}[\text{ATEEC}(\cos\phi)]_{\mu_F}$ defined




Fig. 7 Renormalization scale dependence of the differential distribution $1/\sigma' d\Sigma'/d(\cos\phi)$ and its asymmetry $1/\sigma' d\Sigma'^{asym}/d(\cos\phi)$ in the leading order (upper frames) and the next to leading order (lower frames) as in Fig. 6, but for the high- Q^2 range: $150 \text{ GeV}^2 < Q^2 < 1000 \text{ GeV}^2$ at HERA

below

$$\begin{aligned} \mathcal{R}[\text{TEEC}(\cos\phi)]_{\mu_F} &\equiv \frac{\text{TEEC}(\cos\phi)_{\mu_F=x\mu_0, \mu_R=\mu_0}}{\text{TEEC}(\cos\phi)_{\mu_R=\mu_F=\mu_0}}, \\ x &\in [0.5, 2], \\ \mathcal{R}[\text{ATEEC}(\cos\phi)]_{\mu_F} &\equiv \frac{\text{ATEEC}(\cos\phi)_{\mu_F=x\mu_0, \mu_R=\mu_0}}{\text{ATEEC}(\cos\phi)_{\mu_R=\mu_F=\mu_0}}, \\ x &\in [0.5, 2]. \end{aligned} \quad (16)$$

The μ_F -dependence shown for $\mathcal{R}[\text{TEEC}(\cos\phi)]_{\mu_F}$ and $\mathcal{R}[\text{ATEEC}(\cos\phi)]_{\mu_F}$ is the resulting envelope by varying the scale μ_F in the indicated range and the statistical errors arising from the numerical integration of the phase space. The statistical errors of $\text{TEEC}(\cos\phi)$ and $\text{ATEEC}(\cos\phi)$ are about 3%. The μ_F -dependence of $\text{TEEC}(\cos\phi)$ is small, decreasing for the high- Q^2 range. It is comparatively smaller

for the asymmetry $\text{ATEEC}(\cos\phi)$, except for the last bin, where the numerical integration has a large statistical error.

The μ_R -dependence in the corresponding Q^2 -ranges are shown in Figs. 6 and 7, respectively. Here, we fixed $\mu_F = \mu_0$, and varied μ_R in the range $\mu_R = [0.5, 2]\mu_0$. One notices marked improvement in the μ_R -dependence from the LO to NLO. The μ_R -uncertainty on $\text{TEEC}(\cos\phi)$ and $\text{ATEEC}(\cos\phi)$ are plotted in the lower frames of Figs. 6 and 7 in terms of the ratios $\mathcal{R}[\text{TEEC}(\cos\phi)]_{\mu_R}$, and $\mathcal{R}[\text{ATEEC}(\cos\phi)]_{\mu_R}$, defined in Eq. (17).

$$\begin{aligned} \mathcal{R}[\text{TEEC}(\cos\phi)]_{\mu_R} &\equiv \frac{\text{TEEC}(\cos\phi)_{\mu_R=x\mu_0, \mu_F=\mu_0}}{\text{TEEC}(\cos\phi)_{\mu_R=\mu_F=\mu_0}}, \\ x &\in [0.5, 2], \\ \mathcal{R}[\text{ATEEC}(\cos\phi)]_{\mu_R} &\equiv \frac{\text{ATEEC}(\cos\phi)_{\mu_R=x\mu_0, \mu_F=\mu_0}}{\text{ATEEC}(\cos\phi)_{\mu_R=\mu_F=\mu_0}}, \end{aligned}$$




Fig. 8 Upper frames: Dependence of the differential distribution $1/\sigma' d\Sigma'/d(\cos\phi)$ (left) and its asymmetry $1/\sigma' d\Sigma'^{asym}/d(\cos\phi)$ (right) on the QCD coupling constant $\alpha_s(M_Z^2)$ for three indicated values of $\alpha_s(M_Z^2) = 0.108, 0.118, 0.128$ using the PDFs

of MMHT14 in the low- Q^2 range at HERA setting the scales $\mu_F = \mu_R = \mu_0$. Lower frames: The ratios $\mathcal{R}[\text{TEEC}(\cos\phi)]_{\alpha_s=0.108}$ and $\mathcal{R}[\text{TEEC}(\cos\phi)]_{\alpha_s=0.128}$ and $\mathcal{R}[\text{ATEEC}(\cos\phi)]_{\alpha_s=0.108}$ and $\mathcal{R}[\text{ATEEC}(\cos\phi)]_{\alpha_s=0.128}$, defined in Eq. (18)




Fig. 9 Dependence of the differential distribution $1/\sigma' d\Sigma'/d(\cos\phi)$ (left) and its asymmetry $1/\sigma' d\Sigma'^{asym}/d(\cos\phi)$ (right) on the QCD coupling constant $\alpha_s(M_Z^2)$ as in Fig. 8, but for the high- Q^2 range at HERA

$$x \in [0.5, 2]. \quad (17)$$

Based on these numerical results, we find that the combined uncertainty due to the PDFs, and the μ_F and μ_R -scales, is at about 10% in the TEEC ($\cos\phi$), and smaller in ATEEC ($\cos\phi$).

Further reduction in the scale uncertainty requires additional input, which we anticipate from the NNLO improvements as well as from the fits of the HERA data. This is suggested by the detailed NLO- and NNLO-studies done for the inclusive-jet and dijet data at HERA [50], which can be summarized as follows: The effect of varying μ_F in the range 10–90 GeV on the jet cross sections is small, and this scale

can be fixed to a value within this range without risking a perceptible change elsewhere, which is essentially in line what we find in our analysis. The effect of varying the scale μ_R is found more significant in the HERA jet-analysis. However, the choice $\mu_R = \sqrt{\langle E_T \rangle^2 + Q^2}$ yields a good fit of the jet data in both the NLO and NNLO accuracy. The reduced μ_R -dependence in the NNLO accuracy leads to a factor 2 improvement in the accuracy of $\alpha_s(M_Z^2)$. Following [50], we shall fix the scale μ_R to its nominal value in studying the sensitivity of TEEC ($\cos\phi$) and ATEEC ($\cos\phi$) on $\alpha_s(M_Z^2)$.

We now discuss the sensitivity of TEEC ($\cos\phi$) and ATEEC ($\cos\phi$) on $\alpha_s(M_Z^2)$. The results presented are




Fig. 10 A comparison of the LO and the NLO differential distribution $1/\sigma' d\Sigma'/d(\cos\phi)$ (left) and its asymmetry $1/\sigma' d\Sigma'^{asym}/d(\cos\phi)$ (right) at HERA ($\sqrt{s} = 314$ GeV) in the high- Q^2 range (upper frames) and low- Q^2 range (lower frames), with $\mu_F = \mu_R = \mu_0 = \sqrt{\langle E_T \rangle^2 + Q^2}$ and $\alpha_s(M_Z^2) = 0.118$

obtained by making the nominal choice of the scales $\mu_F = \mu_R = \mu_0$ and the MMHT14 PDFs. Results for three representative values $\alpha_s(M_Z^2) = 0.108, 0.118, 0.128$ are shown, which bracket most other determinations of this quantity, with $\alpha_s(M_Z^2) = 0.118$ being the central value¹ quoted by the Particle Data Group [51]. They are shown in Fig. 8 (low- Q^2 range) and Fig. 9 (high- Q^2 range) at the NLO accuracy. To quantify the $\alpha_s(M_Z^2)$ -sensitivity, we define the following ratios:

$$\mathcal{R}[\text{TEEC}(\cos\phi)]_{\alpha_s} \equiv \frac{\text{TEEC}(\cos\phi)_{\alpha_s(M_Z^2)}}{\text{TEEC}(\cos\phi)_{\alpha_s(M_Z^2)=0.118}}$$

$$\mathcal{R}[\text{ATEEC}(\cos\phi)]_{\alpha_s} \equiv \frac{\text{ATEEC}(\cos\phi)_{\alpha_s(M_Z^2)}}{\text{ATEEC}(\cos\phi)_{\alpha_s(M_Z^2)=0.118}}. \quad (18)$$

¹ The current PDG world average is $\alpha_s(M_Z^2) = 0.1179 \pm 0.0010$.

They are shown in the bottom frames in Fig. 8 (low- Q^2 range) and Fig. 9 (high- Q^2 range) in terms of the ratios $\mathcal{R}[\text{TEEC}(\cos\phi)]_{\alpha_s=0.128}$ and $\mathcal{R}[\text{TEEC}(\cos\phi)]_{\alpha_s=0.108}$. We note that both $\mathcal{R}[\text{TEEC}(\cos\phi)]_{\alpha_s}$ and the corresponding ratio for the asymmetry $\mathcal{R}[\text{ATEEC}(\cos\phi)]_{\alpha_s}$ show a marked sensitivity on $\alpha_s(M_Z^2)$. Hence, these shape functions at HERA offer competitive avenues to determine $\alpha_s(M_Z^2)$, and we urge our experimental colleagues to undertake a detailed data analysis of these variables at HERA.

A comparison of the LO and the NLO $\text{TEEC}(\cos\phi)$ and its asymmetry $\text{ATEEC}(\cos\phi)$ at HERA ($\sqrt{s} = 314$ GeV) in the high- Q^2 range and the low- Q^2 range are shown in Fig. 10. These results are obtained for the choice $\mu_F = \mu_R = \mu_0 = \sqrt{\langle E_T \rangle^2 + Q^2}$, $\alpha_s(M_Z^2) = 0.118$, and MMHT14 set of PDFs. They show that these correlations are remarkably stable against NLO corrections. We conjecture that NNLO corrections are, likewise, small. This remains to be shown and we hope that our work will stimulate working them out.

4 Summary

In this paper, we have studied for the first time, the transverse energy–energy correlations TEEC ($\cos \phi$) and its asymmetry ATEEC ($\cos \phi$) in deep inelastic scattering at the electron–proton collider HERA at the center of mass energy $\sqrt{s} = 314$ GeV, where ϕ is the angle in the Breit frame between two jets defined using a transverse-momentum (k_T) jet algorithm. We use NLOJET++ to calculate these functions in the LO and the NLO approximations in QCD for two ranges in the momentum transfer squared Q^2 . In the LO, these results are checked using the package MadGraph5_aMC@NLO [46] with the MMHT14 PDF set. We show the sensitivity of these functions on the PDFs, factorization (μ_F) and renormalization (μ_R) scales, and on $\alpha_s(M_Z^2)$. With the various cuts in the event generation matched with the ones in the measurements by the H1 collaboration at HERA, these studies are potentially useful in the analysis of the HERA data, including the determination of $\alpha_s(M_Z^2)$ from the shape variables.

An NNLO calculation for these shape variables is still lacking. This has the consequence that significant renormalization-scale dependence which enters in the partonic cross sections remains. At the present theoretical accuracy followed in this paper, this may compromise the precision on $\alpha_s(M_Z^2)$. Theoretical precision can be improved by including the NNLO contribution, as shown for the dijet and inclusive jet cross sections in DIS [56–59]. However, the scale uncertainty could also be reduced by analysing the HERA data for the shape variables by narrowing the allowed range of μ_R for which one gets a good quality fit. This is the case in the analysis of the inclusive-jet and dijet HERA data, in which the choice $\mu_R = \sqrt{\langle E_T \rangle^2 + Q^2}$ accounts well for the H1 measurements, also in the NLO accuracy [50]. For this choice of the μ_R scale, we have shown that the event shape TEEC ($\cos \phi$) and its asymmetry are very sensitive to the value of $\alpha_s(M_Z^2)$. We hope that our case-study for the TEEC and ATEEC at HERA, carried out at the NLO accuracy, will help focus on the analysis of the data on these shape variables with improved theoretical accuracy.

Acknowledgements We thank Fernando Barreiro, Xiao-Hui Liu and Stefan Schmitt for valuable discussions on the HERA measurements of jets. This work is supported in part by the Natural Science Foundation of China under grant No. 11735010, 11911530088, U2032102, by the Natural Science Foundation of Shanghai under grant No. 15DZ2272100, and by MOE Key Lab for Particle Physics, Astrophysics and Cosmology. GL is supported under U.S. Department of Energy contract DE-SC0011095.

Data Availability Statement This manuscript has no associated data or the data will not be deposited. [Authors' comment: Except the results and figures in the presented manuscript, we did not have any additional data.]

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Funded by SCOAP³.

Appendix-A

As a cross check on our calculations, we have also used the program Madgraph to calculate the leading order TEEC and ATEEC functions. To compare with the results obtained using NLOJET++, parton-level events are generated in MadGraph5_aMC@NLO [46] with the MMHT14 PDF set. To that end, the following basic cuts in the lab frame are imposed at the generator level in Madgraph:

$$p_{T,j}^{\text{lab}} > 2\text{GeV}, \quad |\eta_{j,e}^{\text{lab}}| < 5, \quad \Delta R_{jj}^{\text{lab}} > 0.1. \quad (19)$$

In the above, j denotes light-flavor quarks, and the angular distance in the $\eta - \phi$ plane is defined as $\Delta R_{ij} \equiv \sqrt{(\eta_i - \eta_j)^2 + (\phi_i - \phi_j)^2}$ with η_i and ϕ_i being the pseudorapidity and azimuthal angle of particle i , respectively. The momenta of the generated events are defined in the lab frame. After the appropriate Lorentz transformation, the TEEC and ATEEC distributions in the Breit frame can be constructed, and the events are selected in the low and high Q^2 ranges. Given the available choices of the factorization and renormalization scales in Madgraph, we set the scales $\mu_F = \mu_R = E_T$ with E_T being the scalar sum of transverse energies of all jets in both Madgraph and NLOJET++. The transverse energy of each jet in the Breit frame is limited in the range [4.5 GeV, 50 GeV] [49] to reduce the impact of the basic cuts in Eq. (19), apart from the cuts on the transverse energies for dijet and trijet events in Eq. (7). In Fig. 11, a comparison of the LO TEEC ($\cos \phi$) distributions obtained using NLOJET++ and Madgraph is shown, using $\alpha_s(M_Z^2) = 0.118$. The distributions obtained from the two packages agree well in both the low- Q^2 and high- Q^2 ranges. In Fig. 11 the error bars indicate the statistical errors, which are negligible for the distributions obtained from NLOJET++.




Fig. 11 Comparison of the LO differential distribution $1/\sigma' d\Sigma'/d(\cos \phi)$ obtained using Madgraph and NLOJET++ in the low- Q^2 range (left frame) and the high- Q^2 range (right frame)

References

- C.L. Basham, L.S. Brown, S.D. Ellis, S.T. Love, Phys. Rev. Lett. **41**, 1585 (1978). <https://doi.org/10.1103/PhysRevLett.41.1585>
- C.L. Basham, L.S. Brown, S.D. Ellis, S.T. Love, Phys. Rev. D **19**, 2018 (1979). <https://doi.org/10.1103/PhysRevD.19.2018>
- A. Ali, F. Barreiro, Phys. Lett. B **118**, 155 (1982). [https://doi.org/10.1016/0370-2693\(82\)90621-9](https://doi.org/10.1016/0370-2693(82)90621-9)
- A. Ali, F. Barreiro, Nucl. Phys. B **236**, 269 (1984). [https://doi.org/10.1016/0550-3213\(84\)90536-4](https://doi.org/10.1016/0550-3213(84)90536-4)
- D.G. Richards, W.J. Stirling, S.D. Ellis, Phys. Lett. B **119**, 193 (1982). [https://doi.org/10.1016/0370-2693\(82\)90275-1](https://doi.org/10.1016/0370-2693(82)90275-1)
- D.G. Richards, W.J. Stirling, S.D. Ellis, Nucl. Phys. B **229**, 317 (1983). [https://doi.org/10.1016/0550-3213\(83\)90335-8](https://doi.org/10.1016/0550-3213(83)90335-8)
- E.W.N. Glover, M.R. Sutton, Phys. Lett. B **342**, 375 (1995). [https://doi.org/10.1016/0370-2693\(94\)01354-F](https://doi.org/10.1016/0370-2693(94)01354-F). arXiv:hep-ph/9410234
- H.N. Schneider, G. Kramer, G. Schierholz, Z. Phys. C **22**, 201 (1984). <https://doi.org/10.1007/BF01572173>
- N.K. Falck, G. Kramer, Z. Phys. C **42**, 459 (1989). <https://doi.org/10.1007/BF01548452>
- G. Kramer, H. Spiesberger, Z. Phys. C **73**, 495 (1997). <https://doi.org/10.1007/s002880050339>. arXiv:hep-ph/9603385
- S. Catani, M.H. Seymour, Phys. Lett. B **378**, 287 (1996). [https://doi.org/10.1016/0370-2693\(96\)00425-X](https://doi.org/10.1016/0370-2693(96)00425-X). arXiv:hep-ph/9602277
- S. Catani, M.H. Seymour, Nucl. Phys. B **485**, 291 (1997). [https://doi.org/10.1016/S0550-3213\(96\)00589-5](https://doi.org/10.1016/S0550-3213(96)00589-5). arXiv:hep-ph/9605323, Erratum: [Nucl. Phys. B **510**, 503 (1998). [https://doi.org/10.1016/S0550-3213\(98\)81022-5](https://doi.org/10.1016/S0550-3213(98)81022-5)]
- V. Del Duca, C. Duhr, A. Kardos, G. Somogyi, Z. Tršćanski, Phys. Rev. Lett. **117**, 15, 152004 (2016). <https://doi.org/10.1103/PhysRevLett.117.152004>. arXiv:1603.08927 [hep-ph]
- Z. Tulipšnt, A. Kardos, G. Somogyi, Eur. Phys. J. C **77**(11), 749 (2017). <https://doi.org/10.1140/epjc/s10052-017-5320-9>. arXiv:1708.04093 [hep-ph]
- D. de Florian, M. Grazzini, Nucl. Phys. B **704**, 387 (2005). <https://doi.org/10.1016/j.nuclphysb.2004.10.051>. arXiv:hep-ph/0407241
- L.J. Dixon, M.X. Luo, V. Shtabovenko, T.Z. Yang, H.X. Zhu, Phys. Rev. Lett. **120**(10), 102001 (2018). <https://doi.org/10.1103/PhysRevLett.120.102001>. arXiv:1801.03219 [hep-ph]
- M.X. Luo, V. Shtabovenko, T.Z. Yang, H.X. Zhu, JHEP **1906**, 037 (2019). [https://doi.org/10.1007/JHEP06\(2019\)037](https://doi.org/10.1007/JHEP06(2019)037)
- I. Moul, H.X. Zhu, JHEP **1808**, 160 (2018). [https://doi.org/10.1007/JHEP08\(2018\)160](https://doi.org/10.1007/JHEP08(2018)160)
- L.J. Dixon, I. Moul, H.X. Zhu, Phys. Rev. D **100**(1), 014009 (2019). <https://doi.org/10.1103/PhysRevD.100.014009>. arXiv:1905.01310 [hep-ph]
- M. Kologlu, P. Kravchuk, D. Simmons-Duffin, A. Zhiboedov, arXiv:1905.01311 [hep-th]
- G.P. Korchemsky, JHEP **2001**, 008 (2020). [https://doi.org/10.1007/JHEP01\(2020\)008](https://doi.org/10.1007/JHEP01(2020)008)
- A.V. Belitsky, S. Hohenegger, G.P. Korchemsky, E. Sokatchev, A. Zhiboedov, Phys. Rev. Lett. **112**(7), 071601 (2014). <https://doi.org/10.1103/PhysRevLett.112.071601>. arXiv:1311.6800 [hep-th]
- J.M. Henn, E. Sokatchev, K. Yan, A. Zhiboedov, Phys. Rev. D **100**(3), 036010 (2019). <https://doi.org/10.1103/PhysRevD.100.036010>
- C. Berger et al., PLUTO Collaboration. Phys. Lett. **99B**, 292 (1981). [https://doi.org/10.1016/0370-2693\(81\)91128-X](https://doi.org/10.1016/0370-2693(81)91128-X)
- H.J. Behrend et al., CELLO Collaboration. Z. Phys. C **14**, 95 (1982). <https://doi.org/10.1007/BF01495029>
- B. Adeva et al., L3 Collaboration. Phys. Lett. B **257**, 469 (1991). [https://doi.org/10.1016/0370-2693\(91\)91925-L](https://doi.org/10.1016/0370-2693(91)91925-L)
- P. Abreu et al., DELPHI Collaboration. Phys. Lett. B **252**, 149 (1990). [https://doi.org/10.1016/0370-2693\(90\)91097-U](https://doi.org/10.1016/0370-2693(90)91097-U)
- K. Abe, et al. [SLD Collaboration], Phys. Rev. D **50**, 5580 (1994). <https://doi.org/10.1103/PhysRevD.50.5580>. arXiv:hep-ex/9405006
- A. Ali, E. Pietarinen, W.J. Stirling, Phys. Lett. B **141**, 447 (1984). [https://doi.org/10.1016/0370-2693\(84\)90283-1](https://doi.org/10.1016/0370-2693(84)90283-1)
- A. Ali, F. Barreiro, J. Llorente, W. Wang, Phys. Rev. D **86**, 114017 (2012). <https://doi.org/10.1103/PhysRevD.86.114017>
- G. Aad, et al. [ATLAS Collaboration], Phys. Lett. B **750**, 427 (2015). <https://doi.org/10.1016/j.physletb.2015.09.050>
- M. Aaboud, et al. [ATLAS Collaboration], Eur. Phys. J. C **77**, (12), 872 (2017). <https://doi.org/10.1140/epjc/s10052-017-5442-0>
- A. Gao, H.T. Li, I. Moul, H.X. Zhu, Phys. Rev. Lett. **123**(6), 062001 (2019). <https://doi.org/10.1103/PhysRevLett.123.062001>
- M. Dasgupta, G.P. Salam, J. Phys. G **30**, R143 (2004). <https://doi.org/10.1088/0954-3899/30/5/R01>. arXiv:hep-ph/0312283

35. V. Antonelli, M. Dasgupta, G.P. Salam, JHEP **0002**, 001 (2000). <https://doi.org/10.1088/1126-6708/2000/02/001>. [arXiv:hep-ph/9912488](https://arxiv.org/abs/hep-ph/9912488)
36. M. Dasgupta, G.P. Salam, JHEP **0208**, 032 (2002). <https://doi.org/10.1088/1126-6708/2002/08/032>. [arXiv:hep-ph/0208073](https://arxiv.org/abs/hep-ph/0208073)
37. Z.B. Kang, X. Liu, S. Mantry, Phys. Rev. D **90**(1), 014041 (2014). <https://doi.org/10.1103/PhysRevD.90.014041>. [arXiv:1312.0301](https://arxiv.org/abs/1312.0301) [hep-ph]
38. D. Kang, C. Lee, I.W. Stewart, PoS DIS **2015**, 142 (2015). <https://doi.org/10.22323/1.247.0142>
39. T. Gehrmann, A. Huss, J. Mo, J. Niehues, Eur. Phys. J. C **79**(12), 1022 (2019). <https://doi.org/10.1140/epjc/s10052-019-7528-3>. [arXiv:1909.02760](https://arxiv.org/abs/1909.02760) [hep-ph]
40. A. Aktas, et al. [H1 Collaboration], Eur. Phys. J. C **46**, 343 (2006) <https://doi.org/10.1140/epjc/s2006-02493-x>. [arXiv:hep-ex/0512014](https://arxiv.org/abs/hep-ex/0512014)
41. S. Chekanov, et al. [ZEUS Collaboration], Eur. Phys. J. C **27**, 531 (2003) <https://doi.org/10.1140/epjc/s2003-01148-x>. [arXiv:hep-ex/0211040](https://arxiv.org/abs/hep-ex/0211040)
42. P. Newman, M. Wing, Rev. Mod. Phys. **86**, 1037 (2014). <https://doi.org/10.1103/RevModPhys.86.1037>
43. S. Catani, Y.L. Dokshitzer, M.H. Seymour, B.R. Webber, Nucl. Phys. B **406**, 187 (1993). [https://doi.org/10.1016/0550-3213\(93\)90166-M](https://doi.org/10.1016/0550-3213(93)90166-M)
44. Z. Nagy, Phys. Rev. Lett. **88**, 122003 (2002). <https://doi.org/10.1103/PhysRevLett.88.122003>. [arXiv:hep-ph/0110315](https://arxiv.org/abs/hep-ph/0110315)
45. Z. Nagy, Z. Trocsanyi, Phys. Rev. Lett. **87**, 082001 (2001). <https://doi.org/10.1103/PhysRevLett.87.082001>. [arXiv:hep-ph/0104315](https://arxiv.org/abs/hep-ph/0104315)
46. J. Alwall, JHEP **1407**, 079 (2014). [https://doi.org/10.1007/JHEP07\(2014\)079](https://doi.org/10.1007/JHEP07(2014)079)
47. T.J. Hou et al., [arXiv:1912.10053](https://arxiv.org/abs/1912.10053) [hep-ph]
48. L.A. Harland-Lang, A.D. Martin, P. Motylinski, R.S. Thorne, Eur. Phys. J. C **75**(5), 204 (2015). <https://doi.org/10.1140/epjc/s10052-015-3397-6>. [arXiv:1412.3989](https://arxiv.org/abs/1412.3989) [hep-ph]
49. V. Andreev, et al. [H1 Collaboration], Eur. Phys. J. C **77**, (4), 215 (2017) <https://doi.org/10.1140/epjc/s10052-017-4717-9>. [arXiv:1611.03421](https://arxiv.org/abs/1611.03421) [hep-ex]
50. V. Andreev et al. [H1 Collaboration], Eur. Phys. J. C **77**, (11), 791 (2017) <https://doi.org/10.1140/epjc/s10052-017-5314-7>. [arXiv:1709.07251](https://arxiv.org/abs/1709.07251) [hep-ex]
51. P. A. Zyla, et al. [Particle Data Group], Prog. Theor. Exp. Phys. **2020**, (8), 083C01 (2020). <https://doi.org/10.1093/ptep/ptaa104>
52. S.D. Ellis, D.E. Soper, Phys. Rev. D **48**, 3160 (1993). <https://doi.org/10.1103/PhysRevD.48.3160>. [arXiv:hep-ph/9305266](https://arxiv.org/abs/hep-ph/9305266)
53. H.T. Li, I. Vitev, Y.J. Zhu., [arXiv:2006.02437](https://arxiv.org/abs/2006.02437) [hep-ph]
54. X. Liu, F. Ringer, W. Vogelsang, F. Yuan, Phys. Rev. Lett. **122**(19), 192003 (2019). <https://doi.org/10.1103/PhysRevLett.122.192003>. [arXiv:1812.08077](https://arxiv.org/abs/1812.08077) [hep-ph]
55. X. Liu, F. Ringer, W. Vogelsang, F. Yuan, [arXiv:2007.12866](https://arxiv.org/abs/2007.12866) [hep-ph]
56. M. Klasen, G. Kramer, M. Michael, Phys. Rev. D **89**(7), 074032 (2014). <https://doi.org/10.1103/PhysRevD.89.074032>. [arXiv:1310.1724](https://arxiv.org/abs/1310.1724) [hep-ph]
57. T. Biekötter, M. Klasen, G. Kramer, Phys. Rev. D **92**(7), 074037 (2015). <https://doi.org/10.1103/PhysRevD.92.074037>. [arXiv:1508.07153](https://arxiv.org/abs/1508.07153) [hep-ph]
58. J. Currie, T. Gehrmann, J. Niehues, Phys. Rev. Lett. **117**(4), 042001 (2016). <https://doi.org/10.1103/PhysRevLett.117.042001>. [arXiv:1606.03991](https://arxiv.org/abs/1606.03991) [hep-ph]
59. J. Currie, T. Gehrmann, A. Huss, J. Niehues, JHEP **1707**, 018 (2017). [https://doi.org/10.1007/JHEP07\(2017\)018](https://doi.org/10.1007/JHEP07(2017)018). [arXiv:1703.05977](https://arxiv.org/abs/1703.05977) [hep-ph]