

Search for $B^0 \rightarrow \tau^\pm \ell^\mp$ ($\ell = e, \mu$) with a hadronic tagging method at Belle

H. Atmacan,⁷ A. J. Schwartz,⁷ K. Kinoshita,⁷ I. Adachi,^{19,15} K. Adamczyk,⁶⁴ H. Aihara,⁸⁸ S. Al Said,^{81,40} D. M. Asner,³ V. Aulchenko,^{4,67} T. Aushev,²⁰ R. Ayad,⁸¹ V. Babu,⁸ S. Bahinipati,²⁴ M. Bauer,³⁷ P. Behera,²⁷ K. Belous,³¹ J. Bennett,⁵⁵ F. Bernlochner,² M. Bessner,¹⁸ V. Bhardwaj,²³ B. Bhuyan,²⁵ T. Bilka,⁵ J. Biswal,³⁶ A. Bobrov,^{4,67} A. Bozek,⁶⁴ M. Bračko,^{52,36} P. Branchini,³³ T. E. Browder,¹⁸ A. Budano,³³ M. Campajola,^{32,59} L. Cao,² D. Červenkov,⁵ M.-C. Chang,¹⁰ P. Chang,⁶³ V. Chekelian,⁵³ A. Chen,⁶¹ B. G. Cheon,¹⁷ K. Chilikin,⁴⁷ H. E. Cho,¹⁷ K. Cho,⁴² S.-K. Choi,¹⁶ Y. Choi,⁷⁹ S. Choudhury,²⁶ D. Cinabro,⁹² S. Cunliffe,⁸ S. Das,⁵¹ N. Dash,²⁷ G. De Nardo,^{32,59} G. De Pietro,³³ R. Dhamija,²⁶ F. Di Capua,^{32,59} J. Dingfelder,² Z. Doležal,⁵ T. V. Dong,¹¹ S. Dubey,¹⁸ D. Epifanov,^{4,67} T. Ferber,⁸ D. Ferlewicz,⁵⁴ A. Frey,¹⁴ B. G. Fulsom,⁶⁹ R. Garg,⁷⁰ V. Gaur,⁹¹ N. Gabyshev,^{4,67} A. Garmash,^{4,67} A. Giri,²⁶ P. Goldenzweig,³⁷ E. Graziani,³³ D. Greenwald,⁸³ T. Gu,⁷² Y. Guan,⁷ K. Gudkova,^{4,67} C. Hadjivasiliou,⁶⁹ S. Halder,⁸² T. Hara,^{19,15} O. Hartbrich,¹⁸ K. Hayasaka,⁶⁶ H. Hayashii,⁶⁰ W.-S. Hou,⁶³ C.-L. Hsu,⁸⁰ K. Inami,⁵⁸ G. Inguglia,³⁰ A. Ishikawa,^{19,15} R. Itoh,^{19,15} M. Iwasaki,⁶⁸ Y. Iwasaki,¹⁹ W. W. Jacobs,²⁸ S. Jia,¹¹ Y. Jin,⁸⁸ K. K. Joo,⁶ A. B. Kaliyar,⁸² K. H. Kang,⁴⁵ Y. Kato,⁵⁸ T. Kawasaki,⁴¹ C. Kiesling,⁵³ C. H. Kim,¹⁷ D. Y. Kim,⁷⁷ K.-H. Kim,⁹⁴ S. H. Kim,⁷⁶ Y.-K. Kim,⁹⁴ P. Kodyš,⁵ T. Konno,⁴¹ A. Korobov,^{4,67} S. Korpar,^{52,36} E. Kovalenko,^{4,67} P. Križan,^{48,36} R. Kroeger,⁵⁵ P. Krokovny,^{4,67} T. Kuhr,⁴⁹ R. Kulasiri,³⁹ K. Kumara,⁹² Y.-J. Kwon,⁹⁴ Y.-T. Lai,³⁸ J. S. Lange,¹² M. Laurenza,^{33,74} S. C. Lee,⁴⁵ J. Li,⁴⁵ L. K. Li,⁷ Y. B. Li,⁷¹ L. Li Gioi,⁵³ J. Libby,²⁷ K. Lieret,⁴⁹ D. Liventsev,^{92,19} C. MacQueen,⁵⁴ M. Masuda,^{87,73} T. Matsuda,⁵⁶ D. Matvienko,^{4,67,47} M. Merola,^{32,59} F. Metzner,³⁷ K. Miyabayashi,⁶⁰ R. Mizuk,^{47,20} G. B. Mohanty,⁸² R. Mussa,³⁴ M. Nakao,^{19,15} Z. Natkaniec,⁶⁴ A. Natchii,¹⁸ L. Nayak,²⁶ M. Nayak,⁸⁴ M. Niiyama,⁴⁴ N. K. Nisar,³ S. Nishida,^{19,15} S. Ogawa,⁸⁵ H. Ono,^{65,66} Y. Onuki,⁸⁸ P. Oskoin,⁴⁷ P. Pakhlov,^{47,57} G. Pakhlova,^{20,47} S. Pardi,³² H. Park,⁴⁵ S.-H. Park,¹⁹ A. Passeri,³³ S. Paul,^{83,53} T. K. Pedlar,⁵⁰ R. Pestotnik,³⁶ L. E. Piilonen,⁹¹ T. Podobnik,^{48,36} E. Prencipe,²¹ M. T. Prim,² I. Ripp-Baudot,⁷⁸ A. Rostomyan,⁹ N. Rout,²⁷ G. Russo,⁵⁹ D. Sahoo,⁸² S. Sandilya,²⁶ A. Sangal,⁷ L. Santelj,^{48,36} T. Sanuki,⁸⁶ V. Savinov,⁷² G. Schnell,^{1,22} J. Schueler,¹⁸ C. Schwanda,³⁰ Y. Seino,⁶⁶ K. Senyo,⁹³ M. E. Sevier,⁵⁴ M. Shapkin,³¹ C. Sharma,⁵¹ C. P. Shen,¹¹ J.-G. Shiu,⁶³ F. Simon,⁵³ A. Sokolov,³¹ E. Solovieva,⁴⁷ M. Starič,³⁶ Z. S. Stottler,⁹¹ M. Sumihama,¹³ K. Sumisawa,^{19,15} T. Sumiyoshi,⁹⁰ W. Sutcliffe,² U. Tamponi,³⁴ K. Tanida,³⁵ Y. Tao,⁹ F. Tenchini,⁸ K. Trabelsi,⁴⁶ M. Uchida,⁸⁹ T. Uglov,^{47,20} Y. Unno,¹⁷ K. Uno,⁶⁶ S. Uno,^{19,15} P. Urquijo,⁵⁴ S. E. Vahsen,¹⁸ R. Van Tonder,² G. Varner,¹⁸ K. E. Varvell,⁸⁰ A. Vinokurova,^{4,67} E. Waheed,¹⁹ C. H. Wang,⁶² E. Wang,⁷² M.-Z. Wang,⁶³ P. Wang,²⁹ X. L. Wang,¹¹ M. Watanabe,⁶⁶ S. Watanuki,⁴⁶ J. Wiechczynski,⁶⁴ E. Won,⁴³ B. D. Yabsley,⁸⁰ W. Yan,⁷⁵ S. B. Yang,⁴³ H. Ye,⁸ J. Yelton,⁹ J. H. Yin,⁴³ Z. P. Zhang,⁷⁵ V. Zhilich,^{4,67} and V. Zhukova⁴⁷

(The Belle Collaboration)

¹*Department of Physics, University of the Basque Country UPV/EHU, 48080 Bilbao*²*University of Bonn, 53115 Bonn*³*Brookhaven National Laboratory, Upton, New York 11973*⁴*Budker Institute of Nuclear Physics SB RAS, Novosibirsk 630090*⁵*Faculty of Mathematics and Physics, Charles University, 121 16 Prague*⁶*Chonnam National University, Gwangju 61186*⁷*University of Cincinnati, Cincinnati, Ohio 45221*⁸*Deutsches Elektronen-Synchrotron, 22607 Hamburg*⁹*University of Florida, Gainesville, Florida 32611*¹⁰*Department of Physics, Fu Jen Catholic University, Taipei 24205*¹¹*Key Laboratory of Nuclear Physics and Ion-beam Application (MOE)**and Institute of Modern Physics, Fudan University, Shanghai 200443*¹²*Justus-Liebig-Universität Gießen, 35392 Gießen*¹³*Gifu University, Gifu 501-1193*¹⁴*II. Physikalisches Institut, Georg-August-Universität Göttingen, 37073 Göttingen*¹⁵*SOKENDAI (The Graduate University for Advanced Studies), Hayama 240-0193*¹⁶*Gyeongsang National University, Jinju 52828*¹⁷*Department of Physics and Institute of Natural Sciences, Hanyang University, Seoul 04763*¹⁸*University of Hawaii, Honolulu, Hawaii 96822*¹⁹*High Energy Accelerator Research Organization (KEK), Tsukuba 305-0801*²⁰*National Research University Higher School of Economics, Moscow 101000*²¹*Forschungszentrum Jülich, 52425 Jülich*²²*IKERBASQUE, Basque Foundation for Science, 48013 Bilbao*²³*Indian Institute of Science Education and Research Mohali, SAS Nagar, 140306*²⁴*Indian Institute of Technology Bhubaneswar, Satya Nagar 751007*

- ²⁵Indian Institute of Technology Guwahati, Assam 781039
- ²⁶Indian Institute of Technology Hyderabad, Telangana 502285
- ²⁷Indian Institute of Technology Madras, Chennai 600036
- ²⁸Indiana University, Bloomington, Indiana 47408
- ²⁹Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049
- ³⁰Institute of High Energy Physics, Vienna 1050
- ³¹Institute for High Energy Physics, Protvino 142281
- ³²INFN - Sezione di Napoli, I-80126 Napoli
- ³³INFN - Sezione di Roma Tre, I-00146 Roma
- ³⁴INFN - Sezione di Torino, I-10125 Torino
- ³⁵Advanced Science Research Center, Japan Atomic Energy Agency, Naka 319-1195
- ³⁶J. Stefan Institute, 1000 Ljubljana
- ³⁷Institut für Experimentelle Teilchenphysik, Karlsruher Institut für Technologie, 76131 Karlsruhe
- ³⁸Kavli Institute for the Physics and Mathematics of the Universe (WPI), University of Tokyo, Kashiwa 277-8583
- ³⁹Kennesaw State University, Kennesaw, Georgia 30144
- ⁴⁰Department of Physics, Faculty of Science, King Abdulaziz University, Jeddah 21589
- ⁴¹Kitasato University, Sagamihara 252-0373
- ⁴²Korea Institute of Science and Technology Information, Daejeon 34141
- ⁴³Korea University, Seoul 02841
- ⁴⁴Kyoto Sangyo University, Kyoto 603-8555
- ⁴⁵Kyungpook National University, Daegu 41566
- ⁴⁶Université Paris-Saclay, CNRS/IN2P3, IJCLab, 91405 Orsay
- ⁴⁷P.N. Lebedev Physical Institute of the Russian Academy of Sciences, Moscow 119991
- ⁴⁸Faculty of Mathematics and Physics, University of Ljubljana, 1000 Ljubljana
- ⁴⁹Ludwig Maximilians University, 80539 Munich
- ⁵⁰Luther College, Decorah, Iowa 52101
- ⁵¹Malaviya National Institute of Technology Jaipur, Jaipur 302017
- ⁵²Faculty of Chemistry and Chemical Engineering, University of Maribor, 2000 Maribor
- ⁵³Max-Planck-Institut für Physik, 80805 München
- ⁵⁴School of Physics, University of Melbourne, Victoria 3010
- ⁵⁵University of Mississippi, University, Mississippi 38677
- ⁵⁶University of Miyazaki, Miyazaki 889-2192
- ⁵⁷Moscow Physical Engineering Institute, Moscow 115409
- ⁵⁸Graduate School of Science, Nagoya University, Nagoya 464-8602
- ⁵⁹Università di Napoli Federico II, I-80126 Napoli
- ⁶⁰Nara Women's University, Nara 630-8506
- ⁶¹National Central University, Chung-li 32054
- ⁶²National United University, Miao Li 36003
- ⁶³Department of Physics, National Taiwan University, Taipei 10617
- ⁶⁴H. Niewodniczanski Institute of Nuclear Physics, Krakow 31-342
- ⁶⁵Nippon Dental University, Niigata 951-8580
- ⁶⁶Niigata University, Niigata 950-2181
- ⁶⁷Novosibirsk State University, Novosibirsk 630090
- ⁶⁸Osaka City University, Osaka 558-8585
- ⁶⁹Pacific Northwest National Laboratory, Richland, Washington 99352
- ⁷⁰Panjab University, Chandigarh 160014
- ⁷¹Peking University, Beijing 100871
- ⁷²University of Pittsburgh, Pittsburgh, Pennsylvania 15260
- ⁷³Research Center for Nuclear Physics, Osaka University, Osaka 567-0047
- ⁷⁴Dipartimento di Matematica e Fisica, Università di Roma Tre, I-00146 Roma
- ⁷⁵Department of Modern Physics and State Key Laboratory of Particle Detection and Electronics, University of Science and Technology of China, Hefei 230026
- ⁷⁶Seoul National University, Seoul 08826
- ⁷⁷Soongsil University, Seoul 06978
- ⁷⁸Université de Strasbourg, CNRS, IPHC, UMR 7178, 67037 Strasbourg
- ⁷⁹Sungkyunkwan University, Suwon 16419
- ⁸⁰School of Physics, University of Sydney, New South Wales 2006
- ⁸¹Department of Physics, Faculty of Science, University of Tabuk, Tabuk 71451
- ⁸²Tata Institute of Fundamental Research, Mumbai 400005
- ⁸³Department of Physics, Technische Universität München, 85748 Garching
- ⁸⁴School of Physics and Astronomy, Tel Aviv University, Tel Aviv 69978
- ⁸⁵Toho University, Funabashi 274-8510
- ⁸⁶Department of Physics, Tohoku University, Sendai 980-8578
- ⁸⁷Earthquake Research Institute, University of Tokyo, Tokyo 113-0032

⁸⁸Department of Physics, University of Tokyo, Tokyo 113-0033

⁸⁹Tokyo Institute of Technology, Tokyo 152-8550

⁹⁰Tokyo Metropolitan University, Tokyo 192-0397

⁹¹Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061

⁹²Wayne State University, Detroit, Michigan 48202

⁹³Yamagata University, Yamagata 990-8560

⁹⁴Yonsei University, Seoul 03722

We present a search for the lepton-flavor-violating decays $B^0 \rightarrow \tau^\pm \ell^\mp$, where $\ell = (e, \mu)$, using the full data sample of 772×10^6 $B\bar{B}$ pairs recorded by the Belle detector at the KEKB asymmetric-energy e^+e^- collider. We use events in which one B meson is fully reconstructed in a hadronic decay mode. The $\tau^\pm$ lepton is reconstructed indirectly using the momentum of the reconstructed B and that of the $\ell^\mp$ from the signal decay. We find no evidence for $B^0 \rightarrow \tau^\pm \ell^\mp$ decays and set upper limits on their branching fractions at 90% confidence level of $\mathcal{B}(B^0 \rightarrow \tau^\pm \mu^\mp) < 1.5 \times 10^{-5}$ and $\mathcal{B}(B^0 \rightarrow \tau^\pm e^\mp) < 1.6 \times 10^{-5}$.

PACS numbers: 13.25.Hw, 14.40.Nd

I. INTRODUCTION

The lepton-flavor-violating decays $B^0 \rightarrow \tau^\pm \ell^\mp$ [1], where $\ell = (e, \mu)$, are promising modes in which to search for new physics. Recently, there have been indications of possible violation of lepton flavor universality (LFU) in $B^0 \rightarrow D^{(*)-} \tau^+ \nu$ [2] and $B^\pm \rightarrow K^\pm \ell^+ \ell^-$ [3, 4] decays. LFU violation is often accompanied by lepton flavor violation (LFV) in theoretical models [5]. The decay $B^0 \rightarrow \tau^\pm \ell^\mp$, like $B^0 \rightarrow D^{(*)-} \tau^+ \nu$, connects a third-generation quark with a third-generation lepton. The decay can occur in principle via neutrino mixing [6]; however, the rate due to such mixing [7] is considerably below current or future experimental sensitivities. Thus, observing these decays would indicate new physics. Some new physics models give rise to branching fractions of 10^{-9} to 10^{-10} . For example, Pati-Salam vector leptoquarks of mass 86 TeV/ c^2 give branching fractions of 4.4×10^{-9} for $B^0 \rightarrow \tau^\pm \mu^\mp$ and 1.6×10^{-9} for $B^0 \rightarrow \tau^\pm e^\mp$ [8]. The general flavor-universal Minimal Supersymmetric Standard Model predicts branching fractions of up to about 2×10^{-10} [9].

These decay modes have previously been studied by the CLEO [10], BABAR [11], and LHCb [12] experiments. No evidence for these decays has been found. The current most stringent upper limits are $\mathcal{B}(B^0 \rightarrow \tau^\pm \mu^\mp) < 1.2 \times 10^{-5}$ [12] and $\mathcal{B}(B^0 \rightarrow \tau^\pm e^\mp) < 2.8 \times 10^{-5}$ [11], both at 90% confidence level (C.L.). In this Letter we report a search for $B^0 \rightarrow \tau^\pm \ell^\mp$ decays using the full Belle data sample recorded at the $\Upsilon(4S)$ resonance of 711 fb^{-1} . This is the first such search from Belle.

II. DATASET AND DETECTOR DESCRIPTION

Our data sample consists of $(772 \pm 11) \times 10^6$ $B\bar{B}$ pairs produced in $e^+e^- \rightarrow \Upsilon(4S)$ events recorded by the Belle detector at the KEKB asymmetric-energy e^+e^- collider [13]. The Belle detector is a large-solid-angle magnetic spectrometer that consists of a silicon vertex detector (SVD), a 50-layer central drift chamber

(CDC), an array of aerogel threshold Cherenkov counters (ACC), a barrel-like arrangement of time-of-flight scintillation counters (TOF), and an electromagnetic calorimeter comprising CsI(Tl) crystals (ECL). All these detectors are located inside a superconducting solenoid coil that provides a 1.5 T magnetic field. An iron flux return yoke located outside the coil is instrumented with resistive-plate chambers (KLM) to detect K_L^0 mesons and to identify muons. Two inner detector configurations were used: for the first 152×10^6 $B\bar{B}$ pairs, a 2.0 cm radius beam-pipe and a three-layer SVD were used; for the remaining 620×10^6 $B\bar{B}$ pairs [14], a 1.5 cm radius beam-pipe, a four-layer SVD, and a small-cell inner drift chamber were used. A more detailed description of the detector is provided in Ref. [15].

We study properties of signal events, sources of background, and optimize selection criteria using Monte Carlo (MC) simulated events. These samples are generated using the software packages EVTGEN [16] and PYTHIA [17], and final-state radiation is included via PHOTOS [18]. The detector response is simulated using GEANT3 [19]. We produce $B^0 \rightarrow \tau^\pm \ell^\mp$ MC events to calculate signal reconstruction efficiencies. To estimate backgrounds, we use MC samples that describe all $e^+e^- \rightarrow q\bar{q}$ processes. Events containing $e^+e^- \rightarrow B\bar{B}$ with subsequent $b \rightarrow c$ decay, and $e^+e^- \rightarrow q\bar{q}$ ($q = u, d, s, c$) continuum events, are both simulated with five times the integrated luminosity of Belle. Semileptonic $b \rightarrow u\ell\nu$ decays are simulated with 20 times the integrated luminosity. Rare $b \rightarrow s$ and $b \rightarrow u$ decays are simulated with 50 times the integrated luminosity.

III. EVENT SELECTION

Our analysis uses a technique uniquely suited to e^+e^- flavor factory experiments, in which the energy and momentum of the initial state are known. We first reconstruct a B meson decaying hadronically; this is referred to as the “tag-side” B meson (B_{tag}). We use the reconstructed B_{tag} momentum and the e^+e^- initial momen-

tum to infer the momentum of the signal-side B meson (B_{sig}). Because $B^0 \rightarrow \tau^\pm \ell^\mp$ are two-body decays, the momentum of the τ lepton can be inferred from the momentum of B_{sig} and the momentum of $\ell^\mp$; thus the $\tau^\pm$ does not need to be reconstructed. We define the “missing mass” as

$$M_{\text{miss}} = \sqrt{(E_{B_{\text{sig}}} - E_\ell)^2/c^4 - (\vec{p}_{B_{\text{sig}}} - \vec{p}_\ell)^2/c^2}, \quad (1)$$

where $E_{B_{\text{sig}}}$ and $\vec{p}_{B_{\text{sig}}}$ are the energy and momentum, respectively, of B_{sig} , and E_ℓ and $\vec{p}_\ell$ are the corresponding quantities for $\ell^\mp$. The quantity M_{miss} is the invariant mass of the unreconstructed or missing particle and, for $B^0 \rightarrow \tau^\pm \ell^\mp$ decays, should peak at the mass of the τ lepton ($m_\tau = 1.776 \text{ GeV}/c^2$ [20]). To improve the resolution in M_{miss} , we evaluate it in the e^+e^- center-of-mass (c.m.) frame and substitute the beam energy E_{beam} for $E_{B_{\text{sig}}}$. To avoid introducing bias in our analysis, we analyze the data in a “blind” manner, i.e., we finalize all selection criteria before viewing events in a region around m_τ . This blinded region is $[1.65, 1.90] \text{ GeV}/c^2$, which corresponds to approximately 3.8σ in the resolution.

A. Tag-side selection

We first reconstruct B_{tag} candidates in one of 1104 hadronic decay channels using a hierarchical algorithm based on the NeuroBayes neural network package [21]. The quality of B_{tag} is represented by a single classifier output, O_{NN} , which ranges from 0 (background-like) to 1 (signal-like). The output O_{NN} is mainly determined by the B_{tag} reconstruction. It includes event-shape information and significantly suppresses $e^+e^- \rightarrow q\bar{q}$ continuum events. In addition to O_{NN} , two other variables are used for selecting B_{tag} candidates: the energy difference $\Delta E \equiv E_{B_{\text{tag}}} - E_{\text{beam}}$, and the beam-energy-constrained mass $M_{\text{bc}} \equiv \sqrt{E_{\text{beam}}^2/c^4 - |\vec{p}_{B_{\text{tag}}}|^2/c^2}$, where $E_{B_{\text{tag}}}$ and $\vec{p}_{B_{\text{tag}}}$ are the reconstructed energy and momentum, respectively, of B_{tag} . These quantities are evaluated in the e^+e^- c.m. system. The B_{tag} candidate is required to satisfy $|\Delta E| < 0.05 \text{ GeV}$. For each signal mode, we choose selection criteria on O_{NN} and M_{bc} by optimizing a figure-of-merit (FOM). The FOM is defined as $\varepsilon_{\text{MC}}/\sqrt{N_{\text{B}}}$, where ε_{MC} is the reconstruction efficiency of signal events as determined from MC simulation, and N_{B} is the number of background events expected within the signal region $M_{\text{miss}} \in [1.65, 1.90] \text{ GeV}/c^2$. Based on FOM studies, we require $O_{\text{NN}} > 0.082$ for $B^0 \rightarrow \tau^\pm \mu^\mp$, $O_{\text{NN}} > 0.095$ for $B^0 \rightarrow \tau^\pm e^\mp$, and $M_{\text{bc}} > 5.272 \text{ GeV}/c^2$ for both modes.

After all B_{tag} selection criteria are applied, about 10% of $B^0 \rightarrow \tau^\pm \mu^\mp$ events and 8% of $B^0 \rightarrow \tau^\pm e^\mp$ events have multiple B_{tag} candidates. For such events, we select a single B_{tag} by choosing the candidate with the highest value of O_{NN} . This criterion selects the correct candidate 90% of the time, according to MC simulation.

B. Signal-side selection

To reconstruct the signal side, only tracks not associated with B_{tag} are considered. Such tracks are required to originate from the interaction point (IP) and have an impact parameter $|dz| < 4.0 \text{ cm}$ along the z axis, which points opposite the e^+ beam direction. We also require $dr < 2.0 \text{ cm}$ in the x - y plane (transverse to the e^+ beam direction), where $dr = \sqrt{dx^2 + dy^2}$.

Charged tracks are identified by combining information from various subdetectors into a likelihood function $\mathcal{L}_i$, where $i = e, \mu, \pi, K$, or p [22]. Muon candidates are identified based on the response of the CDC and KLM [23]. A track with a likelihood ratio $\mathcal{R}_\mu = \mathcal{L}_\mu/(\mathcal{L}_\mu + \mathcal{L}_\pi + \mathcal{L}_K) > 0.90$ is identified as a muon. The detection efficiency of this requirement is about 89%, and the pion misidentification rate is about 2%. Electron candidates are identified mainly using the ratio of the energy deposited in the ECL to the track momentum, the shower shape in the ECL, and the energy loss in the CDC. A track with a likelihood ratio $\mathcal{R}_e = \mathcal{L}_e/(\mathcal{L}_e + \mathcal{L}_{\text{hadrons}}) > 0.90$ is identified as an electron, where $\mathcal{L}_{\text{hadrons}}$ is the product of probability density functions (PDFs) for hadrons [24]. The efficiency of this requirement is about 94%, and the pion misidentification rate is about 0.3%. We recover electron energy lost due to bremsstrahlung by searching for photons within a cone of radius 50 mrad centered around the electron momentum. If such a photon is found, its four-momentum (assuming it originated at the IP) is added to that of the electron.

We require that M_{miss} be in the range $1.40 \text{ GeV}/c^2$ to $2.20 \text{ GeV}/c^2$. Every muon or electron candidate satisfying this requirement is treated as a B_{sig} candidate. After these selections, we find that less than 1% of $B^0 \rightarrow \tau^\pm \mu^\mp$ and $B^0 \rightarrow \tau^\pm e^\mp$ events have multiple B_{sig} candidates. These fractions are consistent with those from MC simulation. For such events, in order to preserve efficiency we retain all such candidates, i.e., we do not apply a best-candidate selection.

C. Background

After applying all selection criteria, a small amount of background remains. This background is studied using MC simulation and found to originate mainly from $b \rightarrow c$ and $b \rightarrow u\ell\nu$ decays. These backgrounds are smoothly falling in the M_{miss} distribution. However, for $B^0 \rightarrow \tau^\pm \mu^\mp$ candidates, two small peaks are observed: one at $M_{\text{miss}} \approx 1.869 \text{ GeV}/c^2$ and the other at $M_{\text{miss}} \approx 2.010 \text{ GeV}/c^2$. The former corresponds to $B^0 \rightarrow D^-\pi^+$ decays, while the latter corresponds to $B^0 \rightarrow D^{*-}\pi^+$ decays, where in both cases the π^+ is misidentified as μ^+ . These $B^0 \rightarrow D^{(*)-}\pi^+$ decays are taken into account when fitting the M_{miss} distribution for the signal yield (described below).

D. Control samples

We use control samples of $B^0 \rightarrow D^{(*)-}\pi^+$ decays to determine corrections to the shapes of the $B^0 \rightarrow \tau^\pm \ell^\mp$ PDFs used to fit for the signal yields (see Section IV). To identify $B^0 \rightarrow D^{(*)-}\pi^+$ decays, we select pions on the signal side rather than leptons. Pion candidates are identified using dE/dx measured in the CDC, time-of-flight information from the TOF, and the photon yield in the ACC. A track with a likelihood ratio $\mathcal{R}_\pi = \mathcal{L}_\pi/(\mathcal{L}_\pi + \mathcal{L}_K) > 0.90$ is identified as a pion [22]. All other selection criteria are the same as for the $B^0 \rightarrow \tau^\pm \ell^\mp$ search. In addition, we veto leptons by requiring that $\mathcal{R}_\mu < 0.90$ and $\mathcal{R}_e < 0.90$. With the above selection, the pion identification efficiency is about 95%, and the kaon misidentification rate is about 5%.

IV. MAXIMUM LIKELIHOOD FITS

We determine the $B^0 \rightarrow \tau^\pm \ell^\mp$ signal yields by performing an unbinned extended maximum-likelihood fit to the M_{miss} distributions. The PDF used to model correctly reconstructed signal decays are a double Gaussian for $B^0 \rightarrow \tau^\pm \mu^\mp$ and the sum of three Gaussians for $B^0 \rightarrow \tau^\pm e^\mp$. These Gaussians are allowed to have different means. We also model misreconstructed signal decays in which the lepton selected is subject to final-state radiation or is not a direct daughter in the two-body $B^0 \rightarrow \tau^\pm \ell^\mp$ decay, i.e., it originates from $\tau^\pm \rightarrow \ell^\pm \nu \bar{\nu}$ or $\tau^\pm \rightarrow \pi^\pm (\rightarrow \ell^\pm \nu) \bar{\nu}$. This component is referred to as “self-cross-feed” signal, and we model it with a double Gaussian and an exponential function. The fractions of self-cross-feed signal are fixed to the values obtained from MC simulation: $(5.0 \pm 0.2)\%$ for $B^0 \rightarrow \tau^\pm \mu^\mp$ and $(14.0 \pm 0.3)\%$ for $B^0 \rightarrow \tau^\pm e^\mp$. The self-cross-feed fraction is larger for the electron channel due to a larger contribution from $B^0 \rightarrow \tau^\pm e^\mp \gamma$ decays.

The shape parameters of the signal PDFs are obtained from MC simulation. We make corrections to these to account for small differences observed between MC simulation and data. We obtain these correction factors by fitting the M_{miss} distributions of the high-statistics $B^0 \rightarrow D^{(*)-}\pi^+$ control samples. For the $B^0 \rightarrow D^{(*)-}\pi^+$ samples, we fit both data and MC events and record small shifts observed in the means of the PDFs, and nominal differences in the widths. We apply these shifts for the means and scaling factors for the widths to the $B^0 \rightarrow \tau^\pm \ell^\mp$ signal PDFs. The uncertainties in these correction factors are accounted for when evaluating systematic uncertainties.

The PDFs for misidentified $B^0 \rightarrow D^-\pi^+$ and $B^0 \rightarrow D^{*-}\pi^+$ decays are taken to be a double Gaussian and sum of three Gaussians, respectively. Background PDFs of all modes are modeled with exponential functions. The shape parameters for these background PDFs are all floated, along with the background and signal yields.

We validate our fitting procedure and check for fit bias

using MC simulation. We generate large ensembles of simulated experiments, in which the M_{miss} distributions are generated from the PDFs used for fitting. We fit these ensembles and find that the fitted signal yields are consistent with the input values; the mean difference is -0.08 ± 0.05 events for $B^0 \rightarrow \tau^\pm \mu^\mp$ and -0.00 ± 0.05 events for $B^0 \rightarrow \tau^\pm e^\mp$. We include these small potential biases when evaluating systematic uncertainties. We also fit ensembles of fully simulated events and again find the signal yields to be consistent with the input values.

To further check our analysis procedure and acceptance calculation, we measure the branching fractions for the control channels $B^0 \rightarrow D^{(*)-}\pi^+$. The M_{miss} distributions of these decays along with projections of the fit result are shown in Fig. 1. To assess the goodness-of-fit, we calculate a χ^2 statistic from the residuals of the fit result. Dividing by the number of degrees of freedom (n_{dof}) gives $\chi^2/n_{\text{dof}} = 0.89$. The fitted yields are 2136 ± 71 and 2071 ± 74 , and the resulting branching fractions are $\mathcal{B}(B^0 \rightarrow D^-\pi^+) = (2.54 \pm 0.11) \times 10^{-3}$ and $\mathcal{B}(B^0 \rightarrow D^{*-}\pi^+) = (2.67 \pm 0.12) \times 10^{-3}$, where the uncertainties listed are statistical only. These values are in excellent agreement with the current world averages $\mathcal{B}(B^0 \rightarrow D^-\pi^+) = (2.52 \pm 0.13) \times 10^{-3}$ and $\mathcal{B}(B^0 \rightarrow D^{*-}\pi^+) = (2.74 \pm 0.12) \times 10^{-3}$ [20].




FIG. 1. The M_{miss} distribution of $B^0 \rightarrow D^{(*)-}\pi^+$ candidates observed in data (black dots) along with projections of the fit result: the overall fit result (solid blue curve), the background component (dashed red curve), the $B^0 \rightarrow D^-\pi^+$ component (dotted brown curve) and the $B^0 \rightarrow D^{*-}\pi^+$ component (dash-dotted magenta curve). The plot below the distribution shows the residuals divided by the errors (pulls).

The M_{miss} distributions for signal $B^0 \rightarrow \tau^\pm \ell^\mp$ decays along with projections of the fit result are shown in Fig. 2, and χ^2/n_{dof} values are 0.54 and 0.70 for $B^0 \rightarrow \tau^\pm \mu^\mp$ and $B^0 \rightarrow \tau^\pm e^\mp$, respectively. The fitted signal yields are $N_{\text{sig}} = 1.8^{+8.2}_{-7.6}$ for $B^0 \rightarrow \tau^\pm \mu^\mp$ and $N_{\text{sig}} = 0.3^{+8.8}_{-8.2}$ for $B^0 \rightarrow \tau^\pm e^\mp$. Both yields are consistent with zero.




FIG. 2. The M_{miss} distributions of $B^0 \rightarrow \tau^\pm \mu^\mp$ (upper) and $B^0 \rightarrow \tau^\pm e^\mp$ (lower) candidates, along with projections of the fit result. The black dots show the data, the dashed red curve shows the background component, and the solid blue curve shows the overall fit result. The dash-dotted green curve shows the signal PDF, with a normalization corresponding to a branching fraction of 10^{-4} . In the upper plot, the dotted brown curve shows the $B^0 \rightarrow D^- \pi^+$ component. The plots below the distributions show the residuals divided by the errors (pulls).

V. UPPER LIMIT CALCULATION

We calculate upper limits on N_{sig} and the branching fractions at 90% C.L. using a frequentist method. We first generate sets of MC-simulated events, with each set being equivalent to the Belle data sample of $(772 \pm 11) \times 10^6$ $B\bar{B}$ pairs. Both signal and background

events are generated according to their respective PDFs. The number of background events generated is equal to that obtained from the data fit. We vary the number of input signal events, and for each value we generate an ensemble of 10 000 data sets. We fit these data sets and calculate the fraction (f_{sig}) that has a fitted signal yield less than that obtained from the Belle data (1.8 or 0.3 events). Our 90% C.L. upper limit on the number of signal events ($N_{\text{sig}}^{\text{UL}}$) is the number of input signal events that has $f_{\text{sig}} = 0.10$. We convert $N_{\text{sig}}^{\text{UL}}$ to an upper limit on the branching fraction ($\mathcal{B}^{\text{UL}}$) via the formula

$$\mathcal{B}^{\text{UL}} = \frac{N_{\text{sig}}^{\text{UL}}}{2 \times N_{B\bar{B}} \times f^{00} \times \varepsilon}. \quad (2)$$

In this expression, $N_{B\bar{B}} = (772 \pm 11) \times 10^6$ is the number of $B\bar{B}$ pairs; $f^{00} = 0.486 \pm 0.006$ is the fraction that are $B^0 \bar{B}^0$ [20]; and ε is the signal efficiency including tag-side branching fractions and reconstruction efficiencies.

We include systematic uncertainties (discussed below) in $\mathcal{B}^{\text{UL}}$ as follows. We divide all systematic uncertainties into two types (see Table II): those arising from the numerator of Eq. 2 (“additive” uncertainties), and those arising from the denominator of Eq. 2 (“multiplicative” uncertainties). Additive uncertainties arise from fitting for the signal yield, while multiplicative uncertainties correspond to the number of B decays reconstructed. We account for the latter when generating MC data sets in our frequentist procedure. The number of signal events is varied randomly around the nominal input value by the total multiplicative uncertainty. Subsequently, after fitting an MC data set, we adjust the fitted value N_{sig} by a value sampled from a Gaussian distribution with mean zero and a width equal to the total additive uncertainty. As a final step, to include possible fit bias, this value is shifted by an amount obtained by sampling a Gaussian distribution with a mean equal to the fit bias discussed earlier (the central value) and a width equal to the uncertainty in the bias. This final value is used when calculating f_{sig} . The resulting upper limits for $N_{\text{sig}}^{\text{UL}}$ and $\mathcal{B}^{\text{UL}}$ are listed in Table I.

TABLE I. Summary of the fit results for N_{sig} , and the resulting 90% C.L. upper limits $N_{\text{sig}}^{\text{UL}}$ and $\mathcal{B}^{\text{UL}}$ (see text).

Mode	ε ($\times 10^{-4}$)	N_{sig} ($\times 10^{-4}$)	$N_{\text{sig}}^{\text{UL}}$ ($\times 10^{-4}$)	$\mathcal{B}^{\text{UL}}$ ($\times 10^{-5}$)
$B^0 \rightarrow \tau^\pm \mu^\mp$	11.0	$1.8^{+8.2}_{-7.6}$	12.4	1.5
$B^0 \rightarrow \tau^\pm e^\mp$	9.8	$0.3^{+8.8}_{-8.2}$	11.6	1.6

VI. SYSTEMATIC UNCERTAINTIES

The systematic uncertainties in our measurement – aside from potential fit bias, which is treated separately

when setting the upper limits – are listed in Table II. Uncertainties in the shapes of the PDFs used for the signal are evaluated by varying all fixed parameters by $\pm 1\sigma$; the resulting change in the signal yield is taken as the systematic uncertainty. The fixed parameters that are varied include the correction factors to the shapes as obtained from the $B^0 \rightarrow D^{(*)-}\pi^+$ control samples. The fraction of self-cross-feed signal is fixed to the MC value. We vary this fraction by $\pm 50\%$ and take the resulting change in the signal yield as the systematic uncertainty.

The reconstruction efficiency for B_{tag} is evaluated via MC simulation. However, there is uncertainty arising from branching fractions for tagging modes that are not well-measured, and from unknown decay dynamics of (multi-body) hadronic decays. To account for these effects, a correction factor to the reconstruction efficiency is applied. This correction is evaluated as done in Ref. [25], by comparing the number of events containing both a B_{tag} and a semileptonic $B \rightarrow D^{(*)}\ell\nu$ decay in data and MC. As the branching fractions for $B \rightarrow D^{(*)}\ell\nu$ are precisely known, and their reconstruction efficiencies can be separately calculated, the difference between data and MC for B_{tag} reconstruction can be extracted. The resulting correction factor is 0.64 ± 0.03 . The uncertainty in this value is taken as a systematic uncertainty.

The systematic uncertainty due to charged track reconstruction is evaluated using $D^{*+} \rightarrow D^0\pi^+$ decays, with $D^0 \rightarrow K_S^0\pi^+\pi^-$ and $K_S^0 \rightarrow \pi^+\pi^-$. The resulting uncertainty is 0.35% per track. The uncertainty due to lepton identification is evaluated using $e^+e^- \rightarrow e^+e^-\gamma^*\gamma^* \rightarrow e^+e^-\ell^+\ell^-$ events. The resulting uncertainties are 1.6% for muons and 1.8% for electrons.

The systematic uncertainty in the signal reconstruction efficiency due to limited MC statistics is $< 0.1\%$ for both signal modes. The systematic uncertainty arising from the number of $B\bar{B}$ pairs is 1.4%. The world average value of $\mathcal{B}[\Upsilon(4S) \rightarrow B^0\bar{B}^0]$ is $(48.6 \pm 0.6)\%$ [20], which corresponds to a systematic uncertainty of 1.2% in the fraction f^{00} .

The total additive (in number of events) and multiplicative (in percent) systematic uncertainties are obtained by adding in quadrature all systematic uncertainties of that type.

VII. SUMMARY

We have searched for the lepton-flavor-violating decays $B^0 \rightarrow \tau^\pm\ell^\mp$ using the full Belle data set. We find no evidence for these decays and set the following upper limits on the branching fractions at 90% C.L.:

$$\mathcal{B}(B^0 \rightarrow \tau^\pm\mu^\mp) < 1.5 \times 10^{-5} \quad (3)$$

$$\mathcal{B}(B^0 \rightarrow \tau^\pm e^\mp) < 1.6 \times 10^{-5}. \quad (4)$$

Our result for $B^0 \rightarrow \tau^\pm\mu^\mp$ is very similar to a recent result from LHCb [12]. Our result for $B^0 \rightarrow \tau^\pm e^\mp$ is the most stringent limit to date, improving upon the previous

TABLE II. Systematic uncertainties for the branching fraction measurement. Those listed in the upper section (“additive”) arise from fitting for the signal yield and are listed in number of events; those in the lower section (“multiplicative”) arise from the number of reconstructed B decays and are listed in percent.

Source	$B^0 \rightarrow \tau^\pm\mu^\mp$	$B^0 \rightarrow \tau^\pm e^\mp$
PDF shapes	0.7	0.3
Self-cross-feed fraction	< 0.1	0.1
Total (events)	0.7	0.3
B_{tag}	4.5	4.5
Track reconstruction	0.3	0.3
Lepton identification	1.6	1.8
MC statistics	< 0.1	< 0.1
Number of $B\bar{B}$ Pairs	1.4	1.4
f^{00} ($B\bar{B} \rightarrow B^0\bar{B}^0$ fraction)	1.2	1.2
Total (%)	5.1	5.2

limit by almost a factor of two. We find no indication of lepton flavor violation in these decays.

ACKNOWLEDGMENTS

We thank the KEKB group for the excellent operation of the accelerator; the KEK cryogenics group for the efficient operation of the solenoid; and the KEK computer group, and the Pacific Northwest National Laboratory (PNNL) Environmental Molecular Sciences Laboratory (EMSL) computing group for strong computing support; and the National Institute of Informatics, and Science Information NETwork 5 (SINET5) for valuable network support. We acknowledge support from the Ministry of Education, Culture, Sports, Science, and Technology (MEXT) of Japan, the Japan Society for the Promotion of Science (JSPS), and the Tau-Lepton Physics Research Center of Nagoya University; the Australian Research Council including grants DP180102629, DP170102389, DP170102204, DP150103061, FT130100303; Austrian Federal Ministry of Education, Science and Research (FWF) and FWF Austrian Science Fund No. P 31361-N36; the National Natural Science Foundation of China under Contracts No. 11435013, No. 11475187, No. 11521505, No. 11575017, No. 11675166, No. 11705209; Key Research Program of Frontier Sciences, Chinese Academy of Sciences (CAS), Grant No. QYZDJ-SSW-SLH011; the CAS Center for Excellence in Particle Physics (CCEPP); the Shanghai Pujiang Program under Grant No. 18PJ1401000; the Shanghai Science and Technology Committee (STCSM) under Grant No. 19ZR1403000;

the Ministry of Education, Youth and Sports of the Czech Republic under Contract No. LTT17020; Horizon 2020 ERC Advanced Grant No. 884719 and ERC Starting Grant No. 947006 “InterLeptons” (European Union); the Carl Zeiss Foundation, the Deutsche Forschungsgemeinschaft, the Excellence Cluster Universe, and the VolkswagenStiftung; L’Institut National de Physique Nucléaire et de Physique des Particules (IN2P3) du CNRS (France); the Department of Atomic Energy (Project Identification No. RTI 4002) and the Department of Science and Technology of India; the Istituto Nazionale di Fisica Nucleare of Italy; National Research Foundation (NRF) of Korea Grant Nos. 2016R1D1A1B-01010135, 2016R1D1A1B02012900, 2018R1A2B3003643, 2018R1A6A1A06024970, 2018R1D1A1B07047294, 2019K1A3A7A09033840, 2019R1I1A3A01058933; Radiation Science Research Institute, Foreign Large-size

Research Facility Application Supporting project, the Global Science Experimental Data Hub Center of the Korea Institute of Science and Technology Information and KREONET/GLORIAD; the Polish Ministry of Science and Higher Education and the National Science Center; the Ministry of Science and Higher Education of the Russian Federation, Agreement 14.W03.31.0026, and the HSE University Basic Research Program, Moscow; University of Tabuk research grants S-1440-0321, S-0256-1438, and S-0280-1439 (Saudi Arabia); the Slovenian Research Agency Grant Nos. J1-9124 and P1-0135; Ikerbasque, Basque Foundation for Science, Spain; the Swiss National Science Foundation; the Ministry of Education and the Ministry of Science and Technology of Taiwan; and the United States Department of Energy and the National Science Foundation.

-
- [1] Throughout this paper, charge-conjugate modes are implicitly included unless noted otherwise.
 - [2] Y. Amhis et al. (Heavy Flavor Averaging Group), *Eur. Phys. Jour. C* **81**, 226 (2021); and online update at <https://hflav.web.cern.ch/>.
 - [3] R. Aaij et al. (LHCb Collaboration), *Phys. Rev. Lett.* **122**, 191801 (2019).
 - [4] R. Aaij et al. (LHCb Collaboration), [arXiv:2103.11769](https://arxiv.org/abs/2103.11769) (2021).
 - [5] S. L. Glashow, D. Guadagnoli, and K. Lane, *Phys. Rev. Lett.* **114**, 091801 (2015).
 - [6] X.-G. He, G. Valencia, and Y. Wang, *Phys. Rev. D* **70**, 113011 (2004); S. Fukuda et al. (Super-Kamiokande Collaboration), *Phys. Rev. Lett.* **85**, 3999 (2000); Q.R. Ahmad et al. (SNO Collaboration), *Phys. Rev. Lett.* **89**, 011301 (2002); K. Eguchi et al (KamLand Collaboration). *Phys. Rev. Lett.* **90**, 021802 (2003); M. H. Ahn et al., *Phys. Rev. Lett.* **90**, 041801 (2003).
 - [7] Specifically, a tiny neutrino mass in the eV range.
 - [8] A. D. Smirnov, *Mod. Phys. Lett. A* **33**, 1850019 (2018).
 - [9] A. Dedes, J. Ellis, and M. Raidal, *Phys. Lett. B* **549**, 159 (2002).
 - [10] A. Bornheim et al. (CLEO Collaboration), *Phys. Rev. Lett.* **93**, 241802 (2004).
 - [11] B. Aubert et al. (BABAR Collaboration), *Phys. Rev. D* **77**, 091104(R) (2008).
 - [12] R. Aaij et al. (LHCb Collaboration), *Phys. Rev. Lett.* **123**, 211801, (2019).
 - [13] S. Kurokawa and E. Kikutani, *Nucl. Instrum. Methods Phys. Res., Sec. A* **499**, 1 (2003), and other papers included in this volume; T. Abe et al., *Prog. Theor. Exp. Phys.* **2013**, 03A001 (2013) and following articles up to 03A011.
 - [14] Z. Natkaniec et. al. (Belle SVD2 Group), *Nucl. Instrum. Methods Phys. Res., Sec. A* **560**, 1 (2006).
 - [15] A. Abashian et al. (Belle Collaboration), *Nucl. Instr. and Meth. A* **479**, 117 (2002), also see detector section in J. Brodzicka et al., *Prog. Theor. Exp. Phys.* **2012**, 04D001 (2012).
 - [16] D.J. Lange, *Nucl. Instr. and Meth. Sec. A* **462**, 152 (2001).
 - [17] T. Sjöstrand, S. Mrenna and P. Skands, *J. High Energy Phys.* **2006**, 05, 026 (2006).
 - [18] E. Barberio, B. van Eijk, and Z. Was, *Comput. Phys. Commun.* **66**, 115 (1991).
 - [19] R. Brun et al., CERN Report No. DD/EE/84-1 (1984).
 - [20] P.A. Zyla et al. (Particle Data Group), *Prog. Theor. Exp. Phys.* **2020**, 083C01 (2020)
 - [21] M. Feindt, F. Keller, M. Kreps, T. Kuhr, S. Neubauer, D. Zander, and A. Zupanc, *Nucl. Instrum. Methods Phys. Res., Sec. A* **654**, 432 (2011).
 - [22] E. Nakano, *Nucl. Instrum. Methods Phys. Res., Sect. A* **494**, 402 (2002).
 - [23] A. Abashian et al., *Nucl. Instrum. Methods Phys. Res., Sect. A* **491**, 69 (2002).
 - [24] K. Hanagaki et al., *Nucl. Instrum. Methods Phys. Res., Sect. A* **485**, 490 (2002).
 - [25] A. Sibidanov et al. (Belle Collaboration), *Phys. Rev. D* **88**, 032005 (2013).