


# Journal Pre-proof

First record of the spider family Hersiliidae (Araneae) from the Mesozoic of Europe  
(Bakony Mts, Hungary)

Márton Szabó, Jörg U. Hammel, Danilo Harms, Ulrich Kotthoff, Emese Bodor, János Novák, Kristóf Kovács, Attila Ősi



PII: S0195-6671(21)00345-1

DOI: <https://doi.org/10.1016/j.cretres.2021.105097>

Reference: YCRES 105097

To appear in: *Cretaceous Research*

Received Date: 19 June 2021

Revised Date: 19 November 2021

Accepted Date: 19 November 2021

Please cite this article as: Szabó, M., Hammel, J.U., Harms, D., Kotthoff, U., Bodor, E., Novák, J., Kovács, K., Ősi, A., First record of the spider family Hersiliidae (Araneae) from the Mesozoic of Europe (Bakony Mts, Hungary), *Cretaceous Research*, <https://doi.org/10.1016/j.cretres.2021.105097>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 The Author(s). Published by Elsevier Ltd.

Author statement:

**Márton SZABÓ:** Supervision, Conceptualization, Investigation, Writing – Original draft preparation. **Jörg U. HAMMEL:** Software, Writing, Reviewing and Editing. **Danilo HARMS:** Writing, Reviewing and Editing. **Ulrich KOTTHOFF:** Writing, Reviewing and Editing. **Emese BODOR:** Writing, Reviewing and Editing. **János NOVÁK:** Writing, Reviewing and Editing. **Kristóf KOVÁCS:** Software, Writing, Reviewing and Editing, **Attila ÓSI:** Investigation, Writing, Reviewing and Editing

1 First record of the spider family Hersiliidae (Araneae) from the Mesozoic of Europe (Bakony  
2 Mts, Hungary)

3 Márton SZABÓ<sup>\*1, 2</sup>, Jörg U. HAMMEL<sup>3</sup>, Danilo HARMS<sup>4,5</sup>, Ulrich KOTTHOFF<sup>4,5</sup>, Emese BODOR<sup>2,6</sup>,  
4 János NOVÁK<sup>7</sup>, Kristóf KOVÁCS<sup>8</sup>, Attila ŐSI<sup>1, 2</sup>

5  
6 <sup>1</sup> Hungarian Natural History Museum, Department of Paleontology and Geology, Ludovika tér  
7 2., Budapest 1083, Hungary

8 <sup>2</sup> Eötvös Loránd University, Department of Palaeontology, Pázmány Péter sétány 1/C, Budapest  
9 1117, Hungary. E-mail: szaboomarton@ttk.elte.hu, szabo.marton.pisces@gmail.com

10 <sup>3</sup> Helmholtz-Zentrum Hereon, Institute of Materials Physics, Max-Planck-Str. 1, 21502  
11 Geesthacht, Germany

12 <sup>4</sup> Leibniz-Institut zur Analyse des Biodiversitätswandels (LIB), Martin-Luther-King-Platz 3,  
13 20146 Hamburg, Germany

14 <sup>5</sup> Institute for Geology, Universität Hamburg, Bundesstraße 55, 20146 Hamburg

15 <sup>6</sup> Institute for Geological and Geochemical Research, Research Centre for Astronomy and Earth  
16 Sciences, Eötvös Loránd Research Network. Budaörsi út 45, Budapest 1112, Hungary. E-mail:  
17 emesebodor@gmail.com

18 <sup>7</sup> Eötvös Loránd University, Department of Systematic Zoology and Ecology, Pázmány Péter  
19 sétány 1/C, Budapest 1117, Hungary. E-mail: novakjanos01@gmail.com

20 <sup>8</sup> University of Pannonia – Research Centre for Engineering Sciences, Department of Materials  
21 Engineering, Egyetem utca 10, Veszprém 8200, Hungary. E-mail: kris@almos.uni-pannon.hu

22  
23 \*corresponding author

24  
25 key words: Santonian, ajkaite, amber, Hersiliidae, *Hungarosilia*

26  
27  
28  
29  
30  
31  
32  
33 Abstract

An adult male Hersiliidae spider is described from amber that originates from the Upper Cretaceous (Santonian) Ajka Coal Formation (Ajka-Csingervölgy, Hungary), the so-called ajkaite. The spider has elongate posterior spinnerets and a short third pair of legs III, both characteristics of the family, but differs from all known Mesozoic (discovered in burmite) and fossil and Recent European representatives of Hersiliidae so that a new genus and species name is proposed for *Hungarosilia verdesi* gen. et sp. nov.. The new taxon is distinguishable by the following unique combination of characters: pro- and opisthosoma nearly circular in dorsal view; posterior spinnerets with a distal article more than three times longer than the basal article; pedipalpal tibia and patella short and stout but pedipalpal femur as long as the cymbium, cymbium egg-shaped and without cymbial apophysis; bulbus of circular shape in ventral view and slightly flattened in lateral view. The fossil represents the first record of Hersiliidae from the Mesozoic of Europe and establishes the second record of this family in the Mesozoic Era. The estimated paleoclimate and paleoflora of the Ajka coal sub-basin correspond well with habitat preferences of Recent relatives that are often arborical and found in association with tree bark. Overall, our findings highlight the importance of a neglected amber type, the ajkaite, for documenting and studying the European arthropod fauna during the Mesozoic.

## Introduction

Arthropod fossils preserved in amber offer exciting possibilities to study morphology but also aspects of behaviour and ecology because the preservation is three-dimensional, often complete, and at a fine-scale that is unusual for other preservation modes, e.g., compressions fossils. Sporadic occurrences of amber are reported from numerous localities in Hungary that represent various ages (see Csillag and Földvári, 2005; Papp, 2002), but systematically collectable Mesozoic amber with a potential for arthropod-inclusions is only known from two localities: Iharkút (Csehbánya Formation) and Ajka (Ajka Coal Formation). Both ambers are estimated to be of Late Cretaceous (Santonian, 86,3-83,6 Mya) age. Our study focuses on the amber material collected from the Ajka Coal Formation, as despite the large sample size and almost identical geochemical composition (Kovács et al., 2015), arthropod inclusions are yet unknown from Iharkút amber (Hajdu, 2015; Ősi, 2012). The study of Santonian amber of the Ajka area (denominated as *ajkaite*) has a long tradition in Hungary and numerous experts have investigated amber from these deposits. A few hundred pieces have been collected by either scientists or amateur collectors in the last 160 years but, unfortunately, no one made comprehensive, layer-by-layer excavations or collections to increase the number of valuable inclusions. This may be the reason that very few arthropod fossils are still known from this type of amber although it preserves, amongst others, spiders and other arachnids.

The family Hersiliidae is an ecribellate entelegyne spider family that presently includes 183 species belonging to 16 genera (see the online database of the World Spider Catalog, Version 22.5). This family has quite a rich fossil record with roughly 20 species known mostly from fossil resins, i.e. the Cretaceous amber of Myanmar, Baltic amber in Europe from the Eocene, Dominican and Chiapas ambers of Miocene age, and even copal in Madagascar (see Table 1). Also termed “two-tailed spiders” or “tree trunk spiders”, these animals reside on or under the bark of trees where they hunt for prey at night. These spiders use their characteristic, elongate posterior lateral spinnerets for prey capture, usually insects, and rotate around the prey at great speed while holding the spinnerets over the victim so that it is fixed to the substrate with strong silk threads (Dippenaar-Schoeman, 2014; Eberhard, 1967; Murphy and Murphy, 2000; Rheims and Brescovit, 2004a; Yağmur et al., 2008).

The aim of the present study is to describe a male Hersiliidae individual discovered in *ajkaite* and place it in a systematic framework. We also evaluate the fossil within the climate and paleobotanical framework that has been proposed for the Ajka Coal paleoenvironment during the Santonian.

## Previous work on *ajkaite*

The presence of amber in the brown coal deposits of Ajka (Ajka Coal Formation, see below) has been known since the 19<sup>th</sup> century when coal mining in the area of Ajka had begun. Scientific evaluation of the Ajka amber started only in the 1920's (Kozma, 1991). The earliest report about the fossil resins of the Ajka Coal was established by Miksa Hantken (data from 1866, in Szabó, 1871) who mentioned „amber-like resin” from the Upper Cretaceous brown coal deposits. Later, this fossil resin was mentioned as ajkite (Anonymous, 1878) and the term ajkaite (*ajkait* in Hungarian) was first used by Zechmeister (1926). For physical and chemical characteristics of ajkaite see Csillag and Földvári (2005), Hlasiwetz (1871), Kovács et al. (2015), Papp (2002, and references therein), Rozložník (1940) and Zechmeister (1926). The composition is clearly different from the much younger Eocene succinites (Baltic-, Bitterfeld- and Rovno ambers) or rentinite and the most probable origin is copal (Góczán, 1961).

Although amber inclusions are rather poorly documented from Hungary, the arthropod inclusions preserved in ajkaite did attract the interest of some researchers. The earliest report about these inclusions belongs to Tasnádi Kubacska (1957) who mentioned „(...) mites, spiders, mosquitos, flies, tropical and subtropical species”. Koch (1985) wrote the following notes: „Insect inclusions are rather common in the amber pieces coming from here (Ajka)”. The first detailed paleoentomological report belongs to Borkent (1997) who described two new ceratopogonid species, *Leptoconops clava* and *Adelohelea magyarica*, from ajkaite. Czirják (2012), Czirják and Hajdu (2011), Főzy and Sente (2012), Hajdu and Czirják (2012) and Ősi (2012) briefly mentioned Arachnida, Coleoptera, Hymenoptera and Diptera inclusions. In her unpublished MSc thesis, Hajdu (2015) described the ajkaite fauna of flies (Diptera) in detail.

#### **Locality, geological background and age**

The first outcrops of the Ajka Coal Formation were discovered in the 1860's in the Ajka-Csinger Valley (or Ajkacsinger), roughly four kilometres southeast from the city Ajka (Bakony Mountains, western Hungary) (see Hantken, 1878; Szabó, 1871; see also Fig. 1A, B of the present study). Most if not all the available ajkaite specimens come from underground mines or spoil-banks along the Csinger Valley.

The Ajka Coal Formation occurs in the Bakony Mountains (Transdanubian Range) and is sometimes over 100 m thick. It is concentrated in three different carbonate terrain sub-basins (Ajka, Magyarpolány-Devecser, and Gyepükaján) (see Császár and Góczán, 1988; Siegl-Farkas, 1988). With Upper Triassic or, more rarely, Jurassic or Lower Cretaceous limestones forming a geological basement, the Ajka Coal Formation is built up by an alternation of coal or clayey coal beds, marls, sands and sandstone beds, as well as light grey to brownish

carbonaceous to argillaceous pelitic rocks with interbedded molluscan lumachelles (Haas, 1983). The depositional area was characterized by swampy and lacustrine environments resulting in among others at least seven main coal-beds (with the oldest coal bed 6. to the youngest coal bed 0.) with thicknesses ranging from 80-360 cm (Kozma, 1991).

Based on palynological and nannoplankton investigations, the age of the formation is Santonian (Bodor and Baranyi, 2012; Bodrogi et al., 1998; Ősi et al., 2016; Siegl-Farkas and Wagreich, 1996). The Ajka Coal Formation is overlain by either the marine Jákó Marl Formation (Haas et al., 1992), or discontinuously Eocene shallow-marine limestones of the Szóc Limestone Formation (Fig. 1C).

In fact, ajkaite occurs in varying amounts across all coal-bearing layers of the formation. However, the largest quantities come from the strata of the so-called Coal bed 0, also called ‘amber bank’ (*borostyánteleg* or *gyantásteleg* or *nullás telep* in Hungarian) in the late 1800’s (Rozlozsnik, 1940).

Besides amber fossils, a diverse flora and fauna of the Ajka Coal Formation have been described by numerous authors. The fossil flora of the formation has been described in detail by Greguss (1949), Kedves et al. (2000, 2001, 2002), Kedves and Alvarez Ramis (2002), Rákosi (1991), Rákosi and Barbacka (2000), Siegl-Farkas et al. (1988) and Tasnádi Kubacska (1957). Kedves and Párdutz (2002) reported the presence of three different types of morphologically intact bacteria preserved in ajkaite. The Ajka Coal Formation is famous for its extremely rich mollusc fauna, which was documented in several papers (Bandel and Riedel, 1994; Czabalay, 1988; Oppenheim, 1892; Tausch, 1886). Ostracods were investigated by Monostori (1988). Vertebrates are represented by fishes, turtles, mosasauroids, crocodyliforms, and dinosaurs (Ősi et al., 2016).

## Material and methods

Since coal mining in the area of Ajka ceased in 2004, ajkaite is collectable only at the coal refuses nearby the city (e.g. at the so-called Jókai coal refuse), usually as tiny splinters, or opaque, oxidated drops. For this reason, ajkaite is highly valued both by scientists and private collectors. Several hundreds of ajkaite pieces, housed in official and private collections, were inventoried first and then checked for arthropod remains. The present specimen results from this inventory and is deposited in the collection of the Department of Palaeontology and Geology of the Hungarian Natural History Museum (NHMUS, Budapest).

Károly Kozma (1936-2014) donated a large piece of ajkaite to the Paleontological Department of the Eötvös University (Budapest), supervised by Attila Ősi. The specimen came

from an unknown shaft of the Ajka-Csingervölgy coal mine and it was, arguably, the largest piece of ajkaite ever known to science (Fig. 2A). Since it was almost completely opaque and too dark for examination under a light microscope, the piece was scanned by using the NIKON Metrology Model XT H225 ST X-Ray tomograph of the University of Pannonia (Veszprém, Hungary). During the scanning procedure, multiple arthropod inclusions were discovered inside the piece, including a relatively large-sized spider that we describe here (Fig. 2B, C).

For the best available resolution of modelling, the ajkaite specimen was carefully broken into pieces around the spider. To achieve the highest resolution possible, the piece containing the spider was examined further at the Deutsches Elektronen Synchrotron (Hamburg, Germany) where it was studied using synchrotron radiation based micro-computed tomography (SR $\mu$ CT). Imaging was performed at the Imaging Beamline P05 (IBL) (Greving et al., 2014; Haibel et al., 2010; Wilde et al., 2016) operated by the Helmholtz-Zentrum Hereon at the storage ring PETRA III (Deutsches Elektronen Synchrotron – DESY, Hamburg, Germany). Imaging was done at a photon energy of 32.6 keV and a sample to detector distance of 80 mm. Projections were recorded using a custom-developed 20 MP CMOS camera system (Lytaev et al., 2014) with an effective pixel size of 1.28  $\mu$ m. For each tomographic scan 2401 projections at equal intervals between 0 and  $\pi$  have been recorded. Tomographic reconstruction has been done by applying a transport of intensity phase retrieval approach and using the filtered back projection algorithm (FBP) implemented in a custom reconstruction pipeline (Moosmann et al., 2014) using Matlab (Math-Works) and the Astra Toolbox (van Aarle et al., 2015, 2016; Palenstijn et al., 2011). For the processing, raw projections were binned for further processing two times, resulting in an effective pixel size of the reconstructed volume of 2.56  $\mu$ m.

### **Systematic paleontology**

Order Araneae Clerck, 1757

Infraorder Araneomorphae Smith, 1902

Family Hersiliidae Thorell, 1870

Type genus: *Hersilia* Audouin, 1826

Remarks: Based on the elongate, tapering posterior lateral spinnerets and the short third pair of legs, the specimen described below clearly represents the family Hersiliidae (Mirshamsi et al., 2013; Wunderlich, 2004, 2015; Wunderlich and Müller, 2020) although some other diagnostic characters remain equivocal. Based on the listed characteristics below, the Ajka-Csingervölgy specimen cannot be referred to any known Hersiliidae genus from the Mesozoic, nor to any

fossil or extant European representative of the family. A new genus is therefore proposed to accommodate the fossil.

Genus *Hungarosilia* gen. nov.

urn:lsid:zoobank.org:act:A59579E8-0D0D-4E5C-9711-63419C044573

Etymology: After Hungary; combined with the term „*silia*”, part of the family name Hersiliidae.

Type species: *Hungarosilia verdesi* gen. et sp. nov. (by monotypy).

Type horizon, age and locality: Ajka Coal Formation (Santonian, Upper Cretaceous), unknown shaft of the Ajka-Csingervölgy coal mine; Ajka-Csingervölgy (Hungary).

Diagnosis: The diagnosis is given for ♂, the ♀ is unknown. Details of clypeus, eyes, chelicerae, dorsal surface of prosoma and opisthosoma are unknown. Pro- and opisthosoma nearly as long as wide. Legs I, II and IV incomplete; metatarsus of legs III undivided and lacking long trichobothria; tarsi of legs III short without any sign of long trichobothria. Posterior lateral spinnerets elongate and roughly as long as the opisthosoma, their apical segment is more than three times longer than the basal article; anterior spinnerets separated by a triangular colulus. Pedipalpal tibia and patella short and stout but pedipalpal femur long (as long as the cymbium); cymbium egg-shaped, without cymbial apophysis; questionable apical cymbial bristles present; bulbus of circular shape in ventral view and slightly flattened in lateral view.

Remarks: Due to the incompleteness of the only known specimen, the key given by Wunderlich (2004:815) to fossil and extant hersiliid genera cannot be used here. However, the new genus differs from both the extinct and extant genera in a number of characters.

Up to the present study only two Mesozoic genera of the family have been described, namely *Burmesiola* and *Spinasilia* (Wunderlich, 2011a, 2015; Wunderlich and Müller, 2020), both from the Burmese amber of the Cenomanian (ca. 99 Ma) of Myanmar. *Burmesiola* has 5 long trichobothria on metatarsi III, and posterior spinnerets at least 2/3 as long as the opisthosoma, with the apical article at best slightly longer than the basal article (Wunderlich, 2011a:551, fig. 9 and photo 53a,b; 2015:303-304; photo 142; Wunderlich and Müller, 2020, photo 27). By contrast, preserved metatarsi III of *Hungarosilia* gen. nov. possess no signs of long trichobothria. The posterior spinnerets are as long as the opisthosoma and the apical article

is much (more than three times) longer than the basal one. *Hungarosilia* gen. nov. differs from *Spinasilia* in having posterior spinnerets about as long as the opisthosoma (posterior spinnerets of *Spinasilia* are longer than the opisthosoma), in the more egg-shaped cymbium, and in the lack of a cymbial apophysis (present in *Spinasilia*, see in Wunderlich, 2015).

*Gerdia*, a hersiliid genus described from the Eocene Baltic amber, differs from *Hungarosilia* gen. nov. in having an ovally elongate pro- and opisthosoma (Menge, 1869; Wunderlich, 2004). *Gerdiopsis*, another Eocene genus also described from the Baltic amber, possesses pedipalpi with long tibiae (see Wunderlich, 2004, fig. 7). Conversely, in *Hungarosilia* gen. nov. the pedipalpal tibia is short and stout. Unlike in *Hungarosilia* gen. nov., the pedipalp of the Baltic amber genus *Gerdiorum* also has long tibia (see Wunderlich, 2004, fig. 7a).

*Hungarosilia* gen. nov. also differs from extant Hersiliidae genera. According to Foord and Dippenaar-Schoeman (2006), *Hersilia* has armed chelicerae with teeth on the pro- and retromargin. Unfortunately, chelicerae of *Hungarosilia* gen. nov. are not visible in detail. An important difference between *Hungarosilia* gen. nov. and *Hersilia* is that the opisthosoma of *Hersilia* is oval to elongated (Chen, 2007; Pravalikha et al., 2014), sometimes wider in its posterior half (see *Hersilia jajaj* in Rheims & Brescovit, 2004b), while in *Hungarosilia* gen. nov. it is circular. Compared to *Hungarosilia* gen. nov., the extant genera *Bastanius*, *Delshevia*, *Duninia*, *Hersiliola*, *Ovtsharenkoia* have short posterior lateral spinnerets and a pedipalpal cymbium with a long tip (Marusik & Fet, 2009; Mirshamsi et al., 2015). Genera *Iviraiva*, *Neotama*, *Prima*, *Tamopsis*, *Tama* and *Ypyruera* possess pedipalpal characteristics (i.e. proportion of pedipalpal cymbium, tibia, patella and femur; shape of cymbium) very different from those of *Hungarosilia* gen. nov.. (see Rheims & Brescovit, 2004a; Foord & Dippenaar-Schoeman, 2005b; Foord, 2008b). The genus *Tama* also has an opisthosomal shape that is wider in its posterior half and a markedly flattened bulbus (Foord & Dippenaar-Schoeman, 2005a, table 1) while in *Hungarosilia* gen. nov. the opisthosoma is almost circular in dorsal view and the bulbus is flattened only slightly. The pedipalpal cymbium of *Neotama* is digiform but that of *Hungarosilia* gen. nov. is egg-shaped (Foord & Dippenaar-Schoeman, 2005b). The opisthosoma of *Murricia* is often subtriangular to subquadrate (Javed & Tampal, 2010), in contrast to *Hungarosilia* gen. nov. which has an almost circular opisthosoma. *Tyrotama* has short posterior lateral spinnerets (Foord & Dippenaar-Schoeman, 2005a) and differs from *Hungarosilia* gen. nov., that has posterior lateral spinnerets nearly as long as the opisthosoma. *Yabisi* has a somewhat deltoid opisthosoma wider in its posterior half (see figures of Rheims & Brescovit, 2004a) while opisthosoma of *Hungarosilia* gen. nov. is circular in dorsal view. The genus *Promurricia* includes a single species, *P. depressa* (Baehr & Baehr, 1993), which is based

on a female specimen. It shows a wide opisthosoma which is rather triangular but *Hungarosilia* gen. nov., which has a nearly circular opisthosoma.

*Hersiliola* and *Tama* are the only extant representatives of the family in Europe (Nentwig et al., 2021; Wunderlich, 2011e). Note here that the genus *Tama* might be present in the Baltic amber arachnofauna (see the short posterior lateral spinnerets of the figured *Hersiliidae* indet. in Wunderlich, 2004, photos 90, 91).

*Hungarosilia verdesi* gen. et sp. nov.

urn:lsid:zoobank.org:act:2ABE0ED0-FD7E-4E51-ACBB-7EA7875EC9FE

Etymology: In honour of †Gábor Verdes (1937-2010), grandfather of the first author who worked in Ajka for decades as a coal miner.

Type specimen: NHMUS PAL 2021.30.1., adult ♂ (only known specimen).

Type horizon, age and locality: Ajka Coal Formation (Santonian, Upper Cretaceous), unknown shaft of the Ajka-Csingervölgy coal minery; Ajka-Csingervölgy (Hungary).

Diagnosis: See the diagnosis of the genus.

Description: Habitus and legs: Prosoma nearly as wide as long (ca. 1.6 x 1.5 mm), with a closely circular outline in dorsal view, dorsal side is damaged; no characters of the clypeus and eyes are observable; order of legs cannot be given (as all I, II and IV legs are incomplete), however, based on their preserved parts, legs III seem to be the shortest; based on the preserved portions, the legs I, II and IV long and slender (although femur I is incomplete, it is still longer than the anteroposterior length of the prosoma); legs III with unarticulate metatarsi and short tarsi; opisthosoma about as long as wide (ca. 1.7 x 1.8 mm), nearly circular in ventral and dorsal view.

Spinnerets: posterior spinnerets elongate and about as long as the opisthosoma (ca. 1.8 mm, resulting in a body length/ posterior spinneret ratio of 1:1; see Fig. 4A), with feathery hairs medially (Fig. 4, psh), their distal article (ca. 1.4 mm; Fig. 4, daps) tapering and much longer

than the basal article (ca. 0.4 mm; Fig. 4, baps); anterior spinnerets (Fig. 4, as) much smaller than posteriors, separated by a distinct, triangular colulus (Fig. 4, co).

Copulatory organs: Both pedipalpi are preserved, however, several microfractures run across the amber and it hard to identify fine structures (e.g., hairs). Tibia (ca. 0.3 mm long; Fig. 5, t) and patella (ca. 0.25 mm long; Fig. 5, p) stout, femur (ca. 0.7 mm long; Fig. 5, f) elongate; cymbium (Fig. 5, c) with egg-shaped dorsal outline, as long as the pedipalpal femur; bulbus with circular shape in ventral view (Fig. 5E, J); questionable remains of apical cymbial bristles are present; no signs of a cymbial apophysis are visible.

Remarks: We compared *Hungarosilia verdesi* gen. et sp. nov. to chronologically (Late Cretaceous) or geographically close (European) hersiliid taxa. Except for *Hungarosilia verdesi* gen. et sp. nov., four Mesozoic hersiliid species, representing two genera are known, all from Burmese amber: *Burmesiola cretacea*, *B. daviesi*, ?*B. kachinensis* and *Spinasilia dissoluta* (Wunderlich, 2011a, 2015; Wunderlich and Müller, 2020). For a figurative summary of their comparison to *Hungarosilia verdesi* gen. et sp. nov. see Fig. 6. Unlike *Hungarosilia verdesi* gen. et sp. nov., the type specimen of *Burmesiola cretacea* has posterior spinnerets with the distal article slightly (~1.1-1.2x) longer than the basal, while the posterior spinnerets of *B. daviesi* have articles of about the same length (Wunderlich, 2011a, fig. 9; 2015). *Hungarosilia verdesi* gen. et sp. nov. has posterior spinnerets with a distal article more than three times longer than the basal article. The posterior spinnerets of *B. cretacea* are shorter while those of *B. daviesi* are longer than the opisthosoma (Wunderlich, 2011a, 2015). Conversely, the posterior spinnerets of *Hungarosilia verdesi* gen. et sp. nov. are as long as the opisthosoma. The pedipalpal tibiae of ?*Burmesiola kachinensis* (a species tentatively referred into genus *Burmesiola*) are elongate and longer than the cymbium. The prosoma of this species is longer than wide, and the posterior spinnerets are slightly longer than the opisthosoma (Wunderlich and Müller, 2020). By contrast, *Hungarosilia verdesi* gen et sp. nov. has stout pedipalpal tibiae that are shorter than the cymbium, the prosoma is nearly as long as wide, and the posterior spinnerets are as long as the opisthosoma. The type specimen of *Spinasilia dissoluta* is hard to observe in detail due to decomposition and the darkness of the embedding amber matrix (see Wunderlich, 2015). Cymbia of *S. dissoluta* are short and circular (Wunderlich, 2015) while those of *Hungarosilia verdesi* gen. et sp. nov. are egg-shaped. Cymbia of *S. dissoluta* bear a large cymbial apophysis while those of *Hungarosilia verdesi* gen. et sp. nov. lack this character. A further important difference is that posterior spinnerets of *S. dissoluta* are ~1.3x longer than

the opisthosoma while the posterior spinnerets and opisthosoma of *Hungarosilia verdesi* gen. et sp. nov. are more or less of the same length.

Five hersiliid species have been described from the Eocene Baltic amber so far: *Gerdia myura*, *Gardiopsis infringens*, *Gerdiorum inflexum*, *Hersilia longipes* and *Hersilia miranda* (Giebel, 1856; Koch and Berendt, 1854; Menge, 1869; Wunderlich, 2004, 2011b). The prosoma of *Gerdia myura* is elongate to oval (Wunderlich, 2004), unlike that of *Hungarosilia verdesi* gen. et sp. nov., which is almost as wide as long. The pedipalpal tibia of *Gardiopsis infringens* and *Gerdiorum inflexum* is elongate, longer than wide (as long as the cymbium, see Wunderlich, 2004, fig. 7, 7a) while that of *Hungarosilia verdesi* gen. et sp. nov. is short and stout (less than half of the length of the cymbium). The description of *Hersilia longipes* (see Giebel, 1856) is fairly incomplete and lacks illustrations, therefore a detailed comparison of *Hungarosilia verdesi* gen. et sp. nov. with this Baltic amber species is not possible. The description of *Hersilia miranda* does not include important characteristics of the male copulatory organs, but according to Koch and Berendt (1854) the opisthosoma of *Hersilia miranda* is egg-shaped whereas that of *Hungarosilia verdesi* gen. et sp. nov. is circular. Note that the only available illustration of *Hersilia miranda* (Koch and Berendt, 1854, pl. 17, fig. 147) is controversial because legs III are not the shortest (which questions the figured specimen's assignment to Hersiliidae). A revision of the validity of *Hersilia longipes* and *Hersilia miranda* would be helpful but the current location of the holotype specimen is unknown. We also note that Koch & Berendt (1854) did not produce accurate drawings of the appearance of their fossils but rather produced somewhat idealised pictures of their appearance.

*Hersilia aquisextana* has been published from the Oligocene of Aix-en-Provence (France) (Gourret, 1887). The validity of *Hersilia aquisextana* requires a detailed revision because the specimen is poorly illustrated and its relation to Hersiliidae remains dubious (see Gourret, 1887, pl. 21, fig. 14; note the lack of elongate posterior spinnerets and the relative length of legs III). Yet, if *Hersilia aquisextana* is tentatively accepted as member of family Hersiliidae and it differs from *Hungarosilia verdesi* gen. et sp. nov. in having an ovally elongate opisthosoma.

*Hersiliola* and *Tama* are represented in the Recent European arachnofauna by *Hersiliola simoni*, *Hersiliola macullulata* and *Tama edwardsi* (see Nentwig et al., 2021). These species differ from *Hungarosilia verdesi* gen. et sp. nov. in the following characters: *Hersiliola simoni* has an elongate (somewhat deltoid shaped) opisthosoma (Foord and Dippenaar-Schoeman, 2005a, fig. 3e) while that of *Hungarosilia verdesi* gen. et sp. nov. is circular. The pedipalpi of *Hersiliola simoni* possess more tapering cymbia and a more flattened bulbus (see Foord and

Dippenaar-Schoeman, 2005a, fig. 3a, b; Marusik and Fet, 2009, figs. 4.2, 5.2, 6.2-3; Rheims and Brescovit, 2004a, figs 39, 40), compared to *Hungarosilia verdesi* gen et sp. nov.. The same is true of the male copulatory organs of *Hersiliola macullulata* (Marusik and Fet, 2009, figs 4.1, 5.1, 6.1). The pedipalp of *Tama edwardsi* has tibiae that are as long as the cymbia, unlike *Hungarosilia verdesi* gen et sp. nov., that has a stout pedipalpal tibia much shorter than the pedipalpal cymbium.

## Discussion

Up to the present study, fossil spiders from Hungary were known only from the Pula Alginite Formation (Pula, Bakony Mountains, Hungary) of Pliocene age and a single fossil crab spider (*Xysticus* sp.) had been reported from this deposit (Katona et al., 2014, fig. 14).

Upper Cretaceous amber-bearing deposits are rather rare in Europe but some localities in France, Germany and Hungary are known (see Adl et al., 2011; Girard, 2009; Néraudeau et al., 2003, 2020; Perrichot, 2014; Perrichot et al., 2007a, b; Schlüter, 1978; Schmidt et al., 2001; Valentin et al., 2020; present study). Although some of these localities preserve rich assemblages of arthropods, only a few spiders or spider-related inclusions have been identified to date (Néraudeau et al., 2009; Penney, 2014; Saint Martin et al., 2014). The occurrence of Hersiliidae in the Santonian ajkaite establishes the oldest known record for this family in Europe, but also the second record of this family from the Mesozoic because all other fossils from this period are from Burmese amber in Myanmar (Table 1). Nowadays, only two hersiliid genera, namely *Hersiliola* and *Tama*, occur in Europe (Nentwig et al., 2021; Wunderlich, 2011e) and both are ground-dwelling members of the family found under stones where they build irregular webs plastered with small pebbles (Foord and Dippenaar-Schoeman, 2005a; Lawrence, 1964; Smithers, 1945).

The presence of Hersiliidae in the ajkaite arachnofauna corresponds well with the estimated paleoclimate and vegetation of the Ajka Coal paleoenvironment. Extant members of the family are medium-sized, often cryptic spiders that occur in warm climates and have the highest diversity in the tropics or subtropics. Based on drill core samples (Siegl-Farkas, 1988), the climate for the Ajka Coal Formation has been estimated to be tropical monsoon climate with high precipitation (2000–2500 mm per year). The later palynological and paleocarpological studies suggest Normapolles-related forests with fern-dominated underwood, but gymnosperm pollen and branches can be also found in the fossil record (Bodor and Baranyi, 2012). The climate might have been tropical or subtropical according to the ecological needs

of the nearest living relatives of the fossil plant forms (Bodor et al., 2012). Seasonality of the precipitation can be presumed based on the tree rings of fossil woods (L. Rákosi pers. comm.).

Life history traits and habitat choice of extant hersiliid spider species vary greatly between genera and some are arboreal hunters, living on the bark of tree trunks, whilst others are ground-dwelling spiders that prefer rock or boulders as shelters. The new species resembles the typical morphology of extant arboreal hersiliids (e.g., enlarged posterior spinnerets, elongate first, second and fourth pairs of legs, short third pair of legs). An arboreal lifestyle similar to many present-day hersiliid spiders is therefore probable (see in Foord, 2008a; Rheims and Brescovit, 2004a; also in Wunderlich, 2015:306 in fossil species *S. dissoluta*).

The swamp where the Ajka Coal was formed was arboreous and dominated by large angiospermes with possible affinity to Fagales (Normapolles-related; compare Friis et al., 2006). However, Platanaceae-related leaf fragments have also been found in the drill cores from the Ajka Coal Formation and Magnoliaceae-related *Padragkutia* and *Operculispermum* seeds were found in several layers of Ajka Coal Formation (Rákosi unpublished reports; Bodor et al., 2013). Rákosi also reported Araucariaceae trunks from the formation and recent relatives of this gymnosperm group have large trunks (reaching a height of 5–80 metres) with deeply seamed bark (Stockey et al., 2020) which could have been a potential habitat for this species. This variety in the floral assemblage indicates a complex ecosystem in which *Hungarosilia verdesi* gen. et sp. nov. may have found suitable habitat.

## Conclusion

The present study provides the first data on the Santonian arachnofauna of the western Tethyan archipelago. The discovery of *Hungarosilia verdesi* gen. et sp. nov. improves our knowledge on the Mesozoic diversity of family Hersiliidae. The suggested arboreal life style of our new hersiliid genus matches with the climate and paleoflora of the Ajka Coal paleoenvironment. There are further specimens of spiders from the Santonian ajkaite available for study, some of which are better preserved and will most likely reveal a greater diversity of spiders from the Mesozoic of Europe. The occurrence of further arboreal arachnids in ajkaite is expected.

## Acknowledgements

We thank the two anonymous reviewers for the constructive comments, which highly improved an earlier version of the manuscript. The authors are grateful to Zsófia Hajdu and Jörg Wunderlich for helpful discussions. This research was supported by the MTA-ELTE

Lendület Dinosaur Research Group (Grant no. 95102), National Research, Development and Innovation Office (NKFIH K 116665, K 131597, PD 130190, FK 130627), Hungarian Natural History Museum, Eötvös Loránd University, University of Pannonia (Research Centre for Engineering Sciences) and the Hungarian Dinosaur Foundation. We acknowledge DESY (Hamburg, Germany), a member of the Helmholtz Association HGF, for the provision of experimental facilities. Beamtime was allocated for proposal I-20190010. Danilo Harms and Ulrich Kotthoff thank the German Research Foundation (DFG) for financial support (HA 8785/5-1 and KO 3944/10-1).

## References

- Adl, S., Girard, V., Breton, G., Lak, M., Maharning, A., Mills, A., Perrichot, V., Trionnaire, M., Vullo, R., Néraudeau, D., 2011. Reconstructing the soil food web of a 100 million-year-old forest: The case of the mid-Cretaceous fossils in the amber of Charentes (SW France). *Soil Biology & Biochemistry* 43, 726–735., DOI: 10.1016/j.soilbio.2010.12.003
- Anonymous, 1878. Extraits de diverses publications. *Bulletin de la Société Française de Minéralogie et de Crystallographie* 1, 125–146.
- van Aarle, W., Palenstijn, W.J., De Beenhouwer, J., Altantzis, T., Bals, S., Batenburg, K.J., Sijbers, J., 2015. The ASTRA Toolbox: A platform for advanced algorithm development in electron tomography. *Ultramicroscopy* 157, 35–47, DOI: 10.1016/j.ultramic.2015.05.002
- van Aarle, W., Palenstijn, W.J., Cant, J., Janssens, E., Bleichrodt, F., Dabrovolski, A., De Beenhouwer, J., Batenburg, K.J., Sijbers, J., 2016. Fast and flexible X-ray tomography using the ASTRA toolbox. *Optics Express* 24(22), 25129–25147., DOI: 10.1364/OE.24.025129
- Baehr, M., Baehr, B., 1993. The Hersiliidae of the Oriental Region including New Guinea. *Taxonomy, phylogeny, zoogeography (Arachnida, Araneae)*. *Spixiana* 19, 1–96.
- Bandel, K., Riedel, F., 1994. The Late Cretaceous gastropod fauna from Ajka (Bakony Mountains, Hungary): a revision. *Annalen des Naturhistorischen Museums in Wien* 96A, 1–65.
- Bodor, E.R., Baranyi, V., 2012. Palynomorphs of the Normapolles group and related plant mesofossils from the Iharkút vertebrate site, Bakony Mountains (Hungary). *Central European Geology* 55(3), 259–292.
- Bodor, E.R., Baranyi, V., Hermanová, Z., 2012. The earliest Sabiaceae fruit remains of Hungary. *Hantkeniana* 7, 11–18.

- Bodor, E.R., Rákosi, L., Baranyi, V., Barbacka, M., 2013. Plant mesofossil based environmental reconstruction of the vicinity of Iharkút (the Bakony Mts. Hungary). In: Green Planet 400 million years of terrestrial floras: Symposium on the occasion of the 70th birthday of Han (Johanna H. A.) van Konijnenburg van Cittert Leiden, Netherlands (2013) p. 22
- Bodrogi, I., Fogarasi, A., Yazikova, E.A., Sztanó, O., Báldi-Beke, M., 1998. Upper Cretaceous of the Bakony Mts. (Hungary): sedimentology, biostratigraphy, correlation. Zentralblatt für Geologie und Paläontologie Teil I 11/12, 1179–1194.
- Borkent, A., 1997. Upper and Lower Cretaceous biting midges (Ceratopogonidae: Diptera) from Hungarian and Austrian amber and the Koonwarra Fossil Bed of Australia. Stuttgarter Beiträge zur Naturkunde Serie B, 249, pp. 10
- Chen, S.H., 2007. Spiders of the genus *Hersilia* from Taiwan (Araneae: Hersiliidae). Zoological Studies 46, 12–25.
- Császár, G.H., Góczán, F., 1988. A Bakony felső-kréta kőszénkutatás és kőszén láp vizsgálat. (in Hungarian: Upper Cretaceous coal prospecting and peat bog studies in the Bakony Mts). Magyar Állami Földtani Intézet Évi Jelentése 1986-ról, 155-178.
- Csillag, G., Földvári, M., 2005. Felső-triász gyantatöredék a Balaton-felvidékről. (in Hungarian: Upper Triassic amber fragment from the Balaton Upland). Magyar Állami Földtani Intézet Évi Jelentése 2005, 37–46.
- Czabalay, L., 1988. Az Ajkai Kőszén Formáció ökoszisztémái viszonyai a kagyló és csiga fauna alapján. (in Hungarian: Paleoecological study of the Ajka Coal Formation upon bivalves and gastropods). Magyar Állami Földtani Intézet Évi Jelentése 1986-ról, 211–227.
- Czirják, G., 2012. A Kárpát-medence borostyánjai és a borostyánok vizsgálati módszerei. (in Hungarian: Ambers of the Carpathian Basin, and methods for their investigation). BSc Thesis, Eötvös Loránd University, Institute of Geography and Earth Sciences, Budapest, pp. 42
- Czirják, G., Hajdu, Zs. 2011. Borostyánok és zárványaik a felső-kréta Ajkai Kőszén Formációból. (in Hungarian: Ambers and their inclusions from the Late Cretaceous Ajka Coal Formation). Abstract Book of the 14<sup>th</sup> Hungarian Palaeontological Convention (Magyar Őslénytani Vándorgyűlés), Szeged, p. 13–14
- Dippenaar-Schoeman, A.S., 2014. Field Guide to the Spiders of South Africa. LAPA Publishers, 432 p., ISBN: 9780799360189
- Eberhard, B.G., 1967. Attack behavior of digueid spiders and the origin of prey wrapping in spiders. Psyche 74, 173–181.

- 507 Friis, E.M., Raunsgaard Pedersen, K., Schönenberger, J., 2006. Normapolles plants: a  
 508 prominent component of the Cretaceous rosid diversification. *Plant Systematics and*  
 509 *Evolution* 260, 107–140.
- 510 Foord, S.H., 2008a. Cladistic analysis of the family Hersiliidae (Arachnida, Araneae) of the  
 511 Afrotropical Region, with the description of a new genus from Madagascar. *Journal of*  
 512 *Afrotropical Zoology* 4, 111–142.
- 513 Foord, S.H., 2008b. Cladistic analysis of the Afrotropical Hersiliidae (Arachnida, Araneae)  
 514 with the first records of *Murricia* and the description of a new genus from Madagascar.  
 515 *Journal of Afrotropical Zoology* 4, 111–142.
- 516 Foord, S.H., Dippenaar-Schoeman, A.S., 2005a. A revision of the Afrotropical species of  
 517 *Hersiliola* Thorell and *Tama* Simon with the description of a new genus *Tyrotama*  
 518 (Araneae: Hersiliidae). *African Entomology* 13(2), 255–279.
- 519 Foord, S.H., Dippenaar-Schoeman, A.S., 2005b. First records of the genus *Neotama* Baehr &  
 520 Baehr (Araneae: Hersiliidae) from the Afrotropical Region. *African Invertebrates* 46,  
 521 125–132.
- 522 Foord, S.H., Dippenaar-Schoeman, A.S., 2006. A revision of the Afrotropical species of  
 523 *Hersilia* Audouin (Araneae: Hersiliidae). *Zootaxa* 1347, 1–92.
- 524 Főzy, I., Szente, I., 2012. Ősmaradványok. A Kárpát-Pannon térség kövületei. (in Hungarian:  
 525 Fossils of the Carpathian Region). *Geolitera*, Szeged, pp. 584
- 526 Giebel, C.G., 1856. Die Insecten und Spinnen der Vorwelt mit steter Berücksichtigung der  
 527 lebenden Insekten und Spinnen. *Die Fauna der Vorwelt* 2, 1–511.
- 528 Girard, V., 2009. Evidence of Scenedesmeaceae (Chlorophyta) from 100 million-year-old  
 529 amber. *Geodiversitas* 31(1), 145–151.
- 530 Gourret, P., 1887. Recherches sur les Arachnides tertiaires d'Aix en Provence. *Recueil*  
 531 *Zoologique Suisse* 4, 431–496.
- 532 Góczán, F., 1961. Die Palynologie der Senon-Bildungen des Süd-Bakony. *Magyar Állami*  
 533 *Intézet Évi Jelentése* 49, 789–799.
- 534 Greguss, P., 1949. Az ajkai felső-kréta korú barnaköszén fusitzárványának meghatározása  
 535 (*Podocarpoxylon ajkaense* n. sp.). (in Hungarian: Determination of the fusit inclusion  
 536 from the Upper Cretaceous brown coal of Ajka (*Podocarpoxylon ajkaense* n. sp.)).  
 537 *Földtani Közlöny* 79, 9–12.
- 538 Greving, I., Wilde, F., Ogurreck, M., Herzen, J., Hammel, J.U., Hipp, A., Friedrich, F.,  
 539 Lottermoser, L., Dose, T., Burmester, H., Müller, M., Beckmann, F., 2014. P05 imaging  
 540 beamline at PETRA III: first results. In: Stock, R. S. (Ed.), *Proceedings of SPIE –*

- Developments in X-Ray Tomography IX, Vol. 9212, San Diego, California, United States, pp. 92120O-92120O-8.
- Haas, J., 1983. Senonian in the Transdanubian Central Range. *Acta Geologica Hungarica* 26, 21–40.
- Haas, J., Jocha-Edelényi, E., Császár, G., 1992. Upper Cretaceous coal deposits in Hungary. In: McCabe, P., Parris, J.T. (Eds.), *Controls on the distribution and quality of Cretaceous coals*. Geological Society of America, Special Paper 267, 245–262.
- Haibel, A., Ogurreck, M., Beckammn, F., Dose, T., Wilde, Herzon, J., Müller, M., Schreyer, A., Nazmov, V., Simon, M., Last, A., Mohr, J., 2010. Micro- and nano-tomography at the GKSS Imaging Beamline at PETRA III. *SPIE Optical Engineering + Applications*, 7804, SPIE, pp. 8
- Hajdu, Zs., 2015. A késő-kréta Ajkai Kőszén Formáció borostyánjai és kétszárnyú (Insecta: Diptera) faunája. (in Hungarian: Ambers and Diptera inclusions from the Late Cretaceous Ajka Coal Formation). MSc Thesis, Eötvös Loránd University, Institute of Geography and Earth Sciences, Budapest, pp. 56
- Hajdu, Zs., Czirják, G., 2012. A bakonyi kréta borostyánok és zárványaik. (in Hungarian: Cretaceous ambers and their inclusions from the Bakony Mountains). Abstract Book of the 15<sup>th</sup> Hungarian Palaeontological Convention (Magyar Őslénytani Vándorgyűlés), Uzsa, p. 14
- Hantken, M., 1878. A magyar korona országainak széntelepei és szénbányászata. Budapest, Légrády, 331 p.
- Hlasiwetz, H., 1871. Harz aus Braunkohle von Ajka im Veszprimer Comit. *Verhandlungen der Kaiserlich-Königlichen Geologischen Reichsanstalt* 5, 191–192.
- Javed, S.M.M., Tampal, F., 2010. Spiders of the genus *Murricia* Simon, 1882 (Araneae: Hersiliidae) from India. *Acta Zoologica Lituanica* 20(2), 88–97., DOI: 10.2478/v10043-010-0012-9
- Katona, L.T., Kutasi, Cs., Papp, B., Tóth, S., 2014. Further remarkable palaeontological finds at the alginite quarry in Pula. *Annales Historico-Naturales Musei Nationalis Hungarici* 106, 117–140.
- Kedves, M., Alvarez Ramis, C., 2002. Types of sporomorphs from the Ajkaite containing brown coal samples from Ajka (Hungary). *Plant cell biology and development* 14, 7–10.
- Kedves, M., Párdutz, Á., 2002. Ultrastructure of 100 million years old microorganisms of the ajkaite (Upper Cretaceous Amber) from Hungary. *Taiwania* 47, 273–280.

- Kedves, M., Szónoky, M., Madarász, M., Kovács, G., 2000. LM and TEM investigations on the Upper Cretaceous Ajkaite of Hungary I. *Plant cell biology and development* 12, 8–17.
- Kedves, M., Borbola, A., Priskin, K., 2001. LM and TEM investigations on the Upper Cretaceous Ajkaite of Hungary II. *Plant cell biology and development* 13, 22–31.
- Kedves, M., Borbola, A., Priskin, K., 2002. LM and TEM investigations on the Upper Cretaceous Ajkaite of Hungary III. *Plant cell biology and development* 14, 11–16.
- Koch, S., 1985. Magyarország ásványai. (in Hungarian: Minerals of Hungary). Akadémiai Kiadó, Budapest, pp. 562
- Koch, C.L., Berendt, G.C., 1854. Die im Bernstein befindlichen Crustaceen, Myriapoden, Arachniden und Apteren der Vorwelt. In: Berendt, G.C. (Ed.), Die im Bernstein Befindlichen Organischen Reste der Vorwelt Gesammelt in Verbindung mit Mehreren Bearbeitet und Herausgegeben, Berlin, pp. 1–124.
- Kozma, K., 1991. Az ajkai szénbányászat története. (in Hungarian: History of the coal mining at Ajka). Veszprémi Szénbányák Kiadó, Veszprém, pp. 531
- Kovács, I., Udvardi, B., Falus, G., Földvári, M., Fancsik, T., Kónya, P., Bodor, E.R., Mihály, J., Németh, C., Czirják, G., Ósi, A., Vargáné Barna, Z., Bhattoa, H.P., Szekanetz, Z., Turza, S., 2015. Az ATR FTIR spektrometria gyakorlati alkalmazása néhány - elsősorban földtani - esettanulmány bemutatásával. *Földtani Közlöny* 145(2), 173–192.
- Lawrence, R.F., 1964. A conspectus of South African spiders. Department of Agricultural Technical Services. South Africa. Science Bulletin 369, 1–64.
- Lytaev, P., Hipp, A., Lottermoser, L., Herzen, J., Gerving, I., Khokhriakov, I., Meyer-Loges, S., Plewka, J., Burmester, J., Caselle, M., Vogelgesang, M., Chiligaryan, S., Kopmann, A., Balzer, M., Schreyer, A., Beckmann, F., 2014. Characterization of the CCD and CMOS cameras for grating-based phase-contrast tomography. *SPIE Optical Engineering + Applications*, 9212, SPIE, pp. 10
- Marusik, Y. M., Fet, V., 2009. A survey of East Palaearctic *Hersiliola* Thorell, 1870 (Araneae, Hersiliidae), with a description of three new genera. *ZooKeys* 16, 75–114., DOI: 10.3897/zookeys.16.229
- Menge, A., 1869. Ueber einen Scorpion und zwei Spinnen im Bernstein. *Schriften der Naturforschenden Gesellschaft in Danzig (Neue Folge)*, 2(2-10), 1–9.
- Mirshamsi, O., Marusik, Y.M., Jamili, E., 2013. The spider family Hersiliidae Thorell, 1870 (Arachnida: Araneae) in Iran. *Zoology in the Middle East* 59(4), 347–352., DOI: 10.1080/09397140.2013.868140

- Mirshamsi, O., Zamani, A., Marusik, Y.M., 2015. A survey of Hersiliidae (Arachnida: Araneae) of Iran with description of one new genus and two new species. *Journal of Natural History* 50(23-24), 1447–1461., DOI:10.1080/00222933.2015.1113318
- Monostori, M., 1988. Jelentés az ajkai felső-kréta kőszénterület Ostracoda faunáinak vizsgálatáról. (in Hungarian: Report on the ostracod fauna of the Upper Cretaceous Ajka coal subbasin). Unpublished manuscript, 17 pp.
- Moosmann, J., Ershov, A., Weinhardt, V., Baumbach, T., Prasad, M.S., LaBonne, C., Xiao, X., Kashef, J., Hoffmann, R., 2014. Time-lapse X-ray phase-contrast microtomography for in vivo imaging and analysis of morphogenesis. *Nature Protocols* 9, 294–304.
- Murphy, F., Murphy, J., 2000. An Introduction to the Spiders of South East Asia with Notes on all the Genera. Kuala Lumpur: United Selangor Press Sdn Bhd, 665 pp.
- Nentwig, W., Blick, T., Bosmans, R., Gloor, D., Hänggi, A., Kropf, C., 2021, June 03. Spiders of Europe. Version 04.2021. Online at <https://www.araneae.nmbe.ch>, accessed on 04.27.2021, DOI: <https://doi.org/10.24436/1>
- Néraudeau, D., Allain, R., Perrichot, V., Videt, B., De Broin, F., Guillocheau, F., Philippe, M., Rage, J.C., Vullo, R., 2003. Découverte d'un dépôt paralique à bois fossiles, ambre insectifère et restes d'Iguanodontidae (Dinosauria, Ornithopoda) dans le Cénomanién inférieur de Fouras (Charente-Maritime, Sud-Ouest de la France). *Comptes Rendus Palevol* 2, 221–230.
- Néraudeau, D., Vullo, R., Gomez, B., Girard, V., Lak, M., Videt, B., Dépré, É., Perrichot, V., 2009. Amber, plant and vertebrate fossils from the Lower Cenomanian paralitic facies of Aix Island (Charente-Maritime, SW France). *Geodiversitas* 31(1), 13–27.
- Néraudeau, D., Saint Martin, J.-P., Saint Martin, S., Jeanneau, L., Mureau, J.-D., Philippe, M., Polette, F., Gendry, D., Brunet, J., Tréguier, J., 2020. Amber- and plant-bearing deposits from the Cenomanian of Neau (Mayenne, France). *Bulletin de la Société Géologique de France* 191(2), DOI: 10.1051/bsgf/2020039
- Oppenheim, P., 1892. Ueber einige Brackwasser - und Binnenmollusken aus der Kreide und dem Eocän Ungarns. *Zeitschrift der Deutschen Geologischen Gesellschaft* 44, 697–818.
- Ősi, A., 2012. Dinoszauruszok Magyarországon. (in Hungarian: Dinosaurs in Hungary). *Geoliter*, Szeged, 168 pp.
- Ősi, A., Bodor, E.R., Makádi, L., Rabi, M., 2016. Vertebrate remains from the Upper Cretaceous (Santonian) Ajka Coal Formation, western Hungary. *Cretaceous Research* 57, 228–238.

- Palenstijn, W.J., Batenburg, K.J., Sijbers, J., 2011. Performance improvements for iterative electron tomography reconstruction using graphics processing units (GPUs). *Journal of Structural Biology* 176(2), 250–253.
- Papp, G., 2002. A Kárpát-övezetben felfedezett ásványok, kőzetek és fosszilis gyanták története. *Studia Naturalia* 14, 147–163.
- Penney, D., 2006. Tertiary neotropical Hersiliidae (Arthropoda, Araneae) with new combinations for the extant fauna and comments on historical biogeography of the family. *Palaeontology* 49(4), 899–906.
- Penney, D., 2014. A fossil ray spider (Araneae: Theridiostomatidae) in Cretaceous amber from Vendée, France. In: Perrichot, V. (Ed.), *Fossil arthropods in Late Cretaceous Vendean amber (northwestern France)*. Paleontological Contributions, Number 10, p. 5–8., University of Kansas, p. 5–8.
- Perrichot, V., 2014. Fossil arthropods in Late Cretaceous Vendean amber (northwestern France). *Paleontological Contributions*, Number 10, University of Kansas, Paleontological Institute, ISSN 1946-0279 (online), 51 pp.
- Perrichot, V., Néraudeau, D., Nel, A., de Ploëg, G. 2007a. A reassessment of the Cretaceous amber deposits from France and their palaeontological significance. *African Invertebrates* 48(1), 213–227.
- Perrichot, V., Nel, A., Néraudeau, D., Lacau, S., Guyot, T., 2007b. New fossil ants in French Cretaceous amber (Hymenoptera: Formicidae). *Naturwissenschaften* 95(2), 91–97., DOI: 10.1007/s00114-007-302-7
- Petrunkévitch, A., 1963. Chiapas amber spiders. *University of California Publications in Entomology* 31, 1–40.
- Petrunkévitch, A., 1971. Chiapas amber spiders, II. *University of California Publications in Entomology* 63, 1–44.
- Pravalikha, G.B., Srinivasulu, C., Srinivasulu, B., 2014. New species of genus *Hersilia* Audouin, 1826 (Araneae: Hersiliidae) from India. *Journal of Threatened Taxa* 6(3), 5553–5557.
- Rákosi, L., 1991. Paleokarológiai vizsgálatok a Kretaceous és Tertiár Magyarországon. *Őslénytani Viták* 36–37, 127–133.
- Rákosi, L., Barbacka, M., 2000. Upper Cretaceous flora from Ajka (SW Hungary). I. *Thallophyta*. *Studia Botanica Hungarica* 30–31, 27–55.

- Rheims, C.A., Brescovit, A.D., 2004a. Revision and cladistic analysis of the spider family Hersiliidae (Arachnida, Araneae) with emphasis on Neotropical and Nearctic species. *Insect Systematics and Evolution* 35, 189–239.
- Rheims, C.A., Brescovit, A.D., 2004b. Description of four new species of Hersiliidae (Arachnida, Araneae) from Kinabalu National Park, Sabah, Borneo, Malaysia. *Journal of Natural History* 38, 2851–2861.
- Rozlozsnik, P., 1940. A csingervölgyi bányászat múltja, jelene és jövője. (in Hungarian: Past, present and future of the coal mining in Csingervölgy). *Magyar Királyi Állami Földtani Intézet Évi Jelentése az 1933-35 évekről* 3, 1231–1245.
- Saint Martin, J.-P., Saint Martin, S., Bolte, S., Néraudeau, D., 2014. Spider web in Late Cretaceous French amber (Vendée): The contribution of 3D image microscopy. (Les toiles d'araignée préservées dans l'ambre du Crétacé supérieur de France (Vendée): apport de l'imagerie 3D). *Comptes Rendus Palevol* 13(5), 463–472., DOI: 10.1016/j.crpv.2014.03.005
- Schmidt, A.R., von Eynatten, H., Wägreich, M., 2001. The Mesozoic amber of Schliersee (southern Germany) is Cretaceous in age. *Cretaceous Research* 22, 423–428., DOI: 10.1006/cres.2001.0267
- Schlüter, T., 1978. Zur Systematik und Palökologie harzkonservierter Arthropoda einer Taphozönose aus dem Cenomanium von NW-Frankreich. *Berliner Geowissenschaftliche Abhandlungen* A9, 150 p.
- Selden, P.A., Wang, Y., 2014. Fossil spiders (Araneae) from the Eocene Green River Formation of Colorado. *Arthropoda Selecta* 23(2), 207–219.
- Siegl-Farkas, Á., 1988. Az Ajkai Kőszén Formáció palynosztratigráfiája és fejlődéstörténete. *Magyar Állami Földtani Intézet Évi Jelentése 1986-ról*, 179–209.
- Siegl-Farkas, Á., Wägreich, M., 1996. Correlation of palyno- (spores, pollen, dinoflagellates) and calcareous nannofossil zones in the Late Cretaceous of the Northern Calcareous Alps (Austria) and the Transdanubian Central Range (Hungary). *Advances in Austrian–Hungarian Joint Geological Research*, Budapest, 127–135.
- Smithers, R.H.N., 1945. The Hersiliidae (Araneae) of South Africa. *Transactions of the Royal Society of South Africa* 31, 1–18.
- Stockey, R.A., Rothwell, G.W., 2020. Diversification of crown group *Araucaria*: the role of *Araucaria famii* sp. nov. in the Late Cretaceous (Campanian) radiation of *Araucariaceae* in the Northern Hemisphere. *American Journal of Botany* 107(7), 1072–1093.

- Szabó, J., 1871. Az ajkai kőszéntelep a Bakonyban. (in Hungarian: The coal deposits of Ajka in the Bakony Mountains). *Földtani Közlöny* 1, 124–130.
- Tasnádi Kubacska, A., 1957. Magyar ősgyanta. (in Hungarian: Hungarian fossil resin). *Élet és Tudomány*, 1957
- Tausch, L., 1886. Ueber die Fauna der nicht marinen Ablagerungen der oberen Kreide des Csongerthales bei Ajka im Bakony (Veszprimer Comitat, Ungarn) und über einige Conchylien des Gosaumergel von Aigen bei Salzburg. *Abhandlungen der Kaiserlich-Königliche Geologischen Reichsanstalt* 12, 1–32.
- Valentin, X., Garcia, G., Gomez, B., Gomez, V.D., Boiteau, J.-M., Saint Martin, S., Saint Martin, J.-P., 2020. New fossil assemblage with amber, plants and vertebrates from the lower Cenomanian near Châtelleraut (Vienne, western France). *Bulletin de la Société Géologique de France* 191(12), DOI: 10.1051/bsgf/2020034
- Wilde, F., Ogurreck, M., Greving, I., Hammel, J.U., Beckmann, F. Hipp, A., Lottermoser, L., Khokhriakov, I., Lytaev, P., Dose, T., Burmester, H., Müller, M., Schreyer, A., 2016. Micro-Ct at the imaging beamline P05 at PETRA III. *AIP Conference Proceedings* 1741, 030035, DOI: 10.1063/1.4952858.
- World Spider Catalog, 2021. World Spider Catalog. Version 22.5. Natural History Museum Bern, online at <http://wsc.nmbe.ch>, accessed on 2021.09.23. DOI: 10.24436/2
- Wunderlich, J. 1987. *Tama minor* n. sp., eine fossile Spinnenart der Familie Hersiliidae in Dominikanischen Bernstein (Arachnida: Araneae). *Entomologische Zeitschrift* 97, 93–96.
- Wunderlich, J., 1988. Die Fossilen Spinnen im Dominikanischen Bernstein. *Beiträge zur Araneologie* 2, 1–378.
- Wunderlich, J., 2004. On selected higher and lower taxa of fossil and extant spiders of the superfamily Oecobioidea, with a provisional cladogram (Araneae: Hersiliidae and Oecobiidae). *Beiträge zur Araneologie* 3, 809–848.
- Wunderlich, J., 2011a. Some fossil spiders (Araneae) in Cretaceous ambers. In: Wunderlich, J. (Ed.), *Extant and Fossil Spiders (Araneae)*. *Beiträge zur Araneologie* 6, 539–557.
- Wunderlich, J., 2011b. Some fossil spiders (Araneae) in Eocene European ambers. In: Wunderlich, J. (Ed.), *Extant and Fossil Spiders (Araneae)*. *Beiträge zur Araneologie* 6, 472–438.

- Wunderlich, J., 2011c. Some fossil spiders in Dominican amber (Araneae: Hersiliidae, Theridiidae, Gnaphosidae). In: Wunderlich, J. (Ed.), Extant and Fossil Spiders (Araneae). Beiträge zur Araneologie 6, 461–471.
- Wunderlich, J., 2011d. Some subrecent spiders (Araneae) in copal from Madagascar. In: Wunderlich, J. (Ed.), Extant and Fossil Spiders (Araneae). Beiträge zur Araneologie 6, 445–460.
- Wunderlich, J., 2011e. On extant West-Palaearctic (mainly Southern European) spiders (Araneae) of various families, with new descriptions. In: Wunderlich, J. (Ed.), Extant and Fossil Spiders (Araneae). Beiträge zur Araneologie 6, 158–338.
- Wunderlich, J., 2015. On the evolution and the classification of spiders, the Mesozoic spider faunas, and descriptions of new Cretaceous taxa mainly in amber from Myanmar (Burma) (Arachnida: Araneae). In: Wunderlich, J. (Ed.), Mesozoic Spiders (Araneae): Ancient Spider Faunas and Spider Evolution. Beiträge zur Araneologie 9, 21–408.
- Wunderlich, J., Müller, P., 2020. New and already described fossil spiders (Araneae) of 20 families in Mid Cretaceous Burmese amber with notes on spider phylogeny, evolution and classification. In: Wunderlich, J. (Ed.), Five papers on fossil and extant spiders. Beiträge zur Araneologie 6, 22–164.
- Yağmur, E. A., Kunt, K.B., Yalçın, M., 2008. The first record of family Hersiliidae from Turkey (Arachnida: Araneae). Serket 11(2), 62–64.
- Zechmesiter, L., 1926. Adatok az ajkait, egy hazai fosszilis gyanta ismeretéhez. (in Hungarian: Notes on ajkaite, a fossil resin from Hungary). Matematikai és Természettudományi Értesítő 43, 332–341.

Figure and table captions:

Fig. 1: A: Location of Ajka-Csingervölgy and the Ajka Coal area in western Hungary. B: distributional map of the Ajka Coal Formation (light grey) (modified after Császár and Góczán, 1988). C: Simplified geological section of the Ajka Subbasin (modified after Kozma, 1991).

Fig. 2: The possibly largest ever known piece of ajkaite. B, C: CT scan picture of the same specimen, showing the spider inclusions inside (area indicated by a white rectangle in figure B is enlarged in figure C). Pictures of B and C were taken at the University of Pannonia (Veszprém, Hungary). Scale bars: A, B: 3 cm; C: 2 mm

773

774 Fig. 3: *Hungarosilia verdesi* gen. et sp. nov., holotype (NHMUS PAL 2021.30.1.), body  
 775 habitus. A: dorsal view; B: ventral view; C: anterior view; D: right lateral view; E:  
 776 posterior view; F: left lateral view. Scale bar: 2 mm

777

778 Fig. 4: *Hungarosilia verdesi* gen. et sp. nov. (holotype); opisthosoma with spinnerets. A: right  
 779 lateral view, with slight saggital cross-cut on the lateral edge of the opisthosoma; B:  
 780 oblique posterior view; C: ventral view. Abbreviations: as: anterior spinnerets; baps: basal  
 781 article of the posterior spinneret; co: colulus; daps: dorsal article of the posterior  
 782 spinneret; psh: medial feathery hairs of the posterior spinneret. Scale bars: 500  $\mu$ m

783

784 Fig. 5: *Hungarosilia verdesi* gen. et sp. nov. (holotype); palpal morphology. A: left pedipalp in  
 785 lateral view; B: in medial view. C: right pedipalp in lateral view; D: in medial view. E:  
 786 left palpal tarsus (showing the cymbium and the bulbus) in ventral view; F: left cymbium  
 787 in dorsal view; G-I: saggital cross-section (in lateromedial order) of the left palpal tarsus.  
 788 J: right palpal tarsus (showing the cymbium and the bulbus) in ventral view; K: right  
 789 cymbium in dorsal view; L-N: saggital cross-sections (in lateromedial order) of the right  
 790 palpal tarsus. Abbreviations: c: cymbium; ?cb: possible cymbial bristles; b: bulbus; f:  
 791 palpal femur; p: palpal patella; t: palpal tibia. Scale bars: A-D: 750  $\mu$ m; E-N: 500  $\mu$ m


792

793 Fig. 6: Schematic drawings of characteristics of Mesozoic Hersiliidae (depictions of  
 794 *Burmesiola* spp. and *Spinasilia dissoluta* are re-drawn after the text figures and  
 795 photographs in Wunderlich, 2011a, 2015; Wunderlich and Müller, 2020).

796

797 Table 1: Fossil record of family Hersiliidae.


| Fossil Hersiliidae of the World |                                             |                                                     |                    |                                   |
|---------------------------------|---------------------------------------------|-----------------------------------------------------|--------------------|-----------------------------------|
| Horizon                         |                                             | Taxon                                               | Locality           | References                        |
| Mesozoic                        | Cenomanian (Cretaceous)                     | <i>Burmesiola cretacea</i>                          | Myanmar            | Wunderlich, 2011a                 |
|                                 |                                             | <i>Burmesiola daviesi</i>                           |                    | Wunderlich, 2015                  |
|                                 |                                             | <i>?Burmesiola kachinensis</i>                      |                    | Wunderlich & Müller, 2020         |
|                                 |                                             | <i>Spinasilia dissoluta</i>                         |                    | Wunderlich, 2015                  |
|                                 |                                             | Hersiliidae indet.                                  |                    | Wunderlich, 2015                  |
|                                 | <b>Santonian (Cretaceous)</b>               | <b><i>Hungarosilia verdesi</i> gen. et sp. nov.</b> | <b>Hungary</b>     | <b>Present study</b>              |
| Cenozoic                        | Ypresian (Eocene)                           | <i>Hersiliola</i> sp.                               | U.S.A.             | Selden & Wang, 2014               |
|                                 | Lutetian (Eocene)                           | <i>Gerdia myura</i>                                 | Baltic region      | Menge, 1869                       |
|                                 |                                             | <i>Gerdrops infringens</i>                          |                    | Wunderlich, 2004                  |
|                                 |                                             | <i>Gerdiorum inflexum</i>                           |                    | Wunderlich, 2004, 2011b           |
|                                 |                                             | <i>Hersilia longipes</i>                            |                    | Giebel, 1856                      |
|                                 |                                             | <i>Hersilia miranda</i>                             |                    | Koch & Berendt, 1854              |
|                                 |                                             | Hersiliidae indet. 1-3                              |                    | Wunderlich, 2004                  |
|                                 | upper Oligocene                             | <i>Hersilia aquisextana</i>                         | France             | Gourret, 1887                     |
|                                 | upper Oligocene to lower Miocene            | <i>Fictotama extincta</i>                           | Mexico             | Petrunkévitch, 1963; Penney, 2006 |
|                                 |                                             | <i>Perturbator corniger</i>                         |                    | Petrunkévitch, 1971; Penney, 2006 |
|                                 |                                             | <i>Prototama antiqua</i>                            |                    | Petrunkévitch, 1971; Penney, 2006 |
|                                 |                                             | <i>Prototama succinea</i>                           |                    | Petrunkévitch, 1971; Penney, 2006 |
|                                 | upper lower Miocene to lower middle Miocene | <i>Fictotama maculosa</i>                           | Dominican Republic | Wunderlich, 2011c                 |
|                                 |                                             | <i>Prototama maior</i>                              |                    | Wunderlich, 1988; Penney, 2006    |
|                                 |                                             | <i>Prototama media</i>                              |                    | Wunderlich, 1988; Penney, 2006    |
|                                 |                                             | <i>Prototama minor</i>                              |                    | Wunderlich, 1987; Penney, 2006    |
|                                 |                                             | <i>Prototama</i> sp.                                |                    | Wunderlich, 1988                  |
|                                 |                                             | <i>Tama</i> sp.                                     |                    | Schawaller, 1981                  |
|                                 | Quaternary (Pleistocene–Holocene)           | <i>Gorgorops jadis</i>                              | Madagascar         | Wunderlich, 2011d                 |
|                                 |                                             | <i>Hersilia madagascarensis</i>                     |                    | Wunderlich, 2004                  |
|                                 |                                             | <i>Hersiliana brevipes</i>                          |                    | Wunderlich, 2004                  |
|                                 |                                             | Hersiliidae indet.                                  |                    | Wunderlich, 2015                  |




A





B










C









|                                                 | Opisthosoma                                                                          | Palpus                                                                                |
|-------------------------------------------------|--------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------|
| <i>Hungarosilia verdesi</i><br>gen. et sp. nov. |    |    |
| <i>Burmesiola cretacea</i>                      |    | type specimen is<br>juvenile<br>?female                                               |
| <i>Burmesiola daviesi</i>                       |   | type specimen is<br>?juvenile<br>female                                               |
| ? <i>Burmesiola kachinensis</i>                 |  |  |
| <i>Spinasilia dissoluta</i>                     | opsithosoma is strongly<br>deformed                                                  |  |

**Declaration of interests**

☒ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

|  |
|--|
|  |
|--|