

Nonadiabatic charge-transfer within photoexcited Nickel porphyrins

Maria A. Naumova,¹ Gheorghe Paveliuc,² Mykola Biednov,³ Katharina Kubicek,^{3,4,5} Aleksandr Kalinko,¹ Jie Meng,^{6,7} Mingli Liang,⁶ Ahibur Rahaman,^{6,7} Mohamed Abdellah,^{7,8,9} Stefano Checchia,¹⁰ Frederico Alves Lima,³ Peter Zalden,³ Wojciech Gawelda,^{3,11,12,13} Christian Bressler,^{3,4,5} Huifang Geng,¹⁴ Weihua Lin,⁷ Yan Liu,⁷ Qian Zhao,⁶ Qingyin Pan,⁷ Marufa Akter,⁷ Qingyu Kong,¹⁵ Marius Retegan,¹⁰ David J. Gosztola,¹⁶ Matyas Pápai,¹⁷ Dmitry Khakhulin,³ Max Lawson Daku,² Kaibo Zheng,^{6,7} and Sophie E. Canton⁶


1. Deutsches Elektronen-Synchrotron DESY, Notkestr. 85, 22607 Hamburg, Germany
2. Département de Chimie Physique, Université de Genève, Quai E. Ansermet 30, CH-1211 Genève, Switzerland
3. European XFEL, Holzkoppel 4, 22869 Schenefeld, Germany
4. The Hamburg Centre for Ultrafast Imaging, University of Hamburg, Luruper Chaussee 149, 22761 Hamburg, Germany
5. Fachbereich Physik, Universität Hamburg, Notkestraße 9-11, 22607 Hamburg, Germany.
6. Department of Chemistry, Technical University of Denmark, Kongens, Lyngby DK-2800, Denmark
7. Chemical Physics and NanoLund, Lund University
8. Department of Chemistry, United Arab Emirates University, Al Ain P.O. Box 15551, United Arab Emirates
9. Department of Chemistry, Qena Faculty of Science, South Valley University, Qena, Egypt
10. ESRF - The European Synchrotron, 71, avenue des Martyrs, CS 40220, 38043 Grenoble Cedex 9, France
11. Departamento de Química, Universidad Autónoma de Madrid, Madrid 28049, Spain
12. IMDEA-Nanociencia, Calle Faraday 9, Madrid 28049, Spain
13. Faculty of Physics, Adam Mickiewicz University, Poznan 61-614, Poland
14. Department of Physics, Yantai University, 30 Qingquan Road, Yantai 264005, China
15. Synchrotron Soleil, L'Orme des Merisiers, 91190 Saint-Aubin, France
16. Center for Nanoscale Materials, Argonne National Laboratory, 9700 South Cass Avenue, Lemont, IL 60439, USA
17. HUN-REN Wigner Research Center for Physics, P.O. Box 49, Budapest H-1525, Hungary.

This document is the unedited Author's version of a Submitted Work that was subsequently accepted for publication in the Journal of Physical Chemistry Letters, copyright © 2024 American Chemical Society, after peer review. To access the final edited and published work see ACS Articles on Request

<https://pubs.acs.org/page/4authors/benefits/index.html#articles-request>

ABSTRACT: Metalloporphyrins with open d-shell ions can drive biochemical energy cycles. However, their utilization in photoconversion is hampered by rapid deactivation. Mapping the relaxation pathways is essential for elaborating strategies that can favorably alter the charge dynamics through chemical design and photoexcitation conditions. Here, we combine transient optical absorption spectroscopy and transient X-ray emission

spectroscopy with femtosecond resolution to probe directly the coupled electronic and spin dynamics within a photoexcited Nickel porphyrin in solution. Measurements and calculations reveal that a state with charge-transfer character mediates the formation of the thermalized excited state, thereby advancing the description of the photocycle for this important representative molecule. More generally, establishing that intramolecular charge-transfer steps play a role in the photoinduced dynamics of metalloporphyrins with open d-shell sets a conceptual ground for their development as building blocks capable of boosting nonadiabatic photoconversion in functional architectures through “hot” charge transfer down to the attosecond time scale.



KEYWORDS Metalloporphyrins, hot charge transfer, photoconversion, nonadiabatic processes, XFEL radiation, femtosecond X-ray sciences.

Metalloporphyrins (MPs) incorporating 3d transition metal ions with closed or open d-shells are primary reactants in biochemical energy cycles.^{1, 2, 3, 4, 5} Building upon the first reports of porphyrin π -ring synthesis in the 1930s,^{6, 7} modern protocols have expanded the MP family to encompass most of the stable elements.^{8, 9, 10, 11, 12, 13} The so-called “MP periodic table”^{14, 15} rationalizes how the nature of the metal ion imparts unique photophysical properties, placing MPs at the focus of a cross-disciplinary research effort that targets the bio-inspired production, conversion, transduction and storage of renewable energy.^{16, 17, 18, 19, 20} In contrast, the integration of MPs containing transition metal ions with open 3d-shell in photoconversion applications remains limited despite strong prospects associated with their coupled electronic and magnetic activities. This status can be traced back to the fact that, although the initial photoexcitation is delocalized over the porphyrin π -ring as for their closed-shell congeners, the pathways of intramolecular relaxation are rapidly obscured by the participation of intermediate states that involve the d-levels of the metal ions.^{16, 17, 18} Several decades of investigations with electronic spectroscopy have shown that it is rarely possible to unambiguously differentiate (d,d), (π ,d) and (d, π) states in the UV-visible and near-infrared spectral ranges.^{21, 22, 23} Ultrafast techniques with inherent structural sensitivity such as time-resolved Raman spectroscopy (tr-RSS)^{24, 25, 26} and, recently, time-resolved X-ray absorption spectroscopy (tr-XAS) on the picosecond (ps) and femtosecond (fs) time scales^{23, 27, 28, 29, 30, 31, 32} are partially challenged by the lack of direct spin-sensitivity, so that they tend to return highly-correlated rate constants. As a result, for most open d-shell MPs, the description of the photocycle is not exhaustive and the photoredox capabilities of the excited states cannot be fully explored. Here, we combine transient optical absorption spectroscopy (tr-OAS) with transient X-ray emission spectroscopy (tr-XES), both achieving femtosecond resolution, in order to track directly the coupled electronic and spin dynamics

within a photoexcited Nickel (Ni) porphyrin in solution. The measurements reveal a distinct intermediate state with charge-transfer (CT) character in the excited triplet manifold. Supported by time-dependent density functional theory (TD-DFT) and excited-state DFT (e-DFT) theoretical calculations, the observations allow proposing a pathway connecting the singlet Franck-Condon (FC) state with excitation localized on the π -ring (i.e., ligand-centered (LC) character) to the triplet metastable (d,d) state (i.e., metal-centered (MC) character). Besides completing the description of the photocycle for this important representative molecule, the present study also outlines novel directions for utilizing photoexcited Ni porphyrins and other open d-shell MPs to trigger “hot” charge transfer events in large functional architectures down to the attosecond time scale.

The basic Nickel porphyrin molecule is formed through metalation of the porphin dianion (P^{2-}) by Ni(II) cations. The Ni(II) tetrakis (2,4,6-trimethylphenyl) porphyrin (NiTMP) possesses aryl substituents added at the four meso positions. The chemical structure is shown in **Figure 1a**.

The experimental details for the synthesis of the molecule are given in **S.I.1**. The structural optimization of NiTMP with density functional theory (DFT) calculations performed in C_{2v} symmetry delivers the geometry of the $1A_1$ singlet ground state in toluene (**Figure 1b**). The computational details are given in **S.I.2 (S1)**. The value of the average Ni-N bond length, noted R, is 1.953 Å. The molecule departs from planarity as the ring presents clear ruffling. This type of deformation is ascribed to the fact that the Ni-N bonds are shorter than the radius of the N_4 cavity in the planar π -ring of free P^{2-} .³³ The Ni(II) ion is in the $(d_{xy})^2(d_{xz})^2(d_{yz})^2(d_z^2)^2$ ($S=0$) low spin (LS) electronic configuration. The Kohn-Sham orbitals are displayed in **S.I.2 (S2)**.

The optical spectra of closed d-shells MPs are very well described by the so-called Gouterman model involving four orbitals, namely the two nearly-degenerate a_{2u} HOMO, a_{1u} HOMO-1³⁴ (red) and the doubly-degenerate e_g LUMO (purple) of P^{2-} (Figure 2a).^{35, 36} Within this model, pronounced configuration interaction (C.I.) between the excited states originating from the $(a_{1u})^2(a_{2u})^1(e_g)^1$ and $(a_{1u})^1(a_{2u})^2(e_g)^1$ configurations produces two (π, π^*) singlet states, S_1 and S_2 , and two (π, π^*) triplet states, T_1 and T_2 , for which the excitation is localized on the π -ring, i.e., with ligand-centered (LC) character. These states are indicated as solid and dashed black lines in **Figure 1c**.

The d-electrons of the Ni(II) ion perturb these energetics only slightly, with possible formation of $(\pi, d)/(d, \pi)$ states of charge transfer (CT) character, and (d,d) states of metal-centered (MC) character within the singlet and triplet manifolds. These states are indicated as dashed gray lines. The two optically-allowed transitions, $S_0 \rightarrow S_2$ and $S_0 \rightarrow S_1$, yield the strong Soret B band and the weaker Q band (orange arrows). Direct transitions to the triplet states are optically-forbidden in the absence of spin-orbit coupling (SOC).

The UV-vis spectrum of NiTMP in toluene is shown in **Figure 1d**. The B band is centered around 415 nm. The Q band displays a vibronic sub-structure, with Q(1,0) and Q(0,0) at 527 nm and 558 nm (**inset of Figure 1d**). The experimental spectrum is well reproduced by time-dependent DFT (TD-DFT) calculations using the PBE functional (**Figure 1e**). The main transitions between the occupied and virtual natural transition orbitals (NTOs) underlying the Soret band are depicted in **Figure 1f**. They are located on the porphyrin ring and do not implicate any electronic density at the Ni(II) center, as expected from a (π, π^*) transition (**S3** and **S4** in **S.I.2**).

The ultrafast photoinduced dynamics of NiTMP were first monitored with femtosecond tr-OAS in the UV-visible range. The setup is described in **S.I.3**. NiTMP was dissolved in toluene and excited in the B band at 400 nm. Transient optical absorption spectra acquired over the sub-picosecond, tens of picosecond and hundreds of picosecond temporal windows are presented in **Figures 2a, 2b** and **2c** for specific pump-probe delays Δt . The incident fluence was set in the linear regime of photoexcitation (**S.I.3**).

In **Figure 2a**, the GS bleach (GSB) of the Q(1,0) band at 527 nm (feature B) appears within the instrument response function (IRF) defined by a gaussian function of full-width half maximum (FWHM) ~ 120 fs, along with positive excited-state absorption (ESA) signals on the blue side (feature E_1) and the red side (feature E_2) of feature B. A small peak forms around 635 nm (indicated by the red star). Isosbestic points are observed at 444 nm and 577 nm. The inverse of the GS absorbance is shown as the gray trace. In **Figure 2b**, the intensity of E_1 decays rapidly, while E_2 undergoes profound alterations as it narrows and blue-shifts with the small peak localizing towards shorter wavelengths (indicated by the black star). In **Figure 2c**, the transient signal of the lowest excited state exhibits a derivative-like profile with a shoulder at 580 nm (indicated by the black star). It decays on the ~ 200 ps time scale with a clear isosbestic point at 534 nm. Early studies have assigned this metastable state as a $(d_{z^2}, d_{x^2-y^2})$ state with MC character owing to the derivative-like profile. This distinctive spectral lineshape was proposed to result from the superposition of the GSB signal with an ESA signal corresponding to the same GS spectrum, but shifted to lower energy. The spectral red-shift is attributed to the repulsion between the electrons populating the a_{2u} ring-based orbital and the $d_{x^2-y^2}$ metal-based orbital that both place electronic density in the vicinity of the N atoms. Such interaction destabilizes the a_{2u} HOMO, while leaving the e_g LUMO unaffected, thereby only reducing the HOMO-LUMO gap (see **Figure 1c**).^{37, 38} Although the spin multiplicity of the (d,d) state cannot be inferred from the optical measurement, it is conventionally assumed to become triplet through intersystem crossing (ISC), in line with Hund's rule and the absence of detectable stimulated emission in **Figure 2a**. The individual kinetics are expected to be probe-dependent considering the nonadiabatic nature of the dynamics following photoexcitation. Nevertheless, a global-fit analysis based on a simple sequential kinetic model including three exponential functions captures the essential intensity redistribution over extended ranges of wavelengths and times. The three components are displayed in **Figure 2d**, after scaling their minima to -1 for visualization purposes. The best-fit values of the time constants are 560 ± 60 fs (red line), 6 ± 1 ps (orange line) and 210 ± 20 ps (green line). **Figure 2e** shows the transient optical absorption spectrum observed at 50 ps (dotted line). The model trace (solid line) is constructed as the difference between the ground-state UV-visible spectrum red-shifted by ~ 20 nm and the ground-state UV-visible spectrum itself. The excellent agreement between the traces supports the assignment of the metastable state as a (d,d) state with MC character. Selected kinetics at 450 nm, 500 nm, 600 nm and 700 nm are presented in **Figure 2f**, along with their corresponding best-fit traces. The spectral dynamics over the first several tens of ps reflect the formation of the (d,d) state through competing internal conversion (IC), intersystem crossing (ISC) and intramolecular vibrational energy redistribution (IVR).^{26, 39} As noted above, the absence of stimulated emission in the tr-OAS measurements indirectly suggests that the excited-state manifold acquires a triplet spin-multiplicity on the sub-hundred fs time scale through ISC. However, the multi-exponential evolution of the kinetics does not display clear-cut sensitivity towards the extent of the excitation localization at the Ni center in the metastable MC state.

Therefore, femtosecond tr-XES measurements were carried out at the FXE Instrument of the European XFEL facility^{40, 41, 42, 43} in order to track directly the dynamics of the spin-multiplicity at the Ni(II) center during the relaxation across the excited-state manifold. The setup is shown in **S11** of **S.I.4**. The data acquisition and the analysis procedures are described in **S.I.4**. In the hard X-ray regime, the radiative relaxation upon 1s core-ionization produces the emission lines $2p_{1/2,3/2} \rightarrow 1s$ ($K\alpha_{1,2}$) and $3p_{1/2,3/2} \rightarrow 1s$ ($K\beta$). They are sensitive to the coupled changes in spin-multiplicity and electronic structure.⁴⁴ **Figures 3a** and **3b** show the $K\alpha_{1,2}$ and $K\beta$ spectra for NiTMP in toluene in its GS. **Figures 3c** and **3d** present the difference traces [(laser_on)-(laser_off)] delivering the $K\alpha_{1,2}$ and $K\beta$ tr-XES spectra following photoexcitation at 400 nm with a fluence of ~ 4 mJ/cm² (linear regime) for the optical pump-X-ray probe time delays Δt of 1 ps and 50 ps (**S12** in **S.I.5** and **S13, S14** in **S.I.6**). The similarity between the two transient XES spectra demonstrates that the MC character is acquired on the sub-picosecond time scale. Taking the summed absolute value of the integrated areas under the transient difference signals as figure of merit (FOM) and tracking it as a function of pump-probe time delay allows accessing the kinetics of the relaxation with spin-sensitivity.^{44, 45} **Figures 3e** and **3f** show the $K\alpha_{1,2}$ and $K\beta$ kinetics acquired under low power laser excitation, along with their single-exponential decay fits (black solid and dotted lines) where the IRF FWHM is set to 110 fs, delivering a lifetime $\sim 161 \pm 18$ ps and $\sim 143 \pm 34$ ps for the metastable state. The nominal powers and best-fit values for the parameters are given in **S15** and **S16, S17 and S18** in **S.I.7**. The values of the single-exponential time constants are similar for the optical and X-ray spectroscopic measurements, ascertaining that both techniques are probing the decay of the metastable state.

The characteristics of the metastable state are established with theoretical methods. The lowest triplet states obtained from DFT and TD-DFT applying C_{2v} constraint are of 3A_2 symmetry. The optimized DFT / relaxed TD-DFT structure of this state denoted 1^3A_2 exhibits an average Ni-N bond elongation ΔR of 0.072 / 0.086 Å with concurrent planarization of the porphyrin π -ring. The molecular geometries are shown respectively in **S5, S7, S8, S9** of **S.I.2** and **Figure 3g**. Examining the Kohn-Sham orbitals from DFT for the $e1^1A_1$ ground state and 1^3A_2 shown in the **S.I.2** (**S6**) reveals that the two states differ by the promotion of an electron from a d_{z^2} -like MO to a $d_{x^2-y^2}$ -like MO. Examining the NTOs from TD-DFT associated to the relaxed 1^3A_2 confirms its formation via an electronic transition from a d_{z^2} -like MO to a $d_{x^2-y^2}$ -like MO (**Figure 3g**). Overall, both methods establish that the lowest excited triplet state 1^3A_2 displays an MC character with an expanded and planar core compared to the 1^1A_1 ground state correlated to the population of a $d_{x^2-y^2}$ -like MO.

The early spin dynamics were followed with tr-XES on the sub-ps time scale with a fluence of ~ 2 mJ/cm² (**S12** in **S.I.5**). Transient XES spectra were acquired on a fine temporal grid over 2 ps. **Figures 4a** and **4b** display the transient $K\alpha_{1,2}$ and $K\beta$ traces averaged around the central pump-probe delays Δt of -50 fs and 50 fs (binned over a 100 fs time window), 200 fs, 450 fs and 800 fs (binned respectively over a 200, 300 and 400 fs time window) across the energy range where the transient signal is detected. **Figures 4c** and **4d** show the $K\alpha_{1,2}$ and $K\beta$ kinetics, with their fit as single-exponential rise tracking the formation of the MC state. (**S19, S20, S21, S22, S23** in **S.I.7**).

However, close inspection of the profiles reveals a departure from a uniform growth, with clear alterations in spectral weight as a function of Δt . The differences in feature positions and amplitude ratios flag the participation of an intermediate species in the ultrafast formation of the metastable MC species. Focusing on the first 2 ps, the global-fit analysis of the $K\alpha_{1,2}$ and $K\beta$ lines based on a single exponential rise convoluted by a Gaussian IRF of 110 fs FWHM⁴¹ captures the early dynamical evolution (**S.I.6**). The best fit parameters are summarized in **S24**, **S25** and **S26** of **S.I.7**. The components A_1 , A_2 for $K\alpha_{1,2}$ and B_1 , B_2 for $K\beta$ are shown in **Figure 4e** and **4f** with 2-point rebinning. The first components (A_1 for $K\alpha_{1,2}$ and B_1 for $K\beta$, in blue) appear within the IRF of the tr-XES measurement. They evolve towards the second component (A_2 for $K\alpha_{1,2}$ and B_2 for $K\beta$, in red) with a time-constant $\tau_1 \sim 640 \pm 120$ fs ($K\alpha_{1,2}$) and $\tau'_1 \sim 660 \pm 150$ fs ($K\beta$). The lineshapes of components A_2 and B_2 are very similar to the $K\alpha_{1,2}$ and $K\beta$ transient difference spectra acquired at a pump-probe delay Δt of 1 ps and 50 ps displayed in **Figures 3c** and **3d** so that they can be unambiguously identified as the fingerprints of the triplet state (d,d) of MC character. In contrast, the lineshapes of A_1 and B_1 present clear variations. The amplitudes of features a_1 and a_2 in the $K\alpha_1$ range of A_1 are smaller than for A_2 , while the feature a_3 is slightly broader. The amplitudes of all transient features in the $K\alpha_2$ range are smaller. The amplitudes of features b_1 , b_3 and b_4 in the $K\beta$ range of B_1 are smaller than for B_2 , while b_2 appears suppressed. In other words, the intermediate species exhibits transient XES signatures that are distinguishable from the ones of the triplet MC state displayed in **Figures 3c** and **3d**. Moreover, it should be noted here that the vertical transition to the singlet FC state (without geometrical changes) does not involve any appreciable electronic density at the metal ion (**Figure 1f**) so that it could be expected to appear predominantly silent in the tr-XES measurement. Therefore, the intermediate species possesses a CT character. The temporal evolution of the concentration of the CT and MC species using the best-fit parameters extracted from the global-fit analysis of $K\alpha_{1,2}$ for which the formation time is 640 fs are shown in **S27** of **S.I.7**.

The assignment of the intermediate CT state is then investigated with theoretical methods. The excited states of triplet spin-multiplicity lying above the lowest metastable 1^3A_2 triplet state and below the initial excitation energy were first characterized with TD-DFT calculations for each symmetry. The optimized geometries for 1^3A_1 of MLCT character and 2^3A_2 of LMCT character are shown in **Figure 5a** and **5b**, along with the NTO \rightarrow NTO transitions associated with the corresponding electronic excitation. The frequency analysis necessary to determine whether the relaxed geometries correspond to local minima or transition states could not be performed due to the lack of analytical second derivatives in the TD-DFT implementation of the ADF software. The optimizations were then performed with the Gaussian16 software that allows such frequency analysis. The resulting triplet structures obtained with very tight convergence criteria are similar to the ones obtained with ADF and all exhibit imaginary frequencies (**S10** in **S.I.2**). Consequently, the low-lying states of triplet spin-multiplicity for each symmetry were also characterized with e-DFT calculations, wherein the occupation of the MOs is constrained and allowed to be non-Aufbau during the self-consistent (SCF) procedure.⁴⁶ The resulting structures of the 2^3A_2 state with mixed LMCT/MC character and the 3^3A_1 state with LMCT character are shown in **Figures 5c** and **5d**. Frequency analyses performed on the optimized geometries indicate that they are actual local minima, without imaginary frequencies. The calculations for 1^3A_1 and 2^3A_1 converged to extrema corresponding to transition states, with one imaginary frequency. For each state, the MO \rightarrow MO transition depicts the passing from the closed-shell ground state to the triplet state of interest as a constrained spin-flip promotion (**Figures 5c** and **5d**). The corresponding spin-density isosurfaces are displayed in **S.I.2**. Using the D_{4h} notation, which is applicable to planar and quasi-planar geometries, the TD-DFT and e-DFT calculations with C_{2v} constraints identify the 2^3A_2 triplet state of dominant LMCT (TD-DFT)

or mixed LMCT/MC (e-DFT) character that implicate an a_{2u} -like HOMO and a $d_{x^2-y^2}$ -like orbital, thereby noted $(a_{2u})^1(d_{x^2-y^2})^1$. The degenerate 1^3B_1 and 1^3B_2 of MC character obtained with TD-DFT and e-DFT are shown in **S.I.2** for comparison, although this character can be ruled out based on the tr-OAS and tr-XES observations.

To summarize, following the FC singlet excitation of LC character, i.e. localized on the π -ring, the tr-XES measurements establish that the nonadiabatic formation of the metastable triplet state with MC character, where the excitation is localized over the metal, is mediated by the population of a triplet state with CT character, where the excitation is delocalized over the ligand and the metal. This short-lived state presents a ~ 640 fs lifetime. The participation of low-lying CT triplet states to the formation of the MC state is discussed below in connection with the description of the early steps in the photocycle.

By exploiting the synthetic versatility of Nickel MPs that enables extensive rational design, systematic investigations using time-resolved electronic and vibrational spectroscopies have delineated generic patterns in their photocycle, which are now summarized. Following photoexcitation to the S_2 and S_1 state, the system deactivates very rapidly. Depending upon the experimental conditions (including excitation wavelength, solvent) and probing technique, S_2 and S_1 are quenched on the sub-ps time scale, with or without detection of an intermediate species, whose nature (S_1 or CT), population mechanism (IC or ISC) and lifetime remain elusive.^{47, 21, 48, 29, 30, 49, 23, 22, 50} On the ps time scale, a vibrationally-hot (d,d) state, noted (d,d)*, is formed via intramolecular energy relaxation (IVR) and conformational relaxation, as monitored by tr-OAS^{47, 22, 51, 52} and tr-RSS.^{33, 53, 54} This state thermalizes to an equilibrated (d,d) state through vibrational cooling (VC) over tens of ps. The metastable (d,d) state decays back to the ground-state on time scales ranging from few hundreds of ps to several ms, via nonradiative channels that are governed by ring-deformation modes, solute-solvent interactions and temperature.^{44, 22, 51, 52}

Combining the findings from the present femtosecond spectroscopic measurements performed in the UV-visible and X-ray spectral ranges with comparable resolution (120 fs IRF and 110 fs IRF, respectively) yields a comprehensive description of the dynamical processes across the excited-state manifold, along with robust time constants.

Considering first the tr-OAS measurements, the observations follow the general trends outlined above. Photoexcitation into the S_2 state at 400 nm induces a $1^1(\pi, \pi^*)$ transition centered on the π -ring. As seen from **Figure 1f** and **S.I.2**, this transition does not contain appreciable metal-based contribution. In **Figure 2a**, the absence of any stimulated emission feature similar to the one reported for photoexcited closed-shell porphyrins (containing Mg^{2+} or Zn^{2+}) shows that the population of S_1 is not observed with this technique and that the triplet multiplicity of the excited-state manifold is acquired on the sub-100 fs time scale through ISC. The intricate evolution of the tr-OAS spectral lineshapes (**Figure 2a** and **2b**) reflects the nonadiabatic intramolecular dynamics induced by the photoexcitation of the π -ring. This stage results in the emergence of a derivative-like feature with a broad absorbance over the red side of the bleach, which evolves to the known spectral profile unambiguously attributed to the equilibrated (d,d) state reached through the VC of the (d,d)* state (**Figure 2b** and **2c**). The global-fit analysis based on a sequential kinetic model captures this evolution and delivers three time constants of 560 ± 60 fs, 6 ± 1 ps and 210 ± 20 ps (**Figure 2d**).

Considering then the tr-XES measurements, photoexcitation into the S_2 state at 400 nm triggers the IRF-limited population of a triplet state of CT character that presents a well defined spectral lineshape. This state decays to the metastable triplet state of MC character with a time constant of 640 fs. The TD-DFT and e-DFT calculations identify several potential triplet states of CT character (**Figure 5**). **Figure 6** displays the Jablonski diagram of the observable states and the photocycle, where the time constants are established on the basis of the combined (a) optical and (b) X-ray measurements. It highlights the strong complementarity between the two spectroscopies when tracking the formation and the relaxation of the MC state. The energy of the (d,d) state has been estimated in an early measurement based on the transient grating techniques.⁵⁵ **Figure 6** also suggests similarities and differences between the photocycle of Ni porphyrins and the ones of low-Z transition metal complexes (e.g., square planar Ni complexes and octahedral d^6 complexes⁴⁴ (see **S.I.8**)

Additionally, correlating the tr-OAS and tr-XES observations later in the ps delay range provides novel insights into the nature of the (d,d)* state. As predicted by the Gouterman model illustrated in **Figure 1c**, the energetic separation between the HOMO and the HOMO-1 affects the positions and the relative intensities of the B and Q bands through C.I. and further, those of the Q(1,0) and Q(0,0) bands through vibronic coupling. Since the spectrum of the equilibrated (d,d) state is very well reproduced by a rigid shift of the GS spectrum without any spectral weight redistribution (**Figure 2e**), the energetic separation between a_{1u} and a_{2u} , the degree of C.I. and the strength of the vibronic coupling are comparable in the GS and in the equilibrated (d,d) state. From **Figures 3b, 3c** and **Figures 4e, 4f** and **S.I.6**, the tr-XES measurements show that the (d,d)* state is largely formed by 1 ps. This implies that the corresponding ESA in **Figure 3b** observed at this particular time delay can already be approximated by the typical UV-vis spectrum comprising B, Q(1,0) and Q(0,0) bands that are red-shifted. As such, the broadened peak detected at 635 nm (red star in **Figure 2b**) can be identified with the quasi Q(0,0) band of the (d,d)* state. Based on this tentative assignment, the unexpectedly large energy separation and the relative intensities of Q(1,0) and Q(0,0) evolve during the relaxation from (d,d)* to (d,d) (**Figure 2b** and **2c**). This observation then implies that, unlike for the GS and thermalized (d,d) state, the energetic separation between a_{2u} and a_{1u} , the degree of C.I. and the strength of vibronic coupling are dynamically altered in the (d,d)* state. In other words, the (d,d)* state should be viewed as a vibronically-hot state rather than a simpler vibrationally-hot state.

At the outset of the photocycle, the dynamics are driven by concurrent ISC, IC and CT. Elucidating the detailed transition pathway from the singlet FC state to the thermalized triplet (d,d) state will require further experimental studies at higher temporal resolution with assistance from advanced theoretical modeling of the underlying nonadiabatic processes, including vibronic and SOC couplings, while going beyond the sequential kinetic model.

The first key result of this work is that, within a sequential kinetic model, a triplet state of CT character mediates the nonadiabatic intramolecular dynamics leading to the formation of the metastable triplet (d,d) state of MC character. The CT character of the intermediate triplet species indicates that photoexcited Ni MPs can engage in ultrafast intermolecular CT events. Although the lifetime of the CT state is very short, it falls within the characteristic range reported for CT events in MPs arrays^{56, 57, 58} and MPs-containing COFs and MOFs.^{59, 60,}

⁶¹ It is also compatible with the time spans necessary for electron/hole extraction in

organic/inorganic nanohybrids of MPs anchored to surfaces (e.g., functionalized electrodes).^{62, 63}

The second key result of this work is that the thermalization of the (d,d)* state taking place over a few ps involves strong vibronic coupling, which transiently impacts the electronic structure of the molecule, in particular the energetic separation between the a_{2u} and a_{1u} orbitals. This observation suggests that photoexcited NiP could conceivably sustain solvent-assisted CT and even vibrationally-assisted CT under suitable conditions, provided that energy transfer (EET) to the surroundings can be partly overcome.^{64, 65}

Some questions are still hovering over the dynamics of the excited system as it evolves away from the FC region. The sub-100 fs time scale of the CT formation hints at the participation of a conical intersection.^{66, 67} However, it is not possible to determine whether the $^3(\pi,\pi^*)$ is effectively populated or not. In addition, the strength of the JT effects, which are expected to manifest for all electronically-degenerate excited states, cannot yet be assessed.⁶⁸ All these aspects are rooted in the extent of symmetry breaking and the ensuing localization of delocalized electrons and holes, in the initial ground state, intermediate excited states and metastable state⁶⁹ (**Figures 1f**, **Figure 3g**, and **Figure 5**). Such phenomena could promote the role of other low-lying triplet states, thereby complicating the simplified picture based on quasi- D_{4h} representations.

Ultimately, functionalizing the nonadiabatic intramolecular dynamics of photoexcited NiPs and other open-shell MPs for boosting the performances of photoconversion in practical applications will rely on maximizing the yield of hot charge separation, while minimizing energy losses in optimized architectures. Guidelines for the associated physico-chemical engineering will be obtained by establishing structure-property relationships that are applicable on the ultrafast time scale. They will enable correlating the rates of forward/backward hot CT and EET to the non-equilibrated electronic and geometric structures of the short-lived excited species involved in these processes. Owing to the CT character of the manifold mediating the nonadiabatic dynamics, the specific interactions implicating the first solvation shell should be considered for aiding the stabilization of the charge separation.

Elaborating this extended framework will rely on combining ultrafast X-ray emission spectroscopy, X-ray absorption spectroscopy along with diffuse wide-angle and small-angle X-ray scattering in order to provide atomically-resolved and spin-resolved kinetics on multiple length scales and time scales. This knowledge will facilitate the interpretation of the multi-exponential and probe-wavelengths-dependent kinetics that are usually introduced to describe the dynamics of nonadiabatic photoconversion processes in the UV-visible and IR spectral ranges.

Finally, the exploration of the few-ps down to the sub-ps regime is also now within reach for the ultrafast X-ray based techniques implemented at XFEL facilities and table-top setups. These studies will be assisted by the non-linear electronic and vibrational laser-based spectroscopies that already reveal ever-finer details about vibronic couplings,^{70, 71} and by the attosecond methodologies that are currently developed to track charge migration in large organic molecules.⁷²

In conclusion, this study probes the early stages of the photoinduced dynamics within the NiTMP molecule in solution using ultrafast tr-OAS and tr-XES achieving comparable femtosecond resolution. Analyzing the two types of measurements by employing a sequential kinetic model with support from TD-DFT and e-DFT calculations reveals that the population of the thermalized metastable state of MC character proceeds via the

mediation of a short-lived intermediate state of CT character. This feature should be generally observable in open-shell MPs, setting them apart from closed-shell MPs, where the excited-state manifold remains LC. Besides completing the description of the photocycle for this important representative molecule by delivering robust assignment and time constants, the present findings establish the strong potential of photoexcited NiP and other open-shell MPs for joining the classes of low-Z photosensitizers, for which innovative chemical engineering can prolong the lifetime of the CT state. Considering the distinctive formation and deactivation mechanisms outlined in this study,, the principles governing the sub-picosecond interplay between the CT and LC characters are expected to be specific to this family.

Moreover, the vibronically-coupled nature of the thermalization dynamics in the MC manifold brings further opportunities for promoting and stabilizing charge separation through the interaction between the building blocks and with the surroundings (solvent or surface) on longer time scales. The comprehensive exploration of the transient properties afforded by the CT manifold of open-shell MPs will require fostering synergies between complementary ultrafast techniques that should be combined in order to realize an exhaustive mapping of the spin, electronic and structural dynamics on multiple length scales down to the attosecond regime of charge localization and migration. Uncovering the underlying structure-properties relationships for the CT manifold of open-shell MPs will lead to the design of non-equilibrated photoreactants that can engage in hot CT steps. Their functionalization will produce the novel generations of hybrid molecular architectures capable of boosting the performances reachable by photoconversion schemes out of equilibrium.

ASSOCIATED CONTENT

Supporting Information

S.I.1. Synthesis of NiTMP, S.I.2. Theoretical calculations and modeling, S.I.3. Transient optical absorption spectroscopy measurements, S.I.4. Transient X-ray emission spectroscopy measurements, S.I.5. Excitation conditions in the transient X-ray emission spectroscopy measurements, S.I.6. Power dependencies, S.I.7. Global spectral analysis of the photoinduced dynamics in the X-ray range, S.I.8. Observation of CT states in low-Z transition metal complexes.

AUTHOR INFORMATION

Corresponding Author

* socant@kemi.dtu.dk, kaibo.zheng@chemphys.lu.se, max.lawson@unige.ch

Author Contributions

SEC designed the study and the associated experiments. AR, MAK performed the synthesis of the molecular system and AR, MAK, YL, QP performed its chemical characterization. GP and MLD performed the DFT, TD-DFT and e-DFT calculations with the ADF software. MP performed the DFT and TD-DFT calculations with the ORCA software. HG, WL, YL, QZ, KZ conducted the tr-OAS experiments. HG, KZ, SEC performed the tr-OAS analysis. MAN, MB, KK, AK, HG, JM, ML, AR, MAb, SC, FAL, PZ, WG, CB, DK, KZ, SEC conducted the tr-XES experiments. MAN performed the tr-XES analysis with feedback from MB, DK, SEC. SEC

wrote the manuscript, with contributions from MAN, GP, MB, MP, DK, MLD, KZ. All authors have given approval to the final version of the manuscript.

ACKNOWLEDGMENT

The authors gratefully acknowledge European XFEL in Schenefeld, Germany, for provision of X-ray free-electron laser beamtime at FXE and thank the instrument group and facility staff for their expert assistance. This research was supported in part through the Maxwell computational resources operated at Deutsches Elektronen-Synchrotron DESY, Hamburg, Germany. This work received funding from the VILLUM FONDEN (VIL50350), and the Swedish Research Council No. 2021-05319 (KZ). This work received funding from the Hungarian National Research, Development and Innovation Fund under Grant Nos. NKFIH FK 145967 (MP) and TKP2021-NVA-04 (MP). KK gratefully acknowledges funding by the DFG within the program "Sachbeihilfe" project ID 497431350 (KU 4184/1-1). CB and KK acknowledge support by the Deutsche Forschungsgemeinschaft (DFG) via the Cluster of Excellence 'Advanced Imaging of Matter', EXC 2056, Project ID 390715994, and via SFB925 ID 170620586 (TP A4). WG acknowledges funding from Spanish MIU through "Ayudas Beatriz Galindo" (BEAGAL18/00092), Comunidad de Madrid and Universidad Autónoma de Madrid through "Proyecto de I+D para Investigadores del Programa Beatriz Galindo" (SI2/PBG/2020-00003), Spanish MICIU through Proyecto de I+D+i 2019 (PID2019-108678GB-I00) and IMDEA-Nanociencia support from the Severo Ochoa Programme for Centers of Excellence in R&D (MINECO, grant CEX2020-001039-S). W.G. further acknowledges partial support from the National Science Centre in Poland under SONATA BIS 6 Grant 2016/22/E/ST4/00543. DG acknowledges support from the Center for Nanoscale Materials, a U.S. Department of Energy Office of Science User Facility, supported by the U.S. DOE, Office of Basic Energy Sciences, under Contract No. DE-AC02-06CH11357. MP acknowledges support from the János Bolyai Scholarship of the Hungarian Academy of Sciences. Guest access for MP to the central HPC cluster of the Technical University of Denmark is acknowledged. The computations performed by GP and MLD were performed at the University of Geneva on the "Baobab" and "Yggdrasil" HPC clusters.

REFERENCES

- (1) Scheer, H. *Chlorophylls*; Springer, 1991.
- (2) Labbé, R. F.; Vreman, H. J.; Stevenson, D. K. Zinc Protoporphyrin: A Metabolite with a Mission. *Clinical Chemistry* **1999**, 45(12), 2060–2072. DOI:[10.1093/clinchem/45.12.2060](https://doi.org/10.1093/clinchem/45.12.2060).
- (3) Ko, Y. J.; Lee, M.-E.; Cho, B.-H.; Kim, M.; Hyeon, J. E.; Han, J. H.; Han, S. O. Bioproduction of Porphyrins, Phycobilins, and Their Proteins Using Microbial Cell Factories: Engineering, Metabolic Regulations, Challenges, and Perspectives. *Critical Reviews in Biotechnology* **2023**, 1–15. DOI:[10.1080/07388551.2023.2168512](https://doi.org/10.1080/07388551.2023.2168512)
- (4) Batinic-Haberle, I.; Tovmasyan, A.; Spasojevic, I. An Educational Overview of the Chemistry, Biochemistry and Therapeutic Aspects of Mn Porphyrins – from Superoxide Dismutation to H₂O₂-Driven Pathways. *Redox Biology* **2015**, 5, 43–65. DOI:[10.1016/j.redox.2015.01.017](https://doi.org/10.1016/j.redox.2015.01.017).
- (5) Shaik, S.; Kshatresh Dutta Dubey. Nanomachines in Living Matters: The Soft-Robot Cytochrome P450. *Trends in Chemistry* **2023**, 5(10), 763–774. DOI:[10.1016/j.trechm.2023.07.002](https://doi.org/10.1016/j.trechm.2023.07.002).

- (6) Rothemund, P. Formation of Porphyrins from Pyrrole and Aldehydes. *Journal of the American Chemical Society* **1935**, 57 (10), 2010–2011. [DOI:10.1021/ja01313a510](https://doi.org/10.1021/ja01313a510).
- (7) Rothemund, P. A New Porphyrin Synthesis. The Synthesis of Porphin 1. *Journal of the American Chemical Society* **1936**, 58 (4), 625–627. [DOI:10.1021/ja01295a027](https://doi.org/10.1021/ja01295a027).
- (8) Woodward, R. B.; Ayer, W. A.; Beaton, J. M.; Bickelhaupt, F.; Bonnett, R.; Buchschacher, P.; Closs, G. L.; Dutler, H.; Hannah, J.; Hauck, F. P.; Itô, S.; Langemann, A.; Le Goff, E.; Leimgruber, W.; Lwowski, W.; Sauer, J.; Valenta, Z.; Volz, H. The Total Synthesis of Chlorophyll. *Journal of the American Chemical Society* **1960**, 82(14), 3800–3802. [DOI:10.1021/ja01499a093](https://doi.org/10.1021/ja01499a093).
- (9) Heimgartner, H.; Hansen, H.-J. Struktur Und Mechanismus in Der Organischen Chemie Im Spiegel Von Helvetica Chimica Acta. **1992**, 75 (2), 359–437. [DOI:10.1002/hlca.19920750202](https://doi.org/10.1002/hlca.19920750202).
- (10) Eschenmoser, A.; Wintner, C. E. Natural Product Synthesis and Vitamin B 12. *Science* **1977**, 196 (4297), 1410–1420. [DOI:10.1126/science.867037](https://doi.org/10.1126/science.867037).
- (11) Hiroto, S.; Miyake, Y.; Shinokubo, H. Synthesis and Functionalization of Porphyrins through Organometallic Methodologies. *Chemical Reviews* **2016**, 117 (4), 2910–3043. [DOI:10.1021/acs.chemrev.6b00427](https://doi.org/10.1021/acs.chemrev.6b00427).
- (12) Sitte, E.; Senge, M. O. The Red Color of Life Transformed - Synthetic Advances and Emerging Applications of Protoporphyrin IX in Chemical Biology. *European Journal of Organic Chemistry* **2020**, 2020(22), 3171–3191. [DOI:10.1002/ejoc.202000074](https://doi.org/10.1002/ejoc.202000074).
- (13) Senge, M. O.; Sergeeva, N. N.; Hale, K. J. Classic Highlights in Porphyrin and Porphyrinoid Total Synthesis and Biosynthesis. *Chemical Society Reviews* **2021**, 50 (7), 4730–4789. [DOI:10.1039/c7cs00719a](https://doi.org/10.1039/c7cs00719a).
- (14) Ostfeld, D.; Tsutsui, M. Novel Metalloporphyrins. Syntheses and Implications. *Accounts of Chemical Research* **1974**, 7 (2), 52–58. [DOI:10.1021/ar50074a004](https://doi.org/10.1021/ar50074a004).
- (15) J. W. Buchler Synthesis and properties of metalloporphyrins, The Porphyrins VI: Structure and Synthesis, Part A, Part 1 (1978).
- (16) D. Dolphin, The Porphyrins, Academic Press, New York (1978).
- (17) K. M. Kadish, K. M. Smith and R. Guilard, The Porphyrin Handbook, Academic Press, New York, p. 2003 (2000).
- (18) Kadish, K. M.; Smith, K. M.; Guilard, R. *Handbook of Porphyrin Science (Volumes 6 – 10): With Applications to Chemistry, Physics, Materials Science, Engineering, Biology and Medicine*; World Scientific, **2010**.
- (19) Imahori, H.; Mori, Y.; Matano, Y. Nanostructured Artificial Photosynthesis. *Journal of Photochemistry and Photobiology C: Photochemistry Reviews* **2003**, 4 (1), 51–83. [DOI:10.1016/S1389-5567\(03\)00004-2](https://doi.org/10.1016/S1389-5567(03)00004-2).
- (20) Park, J. M.; Hong, K.-I.; Lee, H.; Jang, W.-D. Bioinspired Applications of Porphyrin Derivatives. *Accounts of Chemical Research* **2021**, 54 (9), 2249–2260. [DOI:10.1021/acs.accounts.1c00114](https://doi.org/10.1021/acs.accounts.1c00114).
- (21) Bräm, O.; Cannizzo, A.; Chergui, M. Ultrafast Broadband Fluorescence Up-Conversion Study of the Electronic Relaxation of Metalloporphyrins. *The Journal of Physical Chemistry A* **2019**, 123 (7), 1461–1468. [DOI:10.1021/acs.jpca.9b00007](https://doi.org/10.1021/acs.jpca.9b00007).

(22) Zhang, X.; Wasinger, E. C.; Mureşan, A.; Attenkofer, K.; Jennings, G.; Lindsey, J. S.; Chen, L. X. Ultrafast Stimulated Emission and Structural Dynamics in Nickel Porphyrins. *The Journal of Physical Chemistry A* **2007**, 111 (46), 11736–11742.

[DOI:10.1021/jp0751763](https://doi.org/10.1021/jp0751763).

(23) Pattengale, B.; Liu, Q.; Hu, W.; Yang, S.; He, P.; Sir Lawrence Tender; Wang, Y.; Zhang, X.; Zhou, Z.; Zhang, J.; Huang, J. Selective Excited-State Dynamics in a Unique Set of Rationally Designed Ni Porphyrins. *The Journal of Physical Chemistry C* **2019**, 123 (29), 17994–18000.

[DOI:10.1021/acs.jpcc.9b05413](https://doi.org/10.1021/acs.jpcc.9b05413).

(24) Findsen, E. W.; Shelnutt, J. A.; Ondrias, M. R. Photodynamics of Nickel-Porphyrins in Noncoordinating Solvents: Characterization of d-d Excited States Using Transient Raman Spectroscopy. *The Journal of Physical Chemistry* **1988**, 92 (2), 307–314.

[DOI:10.1021/j100313a014](https://doi.org/10.1021/j100313a014).

(25) Kruglik, S. G.; Mizutani, Y.; Kitagawa, T. Time-Resolved Resonance Raman Study of the Primary Photoprocesses of Nickel(II) Octaethylporphyrin in Solution. *Chemical Physics Letters* **1997**, 266 (3-4), 283–289. [DOI:10.1016/S0009-2614\(97\)00008-0](https://doi.org/10.1016/S0009-2614(97)00008-0).

(26) Mizutani, Y.; Uesugi, Y.; Kitagawa, T. Intramolecular Vibrational Energy Redistribution and Intermolecular Energy Transfer in the (d,d) Excited State of Nickel Octaethylporphyrin. *The Journal of Chemical Physics* 1999, 111 (19), 8950–8962. [DOI:10.1063/1.480253](https://doi.org/10.1063/1.480253).

(27) Chen, L. X. Capturing a Photoexcited Molecular Structure through Time-Domain X-Ray Absorption Fine Structure. *Science* **2001**, 292 (5515), 262–264. [DOI:10.1126/science.1057063](https://doi.org/10.1126/science.1057063).

(28) Chen, L. X.; Zhang, X.; Wasinger, E. C.; Klaus Attenkofer; Jennings, G.; Muresan, A. Z.; Lindsey, J. S. Tracking Electrons and Atoms in a Photoexcited Metalloporphyrin by X-Ray Transient Absorption Spectroscopy. *Journal of the American Chemical Society* **2007**, 129 (31), 9616–9618. [DOI:10.1021/ja072979v](https://doi.org/10.1021/ja072979v).

(29) Shelby, M. L.; Mara, M. W.; Chen, L. X. New Insight into Metalloporphyrin Excited State Structures and Axial Ligand Binding from X-Ray Transient Absorption Spectroscopic Studies. *Coordination Chemistry Reviews* **2014**, 277-278, 291–299. [DOI:10.1016/j.ccr.2014.05.025](https://doi.org/10.1016/j.ccr.2014.05.025).

(30) Shelby, M. L.; Lestrangle, P. J.; Jackson, N. E.; Haldrup, K.; Mara, M. W.; Stickrath, A. B.; Zhu, D.; Lemke, H. T.; Chollet, M.; Hoffman, et al. Ultrafast Excited State Relaxation of a Metalloporphyrin Revealed by Femtosecond X-Ray Absorption Spectroscopy. *Journal of the American Chemical Society* **2016**, 138 (28), 8752–8764. [DOI:10.1021/jacs.6b02176](https://doi.org/10.1021/jacs.6b02176).

(31) Della-Longa, S.; Chen, L. X.; Frank, P.; Hayakawa, K.; Hatada, K.; Benfatto, M. Direct Deconvolution of Two-State Pump-Probe X-Ray Absorption Spectra and the Structural Changes in a 100 Ps Transient of Ni(II)-Tetramesitylporphyrin. *Inorganic Chemistry* **2009**, 48 (9), 3934–3942. [DOI:10.1021/ic8008943](https://doi.org/10.1021/ic8008943).

(32) Ryland, E. S.; Zhang, K.; Vura-Weis, J. Sub-100 Fs Intersystem Crossing to a Metal-Centered Triplet in Ni(II)OEP Observed with M-Edge XANES. *Journal of Physical Chemistry A* **2019**, 123 (25), 5214–5222. [DOI:10.1021/acs.jpca.9b03376](https://doi.org/10.1021/acs.jpca.9b03376).

(33) Schindler, J.; Kupfer, S.; Ryan, A. A.; Flanagan, K. J.; Dietzek, B. Sterically Induced Distortions of Nickel(II) Porphyrins – Comprehensive Investigation by DFT Calculations and Resonance Raman Spectroscopy. *Coordination Chemistry Reviews* 2018, 360, 1–16. [DOI:10.1016/j.ccr.2017.12.014](https://doi.org/10.1016/j.ccr.2017.12.014).

- (34) The relative ordering of a_{2u} and a_{1u} as HOMO and HOMO-1 depends on substituents introduced in the porphyrin ring.
- (35) Gouterman, M. Study of the Effects of Substitution on the Absorption Spectra of Porphin. *The Journal of Chemical Physics* **1959**, 30(5), 1139–1161. [DOI:10.1063/1.1730148](https://doi.org/10.1063/1.1730148).
- (36) Gouterman, M. Spectra of Porphyrins. *Journal of Molecular Spectroscopy* **1961**, 6, 138–163. [DOI:10.1016/0022-2852\(61\)90236-3](https://doi.org/10.1016/0022-2852(61)90236-3).
- (37) Rodriguez, J. J.; Holten, D. Ultrafast Vibrational Dynamics of a Photoexcited Metalloporphyrin. 1989, 91 (6), 3525–3531. DOI:10.1063/1.456883.
- (38) Rodríguez, J.; Kirmaier, C.; Holten, D. Time-Resolved and Static Optical Properties of Vibrationally Excited Porphyrins. *Journal of Chemical Physics* **1991**, 94 (9), 6020–6029. [DOI:10.1063/1.460439](https://doi.org/10.1063/1.460439).
- (39) Mizutani, Y.; Kitagawa, T. A Role of Solvent in Vibrational Energy Relaxation of Metalloporphyrins. *Journal of Molecular Liquids* **2001**, 90 (1-3), 233–242. [DOI:10.1016/s0167-7322\(01\)00126-x](https://doi.org/10.1016/s0167-7322(01)00126-x).
- (40) Galler, A.; Gawelda, W.; Biednov, M.; Bömer, C.; Britz, A.; Brockhauser, S.; Choi, T.-K.; Diez, M.; Frankenberger, P.; French, M. T.; et al. Scientific Instrument Femtosecond X-Ray Experiments (FXE): Instrumentation and Baseline Experimental Capabilities. *Journal of Synchrotron Radiation* **2019**, 26 (5), 1432–1447. [DOI:10.1107/s1600577519006647](https://doi.org/10.1107/s1600577519006647).
- (41) Khakhulin, D.; Otte, F.; Biednov, M.; Bömer, C.; Choi, T.-K.; Diez, M.; Galler, A.; Jiang, Y.; Kubicek, K.; Lima, et al. C. Ultrafast X-Ray Photochemistry at European XFEL: Capabilities of the Femtosecond X-Ray Experiments (FXE) Instrument. *Applied Sciences* **2020**, 10 (3), 995. [DOI:10.3390/app10030995](https://doi.org/10.3390/app10030995).
- (42) Lima, F. A.; Otte, F.; Vakili, M.; Ardana-Lamas, F.; Biednov, M.; F Dall'Antonia; Frankenberger, P.; Gawelda, W.; Gelisio, L.; Han, H. et al. Experimental Capabilities for Liquid Jet Samples at Sub-MHz Rates at the FXE Instrument at European XFEL. *Journal of Synchrotron Radiation* **2023**, 30 (6), 1168–1182. [DOI:10.1107/s1600577523008159](https://doi.org/10.1107/s1600577523008159).
- (43) Biednov, M.; Yousef, H.; Otte, F.; Choi, T.-K. Jiang, Y.; Frankenberger, P.; Knoll, M.; Zalden, P.; Ramilli, M.; Gawelda, W.; et al. Hard X-Ray Emission Spectroscopy in Liquids Using MHz XFEL Source and JUNGFRU Detectors. *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment* **2023**, 1055, 168540–168540. [DOI:10.1016/j.nima.2023.168540](https://doi.org/10.1016/j.nima.2023.168540).
- (44) Sinha, N.; Wenger, O. S. Photoactive Metal-To-Ligand Charge Transfer Excited States in 3d6 Complexes with Cr^0 , Mn^I , Fe^{II} , and Co^{III} . *Journal of the American Chemical Society* **2023**, 145 (9), 4903–4920 [DOI:10.1021/jacs.2c13432](https://doi.org/10.1021/jacs.2c13432).
- (45) Kawai, J.; Suzuki, C.; Adachi, H.; Konishi, T.; Gohshi, Y. Charge-Transfer Effect on the Linewidth of $\text{FeK}\alpha$ X-Ray Fluorescence Spectra. *Physical Review B* **1994**, 50 (16), 11347–11354. [DOI:10.1103/physrevb.50.11347](https://doi.org/10.1103/physrevb.50.11347).
- (46) Cheng, C.-L.; Wu, Q.; Van Voorhis, T. Rydberg Energies Using Excited State Density Functional Theory. *The Journal of Chemical Physics* **2008**, 129 (12), 124112. [DOI:10.1063/1.2977989](https://doi.org/10.1063/1.2977989).
- (47) Michael, C.; Gentemann, S.; Roberts, J. A.; Nelson, N. Y.; Medforth, C. J.; Jia, S.; Smith, K. M.; Fajer, J.; Shelnutt, J. A.; Holten, D. Picosecond to Microsecond Photodynamics of a

Nonplanar Nickel Porphyrin: Solvent Dielectric and Temperature Effects. **1998**, 120 (15), 3781–3791. [DOI:10.1021/ja974101h](https://doi.org/10.1021/ja974101h).

(48) Sorgues; S., Poisson, L.; Raffael, K. D.; Krim, L.; Soep, B. ; Shafizadeh; N. Femtosecond Electronic Relaxation of Excited Metalloporphyrins in the Gas Phase. *Journal of Chemical Physics* **2006**, 124 (11). [DOI:10.1063/1.2176612](https://doi.org/10.1063/1.2176612).

(49) Ryland, E. S.; Zhang, K.; Vura-Weis, J. Sub-100 Fs Intersystem Crossing to a Metal-Centered Triplet in Ni(II)OEP Observed with M-Edge XANES. *Journal of Physical Chemistry A* **2019**, 123 (25), 5214–5222. [DOI:10.1021/acs.jpca.9b03376](https://doi.org/10.1021/acs.jpca.9b03376).

(50) Zamyatin, A. V.; Gusev, A. V.; Michael. Two-Pump–One-Probe Femtosecond Studies of Ni(II) Porphyrins Excited States. *Journal of the American Chemical Society* 2004, 126 (49), 15934–15935. [DOI:10.1021/ja0448059](https://doi.org/10.1021/ja0448059).

(51) Rodriguez, J. J.; Holten, D. Ultrafast Vibrational Dynamics of a Photoexcited Metalloporphyrin. **1989**, 91 (6), 3525–3531. [DOI:10.1063/1.456883](https://doi.org/10.1063/1.456883).

(52) Rodríguez, J.; Kirmaier, C.; Holten, D. Time-Resolved and Static Optical Properties of Vibrationally Excited Porphyrins. *Journal of Chemical Physics* **1991**, 94 (9), 6020–6029. [DOI:10.1063/1.460439](https://doi.org/10.1063/1.460439).

(53) Rush, T. S.; Kozlowski, P. M.; Piffat, C. A.; Ranjit Kumble; and; Spiro, T. G. Computational Modeling of Metalloporphyrin Structure and Vibrational Spectra: Porphyrin Ruffling in NiTPP. *Journal of Physical Chemistry B* **2000**, 104 (20), 5020–5034. [DOI:10.1021/jp000266s](https://doi.org/10.1021/jp000266s).

(54) Huang, Q.; Medforth, C. J.; Reinhard Schweitzer-Stenner. Nonplanar Heme Deformations and Excited State Displacements in Nickel Porphyrins Detected by Raman Spectroscopy at Soret Excitation. *Journal of Physical Chemistry A* **2005**, 109 (46), 10493–10502. [DOI:10.1021/jp052986a](https://doi.org/10.1021/jp052986a).

(55) Brodard, P.; Vauthey, E. Determination of the Energy of the (d,d) Electronic State Involved in the Radiationless Deactivation of Photoexcited Nickel Tetraphenylporphine. *Chemical Physics Letters* **1999**, 309 (3-4), 198–208.

[DOI:10.1016/s0009-2614\(99\)00679-x](https://doi.org/10.1016/s0009-2614(99)00679-x).

(56) Aratani, N.; Atsuhiko Osuka; Cho, H.-J.; Kim, D.-H. Photochemistry of Covalently-Linked Multi-Porphyrinic Systems. 2002, 3(1), 25–52. [DOI:10.1016/s1389-5567\(02\)00003-5](https://doi.org/10.1016/s1389-5567(02)00003-5).

(57) Park, J. M.; Hong, K.-I.; Lee, H.; Jang, W.-D. Bioinspired Applications of Porphyrin Derivatives. *Accounts of Chemical Research* **2021**, 54 (9), 2249–2260. [DOI:10.1021/acs.accounts.1c00114](https://doi.org/10.1021/acs.accounts.1c00114).

(58) Imahori, H. Giant Multiporphyrin Arrays as Artificial Light-Harvesting Antennas. *The Journal of Physical Chemistry B* 2004, 108(20), 6130–6143. [DOI:10.1021/jp038036b](https://doi.org/10.1021/jp038036b).

(59) Chen, R.; Wang, Y.; Ma, Y.; Mal, A.; Gao, X.-Y.; Gao, L.; Qiao, L.; Li, X.-B.; Wu, L.-Z.; Wang, C. Rational Design of Isostructural 2D Porphyrin-Based Covalent Organic Frameworks for Tunable Photocatalytic Hydrogen Evolution. *Nature Communications* **2021**, 12 (1), 1354. [DOI:10.1038/s41467-021-21527-3](https://doi.org/10.1038/s41467-021-21527-3).

(60) Feng, L.; Wang, K.-Y.; Joseph, E.; Zhou, H.-C. Catalytic Porphyrin Framework Compounds. *Trends in Chemistry* **2020**, 2 (6), 555–568. [DOI:10.1016/j.trechm.2020.01.003](https://doi.org/10.1016/j.trechm.2020.01.003).

(61) Zhang, X.; Wasson, M. C.; Shayan, M.; Berdichevsky, E. K.; Ricardo-Noordberg, J.; Singh, Z.; Papazyan, E. K.; Castro, A. J.; Marino, P.; Ajoyan, et al. A Historical Perspective on

- Porphyrin-Based Metal–Organic Frameworks and Their Applications. *Coordination Chemistry Reviews* **2021**, 429, 213615. [DOI:10.1016/j.ccr.2020.213615](https://doi.org/10.1016/j.ccr.2020.213615).
- (62) Tomokazu Umeyama; Hiroshi Imahori. Self-Organization of Porphyrins and Fullerenes for Molecular Photoelectrochemical Devices. *Photosynthesis Research* 2006, 87 (1), 63–71. [DOI:10.1007/s11120-005-4632-z](https://doi.org/10.1007/s11120-005-4632-z).
- (63) Sekhar, A. R.; Chitose, Y.; Janoš, J.; Dangoor, S. I.; Ramundo, A.; Satchi-Fainaro, R.; Slavíček, P.; Klán, P.; Weinstein, R. Porphyrin as a Versatile Visible-Light-Activatable Organic/Metal Hybrid Photoremovable Protecting Group. *Nature Communications* **2022**, 13 (1), 3614. [DOI:10.1038/s41467-022-31288-2](https://doi.org/10.1038/s41467-022-31288-2).
- (64) Wardlaw, D.M. and Marcus, R. A. On the Statistical Theory of Unimolecular Processes. *Adv. Chem. Phys.* **1988**, 70, 231. DOI: [10.1002/9780470141199.ch7](https://doi.org/10.1002/9780470141199.ch7)
- (65) Marcus, R. A. Solvent Dynamics: Modified Rice–Ramsperger–Kassel–Marcus Theory. II. Vibrationally Assisted Case. *Journal of Chemical Physics* 1996, 105 (13), 5446–5454. [DOI:10.1063/1.472385](https://doi.org/10.1063/1.472385).
- (66) Foster, P. W.; Jonas, D. M. Nonadiabatic Conical Nodes Are near but Not at an Elliptical Conical Intersection. *Chemical Physics* **2019**, 520, 108–121. [DOI:10.1016/j.chemphys.2019.01.004](https://doi.org/10.1016/j.chemphys.2019.01.004).
- (67) K. Falahati; Hamerla, C.; Huix-Rotllant, M.; Burghardt, I. Ultrafast Photochemistry of Free-Base Porphyrin: A Theoretical Investigation of B → Q Internal Conversion Mediated by Dark States. *Physical Chemistry Chemical Physics* **2018**, 20 (18), 12483–12492. [DOI:10.1039/c8cp00657a](https://doi.org/10.1039/c8cp00657a).
- (68) Seth, J.; Bocian, D. F. Electron Paramagnetic Resonance Studies of Metalloporphyrin Anion Radicals. Effects of Solvent, Counterion, Temperature, and Isotopic Substitution on the Jahn-Teller Active 2Eg Ground State. *Journal of the American Chemical Society* 1994, 116 (1), 143–153. [DOI:10.1021/ja00080a017](https://doi.org/10.1021/ja00080a017).
- (69) Kuleff, A. I.; Lünemann, S.; Cederbaum, L. S. Ultrafast Reorganization of the Hole Charge Created upon Outer-Valence Ionization of Porphyrins. *Chemical Physics* **2012**, 399, 245–251. [DOI:10.1016/j.chemphys.2011.10.015](https://doi.org/10.1016/j.chemphys.2011.10.015).
- (70) Biswas, S.; Kim, J.; Zhang, X.; Scholes, G. D. Coherent Two-Dimensional and Broadband Electronic Spectroscopies. *Chemical Reviews* **2022**, 122 (3), 4257–4321. [DOI:10.1021/acs.chemrev.1c00623](https://doi.org/10.1021/acs.chemrev.1c00623).
- (71) Prince, R. C.; Frontiera, R. R.; Potma, E. O. Stimulated Raman Scattering: From Bulk to Nano. *Chemical Reviews* **2017**, 117 (7), 5070–5094. [DOI:10.1021/acs.chemrev.6b00545](https://doi.org/10.1021/acs.chemrev.6b00545).
- (72) Wörner, H. J.; Arrell, C. A.; Banerji, N.; Cannizzo, A.; Chergui, M.; Das, A. K.; Hamm, P.; Keller, U.; Kraus, P. M.; Liberatore, et al. Charge Migration and Charge Transfer in Molecular Systems. *Structural Dynamics* **2017**, 4 (6), 061508. [DOI:10.1063/1.4996505.s](https://doi.org/10.1063/1.4996505.s).




FIGURE1 (a) Chemical structure of NiTMP. (b) Geometric structure for the ground state of NiTMP in toluene obtained using DFT optimizations in C_{2v} symmetry with the PBE functional. (c) The Gouterman model and the associated molecular states for P^{2-} and NiTMP. The occupied a_{2u} HOMO and a_{1u} HOMO-1 are shown in red, the unoccupied LUMOs e_g are shown in purple. The ligand centered (LC) states of singlet multiplicity are indicated as solid black lines, the LC states of triplet multiplicity are indicated as dashed black lines. The states of ligand-to-metal charge transfer (LMCT), metal-to-ligand charge transfer (MLCT) character and metal-center (MC) character are indicated as dashed gray lines. The (π, π^*) transitions centered on the ligand π -ring dominate the absorption spectrum in the UV-visible. They are indicated by the orange arrows. (d) Experimental absorption spectrum of NiTMP in toluene. The inset zooms on the Q band. (e) Theoretical absorption spectrum of NiTMP in toluene (using TD-DFT with the PBE functional). The inset zooms on the Q band. (f) Dominant occupied NTO \rightarrow virtual NTO transitions depicting the main electronic transitions underlying the Soret band, along with their weights.




FIGURE2 Transient optical absorption spectra for given time delays Δt across the (a) sub-picosecond, (b) tens of picosecond and (c) sub-nanosecond temporal windows. The negative of the UV-vis absorbance is also displayed after scaling in panel a (gray line). (d) Three components extracted from a global-fit analysis using a sequential model based on three time constants. (e) Transient optical absorption spectrum observed for a time delay Δt of 50 ps (open squares) and the model trace (solid line) (see main text). (f) Kinetics at 450 nm, 500 nm, 600 nm and 700 nm and their fits extracted from the global-fit analysis (see main text).




FIGURE3 The tr-XES measurements for NiTMP in toluene excited at 400 nm. (a) $K\alpha_{1,2}$ spectrum and (b) $K\beta$ spectrum for the ground state of NiTMP toluene. (c) Transient difference $K\alpha_{1,2}$ and (d) $K\beta$ tr-XES spectra for pump-probe delays Δt of 1 ps (green) and 50 ps (blue). The shaded areas around the lines indicate the experimental error bars (rms noise). (e) Normalized $K\alpha_{1,2}$ and (f) $K\beta$ kinetics of the XES FOM on the sub-nanosecond time scale taken over two runs under low fluence laser excitation with their single-exponential decay fit (solid and dotted lines, see main text). (g) Relaxed TD-DFT structure of the lowest triplet state 1^3A_2 . The adiabatic energy difference ΔE and the average bond length elongation ΔR compared to the singlet ground state is $\Delta R = 0.086$ Å.




FIGURE4 (a) Difference $K\alpha_{1,2}$ spectra and (b) $K\beta$ spectra for NiTMP in toluene photoexcited at 400 nm for different pump-probe delays. (c) $K\alpha_{1,2}$ and (d) $K\beta$ kinetics (open squares) and their single exponential fits (solid lines) for the rise (see main text). (e) The A_1 and A_2 components and (f) the B_1 and B_2 components extracted from global fitting with 2-point rebinning (see main text).





FIGURE5 Relaxed TD-DFT structures of the low-lying (a) $1\ ^3A_1$ triplet state of MLCT character and (b) $2\ ^3A_2$ of LMCT character in C_{2v} symmetry. The adiabatic energy difference ΔE and the average bond elongation ΔR are indicated, along with the main NTO \rightarrow NTO transitions and their weights. Relaxed e-DFT structures of the low-lying (c) $3\ ^3A_1$ of dominant LMCT character and (d) $2\ ^3A_2$ triplet state of mixed LMCT / MC in C_{2v} symmetry. The adiabatic energy difference ΔE and the average bond elongation ΔR are indicated, along with the MOs involved in the transitions.

(a) Transient optical absorption spectroscopy



(b) Transient X-ray emission spectroscopy




FIGURE6 (a) Jablonski diagram and photocycle with (a) the tr-OAS and (b) tr-XES $K\alpha_{1,2}$ and $K\beta$ times scales.