

Constraints for twist-two alien operators in QCD

G. Falcioni^{a,b}, F. Herzog^c, S. Moch^d and S. Van Thurenhout^e

^a*Dipartimento di Fisica, Università di Torino,
Via Pietro Giuria 1, Torino 10125, Italy*

^b*Physik-Institut, Universität Zürich,
Winterthurerstrasse 190, Zürich 8057, Switzerland*

^c*Higgs Centre for Theoretical Physics, School of Physics and Astronomy,
The University of Edinburgh,
Edinburgh EH9 3FD, Scotland, U.K.*

^d*II. Institute for Theoretical Physics, Hamburg University,
Luruper Chaussee 149, Hamburg D-22761, Germany*

^e*HUN-REN Wigner Research Centre for Physics,
Konkoly-Thege Miklós u. 29–33, Budapest 1121, Hungary*

E-mail: giulio.falcioni@physik.uzh.ch, ferzog@nikhef.nl,
sven-olaf.moch@desy.de, sam.van.thurenhout@wigner.hun-ren.hu

ABSTRACT: Parton evolution equations in QCD are controlled by the anomalous dimensions of gauge-invariant twist-two spin- N quark and gluon operators. Under renormalization, these mix with gauge-variant operators of the same quantum numbers, referred to as alien operators. Our work addresses the systematic study of these alien operators at arbitrary spin N , using generalized BRST symmetry relations to derive their couplings and Feynman rules at all values of N . We observe how the all- N structure of the generalized (anti-)BRST constraints relates the couplings of alien operators with $n + 1$ gluons to those with n gluons. Realizing a bootstrap, we present all one-loop results necessary for performing the operator renormalization up to four loops in QCD.

KEYWORDS: Higher-Order Perturbative Calculations, Parton Distributions, Renormalization and Regularization

ARXIV EPRINT: [2409.02870](https://arxiv.org/abs/2409.02870)

Contents

1	Introduction	1
2	Setting the stage	2
3	Identities among the alien couplings	8
3.1	Class II couplings	8
3.2	Class III couplings	10
3.3	Class IV couplings	13
4	Feynman rules of alien operators	19
4.1	Ghost operators	19
4.2	Alien gluon operators	21
4.3	Alien quark operators	24
5	Conclusions	25

1 Introduction

The study of twist-two operators of spin- N for quarks and gluons in quantum chromodynamics (QCD) and their renormalization dates to the origins of QCD as the gauge theory of the strong interaction [1–6]. The renormalization of off-shell operator matrix elements (OMEs) in QCD, i.e. Green’s functions with off-shell external momenta and insertions of these quark and gluon operators, gives access to their anomalous dimensions. These coincide with the Mellin transforms of the standard QCD splitting functions, that govern the scale evolution of the parton distribution functions. It is well-known that the twist-two operators of spin- N mix under renormalization with a set of gauge-variant operators of the same quantum numbers, which involve equation-of-motion (EOM) and ghost operators. The latter, often referred to in summary as *alien* operators, can be constructed systematically, by employing a generalized gauge symmetry of the QCD Lagrangian in covariant gauge with the addition of the physical quark and gluon operators [1, 4, 7, 8]. The generalized gauge symmetry can be promoted to a generalized BRST (gBRST) symmetry [4, 8]. This provides an algebraic approach for the derivation of a complete set of operators to be considered in the renormalization of the off-shell OMEs at a given loop order in perturbative QCD in an expansion in the strong coupling g_s , $\alpha_s = g_s^2/(4\pi)$. The complete set of operators required up to four loops has been listed in [8, 9].

Each alien operator features a coupling constant that can be interpreted as the renormalization constant that generates mixing of the gauge-invariant operators into each alien. In order to renormalize the physical OMEs, these coupling constants must be computed order-by-order in perturbation theory. The required couplings to renormalize the two-loop OMEs were computed in [1, 7] in closed form for all values of N . A method to determine the alien counterterms, i.e. the Feynman rules obtained by summing all the alien operators with their associated couplings, was presented in [10] together with results up to the three-loop

level for a covariant gauge and all values of N . From this, the n_f^2 contributions to the pure-singlet splitting functions were obtained at four loops [11]. Beyond three loops, ref. [8] determined a set of all-order constraints on the couplings, induced by gBRST and generalized anti-BRST symmetries [12–14]. In [8, 9], these constraints were solved at arbitrary loop order for fixed $N \leq 20$, leaving the systematic study of the alien operators at arbitrary spin N as an open problem. In this paper, we follow a different strategy. Namely, we will solve the constraints on the alien couplings to leading order in α_s but for *all* values of N . The main results of our study are:

- The all- N structure of the couplings is fixed in terms of a small set of constants. The latter can be determined by explicitly computing the couplings for some fixed values of N .
- The structure of the couplings of alien operators with $n + 1$ gluons is related to the ones with n gluons, allowing for a bootstrap in the determination of complicated higher-order couplings in terms of simpler lower-order ones.

The outline of the article is as follows. In section 2 we set the stage, review the generalized gauge symmetry and provide a brief summary of the set of relevant alien operators. In section 3 we study the identities that exist among the couplings of the alien operators and show how they can be used to restrict the all- N structure of the couplings. The results of this analysis are then used in section 4 to derive the Feynman rules of the alien operators, suitable for the renormalization of OMEs at all N up to four loops in QCD. Finally, in section 5, we summarize our findings and provide an outlook on further developments.

2 Setting the stage

In this section, we review the construction of the alien operators and summarize our conventions. The complete gauge-fixed QCD action is written as

$$S = \int d^D x \, (\mathcal{L}_0 + \mathcal{L}_{\text{GF+G}}) . \quad (2.1)$$

Here \mathcal{L}_0 represents the classical part of the QCD Lagrangian

$$\mathcal{L}_0 = -\frac{1}{4} F_a^{\mu\nu} F_{\mu\nu}^a + \sum_{f=1}^{n_f} \bar{\psi}^f (i \not{D} - m_f) \psi^f , \quad (2.2)$$

with the field strength defined as

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g_s f^{abc} A_\mu^b A_\nu^c . \quad (2.3)$$

f^{abc} are the standard QCD structure constants. The covariant derivative in eq. (2.2) is $\not{D} = \gamma^\mu (\partial_\mu - i g_s T^a A_\mu^a)$ with T^a the generator of the gauge group in the fundamental representation. The gauge-fixing and ghost terms are

$$\mathcal{L}_{\text{GF+G}} = -\frac{1}{2\xi} (\partial^\mu A_\mu^a)^2 - \bar{c}^a \partial^\mu D_\mu^{ab} c^b \quad (2.4)$$

with ξ the covariant gauge parameter and \bar{c}^a and c^a the anti-ghost and ghost fields, respectively. The covariant derivative in the adjoint representation is $D_\mu^{ac} = \partial_\mu \delta^{ac} + g_s f^{abc} A_\mu^b$. The QCD Lagrangian can be extended to also include spin- N gauge-invariant operators of twist two, which we define as

$$\begin{aligned}\mathcal{O}_g^{(N)}(x) &= \frac{1}{2} \text{Tr} \left[F_\nu(x) D^{N-2} F^\nu(x) \right], \\ \mathcal{O}_q^{(N)}(x) &= \frac{1}{2} \text{Tr} \left[\bar{\psi}(x) \not{D}^{N-1} \psi(x) \right].\end{aligned}\tag{2.5}$$

Here Δ_μ is a lightlike vector and we introduced the notation

$$F^{\mu;a} = \Delta_\nu F^{\mu\nu;a}, \quad A^a = \Delta_\mu A^{\mu;a}, \quad D = \Delta_\mu D^\mu, \quad \partial = \Delta_\mu \partial^\mu. \tag{2.6}$$

Under renormalization the operators in eq. (2.5) mix with operators proportional to the (classical) EOM and with BRST-exact operators [1, 4]. Following [8, 9], we begin by presenting the EOM aliens in the form

$$\mathcal{O}_{\text{EOM}}^{(N)} = \left(D \cdot F^a + g_s \bar{\psi} T^a \not{D} \psi \right) \mathcal{G}^a(A^a, \partial A^a, \partial^2 A^a, \dots) \tag{2.7}$$

with $D \cdot F^a = D_\nu F^{\nu;a}$ and \mathcal{G}^a a generic local function of the gauge field and its derivatives. It is convenient to expand \mathcal{G}^a in a series of contributions with an increasing number of gauge fields. This leads to

$$\mathcal{O}_{\text{EOM}}^{(N)} = \mathcal{O}_{\text{EOM}}^{(N),\text{I}} + \mathcal{O}_{\text{EOM}}^{(N),\text{II}} + \mathcal{O}_{\text{EOM}}^{(N),\text{III}} + \mathcal{O}_{\text{EOM}}^{(N),\text{IV}} + \dots \tag{2.8}$$

with

$$\mathcal{O}_{\text{EOM}}^{(N),\text{I}} = \eta(N) \left(D \cdot F^a + g_s \bar{\psi} \not{D} T^a \psi \right) \left(\partial^{N-2} A^a \right), \tag{2.9}$$

$$\mathcal{O}_{\text{EOM}}^{(N),\text{II}} = g_s \left(D \cdot F^a + g_s \bar{\psi} \not{D} T^a \psi \right) \sum_{i+j=N-3} C_{ij}^{abc} (\partial^i A^b) (\partial^j A^c), \tag{2.10}$$

$$\mathcal{O}_{\text{EOM}}^{(N),\text{III}} = g_s^2 \left(D \cdot F^a + g_s \bar{\psi} \not{D} T^a \psi \right) \sum_{i+j+k=N-4} C_{ijk}^{abcd} (\partial^i A^b) (\partial^j A^c) (\partial^k A^d), \tag{2.11}$$

$$\mathcal{O}_{\text{EOM}}^{(N),\text{IV}} = g_s^3 \left(D \cdot F^a + g_s \bar{\psi} \not{D} T^a \psi \right) \sum_{i+j+k+l=N-5} C_{ijkl}^{abcde} (\partial^i A^b) (\partial^j A^c) (\partial^k A^d) (\partial^l A^e). \tag{2.12}$$

The coefficients $C_{i_1 \dots i_{n-1}}^{a_1 \dots a_n}$ appearing in eqs. (2.10)–(2.12) can be written in terms of a set of independent colour tensors, each of them multiplying an associated coupling constant, as follows

$$C_{ij}^{abc} = f^{abc} \kappa_{ij}, \tag{2.13}$$

$$C_{ijk}^{abcd} = (f f)^{abcd} \kappa_{ijk}^{(1)} + d_4^{abcd} \kappa_{ijk}^{(2)} + d_{4ff}^{abcd} \kappa_{ijk}^{(3)}, \tag{2.14}$$

$$C_{ijkl}^{abcde} = (f f f)^{abcde} \kappa_{ijkl}^{(1)} + d_{4f}^{abcde} \kappa_{ijkl}^{(2)} \tag{2.15}$$

with

$$(f f)^{abcd} = f^{abe} f^{cde}, \quad (2.16)$$

$$(f f f)^{abcde} = f^{abm} f^{mcn} f^{nde}, \quad (2.17)$$

$$d_4^{abcd} = \frac{1}{4!} [\text{Tr}(T_A^a T_A^b T_A^c T_A^d) + \text{symmetric permutations}], \quad (2.18)$$

$$d_{4ff}^{abcd} = d_4^{abmn} f^{mce} f^{edn}, \quad (2.19)$$

$$d_{\widehat{4ff}}^{abcd} = d_{4ff}^{abcd} - \frac{1}{3} C_A d_4^{abcd}, \quad (2.20)$$

$$d_{4f}^{abcde} = d_4^{abcm} f^{mde}. \quad (2.21)$$

Here $(T_A)^b_{ac} = i f^{abc}$ are the generators of the adjoint representation of the colour group. We now extend the classical Lagrangian \mathcal{L}_0 in eq. (2.2) to include the gauge-invariant operators of twist two as well as the EOM aliens

$$\mathcal{L}_{\text{GGI}} = \mathcal{L}_0 + w_i \mathcal{O}_i^{(N)} + \mathcal{O}_{\text{EOM}}^{(N)}, \quad (2.22)$$

where w_i is a coupling for the operator \mathcal{O}_i with $i = g, q$, playing the same role as the coefficients $\eta(N), \kappa_{ij}, \dots$ defined in eqs. (2.13)–(2.15). The Lagrangian \mathcal{L}_{GGI} is invariant under the generalized gauge transformation [8] $A_\mu^a \rightarrow A_\mu^a + \delta_\omega A_\mu^a + \delta_\omega^\Delta A_\mu^a$, where

$$\begin{aligned} \delta_\omega A_\mu^a &= D_\mu^{ab} \omega^b(x), \\ \delta_\omega^\Delta A_\mu^a &= -\Delta_\mu \left[\eta(N) \partial^{N-1} \omega^a + g_s \sum_{i+j=N-3} \tilde{C}_{ij}^{aa_1 a_2} \left(\partial^i A^{a_1} \right) \left(\partial^{j+1} \omega^{a_2} \right) \right. \\ &\quad + g_s^2 \sum_{i+j+k=N-4} \tilde{C}_{ijk}^{aa_1 a_2 a_3} \left(\partial^i A^{a_1} \right) \left(\partial^j A^{a_2} \right) \left(\partial^{k+1} \omega^{a_3} \right) \\ &\quad \left. + g_s^3 \sum_{i+j+k+l=N-5} \tilde{C}_{ijkl}^{aa_1 a_2 a_3 a_4} \left(\partial^i A^{a_1} \right) \left(\partial^j A^{a_2} \right) \left(\partial^k A^{a_3} \right) \left(\partial^{l+1} \omega^{a_4} \right) + \mathcal{O}(g_s^4) \right] \end{aligned} \quad (2.23)$$

and

$$\tilde{C}_{ij}^{abc} = f^{abc} \eta_{ij}, \quad (2.24)$$

$$\tilde{C}_{ijk}^{abcd} = (f f)^{abcd} \eta_{ijk}^{(1)} + d_4^{abcd} \eta_{ijk}^{(2)} + d_{\widehat{4ff}}^{abcd} \eta_{ijk}^{(3)}, \quad (2.25)$$

$$\tilde{C}_{ijkl}^{abcde} = (f f f)^{abcde} \eta_{ijkl}^{(1)} + d_{4f}^{abcde} \eta_{ijkl}^{(2a)} + d_{4f}^{aebcd} \eta_{ijkl}^{(2b)}. \quad (2.26)$$

The generalized gauge symmetry implies that the couplings $\eta_{n_1 \dots n_j}^{(k)}$ are related to $\kappa_{n_1 \dots n_j}^{(k)}$ in eqs. (2.9)–(2.12)

$$\eta_{ij} = 2\kappa_{ij} + \eta(N) \binom{i+j+1}{i}, \quad (2.27)$$

$$\eta_{ijk}^{(1)} = 2\kappa_{i(j+k+1)} \binom{j+k+1}{j} + 2 \left[\kappa_{ijk}^{(1)} + \kappa_{kji}^{(1)} \right], \quad (2.28)$$

$$\eta_{ijk}^{(2)} = 3\kappa_{ijk}^{(2)}, \quad (2.29)$$

$$\eta_{ijk}^{(3)} = 2 \left[\kappa_{ijk}^{(3)} - \kappa_{kji}^{(3)} \right], \quad (2.30)$$

$$\eta_{ijkl}^{(1)} = 2 \left[\kappa_{ij(l+k+1)}^{(1)} + \kappa_{(l+k+1)ji}^{(1)} \right] \binom{l+k+1}{k} + 2 \left[\kappa_{ijkl}^{(1)} + \kappa_{ilkj}^{(1)} + \kappa_{likj}^{(1)} + \kappa_{lkij}^{(1)} \right], \quad (2.31)$$

$$\eta_{ijkl}^{(2a)} = 3\kappa_{ij(k+l+1)}^{(2)} \binom{k+l+1}{k} + 2\kappa_{ijkl}^{(2)}, \quad (2.32)$$

$$\eta_{ijkl}^{(2b)} = 2\kappa_{lik}^{(2)}. \quad (2.33)$$

The new gauge transformations in eq. (2.23) are promoted to a nilpotent generalized BRST (gBRST) operator, by replacing the transformation parameter ω^a with the ghost field c^a [8]. In turn the ghost alien operator is generated by the action of such gBRST operator on a suitable ancestor operator [8], giving

$$\mathcal{O}_c^{(N)} = \mathcal{O}_c^{(N),\text{I}} + \mathcal{O}_c^{(N),\text{II}} + \mathcal{O}_c^{(N),\text{III}} + \mathcal{O}_c^{(N),\text{IV}} + \dots \quad (2.34)$$

with

$$\mathcal{O}_c^{(N),\text{I}} = -\eta(N)(\partial \bar{c}^a)(\partial^{N-1} c^a), \quad (2.35)$$

$$\mathcal{O}_c^{(N),\text{II}} = -g_s \sum_{i+j=N-3} \tilde{C}_{ij}^{abc} (\partial \bar{c}^a) (\partial^i A^b) (\partial^{j+1} c^c), \quad (2.36)$$

$$\mathcal{O}_c^{(N),\text{III}} = -g_s^2 \sum_{i+j+k=N-4} \tilde{C}_{ijk}^{astu} (\partial \bar{c}^a) (\partial^i A^s) (\partial^j A^t) (\partial^{k+1} c^u), \quad (2.37)$$

$$\mathcal{O}_c^{(N),\text{IV}} = -g_s^3 \sum_{i+j+k+l=N-5} \tilde{C}_{ijkl}^{abcde} (\partial \bar{c}^a) (\partial^i A^b) (\partial^j A^c) (\partial^k A^d) (\partial^{l+1} c^e). \quad (2.38)$$

Renormalization. The complete Lagrangian, including the twist-two physical and alien operators, can be written as

$$\tilde{\mathcal{L}} = \mathcal{L}_0 + \mathcal{L}_{\text{GF}+\text{G}} + w_i \mathcal{O}_i + \mathcal{O}_{\text{EOM}}^{(N)} + \mathcal{O}_c^{(N)} = \mathcal{L}_0(A_\mu^a, g_s) + \mathcal{L}_{\text{GF}+\text{G}}(A_\mu^a, c^a, \bar{c}^a, g_s, \xi) + \sum_k \mathcal{C}_k \mathcal{O}_k, \quad (2.39)$$

where \mathcal{C}_k labels all the distinct couplings of the operators, e.g. $\mathcal{C}_k = (w_i, \eta(N), \kappa_{01}, \kappa_{12}, \dots)$. The ultraviolet (UV) singularities associated with the QCD Lagrangian are absorbed by introducing the bare fields and parameters

$$A_\mu^{a;\text{bare}}(x) = \sqrt{Z_3} A_\mu^a(x), \quad c^{a;\text{bare}}(x) = \sqrt{Z_c} c^a(x), \quad \bar{c}^{a;\text{bare}}(x) = \sqrt{Z_c} \bar{c}^a(x), \quad (2.40)$$

$$g_s^{\text{bare}} = \mu^\epsilon Z_g g_s, \quad \xi^{\text{bare}} = \sqrt{Z_3} \xi. \quad (2.41)$$

We renormalize the singularities originating from the insertion of the composite operators using

$$\mathcal{O}_i^{\text{ren}}(x) = Z_{ij} \mathcal{O}_j^{\text{bare}}(x), \quad (2.42)$$

where $\mathcal{O}_j^{\text{bare}}$ indicates the operators in eqs. (2.5), (2.8) and (2.34) written in terms of the bare fields. Note that throughout this work we use $D = 4 - 2\epsilon$ dimensional regularization, combined

with the $\overline{\text{MS}}$ renormalization scheme. Z_{ij} is the renormalization matrix of the operators, which makes the OMEs featuring an insertion of $\mathcal{O}_i^{\text{ren}}$ finite. The renormalized Lagrangian becomes

$$\tilde{\mathcal{L}} = \mathcal{L}_0(A_\mu^{a;\text{bare}}, g_s^{\text{bare}}) + \mathcal{L}_{\text{GF+G}}(A_\mu^{a;\text{bare}}, c^{a;\text{bare}}, \bar{c}^{a;\text{bare}}, g_s^{\text{bare}}, \xi^{\text{bare}}) + \sum_k \mathcal{C}_k^{\text{bare}} \mathcal{O}_k^{\text{bare}}, \quad (2.43)$$

$$\mathcal{C}_i^{\text{bare}} = \sum_k \mathcal{C}_k Z_{ki}, \quad (2.44)$$

where \mathcal{C}_k is the (finite) renormalized coupling of the operator \mathcal{O}_k . The UV-finite OMEs featuring a single insertion of $\mathcal{O}_g^{\text{ren}}$ are computed by setting the renormalized couplings $\mathcal{C}_i = \delta_{ig}$ in eq. (2.43), which gives

$$\mathcal{C}_i^{\text{bare}} = Z_{gi}. \quad (2.45)$$

Similarly, the renormalized OMEs with an insertion of \mathcal{O}_q are obtained with $\mathcal{C}_i^{\text{bare}} = Z_{qi}$. Therefore, the couplings of the bare operators $\eta^{\text{bare}}(N), \dots$ are interpreted as the renormalization constants that mix the physical operators into the aliens. These quantities can be extracted from the direct calculation of the singularities of the OMEs with an insertion of $\mathcal{O}_g^{\text{bare}}$ ($\mathcal{O}_q^{\text{bare}}$). For instance, the coupling $\eta^{\text{bare}}(N)$, which is associated to an operator with a two-point vertex, was determined in [1, 7] from the renormalization of the OMEs of \mathcal{O}_g with two external ghosts and it was found to be¹

$$\eta^{\text{bare}}(N) = Z_{gc} = -\frac{a_s}{\epsilon} \frac{C_A}{N(N-1)} + \mathcal{O}(a_s^2), \quad (2.46)$$

where C_A is the quadratic Casimir in the adjoint representation and $a_s = \alpha_s/(4\pi) = g_s^2/(4\pi)^2$. The value of η was determined at two loops in [7] and at three loops in [10]. Throughout this paper we will mainly be interested in the one-loop alien couplings. As such, it will be convenient to select just the N -dependent part of the one-loop result of $\eta^{\text{bare}}(N)$, which in the following we simply denote by $\eta(N)$, i.e.

$$\eta(N) = \frac{-1}{N(N-1)}. \quad (2.47)$$

The couplings of the operators featuring multiple fields, e.g., the couplings κ_{ij} multiply *at least* three fields, are determined by renormalizing OMEs with the corresponding external fields. Recently, a method to compute the counterterms of the OMEs with insertions of the gauge-invariant operators as a function of the spin N was put forward in ref. [10]. The result of that paper can be used to extract the coefficients $\kappa_{ij}^{\text{bare}}$ up to $\mathcal{O}(a_s^2)$ and those of $\kappa_{ijk}^{(p);\text{bare}}$, for $p = 1, 2$, at $\mathcal{O}(a_s)$, finding agreement with the low- N values reported in [8]. In addition, the calculation of the five-point counterterms at $\mathcal{O}(a_s)$, which can be used to determine $\kappa_{ijkl}^{(p)}$, for $p = 1, 2$, has been announced recently [15].

In this paper, we would like to determine the renormalization constants Z_{gi} by solving the constraints on the couplings $\mathcal{C}_i^{\text{bare}}$, which are imposed by the symmetries of eq. (2.43).

¹Note that the expression for η in [7] has an additional factor of 2. This is a consequence of the chosen conventions for dimensional regularization. In particular, we use $D = 4 - 2\epsilon$ while [7] employs $D = 4 + \epsilon$.

The latter is the Lagrangian in eq. (2.39) evaluated with bare fields and couplings constants. Therefore the two Lagrangians share the same symmetry properties, with the obvious substitutions. For simplicity, in the rest of this paper we drop the superscript ‘bare’, wherever it does not create any ambiguity.

Independent operators and couplings. The symmetry constraints on the couplings in eq. (2.43) have been derived in ref. [8]. Without repeating the derivation of that paper, we distinguish three types of relations, which follow from the way we have constructed the operators at the beginning of this section.

First of all, the couplings introduced in the EOM operators, see eqs. (2.9)–(2.12) and (2.13)–(2.15), are chosen to inherit the properties of the colour structures they multiply. For example, because of the anti-symmetry of the structure constants, we take

$$\kappa_{ij} = -\kappa_{ji}. \quad (2.48)$$

This implies, e.g., that at spin $N = 4$, where $i, j = 0, 1$, there is only one independent coupling, e.g., κ_{01} .

The second type of constraints regards the couplings that enter the ghost operators, eqs. (2.35)–(2.38). Because these operators were constructed directly from the EOM ones using gBRST, the η couplings are connected to the κ ones. The relevant identities have been listed in eqs. (2.27)–(2.33).

Finally, we impose the invariance of eq. (2.43) under the generalized transformations of anti-BRST type [12–14], which stem from eq. (2.23), by replacing the transformation parameter ω^a with the anti-ghost field \bar{c}^a . This implies the following condition on the ghost operator $\mathcal{O}_c^{(N)}$ defined in eq. (2.34)

$$O_c^{(N)}(A_\mu^a, c^a, \bar{c}^a) = O_c^{(N)}(A_\mu^a, \bar{c}^a, c^a), \quad (2.49)$$

which translates into a set of constraints on the couplings in eqs. (2.35)–(2.38) and, in turn, on those of the EOM operators. Taking the example of $N = 4$, the anti-BRST relation imposes $\kappa_{01} = 2\eta(4)$, thus reducing the number of independent couplings even further [8].

It is highly non-trivial to find all- N solutions for all the constraints. In refs. [8, 9], they were solved only for fixed values of N , in order to fix bases of independent alien operators up to $N = 20$. In the following sections, we solve the relations with exact N dependence. This is done by setting up an ansatz for the function space that enters to leading order in a_s . The construction of this ansatz is primarily based on constraints from (anti-)gBRST. We will see below that the latter allow one to bootstrap the functional form of higher-order couplings from that of the lower-order ones. The determination of the unknown parameters in the ansatz is then performed by using the full set of colour, gBRST and anti-gBRST relations. As will become clear below, this allows one to fix most, but not all, free parameters. The few that remain then need to be determined from the explicit renormalization of a limited number of fixed- N operator matrix elements. This is particularly important for finding any overall N -dependent function.

3 Identities among the alien couplings

In this section we will discuss in detail the identities between the couplings coming from the (anti)-gBRST relations. In particular, we will show that they allow one to restrict the function space of the couplings and hence constrain their generic N -dependence.

3.1 Class II couplings

The class II operators are defined in terms of two couplings, κ_{ij} and η_{ij} , which obey the following relations

$$\kappa_{ij} + \kappa_{ji} = 0, \quad [\text{anti-symmetry of } f] \quad (3.1)$$

$$\eta_{ij} = 2\kappa_{ij} + \eta(N) \binom{i+j+1}{i}, \quad [\text{gBRST}] \quad (3.2)$$

$$\eta_{ij} + \sum_{s=0}^i (-1)^{s+j} \binom{s+j}{j} \eta_{(i-s)(j+s)} = 0. \quad [\text{anti-gBRST}] \quad (3.3)$$

Note that one can generate an equation for the ghost coupling alone by combining the anti-symmetry of κ_{ij} , eq. (3.1), with the gBRST relation, eq. (3.2),

$$\eta_{ij} + \eta_{ji} = \eta(N) \left[\binom{i+j+1}{i} + \binom{i+j+1}{j} \right]. \quad (3.4)$$

The one-loop value of this coupling was first computed in [1] and later corrected in [7]. In our conventions it reads²

$$\eta_{ij} = -\frac{\eta(N)}{4} \left[(-1)^j - 3 \binom{N-2}{i+1} - \binom{N-2}{i} \right] \quad (3.5)$$

which implies

$$\kappa_{ij} = -\frac{\eta(N)}{8} \left[(-1)^j + 3 \binom{N-2}{i} - 3 \binom{N-2}{i+1} \right]. \quad (3.6)$$

The power of the relations described above is that they can be used to gain valuable information about the structure of the couplings at arbitrary N . For example, one can use eq. (3.4) to write down an ansatz for η_{ij} of the form

$$\eta_{ij} = \eta(N) \left[c_1 \binom{i+j+1}{i} + c_2 \binom{i+j+1}{j} \right]. \quad (3.7)$$

Here c_1 and c_2 are constants to be determined. We assume here that the dependence on $\eta(N)$ is factorized at leading order, as suggested by eq. (3.4) and observed in eq. (3.5).

This ansatz can then be substituted in the anti-gBRST consistency relation, eq. (3.3), yielding

$$\eta_{ij} + \sum_{s=0}^i (-1)^{s+j} \binom{s+j}{j} \eta_{(i-s)(j+s)} = c_1 \eta(N) \left[(-1)^j + \binom{i+j+1}{i} \right] \quad (3.8)$$

²Note that there are typos in the corresponding expression in [8]. In particular, the right-hand side of eq. (4.38) in [8] should be replaced by the right-hand side of eq. (3.5) here.

for even values of N . Hence we find a consistent solution when $c_1 = 0$, while c_2 is unconstrained. Assuming furthermore that κ_{ij} lives in the same function space as η_{ij} , which is motivated by their close relationship due to the generalized BRST symmetry, we posit

$$\kappa_{ij} = \eta(N) \left[b_1 \binom{i+j+1}{i} + b_2 \binom{i+j+1}{j} \right]. \quad (3.9)$$

Then, requiring the anti-symmetry relation in eq. (3.1) and the gBRST one in eq. (3.2) to hold actually produces a unique solution for both couplings, which reads

$$\eta_{ij} = \eta(N) \binom{N-2}{j}, \quad (3.10)$$

$$\kappa_{ij} = \frac{\eta(N)}{2} \left[\binom{N-2}{j} - \binom{N-2}{i} \right]. \quad (3.11)$$

Note that we used $i+j = N-3$. Next, we compare with some fixed- N evaluation to check our result. While it is obvious from the actual solution in eq. (3.5) that the result in eq. (3.10) is in fact incorrect, there is numerical agreement between both for $N = 4$. So, to determine that eq. (3.10) does not represent the physical solution, we actually need to evaluate one more moment, say $N = 6$, for which the disagreement does become obvious. Hence, we now need to extend the ansatz, for which we can use the anti-gBRST relation. In particular, note that the right-hand side of eq. (3.8) suggests the inclusion of a term proportional to $(-1)^j$ to the ansatz,

$$\eta_{ij} = \eta(N) \left[c_1 \binom{i+j+1}{i} + c_2 \binom{i+j+1}{j} + c_3 (-1)^j \right]. \quad (3.12)$$

The anti-gBRST relation now becomes

$$\eta_{ij} + \sum_{s=0}^i (-1)^{s+j} \binom{s+j}{j} \eta_{(i-s)(j+s)} = \eta(N) (c_1 + c_3) \left[(-1)^j + \binom{i+j+1}{i} \right] \quad (3.13)$$

such that $c_3 = -c_1$ is a consistent solution. If we now impose also eq. (3.4) then we obtain the relation $c_1 + c_2 = 1$, leaving just one free parameter unconstrained. Hence

$$\eta_{ij} = \eta(N) \left[c_1 \left[\binom{i+j+1}{i} - (-1)^j \right] + (1 - c_1) \binom{i+j+1}{j} \right]. \quad (3.14)$$

It should be noted that, if an ansatz is generated using (anti-)gBRST relations, one is in principle free to add non-zero functions that live in the kernel of these relations. For example, if one adds a term of the form

$$-\frac{f(N)}{4} \left((-1)^j + \binom{N-2}{i+1} - \binom{N-2}{i} \right) \quad (3.15)$$

to eq. (3.5), the corresponding expression for η_{ij} still obeys the constraints. Here $f(N)$ represents an arbitrary function of N , with the actual solution being recovered by setting $f(N) = 0$ for even values of N . In particular, substituting eq. (3.15) in the constraint coming from anti-symmetry and gBRST, cf. eq. (3.4), one finds

$$\left[(-1)^i + (-1)^j \right] f(N) = 0. \quad (3.16)$$

The left-hand side of this expression always vanishes for all physical (even) values of N , independent of the functional form of $f(N)$. In general, the exclusion of this type of function can only be confirmed by comparison with fixed- N computations.

An important consequence is that now we have recovered the full function space of the actual solution, eq. (3.5), using only the symmetry relations of the couplings. More generally, note that eq. (3.3) is an example of a *conjugation* relation, in the sense that a second application of the sum leads to

$$\sum_{t=0}^i (-1)^{t+j} \binom{t+j}{j} \eta_{(i-t)(j+t)} = - \sum_{t=0}^i (-1)^{t+j} \binom{t+j}{j} \sum_{s=0}^{i-t} (-1)^{s+j+t} \binom{s+j+t}{j+t} \eta_{(i-t-s)(j+t+s)} \quad (3.17)$$

and hence

$$\eta_{ij} = \sum_{t=0}^i \binom{t+j}{j} \sum_{s=0}^{i-t} (-1)^s \binom{s+j+t}{j+t} \eta_{(i-t-s)(j+t+s)}. \quad (3.18)$$

The latter identity is actually *always* true for *any* discrete two-variable function η_{ij} . This type of conjugation relation has already been encountered in the computation of the anomalous dimensions of twist-two operators in non-forward kinematics, see e.g. [16, 17], and holds great predictive power. In particular, it provides valuable information about the function space of the object at hand. To take full advantage of such relations, one needs to be able to evaluate them analytically. This is possible by using principles of symbolic summation, in particular by application of the creative telescoping algorithm [18]. The latter is a generalization of classical telescoping and attempts to evaluate the sum of interest by rewriting it as a recursion relation using Gosper's algorithm [19]. The closed-form expression of the sum then corresponds to the linear combination of the solutions of the recursion that has the same initial values as the sum. This methodology is neatly implemented in the **Mathematica** package **Sigma** [20, 21]. For the class III and IV couplings to be described below we will also encounter identities involving multiple sums, for which the package **EvaluateMultiSums** [22, 23] can be used.

3.2 Class III couplings

3.2.1 $\kappa_{ijk}^{(1)}$ and $\eta_{ijk}^{(1)}$

The couplings $\eta_{ijk}^{(1)}$ and $\kappa_{ijk}^{(1)}$ can be thought of as direct generalizations of η_{ij} and κ_{ij} in the class II operators. They obey the following relations

$$\kappa_{ijk}^{(1)} + \kappa_{ikj}^{(1)} = 0, \quad [\text{anti-symmetry of } f] \quad (3.19)$$

$$\kappa_{ijk}^{(1)} + \kappa_{jki}^{(1)} + \kappa_{kij}^{(1)} = 0, \quad [\text{Jacobi identity}] \quad (3.20)$$

$$\eta_{ijk}^{(1)} = 2\kappa_{i(j+k+1)} \binom{j+k+1}{j} + 2[\kappa_{ijk}^{(1)} + \kappa_{kji}^{(1)}], \quad [\text{gBRST}] \quad (3.21)$$

$$\eta_{ijk}^{(1)} = \sum_{m=0}^i \sum_{n=0}^j \frac{(m+n+k)!}{m!n!k!} (-1)^{m+n+k} \eta_{(j-n)(i-m)(k+m+n)}^{(1)}. \quad [\text{anti-gBRST}] \quad (3.22)$$

Note that now the indices are constrained as $i + j + k = N - 4$. As before, one can combine the relations of the EOM coupling with the gBRST relation to connect $\eta_{ijk}^{(1)}$ to $\kappa_{ijk}^{(1)}$. In

particular we find

$$\eta_{ijk}^{(1)} + \eta_{ikj}^{(1)} = 2\kappa_{i(j+k+1)} \binom{j+k+2}{j+1} + 2[\kappa_{kji}^{(1)} + \kappa_{jki}^{(1)}] \quad (3.23)$$

when combining the anti-symmetry property of $\kappa_{ijk}^{(1)}$ with the gBRST identity. Similarly the combination of the Jacobi identity with gBRST leads to

$$\eta_{ijk}^{(1)} + \eta_{kij}^{(1)} + \eta_{jki}^{(1)} = 2\kappa_{i(j+k+1)} \binom{j+k+1}{j} + 2\kappa_{k(i+j+1)} \binom{i+j+1}{i} + 2\kappa_{j(i+k+1)} \binom{i+k+1}{k}. \quad (3.24)$$

The latter identity relates the class III coupling $\eta_{ijk}^{(1)}$, which is $\mathcal{O}(g_s^2)$, to the class II coupling κ_{ij} of $\mathcal{O}(g_s)$, i.e. at one order lower in perturbation theory. As such, we can use it to determine the function space of $\eta_{ijk}^{(1)}$. Taking into account all independent permutations of i, j and k , we find that this function space is 18-dimensional

$$\left\{ (-1)^{i+j} \binom{i+j+1}{i}, \binom{N-2}{k+1} \binom{i+j+1}{i}, \binom{N-2}{k} \binom{i+j+1}{i}, (-1)^{j+k} \binom{j+k+1}{j}, \right. \\ \left. \binom{N-2}{i+1} \binom{j+k+1}{j}, \binom{N-2}{i} \binom{j+k+1}{j}, (-1)^{i+k} \binom{i+k+1}{k}, \binom{N-2}{j+1} \binom{i+k+1}{k}, \right. \\ \left. \binom{N-2}{j} \binom{i+k+1}{k} + \text{independent permutations of } i, j \text{ and } k \right\}. \quad (3.25)$$

Furthermore, due to the close relationship between $\eta_{ijk}^{(1)}$ and $\kappa_{ijk}^{(1)}$, we assume that the functional form of the latter is constructed from the same functions. Hence in total we have 36 free parameters. Using the relations described above, cf. eqs. (3.19)–(3.22), we are able to fix 34 of these. The final two free parameters are then determined using the one-loop results $\kappa_{110}^{(1)} = 0$ and $\kappa_{121}^{(1)} = 13 C_A/336$, which follow from the explicit operator renormalization for $N = 6$ and $N = 8$ respectively. Our final result for $\kappa_{ijk}^{(1)}$ then becomes

$$\kappa_{ijk}^{(1)} = \frac{\eta(N)}{48} \left\{ 2(-1)^{i+j} \binom{i+j+1}{i} + (-1)^{i+k} \binom{i+k+1}{k} + 3(-1)^{j+k+1} \binom{j+k+1}{j} \right. \\ \left. + \binom{i+k+1}{i} \left[2(-1)^{i+k+1} + 5 \binom{N-1}{j+1} \right] + \binom{j+k+1}{k} \left[3(-1)^{j+k} - 10 \binom{N-2}{i} \right] \right. \\ \left. + 4 \binom{N-2}{i+1} \right] + \binom{i+j+1}{j} \left[(-1)^{i+j+1} + 5 \binom{N-2}{k} - 9 \binom{N-2}{k+1} \right] \right\}. \quad (3.26)$$

To verify this expression, agreement with explicitly computed fixed- N values has been established up to $N = 20$. The necessary direct computations at fixed values of N of Feynman diagrams for the OMEs with (physical or alien) spin- N twist-two operators $\mathcal{O}^{(N)}$ inserted in Green's functions with off-shell quarks, gluons or ghosts are performed with the setup used and described in [9, 24–27] for the computation of moments of four-loop QCD splitting functions. In particular, the **Forcer** package [28], written in **Form** [29–31], is used for the

parametric reductions of the two-point functions up to four loops for fixed even integer values of N . Substituting our result for $\kappa_{ijk}^{(1)}$ into the gBRST relation, eq. (3.21), allows one to also reconstruct the full N -dependence of $\eta_{ijk}^{(1)}$

$$\begin{aligned} \eta_{ijk}^{(1)} = & -\frac{\eta(N)}{24} \left\{ 5(-1)^{i+j+1} \binom{i+j+1}{i} + (-1)^{i+k} \binom{i+k+1}{k} + 2(-1)^{j+k+1} \binom{j+k+1}{j} \right. \\ & + \binom{i+k+1}{i} \left[(-1)^{i+k} + 4 \binom{N-2}{j+1} \right] + \binom{j+k+1}{k} \left[5(-1)^{j+k+1} - 3 \binom{N-2}{i} \right] \\ & \left. + \binom{N-2}{i+1} \right] + \binom{i+j+1}{j} \left[4(-1)^{i+j} - 15 \binom{N-2}{k} - 5 \binom{N-2}{k+1} \right] \Big\}. \end{aligned} \quad (3.27)$$

We have verified that eqs. (3.26) and (3.27) are in agreement with the results of ref. [10], as explained in section 4 below.

3.2.2 $\kappa_{ijk}^{(2)}$ and $\eta_{ijk}^{(2)}$

The next alien couplings we consider are $\kappa_{ijk}^{(2)}$ and $\eta_{ijk}^{(2)}$, which obey the following relations

$$\kappa_{ijk}^{(2)} = \kappa_{jik}^{(2)} = \kappa_{ikj}^{(2)} = \kappa_{kji}^{(2)} = \kappa_{jki}^{(2)} = \kappa_{kij}^{(2)}, \quad [\text{symmetry of } d_4] \quad (3.28)$$

$$\eta_{ijk}^{(2)} = 3\kappa_{ijk}^{(2)}, \quad [\text{gBRST}] \quad (3.29)$$

$$\eta_{ijk}^{(2)} = \sum_{m=0}^i \sum_{n=0}^j (-1)^{m+n+k} \frac{(m+n+k)!}{m! n! k!} \eta_{(i-m)(j-n)(m+n+k)}^{(2)}. \quad [\text{anti-gBRST}] \quad (3.30)$$

As the anti-gBRST equation has a similar form as the one for $\eta_{ijk}^{(1)}$, cf. eq. (3.22), we assume the function space for $\eta_{ijk}^{(2)}$ and $\kappa_{ijk}^{(2)}$ to be the same as above, cf. eq. (3.25). Imposing eqs. (3.28)–(3.30) then allows one to fix all but one of the unknowns. Hence we find expressions $\eta_{ijk}^{(2)}$ and $\kappa_{ijk}^{(2)}$ with only one (overall) free parameter

$$\kappa_{ijk}^{(2)} = c \left\{ (-1)^{i+j} \binom{i+j+2}{i+1} + (-1)^{i+k} \binom{i+k+2}{i+1} + \binom{j+k+2}{j+1} \left[(-1)^{j+k} + \binom{N-1}{i+1} \right] \right\}, \quad (3.31)$$

$$\eta_{ijk}^{(2)} = 3\kappa_{ijk}^{(2)}. \quad (3.32)$$

Note that the c parameter can a priori be some N -dependent function. A computation of the OMEs at a few fixed values of N with the procedure outlined in section 3.2.1 for the renormalization of the respective operators fixes $c = 1/N/(N-1)$ such that

$$\kappa_{ijk}^{(2)} = \frac{1}{N(N-1)} \left\{ (-1)^{i+j} \binom{i+j+2}{i+1} + (-1)^{i+k} \binom{i+k+2}{i+1} + \binom{j+k+2}{j+1} \left[(-1)^{j+k} + \binom{N-1}{i+1} \right] \right\}, \quad (3.33)$$

$$\eta_{ijk}^{(2)} = 3\kappa_{ijk}^{(2)}. \quad (3.34)$$

Noting that

$$\frac{1}{N(N-1)} = -\eta(N) \quad (3.35)$$

we see that also for these two couplings $\eta(N)$ factorizes, which is not expected a priori from the constraints.

3.2.3 $\kappa_{ijk}^{(3)}$ and $\eta_{ijk}^{(3)}$

The last set of couplings in the class III alien operators, $\kappa_{ijk}^{(3)}$ and $\eta_{ijk}^{(3)}$, obey the following relations

$$\kappa_{ijk}^{(3)} = \kappa_{ikj}^{(3)}, \quad [\text{symmetry}] \quad (3.36)$$

$$\kappa_{ijk}^{(3)} + \kappa_{kij}^{(3)} + \kappa_{jki}^{(3)} = 0, \quad [\text{generalized Jacobi identity}] \quad (3.37)$$

$$\eta_{ijk}^{(3)} = 2(\kappa_{ijk}^{(3)} - \kappa_{kji}^{(3)}), \quad [\text{gBRST}] \quad (3.38)$$

$$\eta_{ijk}^{(3)} = \sum_{m=0}^i \sum_{n=0}^j (-1)^{m+n+k} \frac{(m+n+k)!}{m! n! k!} \eta_{(j-n)(i-m)(m+n+k)}^{(3)}. \quad [\text{anti-gBRST}] \quad (3.39)$$

As before, we suggest the same function space as for $\kappa_{ijk}^{(1)}$ and $\eta_{ijk}^{(1)}$, cf. eq. (3.25). The above relations then only leave two parameters unfixed, such that we have

$$\begin{aligned} \kappa_{ijk}^{(3)} = & c_1 (-1)^{i+j} \binom{i+j+1}{i} + c_2 (-1)^{i+k} \binom{i+k+1}{k} + \binom{j+k+1}{j} \left[(c_1 + c_2) (-1)^{j+k+1} \right. \\ & + (c_1 + c_2) (-1)^{j+k+1} + c_1 \binom{N-2}{i+1} \left. \right] + \binom{i+k+1}{i} \left[c_1 (-1)^{i+k} + (2c_1 + c_2) \binom{N-2}{j} \right. \\ & \left. + c_2 \binom{N-2}{j+1} \right] + \binom{i+j+1}{j} \left[c_2 (-1)^{i+j} - (2c_1 + c_2) \binom{N-2}{k} - (c_1 + c_2) \binom{N-2}{k+1} \right] \end{aligned} \quad (3.40)$$

with c_1, c_2 to be determined. We emphasize that, as before, these could be N -dependent functions.³ The corresponding expression for $\eta_{ijk}^{(3)}$ depends on the same parameters through the gBRST relation, cf. eq. (3.38). Since the couplings $\kappa_{ijk}^{(3)}$ and $\eta_{ijk}^{(3)}$ do not appear through operator mixing in the renormalization of physical OMEs up to four loops, we leave the two free parameters c_1, c_2 in eq. (3.40) undetermined, for the time being. We will address this issue again when extending the computation of low- N non-singlet anomalous dimensions at five loops [32] to the flavor-singlet sector.

3.3 Class IV couplings

3.3.1 $\kappa_{ijkl}^{(1)}$ and $\eta_{ijkl}^{(1)}$

We have the following set of relations

$$\kappa_{ijkl}^{(1)} + \kappa_{ijlk}^{(1)} = 0, \quad [\text{anti-symmetry}] \quad (3.41)$$

³In this case we expect $c_1 \sim c_2 \sim \eta(N)$.

$$\kappa_{ijkl}^{(1)} + \kappa_{iklj}^{(1)} + \kappa_{iljk}^{(1)} = 0, \quad [\text{Jacobi}] \quad (3.42)$$

$$\kappa_{ijkl}^{(1)} + \kappa_{jilk}^{(1)} + \kappa_{lkji}^{(1)} + \kappa_{klij}^{(1)} = 0, \quad [\text{double Jacobi}] \quad (3.43)$$

$$\begin{aligned} \eta_{ijkl}^{(1)} = & 2[\kappa_{ij(l+k+1)}^{(1)} + \kappa_{(l+k+1)ji}^{(1)}] \binom{l+k+1}{k} \\ & + 2[\kappa_{ijkl}^{(1)} + \kappa_{ilkj}^{(1)} + \kappa_{likj}^{(1)} + \kappa_{lkij}^{(1)}], \end{aligned} \quad [\text{gBRST}] \quad (3.44)$$

$$\begin{aligned} \eta_{ijkl}^{(1)} = & - \sum_{s_1=0}^i \sum_{s_2=0}^j \sum_{s_3=0}^k \frac{(s_1+s_2+s_3+l)!}{s_1! s_2! s_3! l!} (-1)^{s_1+s_2+s_3+l} \\ & \times \eta_{(k-s_3)(j-s_2)(i-s_1)(s_1+s_2+s_3+l)}^{(1)} \end{aligned} \quad [\text{anti-gBRST}] \quad (3.45)$$

with now $i + j + k + l = N - 5$. Combining the double Jacobi identity, eq. (3.43), with the gBRST one, eq. (3.39), allows one to write $\eta_{ijkl}^{(1)}$ in terms of $\kappa_{ijk}^{(1)}$ appearing already in the class III operators at one order in perturbation theory lower,

$$\begin{aligned} \eta_{ijkl}^{(1)} + \eta_{jikl}^{(1)} + \eta_{lkji}^{(1)} + \eta_{klij}^{(1)} = & 2[\kappa_{ij(k+l+1)}^{(1)} + \kappa_{(k+l+1)ji}^{(1)}] \binom{k+l+1}{k} \\ & + 2[\kappa_{ji(k+l+1)}^{(1)} + \kappa_{(k+l+1)ij}^{(1)}] \binom{k+l+1}{l} \\ & + 2[\kappa_{lk(i+j+1)}^{(1)} + \kappa_{(i+j+1)kl}^{(1)}] \binom{i+j+1}{j} \\ & + 2[\kappa_{kl(i+j+1)}^{(1)} + \kappa_{(i+j+1)lk}^{(1)}] \binom{i+j+1}{i}. \end{aligned} \quad (3.46)$$

As such, we can use the expression we have computed for $\kappa_{ijk}^{(1)}$, cf. eq. (3.26), to determine the function space of $\eta_{ijkl}^{(1)}$. Taking into account all the independent permutations of the indices i, k, j and l this space is now 264-dimensional. Assuming that the functional form of $\kappa_{ijkl}^{(1)}$ is similar to the one of $\eta_{ijkl}^{(1)}$ then implies that in total we now have 528 parameters to fix. However, after implementing all of the above relations, only 8 remain in the end. The latter can again be fixed from the explicit renormalization of a few fixed- N matrix elements. More specifically we extracted them by performing a small momentum expansion around the limit $p_3, p_4, p_5 \rightarrow 0$ of the OME

$$\langle O_g^{(N)}; \bar{c}(p_1)c(p_2)g(p_3)g(p_4)g(p_5) \rangle. \quad (3.47)$$

This expansion is achieved on a diagram-by-diagram basis using the expansion-by-subgraph method [33–35] to second order at $N = 10$ and to third order at $N = 12$. By expanding sequentially in the external gluon momenta p_3, p_4 and p_5 the integrals are reduced to simple one-scale propagator integrals. We have implemented the expansion-by-subgraph in **Maple** [36] and then subsequently evaluated the expressions in **Form**. This methodology was also used to cross-check the expressions for $\kappa_{ij}^{(1)}$ and $\kappa_{ijk}^{(r=1,2)}$ up to $N = 20$. At one loop the poles of the

OME, eq. (3.47), are generated purely by the ghost alien operator $O_c^{(N),IV}$ allowing for a clean extraction of $\eta_{ijkl}^{(1)}$ renormalization constants, from which the $\kappa_{ijkl}^{(1)}$ values can be obtained. In particular, in order to determine the remaining constants in the all- N ansatz for $\kappa_{ijkl}^{(1)}$, we use

$$\kappa_{0210}^{(1)} = -\frac{1}{128}C_A, \quad \kappa_{0050}^{(1)} = \frac{109}{1440}C_A, \quad \kappa_{0104}^{(1)} = -\frac{935}{6912}C_A, \quad \kappa_{1006}^{(1)} = -\frac{2537}{16896}C_A. \quad (3.48)$$

We then find

$$\begin{aligned} \kappa_{ijkl}^{(1)} = & -\frac{\eta(N)}{384} \left\{ \left[6(-1)^{j+k} \binom{i+l+1}{i} - 3(-1)^{j+k} \binom{i+l+1}{l} + 7(-1)^{j+k+l} \binom{j+k+l+2}{l} \right] \right. \\ & + 7(-1)^{j+k+l} \binom{j+k+l+2}{j+k+1} - 27 \binom{N-1}{i+1} \binom{j+k+l+2}{j+k+1} + 2 \binom{i+j+k+2}{j+k+1} \\ & \times \left[2(-1)^{i+j+k} + 9 \binom{N-2}{l} \right] - 2 \binom{i+j+k+2}{i} \left[4(-1)^{i+j+k} + 15 \binom{N-2}{l+1} \right] \Big] \\ & \times \binom{j+k+1}{j} - \left[5(-1)^{j+k} \binom{i+l+1}{i} - 4(-1)^{j+k} \binom{i+l+1}{l} \right. \\ & \left. - 14(-1)^{j+k+l} \binom{j+k+l+2}{l} + 7(-1)^{j+k+l} \binom{j+k+l+2}{j+k+1} \right] \\ & + 54 \binom{N-2}{i} \binom{j+k+l+2}{j+k+1} + \binom{i+j+k+2}{j+k+1} \left[-3(-1)^{i+j+k} + 4 \binom{N-1}{l+1} \right] \\ & + \binom{i+j+k+2}{i} \left[3(-1)^{i+j+k} + 13 \binom{N-1}{l+1} \right] \Big] \binom{j+k+1}{k} \\ & + (-1)^{i+j+k+1} \binom{i+k+1}{k} \binom{i+j+k+2}{j} - 2(-1)^{i+j+k} \binom{i+j+1}{j} \binom{i+j+k+2}{k} \\ & - 6(-1)^{j+l} \binom{i+k+1}{i} \binom{j+l+1}{j} + 3(-1)^{j+l} \binom{i+k+1}{k} \binom{j+l+1}{j} \\ & + 5(-1)^{j+l} \binom{i+k+1}{i} \binom{j+l+1}{l} - 4(-1)^{j+l} \binom{i+k+1}{k} \binom{j+l+1}{l} \\ & + 30 \binom{N-2}{k+1} \binom{j+l+1}{j} \binom{i+j+l+2}{i} - 5(-1)^{i+j+l} \binom{j+l+1}{l} \binom{i+j+l+2}{i} \\ & + 13 \binom{N-1}{k+1} \binom{j+l+1}{l} \binom{i+j+l+2}{i} - 4(-1)^{i+j+l} \binom{i+l+1}{i} \binom{i+j+l+2}{j} \\ & + 12 \binom{N-2}{k} \binom{i+l+1}{i} \binom{i+j+l+2}{j} + 24 \binom{N-2}{k+1} \binom{i+l+1}{i} \binom{i+j+l+2}{j} \\ & - 3(-1)^{i+j+l} \binom{i+l+1}{l} \binom{i+j+l+2}{j} + 49 \binom{N-1}{k+1} \binom{i+l+1}{l} \binom{i+j+l+2}{j} \\ & \left. - 30 \binom{N-1}{k+1} \binom{i+j+1}{i} \binom{i+j+l+2}{l} + 2(-1)^{i+j+l} \binom{i+j+1}{j} \binom{i+j+l+2}{l} \right\} \end{aligned}$$

$$\begin{aligned}
 & -60 \binom{N-2}{k+1} \binom{i+j+1}{j} \binom{i+j+l+2}{l} + 8(-1)^{i+j+l} \binom{i+l+1}{i} \binom{i+j+l+2}{i+l+1} \\
 & -6 \binom{N-2}{k+1} \binom{i+l+1}{i} \binom{i+j+l+2}{i+l+1} - 3(-1)^{i+j+l} \binom{i+l+1}{l} \binom{i+j+l+2}{i+l+1} \\
 & + 71 \binom{N-1}{k+1} \binom{i+l+1}{l} \binom{i+j+l+2}{i+l+1} - 11(-1)^{k+l} \binom{i+j+1}{i} \binom{k+l+1}{k} \\
 & + 7(-1)^{k+l} \binom{i+j+1}{j} \binom{k+l+1}{k} + 11(-1)^{k+l} \binom{i+j+1}{i} \binom{k+l+1}{l} \\
 & - 7(-1)^{k+l} \binom{i+j+1}{j} \binom{k+l+1}{l} + 60 \binom{N-2}{j+1} \binom{k+l+1}{k} \binom{i+k+l+2}{i} \\
 & - 10(-1)^{i+k+l} \binom{k+l+1}{l} \binom{i+k+l+2}{i} + 26 \binom{N-1}{j+1} \binom{k+l+1}{l} \binom{i+k+l+2}{i} \\
 & + (-1)^{i+k+l+1} \binom{i+l+1}{i} \binom{i+k+l+2}{k} - 4 \binom{N-1}{j+1} \binom{i+l+1}{i} \binom{i+k+l+2}{k} \\
 & - 2(-1)^{i+k+l} \binom{i+l+1}{l} \binom{i+k+l+2}{k} - 44 \binom{N-2}{j} \binom{i+l+1}{l} \binom{i+k+l+2}{k} \\
 & - 26 \binom{N-2}{j+1} \binom{i+l+1}{l} \binom{i+k+l+2}{k} - 15 \binom{N-1}{j+1} \binom{i+k+1}{i} \binom{i+k+l+2}{l} \\
 & + (-1)^{i+k+l} \binom{i+k+1}{k} \binom{i+k+l+2}{l} - 30 \binom{N-2}{j+1} \binom{i+k+1}{k} \binom{i+k+l+2}{l} \\
 & + 5(-1)^{i+k+l} \binom{i+l+1}{i} \binom{i+k+l+2}{i+l+1} - 10 \binom{N-1}{j+1} \binom{i+l+1}{i} \binom{i+k+l+2}{i+l+1} \\
 & + (-1)^{i+k+l} \binom{i+l+1}{l} \binom{i+k+l+2}{i+l+1} - 18 \binom{N-2}{j+1} \binom{i+l+1}{l} \binom{i+k+l+2}{i+l+1} \\
 & - 14(-1)^{j+k+l} \binom{k+l+1}{l} \binom{j+k+l+2}{j} - 7(-1)^{j+k+l} \binom{j+l+1}{l} \binom{j+k+l+2}{k} \Big\} \\
 & \tag{3.49}
 \end{aligned}$$

and

$$\begin{aligned}
 \eta_{ijkl}^{(1)} = & -\frac{\eta(N)}{96} \left\{ \left[\left[\left(\binom{j+k+l+2}{l} + 5 \binom{j+k+l+2}{j+k+1} \right) (-1)^{l+1} + 3 \binom{i+l+1}{i} \right. \right. \right. \\
 & \left. \left. + 3 \binom{i+l+1}{l} \right] (-1)^{j+k} - 17(-1)^{i+j+k} \binom{i+j+k+2}{i} + \binom{i+j+k+2}{j+k+1} \right. \\
 & \left. \times \left[13(-1)^{i+j+k} + 54 \binom{N-2}{l} \right] \right] \binom{j+k+1}{j} + \left[-3(-1)^{j+k} \binom{i+l+1}{i} \right. \\
 & \left. - 3(-1)^{j+k} \binom{i+l+1}{l} + 17(-1)^{j+k+l} \binom{j+k+l+2}{l} + 7(-1)^{j+k+l} \binom{j+k+l+2}{j+k+1} \right]
 \end{aligned}$$

$$\begin{aligned}
 & +6 \binom{N-2}{i} \binom{j+k+l+2}{j+k+1} + \binom{i+j+k+2}{i} \left[(-1)^{i+j+k} + 6 \binom{N-1}{l+1} \right] \\
 & + \binom{i+j+k+2}{j+k+1} \left[(-1)^{i+j+k} + 6 \binom{N-1}{l+1} \right] \binom{j+k+1}{k} \\
 & -12(-1)^{i+j+k} \binom{i+j+1}{j} \binom{i+j+k+2}{k} + (-1)^{j+l+1} \binom{i+k+1}{i} \binom{j+l+1}{j} \\
 & + (-1)^{j+l+1} \binom{i+k+1}{k} \binom{j+l+1}{j} + (-1)^{j+l+1} \binom{i+k+1}{i} \binom{j+l+1}{l} \\
 & + (-1)^{j+l+1} \binom{i+k+1}{k} \binom{j+l+1}{l} - 3(-1)^{i+j+l} \binom{i+l+1}{i} \binom{i+j+l+2}{j} \\
 & -18 \binom{N-2}{k} \binom{i+l+1}{i} \binom{i+j+l+2}{j} + 18 \binom{N-2}{k+1} \binom{i+l+1}{i} \binom{i+j+l+2}{j} \\
 & -3(-1)^{i+j+l} \binom{i+l+1}{l} \binom{i+j+l+2}{j} + 3 \binom{N-1}{k+1} \binom{i+l+1}{l} \binom{i+j+l+2}{j} \\
 & +18 \binom{N-1}{k+1} \binom{i+j+1}{i} \binom{i+j+l+2}{l} - 6(-1)^{i+j+l} \binom{i+j+1}{j} \binom{i+j+l+2}{l} \\
 & -30 \binom{N-2}{k+1} \binom{i+j+1}{j} \binom{i+j+l+2}{l} + 3(-1)^{i+j+l} \binom{i+l+1}{i} \binom{i+j+l+2}{i+l+1} \\
 & +12 \binom{N-2}{k+1} \binom{i+l+1}{i} \binom{i+j+l+2}{i+l+1} + 3(-1)^{i+j+l} \binom{i+l+1}{l} \binom{i+j+l+2}{i+l+1} \\
 & +3 \binom{N-1}{k+1} \binom{i+l+1}{l} \binom{i+j+l+2}{i+l+1} + 7(-1)^{k+l} \binom{i+j+1}{i} \binom{k+l+1}{k} \\
 & -5(-1)^{k+l} \binom{i+j+1}{j} \binom{k+l+1}{k} + 17(-1)^{k+l} \binom{i+j+1}{i} \binom{k+l+1}{l} \\
 & -13(-1)^{k+l} \binom{i+j+1}{j} \binom{k+l+1}{l} - 18 \binom{N-2}{j+1} \binom{k+l+1}{k} \binom{i+k+l+2}{i} \\
 & -2(-1)^{i+k+l} \binom{k+l+1}{l} \binom{i+k+l+2}{i} - 10 \binom{N-1}{j+1} \binom{k+l+1}{l} \binom{i+k+l+2}{i} \\
 & + (-1)^{i+k+l+1} \binom{i+l+1}{i} \binom{i+k+l+2}{k} - 5 \binom{N-1}{j+1} \binom{i+l+1}{i} \binom{i+k+l+2}{k} \\
 & + (-1)^{i+k+l+1} \binom{i+l+1}{l} \binom{i+k+l+2}{k} + 10 \binom{N-2}{j} \binom{i+l+1}{l} \binom{i+k+l+2}{k}
 \end{aligned}$$

$$\begin{aligned}
 & -26 \binom{N-2}{j+1} \binom{i+l+1}{l} \binom{i+k+l+2}{k} + (-1)^{i+k+l+1} \binom{i+l+1}{i} \binom{i+k+l+2}{i+l+1} \\
 & -5 \binom{N-1}{j+1} \binom{i+l+1}{i} \binom{i+k+l+2}{i+l+1} - 3(-1)^{i+k+l} \binom{i+l+1}{l} \binom{i+k+l+2}{i+l+1} \\
 & -24 \binom{N-2}{j+1} \binom{i+l+1}{l} \binom{i+k+l+2}{i+l+1} + 4(-1)^{j+k+l} \binom{k+l+1}{l} \binom{j+k+l+2}{j} \Big\}.
 \end{aligned} \tag{3.50}$$

We have checked the correctness of these expressions by comparing with fixed- N computations up to $N = 14$.

3.3.2 $\kappa_{ijkl}^{(2)}$, $\eta_{ijkl}^{(2a)}$ and $\eta_{ijkl}^{(2b)}$

For this final set of couplings we have the following relations

$$\kappa_{ijkl}^{(2)} + \kappa_{ijlk}^{(2)} = 0, \quad [\text{anti-symmetry}] \tag{3.51}$$

$$\kappa_{ijkl}^{(2)} = \kappa_{jikl}^{(2)}, \quad [\text{symmetry of } d_4] \tag{3.52}$$

$$\eta_{ijkl}^{(2a)} = 3\kappa_{ij(k+l+1)}^{(2)} \binom{k+l+1}{k} + 2\kappa_{ijkl}^{(2)}, \quad [\text{gBRST (a)}] \tag{3.53}$$

$$\eta_{ijkl}^{(2b)} = 2\kappa_{lik}^{(2)}, \quad [\text{gBRST (b)}] \tag{3.54}$$

$$\begin{aligned}
 \eta_{ijkl}^{(2a)} = & - \sum_{s_1=0}^i \sum_{s_2=0}^j \sum_{s_3=0}^k \frac{(s_1+s_2+s_3+l)!}{s_1! s_2! s_3! l!} (-1)^{s_1+s_2+s_3+l} \\
 & \times \eta_{(i-s_1)(j-s_2)(k-s_3)(s_1+s_2+s_3+l)}^{(2a)}, \quad [\text{anti-gBRST (a)}]
 \end{aligned} \tag{3.55}$$

$$\begin{aligned}
 \eta_{ijkl}^{(2b)} = & \eta_{ikjl}^{(2a)} - \eta_{ijkl}^{(2a)} + \sum_{s_1=0}^i \sum_{s_2=0}^j \sum_{s_3=0}^k \frac{(s_1+s_2+s_3+l)!}{s_1! s_2! s_3! l!} (-1)^{s_1+s_2+s_3+l} \\
 & \times \eta_{(i-s_1)(j-s_2)(k-s_3)(s_1+s_2+s_3+l)}^{(2b)}. \quad [\text{anti-gBRST (b)}]
 \end{aligned} \tag{3.56}$$

Note that eqs. (3.51) and (3.53) can be combined to express $\eta_{ijkl}^{(2a)}$ in terms of the class III coupling $\kappa_{ijk}^{(2)}$ as

$$\eta_{ijkl}^{(2a)} + \eta_{ijlk}^{(2a)} = 3\kappa_{ij(k+l+1)}^{(2)} \binom{k+l+2}{k+1}. \tag{3.57}$$

with

$$\varepsilon^{ab} = \frac{1+(-1)^N}{2} i^N \eta(N) \delta^{ab} (\Delta \cdot p_1)^N, \quad (4.2)$$

$$\varepsilon_{\mu}^{ab,c_1} = \frac{1+(-1)^N}{2} i^{N-1} \Delta_{\mu} f^{ac_1b} \sum_{i+j=N-3} \eta_{ij}(\Delta \cdot p_1) (\Delta \cdot p_3)^i (\Delta \cdot p_2)^{j+1}, \quad (4.3)$$

$$\begin{aligned} \varepsilon_{\mu\nu}^{ab,c_1c_2}(p_1, p_2, p_3, p_4) &= \frac{1+(-1)^N}{2} i^N \Delta_{\mu} \Delta_{\nu} \\ &\times \left\{ (ff)^{ac_1c_2b} \sum_{i+j+k=N-4} \eta_{ijk}^{(1)}(\Delta \cdot p_1) (\Delta \cdot p_3)^i (\Delta \cdot p_4)^j (\Delta \cdot p_2)^{k+1} \right. \\ &+ d_4^{ac_1c_2b} \sum_{i+j+k=N-4} \eta_{ijk}^{(2)}(\Delta \cdot p_1) (\Delta \cdot p_3)^i (\Delta \cdot p_4)^j (\Delta \cdot p_2)^{k+1} \\ &+ d_{4ff}^{ac_1c_2b} \sum_{i+j+k=N-4} \eta_{ijk}^{(3)}(\Delta \cdot p_1) (\Delta \cdot p_3)^i (\Delta \cdot p_4)^j (\Delta \cdot p_2)^{k+1} \Big\} \\ &+ [(p_3, \mu, c_1) \leftrightarrow (p_4, \nu, c_2)], \end{aligned} \quad (4.4)$$

$$\begin{aligned} \varepsilon_{\mu\nu\rho}^{ab,c_1c_2c_3}(p_1, p_2, p_3, p_4, p_5) &= -\frac{1+(-1)^N}{2} i^{N-1} \Delta_{\mu} \Delta_{\nu} \Delta_{\rho} \\ &\times \left\{ (fff)^{ac_1c_2c_3b} \sum_{i+j+k+l=N-5} \eta_{ijkl}^{(1)}(\Delta \cdot p_1) (\Delta \cdot p_3)^i (\Delta \cdot p_4)^j (\Delta \cdot p_5)^k (\Delta \cdot p_2)^{l+1} \right. \\ &+ d_4^{ac_1c_2c_3b} \sum_{i+j+k+l=N-5} \eta_{ijkl}^{(2a)}(\Delta \cdot p_1) (\Delta \cdot p_3)^i (\Delta \cdot p_4)^j (\Delta \cdot p_5)^k (\Delta \cdot p_2)^{l+1} \\ &+ d_{4f}^{abc_1c_2c_3} \sum_{i+j+k+l=N-5} \eta_{ijkl}^{(2b)}(\Delta \cdot p_1) (\Delta \cdot p_3)^i (\Delta \cdot p_4)^j (\Delta \cdot p_5)^k (\Delta \cdot p_2)^{l+1} \Big\} \\ &+ \text{permutations} \end{aligned} \quad (4.5)$$

where the ‘+ permutations’ in the $O(g_s^3)$ rule in eq. (4.5) denotes the fact that all permutations of the gluonic quantities (momenta, Lorentz and colour indices) have to be added. Note that p_2 in ε^{ab} in eq. (4.2) was eliminated using momentum conservation, $p_2 = -p_1$. This then agrees with eq. (5.20) in [10] after dividing the latter by i^N , as discussed above. Similarly, after performing the summation, the $O(g_s)$ rule exactly matches eq. (5.21) in [10]. At $O(g_s^2)$, our expression for $\varepsilon_{\mu\nu}^{ab,c_1c_2}(p_1, p_2, p_3, p_4)$ should be compared against eq. (5.22) in [10]. With $\eta_{ijk}^{(1)}$ given by eq. (3.27) and $\eta_{ijk}^{(2)}$ by eq. (3.34), we find exact agreement with that expression.⁵ Finally the Feynman rule for the d_{4ff} part of the ghost operator is computed using

$$\begin{aligned} \eta_{ijk}^{(3)} &= 2 \left\{ (c_1 - c_2)(-1)^{i+k} \left[\binom{i+k+1}{i} - \binom{i+k+1}{k} \right] + (-1)^{j+k+1} \left[(c_1 + 2c_2) \binom{j+k+1}{j} \right. \right. \\ &\quad \left. \left. + (2c_1 + c_2) \binom{j+k+1}{k} \right] + \binom{i+j+1}{i} \left[(2c_1 + c_2)(-1)^{i+j} - c_1 \binom{N-2}{k+1} \right] \right\} \end{aligned}$$

⁵The term in our expression proportional to $(ff)^{ac_1c_2b}$ should be compared to the $f^{a_1a_3a} f^{a_2a_4a}$ part of eq. (5.22) in [10] while our $d_4^{ac_1c_2b}$ rule should be compared to the one proportional to $d_4^{a_1a_2a_3a_4}/C_A$.

$$\begin{aligned}
 & + \binom{i+j+1}{j} \left[c_1(-1)^{i+j} + 2c_2(-1)^{i+j} - 2c_1 \binom{N-2}{k} - c_1 \binom{N-2}{k+1} \right. \\
 & - c_2 \binom{N-1}{k+1} \left. \right] + \binom{i+k+1}{i} \left[2c_1 \binom{N-2}{j} + c_2 \binom{N-1}{j+1} \right] \\
 & - \binom{i+k+1}{k} \left[2c_1 \binom{N-2}{j} + c_2 \binom{N-1}{j+1} \right] + \binom{j+k+1}{j} \left[2c_1 \binom{N-2}{i} \right. \\
 & \left. + c_1 \binom{N-2}{i+1} + c_2 \binom{N-1}{i+1} \right] + c_1 \binom{N-2}{i+1} \binom{j+k+1}{k} \left. \right\}
 \end{aligned} \tag{4.6}$$

which follows from eqs. (3.38) and (3.40). As discussed in section 3.2.3, the free parameters c_1, c_2 can be determined by a computation of fixed- N OMEs.

4.2 Alien gluon operators

Next we derive the Feynman rules for the gluonic EOM operator, whose perturbative expansion can be written as

with

$$\begin{aligned}
 \mathcal{G}_{\mu\nu}^{c_1 c_2}(p_1, p_2) &= \frac{1 + (-1)^N}{2} i^N \eta(N) \delta^{c_1 c_2} (\Delta \cdot p_1)^{N-2} \\
 &\quad \times [2p_1^2 \Delta_\mu \Delta_\nu - (\Delta \cdot p_1)(\Delta_\mu p_{1\nu} + \Delta_\nu p_{1\mu})],
 \end{aligned} \tag{4.7}$$

$$\begin{aligned}
 \mathcal{G}_{\mu\nu\rho}^{c_1 c_2 c_3}(p_1, p_2, p_3) &= -\frac{1 + (-1)^N}{2} i^{N-1} f^{c_1 c_2 c_3} \\
 &\quad \times \left\{ \eta(N) (\Delta \cdot p_1)^{N-2} \Delta_\mu [p_{3\nu} \Delta_\rho - g_{\nu\rho} (\Delta \cdot p_3) \right. \\
 &\quad \left. + \Delta_\rho (p_2 + p_3)_\nu] + \Delta_\nu \Delta_\rho [p_1^2 \Delta_\mu - p_{1\mu} (\Delta \cdot p_1)] \right. \\
 &\quad \left. \times \sum_{i+j=N-3} \kappa_{ij} (\Delta \cdot p_2)^i (\Delta \cdot p_3)^j \right\} + \text{permutations},
 \end{aligned} \tag{4.8}$$

$$\begin{aligned}
 \mathcal{G}_{\mu\nu\rho\sigma}^{c_1 c_2 c_3 c_4}(p_1, p_2, p_3, p_4) &= \frac{1 + (-1)^N}{2} i^{N-2} f^{c_1 c_2 x} f^{x c_3 c_4} \left\{ [\Delta_\nu \Delta_\rho \Delta_\sigma (p_1 + 2p_2)_\mu \right. \\
 &\quad \left. - g_{\mu\nu} \Delta_\rho \Delta_\sigma (\Delta \cdot p_2)] \sum_{i+j=N-3} \kappa_{ij} (\Delta \cdot p_3)^i (\Delta \cdot p_4)^j \right. \\
 &\quad \left. - g_{\nu\rho} \Delta_\mu \Delta_\sigma (\Delta \cdot p_1)^{N-2} + [p_1^2 \Delta_\mu - p_{1\mu} (\Delta \cdot p_1)] \Delta_\nu \Delta_\rho \Delta_\sigma \right. \\
 &\quad \left. \times \sum_{i+j+k=N-4} \kappa_{ijk}^{(1)} (\Delta \cdot p_2)^i (\Delta \cdot p_3)^j (\Delta \cdot p_4)^k \right\}
 \end{aligned}$$

$$\begin{aligned}
 & + \frac{1 + (-1)^N}{2} [p_1^2 \Delta_\mu - p_{1\mu} (\Delta \cdot p_1)] \Delta_\nu \Delta_\rho \Delta_\sigma \\
 & \times \left\{ d_{4ff}^{c_1 c_2 c_3 c_4} \sum_{i+j+k=N-4} \kappa_{ijk}^{(2)} (\Delta \cdot p_2)^i (\Delta \cdot p_3)^j (\Delta \cdot p_4)^j \right. \\
 & \quad \left. + d_{4ff}^{c_1 c_2 c_3 c_4} \sum_{i+j+k=N-4} \kappa_{ijk}^{(3)} (\Delta \cdot p_2)^i (\Delta \cdot p_3)^j (\Delta \cdot p_4)^j \right\} \\
 & + \text{permutations}, \tag{4.9} \\
 \mathcal{G}_{\mu\nu\rho\sigma\tau}^{c_1 c_2 c_3 c_4 c_5}(p_1, p_2, p_3, p_4, p_5) = & \frac{1 + (-1)^N}{2} i^{N-1} f^{c_1 c_2 x} f^{x c_3 y} f^{y c_4 c_5} \\
 & \times \left\{ -g_{\mu\rho} \Delta_\nu \Delta_\sigma \Delta_\tau \sum_{i+j=N-3} \kappa_{ij} (\Delta \cdot p_4)^i (\Delta \cdot p_5)^j \right. \\
 & + \Delta_\rho \Delta_\sigma \Delta_\tau [(p_1 + 2p_2)_\mu \Delta_\nu - (\Delta \cdot p_2) g_{\mu\nu}] \\
 & \times \sum_{i+j+k=N-4} \kappa_{ijk}^{(1)} (\Delta \cdot p_3)^i (\Delta \cdot p_4)^j (\Delta \cdot p_5)^k \\
 & + [p_1^2 \Delta_\mu - p_{1\mu} (\Delta \cdot p_1)] \Delta_\nu \Delta_\rho \Delta_\sigma \Delta_\tau \\
 & \times \sum_{i+j+k+l=N-5} \kappa_{ijkl}^{(1)} (\Delta \cdot p_2)^i (\Delta \cdot p_3)^j (\Delta \cdot p_4)^k (\Delta \cdot p_5)^l \Big\} \\
 & + \frac{1 + (-1)^N}{2} i^{N-1} d_{4f}^{c_1 c_2 c_3 c_4 c_5} \left\{ \Delta_\mu \Delta_\nu \Delta_\rho [(p_4 + 2p_5)_\sigma \Delta_\tau \right. \\
 & - (\Delta \cdot p_5) g_{\sigma\tau}] \sum_{i+j+k=N-4} \kappa_{ijk}^{(2)} (\Delta \cdot p_1)^i (\Delta \cdot p_2)^j (\Delta \cdot p_3)^k \\
 & + [p_1^2 \Delta_\mu - p_{1\mu} (\Delta \cdot p_1)] \Delta_\nu \Delta_\rho \Delta_\sigma \Delta_\tau \\
 & \times \sum_{i+j+k+l=N-5} \kappa_{ijkl}^{(2)} (\Delta \cdot p_2)^i (\Delta \cdot p_3)^j (\Delta \cdot p_4)^k (\Delta \cdot p_5)^l \Big\} \\
 & + \text{permutations}, \tag{4.10}
 \end{aligned}$$

where again all permutations of gluon momenta, Lorentz and colour indices have to be added, if indicated by ‘+ permutations’. Note that p_2 in $\mathcal{G}_{\mu\nu}^{c_1 c_2}(p_1, p_2)$ in eq. (4.7) was again eliminated using momentum conservation. This then agrees with eq. (5.23) in [10] and eq. (243) in [38] after dividing the latter rules by i^N to match to our conventions.

For the $O(g_s)$ EOM vertex three contributions need to be taken into account,

- the non-Abelian part of the field strength in the class I operator with $D \rightarrow \partial$,
- the $O(g_s)$ part of the covariant derivative acting on the Abelian part of the field strength in the class I operator and
- the class II operator, cf. eq. (2.10), with $D \rightarrow \partial$ and keeping only the Abelian part of the field strength.

Our result matches the corresponding rules in the literature, cf. eq. (5.24) in [10] and eq. (244) in [38] respectively (again after dividing by the overall i^N).

Next the four-gluon vertex gets four contributions,

- the $O(g_s)$ part of the covariant derivative acting on the non-Abelian part of the field strength in the class I operator,
- the non-Abelian part of the field strength in the class II operator with $D \rightarrow \partial$,
- the $O(g_s)$ part of the covariant derivative acting on the Abelian part of the field strength in the class II operator and
- the class III operator, cf. eq. (2.11), with $D \rightarrow \partial$ and keeping only the Abelian part of the field strength.

The second and third contributions depend on the lower-order coupling κ_{ij} , while the fourth one is written in terms of the couplings $\kappa_{ijk}^{(1)}$, $\kappa_{ijk}^{(2)}$ and $\kappa_{ijk}^{(3)}$ given by eqs. (3.26), (3.33) and (3.40) respectively. The $(f f)$ and d_4 parts of our rule agree with eq. (5.25) in [10],⁶ while the d_{4ff} part is new.

Finally, as a new result,⁷ we consider the five-gluon vertex $\mathcal{G}_{\mu\nu\rho\sigma\tau}^{c_1 c_2 c_3 c_4 c_5}(p_1, p_2, p_3, p_4, p_5)$ in eq. (4.10). Again we need to take into account higher-order contributions of the lower-point vertices. In particular, the $(f f f)$ part of the five-gluon rule gets four contributions,

- the $O(g_s)$ part of the covariant derivative acting on the non-Abelian part of the field strength in the class II operator,
- the non-Abelian part of the field strength in the class III operator with $D \rightarrow \partial$,
- the $O(g_s)$ part of the covariant derivative acting on the Abelian part of the field strength in the class III operator,
- the class IV operator, cf. eq. (2.12), with $D \rightarrow \partial$ and keeping only the Abelian part of the field strength.

On the other hand the d_{4f} part only gets three contributions,

- the non-Abelian part of the field strength in the class III operator with $D \rightarrow \partial$,
- the $O(g_s)$ part of the covariant derivative acting on the Abelian part of the field strength in the class III operator and
- the class IV operator, cf. eq. (2.12), with $D \rightarrow \partial$ and keeping only the Abelian part of the field strength.

⁶Note however that our result proportional to d_4 needs to be multiplied by a symmetry factor of $1/4!$ to match eq. (5.25) in [10], which is again a consequence of differing conventions.

⁷The corresponding result within the framework of ref. [10] was recently announced in a conference talk [15].

4.3 Alien quark operators

Finally in this section we provide the Feynman rules for the alien quark operators presented in eqs. (2.9)–(2.11). As mentioned above, these operators are written in terms of the same couplings as those in the gluon EOM operators. Assuming the momenta of the external quark fields to be p_1 and p_2 we have the following perturbative expansion

with

$$\mathcal{Q}(p_1, p_2) = 0, \quad (4.11)$$

$$\mathcal{Q}_\mu^{c_1}(p_1, p_2, p_3) = -\frac{1+(-1)^N}{2} i^N \eta(N) T^{c_1} \Delta_\mu \not{\Delta} (\Delta \cdot p_3)^{N-2}, \quad (4.12)$$

$$\mathcal{Q}_{\mu\nu}^{c_1 c_2}(p_1, p_2, p_3, p_4) = [1 + (-1)^N] i^{N-1} T^a f^{a c_1 c_2} \Delta_\mu \Delta_\nu \not{\Delta} \sum_{i+j=N-3} \kappa_{ij}^{(1)} (\Delta \cdot p_3)^i (\Delta \cdot p_4)^j, \quad (4.13)$$

$$\begin{aligned} \mathcal{Q}_{\mu\nu\rho}^{c_1 c_2 c_3}(p_1, p_2, p_3, p_4, p_5) = & [1 + (-1)^N] i^N T^a \Delta_\mu \Delta_\nu \Delta_\rho \not{\Delta} \left\{ f^{a c_1 x} f^{c_2 c_3 x} \sum_{i+j+k=N-4} \kappa_{ijk}^{(1)} (\Delta \cdot p_3)^i (\Delta \cdot p_4)^j (\Delta \cdot p_5)^k \right. \\ & + d_4^{a c_1 c_2 c_3} \sum_{i+j+k=N-4} \kappa_{ijk}^{(2)} (\Delta \cdot p_3)^i (\Delta \cdot p_4)^j (\Delta \cdot p_5)^k \\ & \left. + d_{4ff}^{a c_1 c_2 c_3} \sum_{i+j+k=N-4} \kappa_{ijk}^{(3)} (\Delta \cdot p_3)^i (\Delta \cdot p_4)^j (\Delta \cdot p_5)^k \right\} + [(p_3, \mu, c_1) \leftrightarrow (p_4, \nu, c_2)] \\ & + [(p_3, \mu, c_1) \rightarrow (p_5, \rho, c_3) \rightarrow (p_4, \nu, c_2) \rightarrow (p_3, \mu, c_1)]. \end{aligned} \quad (4.14)$$

The vertices with up to two additional gluons can be compared against the results presented in eqs. (5.17)–(5.19) of [10]. Dividing the latter by i^N to match to our conventions, we find exact agreement. Note that eqs. (4.13)–(4.14) contain an additional factor of two coming from the $[(p_4, \nu, c_2) \leftrightarrow (p_5, \rho, c_3)]$ permutation. This directly follows from the (anti-)symmetry properties of the κ -couplings, cf. eqs. (3.19), (3.28) and (3.36). Finally, because the κ -couplings enter the quark operator at one order in the strong coupling lower than in the gluon EOM one, we can push the perturbative expansion of the quark operator to one order higher. Consequently we also present the quark operator vertex at $\mathcal{O}(g_s^4)$ with four additional gluons

We find

$$\begin{aligned}
 & \mathcal{Q}_{\mu\nu\rho\sigma}^{c_1 c_2 c_3 c_4}(p_1, p_2, p_3, p_4, p_5, p_6) \\
 &= -\frac{1+(-1)^N}{2} i^{N-1} T^a \Delta_\mu \Delta_\nu \Delta_\rho \Delta_\sigma \not{X} \left\{ (f f f)^{a c_1 c_2 c_3 c_4} \sum_{i+j+k+l=N-5} \kappa_{ijkl}^{(1)} \right. \\
 &\quad \times (\Delta \cdot p_1)(\Delta \cdot p_3)^i (\Delta \cdot p_4)^j (\Delta \cdot p_5)^k (\Delta \cdot p_2)^{l+1} \\
 &\quad \left. + d_{4f}^{a c_1 c_2 c_3 c_4} \sum_{i+j+k+l=N-5} \kappa_{ijkl}^{(2)} (\Delta \cdot p_1)(\Delta \cdot p_3)^i (\Delta \cdot p_4)^j (\Delta \cdot p_5)^k (\Delta \cdot p_2)^{l+1} \right\} + \text{permutations}
 \end{aligned} \tag{4.15}$$

with all permutations of the gluonic quantities (momenta, Lorentz and colour indices) to be added.

5 Conclusions

The kernels for parton evolution equations in QCD, i.e. splitting functions or the corresponding anomalous dimensions as their Mellin transforms, can be conveniently determined from the ultraviolet singularities of off-shell Green's functions with insertions of gauge-invariant twist-two spin- N operators. The renormalization of these OMEs, though, requires the computation of unphysical counterterms for the associated set of alien operators, which effectively describe vertices of two gluons, ghosts or quarks with any number $n \geq 0$ of additional gluons. The couplings of these alien operators (EOM and ghost operators) are restricted by the fundamental symmetries, particularly the gBRST relations, which reflect the gauge theory characteristics of QCD.

The set of constraints for these couplings admits explicit solutions, valid for any spin N , which can be obtained using algorithms for symbolic summation to solve the recurrence relations. A small number of boundary conditions in these solutions can be derived from the computation of the relevant OMEs at specific fixed values of N . In addition, we have observed that the constraints contain a hierarchy, such that the function space of the couplings of alien operators with $n+1$ gluons can be derived from that of the n -gluon aliens. Thus, the basic ingredients in this bootstrap turn out to be the EOM and ghost operators with the smallest number of additional gluons at a given loop order.

We have provided results for all one-loop alien operator couplings needed in the renormalization of OMEs with physical (gauge-invariant) operators up to four loops, which represents the current frontier in splitting function computations. This includes in particular the gluon EOM operator with five gluons attached, which is a new result. The all- N solutions for the couplings that we have obtained can all be related to the fundamental one-loop counterterm $\eta(N)$ for the EOM and ghost operators of class I involving only two gluons or ghosts. We have also derived the corresponding Feynman rules and, whenever possible, compared them to those in the literature, finding full agreement.

A `Mathematica` file with our results for the all- N couplings necessary for the renormalization up to four loops is made available as supplementary material. We note that the expressions collected in this file have the fundamental one-loop counterterm $\eta(N)$ divided out.

The symmetries and the structure of the alien operators, that we have exploited in this study, are independent of the order of perturbation theory. Thus, we expect also analytic all- N solutions beyond one loop for the couplings of the alien operators of class II and higher. We leave this task to future studies.

Acknowledgments

The Feynman diagrams in this work are drawn using *FeynGame* [54, 55].

This work has been supported by the EU's Marie Skłodowska-Curie grant 101104792, *QCDchallenge*; the DFG through the Research Unit FOR 2926, *Next Generation pQCD for Hadron Structure: Preparing for the EIC*, project number 40824754, DFG grant MO 1801/4-2, the ERC Advanced Grant 101095857 *Conformal-EIC*; and by grant K143451 of the National Research, Development and Innovation Fund in Hungary.

Data Availability Statement. This article has no associated data or the data will not be deposited.

Code Availability Statement. This article has no associated code or the code will not be deposited.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] J.A. Dixon and J.C. Taylor, *Renormalization of wilson operators in gauge theories*, *Nucl. Phys. B* **78** (1974) 552 [[INSPIRE](#)].
- [2] H. Kluberg-Stern and J.B. Zuber, *Renormalization of Non-Abelian Gauge Theories in a Background Field Gauge. Part 1. Green Functions*, *Phys. Rev. D* **12** (1975) 482 [[INSPIRE](#)].
- [3] H. Kluberg-Stern and J.B. Zuber, *Renormalization of Non-Abelian Gauge Theories in a Background Field Gauge. Part 2. Gauge Invariant Operators*, *Phys. Rev. D* **12** (1975) 3159 [[INSPIRE](#)].
- [4] S.D. Joglekar and B.W. Lee, *General Theory of Renormalization of Gauge Invariant Operators*, *Ann. Phys.* **97** (1976) 160 [[INSPIRE](#)].
- [5] S.D. Joglekar, *Local Operator Products in Gauge Theories. Part 1*, *Ann. Phys.* **108** (1977) 233 [[INSPIRE](#)].
- [6] S.D. Joglekar, *Local Operator Products in Gauge Theories. Part 2*, *Ann. Phys.* **109** (1977) 210 [[INSPIRE](#)].
- [7] R. Hamberg and W.L. van Neerven, *The Correct renormalization of the gluon operator in a covariant gauge*, *Nucl. Phys. B* **379** (1992) 143 [[INSPIRE](#)].
- [8] G. Falcioni and F. Herzog, *Renormalization of gluonic leading-twist operators in covariant gauges*, *JHEP* **05** (2022) 177 [[arXiv:2203.11181](#)] [[INSPIRE](#)].
- [9] G. Falcioni, F. Herzog, S. Moch, A. Pelloni and A. Vogt, *Four-loop splitting functions in QCD — The quark-to-gluon case*, *Phys. Lett. B* **856** (2024) 138906 [[arXiv:2404.09701](#)] [[INSPIRE](#)].

- [10] T. Gehrmann, A. von Manteuffel and T.-Z. Yang, *Renormalization of twist-two operators in covariant gauge to three loops in QCD*, *JHEP* **04** (2023) 041 [[arXiv:2302.00022](#)] [[INSPIRE](#)].
- [11] T. Gehrmann, A. von Manteuffel, V. Sotnikov and T.-Z. Yang, *Complete N_f^2 contributions to four-loop pure-singlet splitting functions*, *JHEP* **01** (2024) 029 [[arXiv:2308.07958](#)] [[INSPIRE](#)].
- [12] G. Curci and R. Ferrari, *On a Class of Lagrangian Models for Massive and Massless Yang-Mills Fields*, *Nuovo Cim. A* **32** (1976) 151 [[INSPIRE](#)].
- [13] I. Ojima, *Another BRS Transformation*, *Prog. Theor. Phys.* **64** (1980) 625 [[INSPIRE](#)].
- [14] L. Baulieu and J. Thierry-Mieg, *The Principle of BRS Symmetry: An Alternative Approach to Yang-Mills Theories*, *Nucl. Phys. B* **197** (1982) 477 [[INSPIRE](#)].
- [15] T. Gehrmann, A. von Manteuffel and T.-Z. Yang, *Leading Twist-Two Gauge-Variant Counterterms*, *PoS LL2024* (2024) 087 [[arXiv:2409.10303](#)] [[INSPIRE](#)].
- [16] S. Moch and S. Van Thurenhout, *Renormalization of non-singlet quark operator matrix elements for off-forward hard scattering*, *Nucl. Phys. B* **971** (2021) 115536 [[arXiv:2107.02470](#)] [[INSPIRE](#)].
- [17] S. Van Thurenhout, *Basis transformation properties of anomalous dimensions for hard exclusive processes*, *Nucl. Phys. B* **1000** (2024) 116464 [[arXiv:2309.16236](#)] [[INSPIRE](#)].
- [18] D. Zeilberger, *The method of creative telescoping*, *J. Symb. Comput.* **11** (1991) 195.
- [19] R.W. Gosper, *Decision procedure for indefinite hypergeometric summation*, *Proc. Natl. Acad. Sci. U.S.A.* **75** (1978) 40.
- [20] C. Schneider, *The summation package sigma: Underlying principles and a rhombus tiling application*, *Discrete Math. Theor. Comput. Sci.* **6** (2004) 365.
- [21] C. Schneider, *Symbolic summation assists combinatorics*, *Sémin. Lothar. Comb.* **56** (2007) 1.
- [22] C. Schneider, *Simplifying Multiple Sums in Difference Fields*, in the proceedings of the LHCPhenoNet School: Integration, Summation and Special Functions in Quantum Field Theory, Linz, Austria, 9–13 July 2012, pp. 325–360 [[DOI:10.1007/978-3-7091-1616-6_14](#)] [[arXiv:1304.4134](#)] [[INSPIRE](#)].
- [23] C. Schneider, *Modern Summation Methods for Loop Integrals in Quantum Field Theory: The Packages Sigma, EvaluateMultiSums and SumProduction*, *J. Phys. Conf. Ser.* **523** (2014) 012037 [[arXiv:1310.0160](#)] [[INSPIRE](#)].
- [24] S. Moch, B. Ruijl, T. Ueda, J.A.M. Vermaseren and A. Vogt, *Four-Loop Non-Singlet Splitting Functions in the Planar Limit and Beyond*, *JHEP* **10** (2017) 041 [[arXiv:1707.08315](#)] [[INSPIRE](#)].
- [25] G. Falcioni, F. Herzog, S. Moch and A. Vogt, *Four-loop splitting functions in QCD — The quark-quark case*, *Phys. Lett. B* **842** (2023) 137944 [[arXiv:2302.07593](#)] [[INSPIRE](#)].
- [26] G. Falcioni, F. Herzog, S. Moch and A. Vogt, *Four-loop splitting functions in QCD — The gluon-to-quark case*, *Phys. Lett. B* **846** (2023) 138215 [[arXiv:2307.04158](#)] [[INSPIRE](#)].
- [27] S. Moch, B. Ruijl, T. Ueda, J. Vermaseren and A. Vogt, *Additional moments and x -space approximations of four-loop splitting functions in QCD*, *Phys. Lett. B* **849** (2024) 138468 [[arXiv:2310.05744](#)] [[INSPIRE](#)].
- [28] B. Ruijl, T. Ueda and J.A.M. Vermaseren, *Forcer, a FORM program for the parametric reduction of four-loop massless propagator diagrams*, *Comput. Phys. Commun.* **253** (2020) 107198 [[arXiv:1704.06650](#)] [[INSPIRE](#)].
- [29] J.A.M. Vermaseren, *New features of FORM*, [math-ph/0010025](#) [[INSPIRE](#)].

- [30] J. Kuipers, T. Ueda, J.A.M. Vermaseren and J. Vollinga, *FORM version 4.0*, *Comput. Phys. Commun.* **184** (2013) 1453 [[arXiv:1203.6543](#)] [[INSPIRE](#)].
- [31] B. Ruijl, T. Ueda and J. Vermaseren, *FORM version 4.2*, [arXiv:1707.06453](#) [[INSPIRE](#)].
- [32] F. Herzog, S. Moch, B. Ruijl, T. Ueda, J.A.M. Vermaseren and A. Vogt, *Five-loop contributions to low- N non-singlet anomalous dimensions in QCD*, *Phys. Lett. B* **790** (2019) 436 [[arXiv:1812.11818](#)] [[INSPIRE](#)].
- [33] V.A. Smirnov, *Asymptotic expansions in limits of large momenta and masses*, *Commun. Math. Phys.* **134** (1990) 109 [[INSPIRE](#)].
- [34] V.A. Smirnov, *Asymptotic expansions in momenta and masses and calculation of Feynman diagrams*, *Mod. Phys. Lett. A* **10** (1995) 1485 [[hep-th/9412063](#)] [[INSPIRE](#)].
- [35] V.A. Smirnov, *Applied asymptotic expansions in momenta and masses*, in *Springer Tracts in Modern Physics* **177**, Springer (2002) [[DOI:10.1007/3-540-44574-9](#)] [[INSPIRE](#)].
- [36] Maplesoft, a division of Waterloo Maple Inc., *Maple*, version 2019 (2019).
- [37] Y. Matiounine, J. Smith and W.L. van Neerven, *Two loop operator matrix elements calculated up to finite terms*, *Phys. Rev. D* **57** (1998) 6701 [[hep-ph/9801224](#)] [[INSPIRE](#)].
- [38] J. Blümlein, P. Marquard, C. Schneider and K. Schönwald, *The two-loop massless off-shell QCD operator matrix elements to finite terms*, *Nucl. Phys. B* **980** (2022) 115794 [[arXiv:2202.03216](#)] [[INSPIRE](#)].
- [39] E.G. Floratos, D.A. Ross and C.T. Sachrajda, *Higher Order Effects in Asymptotically Free Gauge Theories: The Anomalous Dimensions of Wilson Operators*, *Nucl. Phys. B* **129** (1977) 66 [[INSPIRE](#)].
- [40] E.G. Floratos, D.A. Ross and C.T. Sachrajda, *Higher Order Effects in Asymptotically Free Gauge Theories. Part 2. Flavor Singlet Wilson Operators and Coefficient Functions*, *Nucl. Phys. B* **152** (1979) 493 [[INSPIRE](#)].
- [41] R. Mertig and W.L. van Neerven, *The Calculation of the two loop spin splitting functions $P_{ij}^{(1)}(x)$* , *Z. Phys. C* **70** (1996) 637 [[hep-ph/9506451](#)] [[INSPIRE](#)].
- [42] S. Kumano and M. Miyama, *Two loop anomalous dimensions for the structure function h_1* , *Phys. Rev. D* **56** (1997) R2504 [[hep-ph/9706420](#)] [[INSPIRE](#)].
- [43] A. Hayashigaki, Y. Kanazawa and Y. Koike, *Next-to-leading order q^2 evolution of the transversity distribution $h_1(x, q^2)$* , *Phys. Rev. D* **56** (1997) 7350 [[hep-ph/9707208](#)] [[INSPIRE](#)].
- [44] I. Bierenbaum, J. Blümlein and S. Klein, *Mellin Moments of the $O(\alpha_s^3)$ Heavy Flavor Contributions to unpolarized Deep-Inelastic Scattering at $Q^2 \gg m^2$ and Anomalous Dimensions*, *Nucl. Phys. B* **820** (2009) 417 [[arXiv:0904.3563](#)] [[INSPIRE](#)].
- [45] S.W.G. Klein, *Mellin Moments of Heavy Flavor Contributions to $F_2(x, Q^2)$ at NNLO*, Ph.D. Thesis, Technical University of Dortmund, Dortmund, Germany (2009) [[DOI:10.1007/978-3-642-23286-2](#)] [[arXiv:0910.3101](#)] [[INSPIRE](#)].
- [46] J. Blümlein, *On the anomalous dimension of the transversity distribution $h_1(x, Q^2)$* , *Eur. Phys. J. C* **20** (2001) 683 [[hep-ph/0104099](#)] [[INSPIRE](#)].
- [47] V.N. Velizhanin, *Four loop anomalous dimension of the second moment of the non-singlet twist-2 operator in QCD*, *Nucl. Phys. B* **860** (2012) 288 [[arXiv:1112.3954](#)] [[INSPIRE](#)].
- [48] V.N. Velizhanin, *Four-loop anomalous dimension of the third and fourth moments of the nonsinglet twist-2 operator in QCD*, *Int. J. Mod. Phys. A* **35** (2020) 2050199 [[arXiv:1411.1331](#)] [[INSPIRE](#)].

- [49] S. Moch, B. Ruijl, T. Ueda, J.A.M. Vermaseren and A. Vogt, *Low moments of the four-loop splitting functions in QCD*, *Phys. Lett. B* **825** (2022) 136853 [[arXiv:2111.15561](#)] [[INSPIRE](#)].
- [50] G. Falcioni, F. Herzog, S. Moch, J. Vermaseren and A. Vogt, *The double fermionic contribution to the four-loop quark-to-gluon splitting function*, *Phys. Lett. B* **848** (2024) 138351 [[arXiv:2310.01245](#)] [[INSPIRE](#)].
- [51] T. Gehrmann, A. von Manteuffel, V. Sotnikov and T.-Z. Yang, *The $N_f C_F^3$ contribution to the non-singlet splitting function at four-loop order*, *Phys. Lett. B* **849** (2024) 138427 [[arXiv:2310.12240](#)] [[INSPIRE](#)].
- [52] B.A. Kniehl and V.N. Velizhanin, *Anomalous dimensions of twist-two operators in extended $N = 2$ and $N = 4$ super Yang-Mills theories*, *Nucl. Phys. B* **1001** (2024) 116511 [[arXiv:2312.05888](#)] [[INSPIRE](#)].
- [53] G. Somogyi and S. Van Thurenhout, *All-order Feynman rules for leading-twist gauge-invariant operators in QCD*, *Eur. Phys. J. C* **84** (2024) 740 [[arXiv:2403.12623](#)] [[INSPIRE](#)].
- [54] R.V. Harlander, S.Y. Klein and M. Lipp, *FeynGame*, *Comput. Phys. Commun.* **256** (2020) 107465 [[arXiv:2003.00896](#)] [[INSPIRE](#)].
- [55] R. Harlander, S.Y. Klein and M.C. Schaaf, *FeynGame-2.1 — Feynman diagrams made easy*, *PoS EPS-HEP2023* (2024) 657 [[arXiv:2401.12778](#)] [[INSPIRE](#)].