

# Towards a Global Search for New Physics with Isotope Shifts

Elina Fuchs\*

*Deutsches Elektronen-Synchrotron DESY, Notkestr. 85, 22607 Hamburg, Germany  
Institut für Theoretische Physik, Leibniz Universität Hannover, Appelstraße 2, 30167 Hannover, Germany and  
Physikalisch-Technische Bundesanstalt, Bundesallee 100, 38116 Braunschweig, Germany*

Fiona Kirk†

*Physikalisch-Technische Bundesanstalt, Bundesallee 100, 38116 Braunschweig, Germany and  
Institut für Theoretische Physik, Leibniz Universität Hannover, Appelstraße 2, 30167 Hannover, Germany*

Agnese Mariotti‡

*Institut für Theoretische Physik, Leibniz Universität Hannover, Appelstraße 2, 30167 Hannover, Germany*

Jan Richter§

*Physikalisch-Technische Bundesanstalt, Bundesallee 100, 38116 Braunschweig, Germany*

Matteo Robbiati¶

*TIF Lab, Dipartimento di Fisica, Università degli Studi di Milano, Italy and  
European Organization for Nuclear Research (CERN), Geneva 1211, Switzerland*

Isotope shifts have emerged as a sensitive probe of new bosons that couple to electrons and neutrons, and of nuclear structure. The recent Hz- or even sub-Hz-level isotope shift measurements across different elements call for a global assessment of all available data. In this work, we present the fit framework `kifit` that for the first time enables a combined analysis of isotope shift data from several elements, taking into account correlations. We provide a thorough comparison of analytical methods and the fit to analyse linear and nonlinear King plots and quantify their uncertainties. Finally, we provide recommendations for future measurements that could enhance the sensitivity to new physics and offer new insights into nuclear structure.

## CONTENTS

I. Introduction	2	V. Conclusions and Outlook	19
II. Searching for New Physics with King Plots	3	Acknowledgments	20
A. Algebraic Methods for Linear King Plots	3	References	20
B. Algebraic Methods for Nonlinear King Plots	6	A. Calculation of Electronic Coefficients in $\text{Ca}^+$	24
III. The King Plot Fit	9	B. Sensitivity to New Physics	25
A. Geometric Construction (Build Phase)	9	C. Projection of Uncertainties	27
B. Determining the Search Window (Search Phase)	12	D. The <code>kifit</code> Package	28
C. Estimating the Confidence Interval (Experiment Phase)	12	1. Structure of the <code>kifit</code> Package	28
D. Producing Results (Consolidation Phase)	12	2. How to Use <code>kifit</code>	28
E. Comparison to a Previous King Plot Fit	15	3. Validation of the Algorithm	29
IV. Comparison of Fit and Algebraic Methods	16	E. The Impact of Data Sparsity on the Fit	32
A. Blind Directions	17	F. Isotope Shift and Atomic/Nuclear Mass Data	34
B. Geometric Construction & Form of Data Sets	17		

\* elina.fuchs@itp.uni-hannover.de

† fiona.kirk@itp.uni-hannover.de

‡ agnese.mariotti@itp.uni-hannover.de

§ jan.richter@ptb.de

¶ matteo.robbiati@cern.ch

## I. INTRODUCTION

Atomic precision spectroscopy has emerged as a powerful tool for probing the Standard Model of particle physics (SM) and its extensions, which address long-standing questions such as the properties of dark matter [1–3]. The groundbreaking developments of laser cooling [4–7], optical frequency combs [8, 9], optical tweezers [10–12], optical lattices [13–16] and quantum logic spectroscopy [17–20] have increased the relative precision of optical atomic clocks to the level of  $10^{-18}$  [21, 22]. As a result of this unparalleled precision, clock comparisons and other differential spectroscopic measurements probe relevant parts of the parameter space of light dark matter and dark portal models.

A prime example for such developments is isotope-shift spectroscopy, a well-established technique commonly used to determine nuclear charge radii [23–27], but more recently proposed as a method to search for hypothetical new bosons mediating an additional interaction between neutrons and electrons [28, 29]. Isotope shifts can probe bosons in the eV to MeV mass range, allowing them to bridge the gap between fifth force searches via the Casimir effect [30], beam dump experiments [31–33] and searches for exotic meson decays at colliders [34, 35]. New vector or scalar bosons, such as the ones probed by isotope shift spectroscopy [36], are predicted by a wide range of extensions of the SM. These include models that gauge baryon minus lepton number ( $B - L$ ), which result in a new vector boson  $Z'$ , or so-called portal models that introduce new light scalar or vector mediators between the SM and dark matter [37].

The new physics search proposed in Refs. [28, 29] is based on the so-called King plot method [23, 24], which at leading order predicts a linear relation between isotope shifts in different electronic transitions. Using King plots reduces the reliance on atomic structure calculations, which are limited in precision by non-perturbative and many-body effects, particularly within the nucleus. This approach allows new physics signals to be constrained using isotope shift data.

Given the long history of remarkably linear King plots, King plot searches for new physics attracted a lot of attention and significant progress was made in improving the precision of the isotope shift and nuclear mass measurements, with notable advancements in neutral Ca [38, 39],  $\text{Ca}^+$  [26, 40–43] and  $\text{Ca}^{14+}$  [43], as well as in neutral Yb [44–47] and  $\text{Yb}^+$  [48–50]. The first observation of a nonlinear King plot was reported in Ref. [48], which employed the  $^2S_{1/2} \rightarrow ^2D_{3/2}$  vs.  $^2S_{1/2} \rightarrow ^2D_{5/2}$  transitions in  $\text{Yb}^+$ , measured at a precision of about 300 Hz. This so-called “King nonlinearity” was confirmed by subsequent measurements [49, 50] and is currently found to be at the level of 20.17(2) kHz [50]. Recently, nonlinear isotope shifts have also been observed in the  $^3P_0 \rightarrow ^3P_1$  transition, measured at sub-Hz precision in  $\text{Ca}^{14+}$ , and in the  $^2S_{1/2} \rightarrow ^2D_{5/2}$  transition, measured in  $\text{Ca}^+$ , combined with nuclear mass ratios with relative uncertainties

below  $4 \times 10^{-11}$  [43].

Although the presence of nonlinearities complicates King plot searches for new physics, it is by no means a show-stopper: Progress in atomic and nuclear structure theory has facilitated the identification of the leading higher-order effects in Yb [49–51] and in Ca King plots [43, 52], whereas the development of the Generalised King Plot [53] showed that the data-driven King plot approach can be generalised so as to provide constraints on new physics, even in the presence of higher-order nuclear effects. For a recent review of the relevance of (nonlinear) King plots for the search for new (nuclear) physics, see Ref. [54].

The King plot analysis is often restricted to even isotopes to avoid the effects of the nuclear spin, namely hyperfine interactions, which are expected to introduce additional nonlinear effects in the King plot [55]. Ref. [56] discusses the challenges associated with using odd Yb isotopes in King plot analyses. Another example is provided by Refs. [57, 58]: although the King plots for Cd and Zn are linear, the hyperfine interactions would potentially be observable at Hz precision.

With a wealth of high-precision isotope shift data now available for different elements and charged states, a systematic approach is needed to assess the constraints on new physics contributions, which are expected to be governed by the same couplings to electrons and to neutrons, irrespective of the element under consideration. In this work we first review the (generalised) King plot [29, 53], the no-mass (generalised) King plot [53] and the projection [41] methods, which follow a purely algebraic approach (Section II), before presenting our code `kifit`, which provides for the first time a framework to combine all linear King plots across elements in one global constraint on new physics (Section III). The framework provided by `kifit` is based on the fit presented in Ref. [36], but was significantly extended to handle contemporary high-precision isotope shift data. `kifit` performs a fit to linear King plots plus new physics, while the incorporation of higher-order nuclear effects, that would be necessary to analyse nonlinear King plots, is left to future work. Nonetheless, `kifit` presents an important step towards a global view of isotope shift data. In Section IV we compare `kifit` with the algebraic methods introduced in Section II, before concluding in Section V. In the appendices we provide further details on the electronic structure calculations with `AMBiT` [59] (Appendix A), how the choice of transitions used in King plots affects the sensitivity to new physics (Appendix B), the impact of experimental uncertainties on the margin for new physics (Appendix C), a short manual for our code `kifit` and a list of implemented validation checks (Appendix D), an analysis of the impact of data sparsity on the fit (Appendix E), as well as a summary of the available state-of-the-art isotope shift data and isotope masses (Appendix F).

## II. SEARCHING FOR NEW PHYSICS WITH KING PLOTS

An isotope shift  $\nu_i^{AA'} \equiv \nu_i^A - \nu_i^{A'}$  corresponds to the frequency difference between the electronic transition  $i$  measured in isotopes  $A$  and  $A'$ . Isotope shifts are dominated by two effects, both of which can be described by a product of an electronic coefficient ( $K_i$ ,  $F_i$ ) and a nuclear quantity ( $\mu^{AA'}$ ,  $\delta\langle r^2 \rangle^{AA'}$ ), respectively [60–63],

$$\nu_i^{AA'} \approx K_i \mu^{AA'} + F_i \delta\langle r^2 \rangle^{AA'}. \quad (1)$$

The first term in Eq. (1), which is proportional to the difference of inverse nuclear masses<sup>1</sup>,  $\mu^{AA'} \equiv \frac{1}{m^A} - \frac{1}{m^{A'}}$ , is known as the first-order *mass shift* (MS) and describes the nuclear-recoil correction to the electron kinetic energy [64]. The second term, referred to as the first-order *field shift* (FS), accounts for the nuclear charge radius variance  $\delta\langle r^2 \rangle^{AA'} \equiv \langle r^2 \rangle^A - \langle r^2 \rangle^{A'}$  and describes the energy shift due to changes in the nuclear charge distribution between the different isotopes [65].

Since we will mostly be dealing with isotope pairs in the following, we introduce the isotope pair index  $a = AA'$ .

Thanks to the factorisation of the isotope shifts into electronic and nuclear quantities at leading order, we can combine isotope shift measurements for two different transitions to eliminate the nuclear charge radius, which is neither experimentally nor theoretically precisely determined [61, 66]. We obtain a linear relation between the isotope shifts of the two transitions labelled 1 and 2 [61],<sup>2</sup>

$$\tilde{\nu}_2^a|_{\text{lin.}} = K_{21} + F_{21}\tilde{\nu}_1^a \quad (3)$$

where we defined the so-called mass-normalised isotope shifts [63, 67, 68]  $\tilde{\nu}_i^a \equiv \frac{\nu_i^a}{\nu_1^a}$  and introduced the electronic coefficients

$$F_{21} \equiv \frac{F_2}{F_1}, \quad K_{21} \equiv K_2 - F_{21}K_1. \quad (4)$$

<sup>1</sup> The measured mass  $m^{A^0}$  of a neutral atom  $A^0$  can be translated into a nuclear mass  $m^A$  by subtracting the masses  $m_e$  and binding energies  $E_i^b$  of the  $N_e$  electrons [41]:

$$m^A = m^{A^0} - N_e m_e + \sum_{i=1}^{N_e} E_i^b.$$

<sup>2</sup> Instead of the mass-normalised isotope shifts in Eq. (3), it is possible to normalise the isotope shifts by those of the reference transition  $\nu_1^a$ . The King relation for *frequency-normalised* isotope shifts takes the form

$$\bar{\nu}_2^a \equiv \frac{\nu_2^a}{\nu_1^a} = F_{21} + K_{21}\bar{\mu}^a, \quad (2)$$

where  $\bar{\mu}^a \equiv \mu^a/\nu_1^a$ , while the electronic coefficients appear in the same combinations as in Eq. (4).




FIG. 1. Left: Linear 2-dimensional King plot. The isotope shifts follow the relation given in Eq. (1). Right: In the presence of higher-order nuclear or new physics contributions to the isotope shifts, the data points deviate from the King line and define a non-zero volume.




FIG. 2. Illustration of the “King plane” spanned by the vectors  $\nu_1$  and  $\mathbf{1}$  in isotope pair space. If Eq. (5) holds,  $\nu_2$  lies in this plane, while a nonlinearity in the King plot would lead to an out-of-plane component for  $\nu_2$ . Both new physics, here assumed to be proportional to  $\tilde{\gamma}$ , and higher-order SM terms, represented by the vector  $\tilde{\eta}$  (see Eq. (22)) can lead to this kind of effect.

The subscript “lin.” indicates that Eq. (3) only describes the leading linear behaviour.

If multiple isotope pairs ( $a = 1, \dots, n$ ) can be probed, it is useful to arrange the isotope shifts  $\tilde{\nu}_i^a$ ,  $i = 1, 2$ , into vectors in isotope-pair space. Defining the  $n$ -vectors  $\tilde{\nu}_i = (\tilde{\nu}_i^1, \dots, \tilde{\nu}_i^n)$  and  $\mathbf{1} = (1, \dots, 1)$ , one obtains

$$\tilde{\nu}_2|_{\text{lin.}} = K_{21}\mathbf{1} + F_{21}\tilde{\nu}_1. \quad (5)$$

Eq. (5) can be visualised in a so-called King plot [61], illustrated in the left half of Fig. 1. The electronic coefficients  $K_{21}$  and  $F_{21}$ , corresponding to the intercept and slope of the *King line*, can be determined via a linear fit to the isotope shift data. Alternatively, the isotope shift data can be arranged in isotope pair space. Fig. 2 shows the *plane of King linearity*, which is spanned by the vectors  $\mathbf{1}$  and  $\tilde{\nu}_1$ . If  $\tilde{\nu}_2$  is described by Eq. (5), it will lie in this plane.

### A. Algebraic Methods for Linear King Plots

In the following, we will distinguish linear and non-linear King plots. As King nonlinearities we understand

any deviation from the linear King relation in Eq. (5). As long as these are smaller or comparable to the experimental uncertainties, the King plot is considered linear.

Nonlinearities may be caused by higher-order nuclear or electronic corrections, or by new physics contributions. As long as the higher-order SM contributions are negligible or can be determined by additional experimental and theoretical input, King plots can be used to place constraints on the mass and couplings of a new boson  $\phi$  inducing a new interaction between the bound electrons and the neutrons inside the nucleus [29]. Assuming  $\phi$  couples linearly to neutrons and electrons, this effect can be described by a Yukawa potential [28, 69]

$$V_{\text{NP}}(r, m_\phi) = -\alpha_{\text{NP}}(A - Z) \frac{e^{-m_\phi r}}{r}, \quad (6)$$

expressed here in relativistic units ( $c = \hbar = 1$ ). Here,  $\alpha_{\text{NP}} = (-1)^s \frac{y_e y_n}{4\pi}$  is defined as the product of the new physics couplings to electrons ( $y_e$ ) and to neutrons ( $y_n$ ), while  $s = 0, 1, 2$  denotes the spin,  $m_\phi$  corresponds to the mass of the new light boson  $\phi$  and  $A$  and  $Z$  are respectively the mass and atomic numbers of the considered isotope. This introduces a new term in the isotope shifts [29]

$$\nu_i^a|_{\text{pred.}} = K_i \mu^a + F_i \delta \langle r^2 \rangle^a + \frac{\alpha_{\text{NP}}}{\alpha_{\text{EM}}} X_i \gamma^a \quad (7)$$

where we introduced the subscript “pred” to indicate the predicted isotope shift in the presence of new physics. Assuming a linear King plot, we can use Eq. (7) to set a bound on the new physics coupling  $\alpha_{\text{NP}}$ , expressed in units of the fine structure constant  $\alpha_{\text{EM}}$ <sup>3</sup>. The factor  $\gamma^a \equiv A - A'$  is the difference in neutron number between the isotopes  $A$  and  $A'$ , while the electronic coefficients  $X_i$  quantify the sensitivity of the transition frequency  $\nu_i$  to shifts induced by the new physics potential in Eq. (6). The leading behaviour can be understood in a perturbative approach, in which the radial wave functions  $\Psi_i^{\text{init,fin}}(r)$  of the states involved in the transition  $i$  are assumed to be unchanged compared to the SM. In this case, the coefficient  $X_i$  corresponds to the difference between the overlap of the new physics potential (Eq. (6)) with the final- and initial-state wave functions:

$$X_i(m_\phi) = \int \frac{e^{-m_\phi r}}{r} [|\Psi_i^{\text{fin}}(r)|^2 - |\Psi_i^{\text{init}}(r)|^2] dr. \quad (8)$$

The overlap with the new physics potential is particularly pronounced for S-state wave functions. Consequently narrow transitions involving S-states are favourable for isotope shift-based new physics searches.

As discussed in Appendix A, for the numerical calculation of the new physics electronic coefficients, we follow Ref. [29] and use the public code **AMBiT** [59], which applies a non-perturbative finite field method that adds the new physics potential directly to the Hamiltonian in the Dirac equation.

Combining Eq. (7) for the transitions  $i = 1, 2$ , we obtain the modified King relation

$$\tilde{\nu}_2|_{\text{pred.}} = K_{21} \mathbf{1} + F_{21} \tilde{\nu}_1 + \frac{\alpha_{\text{NP}}}{\alpha_{\text{EM}}} X_{21} \tilde{\gamma}, \quad (9)$$

where we defined  $X_{21} \equiv X_2 - F_{21} X_1$  and  $\tilde{\gamma}$  with elements  $\tilde{\gamma}^a \equiv \frac{\gamma^a}{\mu^a}$ . Eq. (9) implies that  $\tilde{\nu}_2|_{\text{pred.}}$  has a component along  $\tilde{\gamma}$ , which pushes it out of the King plane, as visualised in Fig. 2.

If  $n = 3$  isotope pairs are probed, the system of isotope shift equations Eq. (9) can be solved for the new physics coupling [29]:

$$\frac{\alpha_{\text{NP}}}{\alpha_{\text{EM}}} = \frac{\det(\tilde{\nu}_1, \tilde{\nu}_2, \mathbf{1})}{\varepsilon_{ij} \det(X_i \tilde{\gamma}, \tilde{\nu}_j, \mathbf{1})} = \frac{V_{\text{dat}}}{V_{\text{pred}}}, \quad (10)$$

where  $\varepsilon_{ij}$  is the 2-dimensional Levi-Civita symbol and summation over repeated indices is implied.  $\det(\mathbf{u}, \mathbf{v}, \mathbf{w})$  denotes the determinant of the square matrix whose columns are the vectors  $\mathbf{u}$ ,  $\mathbf{v}$  and  $\mathbf{w}$ . In the following, we will refer to Eq. (10) as the *Minimal King Plot (KP)* formula<sup>4</sup>. Note that the normalisation of the vectors entering Eq. (10) does not affect the result, meaning that  $(\tilde{\nu}_1, \tilde{\nu}_2, \mathbf{1})$  can be replaced by  $(\nu_1, \nu_2, \mu)$  or by frequency-normalised counterparts. Geometrically,  $V_{\text{dat}}$  can be visualised either as the volume spanned by data points in the King plot (see Fig. 1) or as the volume of the parallelepiped spanned by  $\tilde{\nu}_1$ ,  $\tilde{\nu}_2$  and  $\mathbf{1}$  (see Fig. 2). Similarly,  $V_{\text{pred}}$  measures the predicted volume spanned by the data vectors in the presence of new physics contributions.

It is instructive to split  $V_{\text{dat}}$  and  $V_{\text{pred}}$  into their respective electronic and nuclear contributions. Assuming the isotope shifts satisfy Eq. (7) and defining

$$\mathcal{M} = \begin{pmatrix} X_1 & F_1 \\ X_2 & F_2 \end{pmatrix}, \quad \mathcal{N} = \begin{pmatrix} \tilde{\gamma}^1 & \delta \langle r^2 \rangle^1 & 1 \\ \tilde{\gamma}^2 & \delta \langle r^2 \rangle^2 & 1 \\ \tilde{\gamma}^3 & \delta \langle r^2 \rangle^3 & 1 \end{pmatrix}, \quad (11)$$

where  $\delta \langle r^2 \rangle^a \equiv \delta \langle r^2 \rangle^a / \mu^a$  denotes the mass-normalised charge radius variance, we can rewrite  $V_{\text{pred}}$  as

$$V_{\text{pred}} = \varepsilon_{ij} \det(X_i \tilde{\gamma}, \tilde{\nu}_j, \mathbf{1}) = \det(\mathcal{M}) \det(\mathcal{N}) \\ = (F_1 X_2 - F_2 X_1) \det(\tilde{\gamma}, \delta \langle r^2 \rangle, \mathbf{1}). \quad (12)$$

<sup>3</sup> Note that alternative conventions exist in the literature. An equivalent relation in terms of  $y_e y_n$  rather than  $\alpha_{\text{NP}}/\alpha_{\text{EM}}$  can be obtained by rescaling the  $X$  coefficients (see Appendix A).

<sup>4</sup> In the **kfit** code, Eq. (10) corresponds to the 3-dimensional case of the generalised King plot formula, presented in Sec. II B. Here 3 refers to the number of isotope pairs, i.e. to the number of data points in the King plot.

This equation explicitly shows that King plots are sensitive to the new physics coupling when the nuclear and electronic quantities simultaneously “open up” new dimensions in isotope-pair space and in transition space.

The solution of  $\alpha_{\text{NP}}$  in Eq. (10) corresponds to the value of the coupling required to reproduce the central values of the King plot points  $\tilde{\nu}_i^a$ ,  $i = 1, 2$ ,  $a = 1, 2, 3$ . However, the King plot method cannot exclude higher-order SM contributions to the isotope shifts and thus cannot identify an observed nonlinearity as being of pure new physics origin. In other words, it is not a discovery tool, but can only be used to set bounds on the coupling  $\alpha_{\text{NP}}$ . The  $N\sigma$  bounds on  $\alpha_{\text{NP}}$  can be estimated as

$$\langle \alpha_{\text{NP}} \rangle + N\sigma[\alpha_{\text{NP}}]. \quad (13)$$

With  $\langle \alpha_{\text{NP}} \rangle$  we denote the absolute value of  $\alpha_{\text{NP}}$  obtained for the experimental central values of the input parameters (isotope shifts  $\nu_i^a$  and nuclear masses  $m^A$ ). The uncertainty on  $\alpha_{\text{NP}}$ ,  $\sigma[\alpha_{\text{NP}}]$ , can either be estimated using (linear) error propagation or with a Monte Carlo approach. The latter involves the generation of samples<sup>5</sup> of the input parameters from a normal distributions fixed by the respective experimental central values and uncertainties:

$$\nu_i^a \sim \mathcal{N}(\langle \nu_i^a \rangle, \sigma[\nu_i^a]), \quad m^A \sim \mathcal{N}(\langle m^A \rangle, \sigma[m^A]) \quad (14)$$

and similarly for the reference isotope masses  $m^{A'}$ . Then,  $\sigma[\alpha_{\text{NP}}]$  is estimated as the standard deviation of the corresponding  $\alpha_{\text{NP}}$  values. In this case one should pay attention whether the distribution of the  $\alpha_{\text{NP}}$  values can be accurately described by a Gaussian.

As already discussed in Ref. [29], the  $m_\phi$ -range that King plots are particularly sensitive to is dictated by the  $m_\phi$ -dependence of the  $X$  coefficients (see Eqs. (6) and (8) or Fig. 15): In the “massless” limit, the Yukawa potential (Eq. (6)) takes the form  $V_{\text{NP}} \propto 1/r$  and the electronic coefficients  $X$ , and consequently the bounds on  $\alpha_{\text{NP}}$ , become independent of  $m_\phi$  (see Figs. 15 and 16, respectively). This corresponds to the case where the interaction range of the mediator  $\phi$  (Compton wavelength  $\lambda_\phi = h/(m_\phi c)$ ) exceeds the size of the atom or ion, and the overlap of the Yukawa potential with the electronic wave functions saturates at maximal King plot sensitivity to new physics.

In the *intermediate  $m_\phi$  region*, the values of the  $X$  coefficients are sensitive to the value of  $m_\phi$  and the combination of electronic coefficients that enters  $V_{\text{pred}}$  (see Eq. (12)) may cancel for specific values of  $m_\phi$ , leading to the characteristic peaks that are visible e.g. in Fig. 16.

In the *large- $m_\phi$  limit*, the Yukawa potential can be approximated by  $V_{\text{NP}} \propto \delta(r)/(m_\phi r)^2$ . This limit corresponds to the case where the Compton wavelength of  $\phi$  is smaller than the nuclear charge radius and the interaction mediated by  $\phi$  becomes a contact interaction. In this case the King plot method is no longer able to distinguish the new physics shift from the field shift induced by changes in the nuclear radius. The  $X$  coefficients align themselves with the electronic field shift coefficients<sup>6</sup>  $F_i \propto |\Psi_i^{\text{fin}}(0)| - |\Psi_i^{\text{init}}(0)|$  in such a way that  $X_i/X_j \rightarrow F_i/F_j$ , and consequently the new physics parameters  $\frac{\alpha_{\text{NP}}}{\alpha_{\text{EM}}} \gamma^a$  can no longer be distinguished from the charge radius variance  $\delta\langle r^2 \rangle^a$ . The resulting suppression of the sensitivity to new physics can be observed e.g. in Fig. 16.

In Appendix B, a King plot analysis is applied to selected transitions in  $\text{Ca}^+$  to illustrate the behaviour of the electronic part of  $V_{\text{pred}}$  across different transitions. Moreover, the impact of the uncertainties on the  $X$  coefficients is briefly discussed therein.

### Additional Algebraic Methods

For completeness, let us briefly introduce the *No-Mass King Plot* and the *Projection Method*, which provide alternatives to Eq. (10) that can be beneficial, respectively, in the case of large nuclear mass uncertainties and in the case of isotope shift data sets for two transitions but more than 3 isotope pairs.

*No-Mass King Plot* — The *No-Mass King Plot (NMKP)* uses isotope shift data not only to eliminate the charge radius variance  $\delta\langle r^2 \rangle$  from Eq. (7), but also the nuclear masses. This requires isotope shift measurements for an additional transition  $\nu_3$ , yielding [53]<sup>7</sup>:

$$\frac{\alpha_{\text{NP}}}{\alpha_{\text{EM}}} = \frac{2\det(\nu_1, \nu_2, \nu_3)}{\varepsilon_{ijk}\det(X_i\gamma, \nu_j, \nu_k)}. \quad (15)$$

The denominator takes a particularly symmetric form in this case:

$$V_{\text{pred}} = \frac{1}{2}\varepsilon_{ijk}\det(X_i\gamma, \nu_j, \nu_k) = \det(\mathcal{M})\det(\mathcal{N}) \quad (16)$$

with

$$\mathcal{M} = \begin{pmatrix} X_1 & K_1 & F_1 \\ X_2 & K_2 & F_2 \\ X_3 & K_3 & F_3 \end{pmatrix}, \quad \mathcal{N} = \begin{pmatrix} \gamma^1 & \mu^1 & \delta\langle r^2 \rangle^1 \\ \gamma^2 & \mu^2 & \delta\langle r^2 \rangle^2 \\ \gamma^3 & \mu^3 & \delta\langle r^2 \rangle^3 \end{pmatrix}. \quad (17)$$

<sup>6</sup> Note that uncertainties in the  $X$  coefficients, which are predicted by means of atomic structure calculations, generally spoil this behaviour. However, it can easily be restored by rescaling the  $X_i$  coefficients in such a way that  $X_i/X_j \rightarrow F_{ij}|_{\text{dat}}$ , where  $F_{ij}|_{\text{dat}}$  is the slope of best fit line to the King plot data.

<sup>7</sup> In the *kifit* code this formula is referred to as the “dimension 3 No-Mass Generalised King Plot”.

<sup>5</sup> In the *kifit* code, the number of samples `num_det_samples` is a hyperparameter that can be set by the user.



The No-Mass King Plot formula allows to mitigate uncertainties on the nuclear masses when their effect on the uncertainty estimation in Eq. (13) dominates over that of the isotope shift uncertainties.

*Projection Method* — The *Projection Method* presented in Ref. [41], yields a bound on new physics using isotope shift data for 2 transitions measured in  $n$  isotope pairs. Introducing the  $(n \times 2)$  matrix

$$D_{\mathbf{w}} \equiv (\tilde{\nu}_1, \mathbf{w}), \quad (18)$$

where  $\mathbf{w}$  is a generic  $n$ -vector in isotope-pair space, we can define the projection of the vector  $\mathbf{1}$  on the plane spanned by the vectors  $\tilde{\nu}_1$  and  $\mathbf{w}$  as

$$\mathbf{p}_{\mathbf{w}} = D_{\mathbf{w}} (D_{\mathbf{w}}^{\top} D_{\mathbf{w}})^{-1} D_{\mathbf{w}}^{\top} \mathbf{1}, \quad (19)$$

where  $\top$  denotes the transpose. Then,  $\alpha_{\text{NP}}$  can be expressed as

$$\left| \frac{\alpha_{\text{NP}}}{\alpha_{\text{EM}}} \right| = \frac{V(\mathbf{1}, \tilde{\nu}_1, \tilde{\nu}_2)}{|X_{21}|V(\mathbf{1}, \tilde{\nu}_1, \tilde{\gamma})}, \quad (20)$$

where

$$V(\mathbf{1}, \tilde{\nu}_1, \mathbf{w}) = \|\mathbf{1} - \mathbf{p}_{\mathbf{w}}\| \sqrt{\|\tilde{\nu}_1\|^2 \|\mathbf{w}\|^2 - (\tilde{\nu}_1 \cdot \mathbf{w})^2}, \quad (21)$$

is the volume of the parallelepiped spanned by the vectors  $\mathbf{1}$ ,  $\tilde{\nu}_1$  and  $\mathbf{w}$ . Notice that Eq. (20) reduces to an equation similar to the Minimal King Plot formula (Eq. (10)), for  $n = 3$  isotope pairs, however, it is insensitive to the sign of  $\alpha_{\text{NP}}$ . When sampling  $\alpha_{\text{NP}}$  (see discussion around Eq. (13)), this insensitivity to the sign can lead to a distortion of the distribution.

## B. Algebraic Methods for Nonlinear King Plots

In the previous section, we discussed how isotope shifts that produce linear King plots can be used to set bounds on the new physics coupling. However, the increasing sensitivity of isotope shift and nuclear mass measurements has resolved deviations from Eq. (1) (see e.g. samarium (Sm) [61, 70, 71] and ytterbium (Yb) [47–50, 72, 73] King plots, and the combination of singly and highly-charged calcium (Ca) [43]<sup>8</sup>), calling for a more detailed analysis of the SM background. Assuming factorisability of the electronic and nuclear contributions, we denote the next-to-leading SM contribution by  $H_i \eta^a$ , such that

$$\nu_i^a = K_i \mu^a + F_i \delta \langle r^2 \rangle^a + H_i \eta^a + \dots + \frac{\alpha_{\text{NP}}}{\alpha_{\text{EM}}} X_i \gamma^a, \quad (22)$$

<sup>8</sup> After a nonlinearity had been measured recently in cadmium (Cd) [74], linearity was restored in [57].

where the ellipsis is a placeholder for unresolved higher-order SM contributions. In the presence of King plot nonlinearities, methods beyond the ones discussed in the previous section are needed. In the following section we discuss how the main sources of nonlinearity can be identified and eliminated, either in a data-driven way, or using additional theoretical input, and why King plots remain a useful tool to search for new physics.

### The Nonlinearity Decomposition

If at least four isotope pairs are probed, there is a clear hierarchy between the King nonlinearities, and predictions for their nuclear structure are available, the *Nonlinearity Decomposition Plot* [48, 49] can provide useful insights into the origin of the dominant King nonlinearity. The main idea is to project the isotope shift data onto the  $n$  basis vectors  $(\mathbf{1}, \tilde{\nu}_1, \Lambda^1, \dots, \Lambda^{n-2})$ , where  $\mathbf{1}$  and  $\tilde{\nu}_1$  define the plane of “King linearity”, and the linearly independent vectors  $\{\Lambda^\ell\}_{\ell=1}^{(n-2)}$ , which can be chosen to be orthogonal<sup>9</sup> to  $\mathbf{1}$  and  $\tilde{\nu}_1$ , define a basis for the “King nonlinearity space”:

$$\tilde{\nu}_i = K_{i1} \mathbf{1} + F_{i1} \tilde{\nu}_1 + \sum_{\ell=1}^{n-2} \lambda_{i1}^\ell \Lambda^\ell. \quad (23)$$

In the minimal case of 4 isotope pairs, as in Refs. [48–50] and Ref. [43], which considered the 5 even stable isotopes of Yb and Ca, respectively, the basis vectors of the King nonlinearity space can be expressed as [49]

$$\begin{aligned} \Lambda^+ &\sim (\tilde{\nu}_1^3 - \tilde{\nu}_1^2, \tilde{\nu}_1^1 - \tilde{\nu}_1^4, \tilde{\nu}_1^4 - \tilde{\nu}_1^1, \tilde{\nu}_1^2 - \tilde{\nu}_1^3), \\ \Lambda^- &\sim (\tilde{\nu}_1^4 - \tilde{\nu}_1^2, \tilde{\nu}_1^1 - \tilde{\nu}_1^3, \tilde{\nu}_1^2 - \tilde{\nu}_1^4, \tilde{\nu}_1^3 - \tilde{\nu}_1^1), \end{aligned} \quad (24)$$

and Eq. (23) reads

$$\tilde{\nu}_i = K_{i1} \mathbf{1} + F_{i1} \tilde{\nu}_1 + \lambda_{i1}^+ \Lambda^+ + \lambda_{i1}^- \Lambda^-. \quad (25)$$

The projections  $\lambda_{i1}^+$  and  $\lambda_{i1}^-$  are used as coordinates in the Nonlinearity Decomposition Plot, which in the case of 4 isotope pairs is 2-dimensional, as illustrated in Fig. 3: Each point  $(\lambda_{i1}^+, \lambda_{i1}^-)$ ,  $i = 2, 3$  in the Nonlinearity Decomposition Plot represents the nonlinearities in the 2-dimensional King plot constructed from the transitions  $i$  and 1. If its uncertainty ellipse contains the origin of the  $(\lambda_{i1}^+, \lambda_{i1}^-)$  plane, the associated King plot can be considered to be linear, else additional terms need to be added to the isotope shift equations to describe the data.

The Nonlinearity Decomposition Plot is particularly useful when there is a strong hierarchy between the nonlinearities. In the case of one dominant King nonlinearity that can be factorised into electronic and nuclear

<sup>9</sup> The results in this section can, of course, be generalised to the case where the vectors  $\Lambda^\ell$  are not orthogonal to the basis vectors  $\mathbf{1}$  and  $\tilde{\nu}_1$ . In this case the projections of the higher-order terms such as  $H_i \eta^a$  onto  $\mathbf{1}$  and  $\tilde{\nu}_1$  need to be taken into account.

contributions (as e.g.  $H_i$  and  $\eta^a$  in Eq. (22)), the data points approximately lie on one line in the Nonlinearity Decomposition Plot. Indeed, in this case the slope  $\lambda_{i1}^-/\lambda_{i1}^+$  defined by the projections onto the space of King nonlinearity is transition independent

$$\frac{\lambda^-}{\lambda^+} \equiv \frac{\lambda_{i1}^-}{\lambda_{i1}^+} = \frac{(\tilde{\eta} \cdot \Lambda^-)(\Lambda^+ \cdot \Lambda^+) - (\tilde{\eta} \cdot \Lambda^+)(\Lambda^- \cdot \Lambda^+)}{(\tilde{\eta} \cdot \Lambda^+)(\Lambda^- \cdot \Lambda^-) - (\tilde{\eta} \cdot \Lambda^-)(\Lambda^- \cdot \Lambda^+)} \quad (26)$$

and a comparison with predictions for different nuclear parameters  $\tilde{\eta}$  can be used to identify the origin of the dominant nonlinearity in the data.

The purple line through the origin in Fig. 3 illustrates how the new physics hypothesis introduced in Eq. (7) can be tested with the help of the Nonlinearity Decomposition Plot: If all uncertainty ellipses overlap with this purple line, the observed nonlinearities are compatible with the new physics hypothesis (or any other effect that is proportional to the neutron number). A similar statement can be made for higher-order nuclear contributions. The slopes of the corresponding lines can either be determined via nuclear structure calculations or using complementary experimental input.

Note that more than one source of nonlinearity, as well as input for their electronic structure, is needed to describe the data that does not lie on a line through the origin of the  $(\lambda_{i1}^+, \lambda_{i1}^-)$  plane.

*Generalisation to  $n$  isotope pairs* If isotope shift data for  $n$  isotope pairs is available, more than one nonlinearity can be identified in this data-driven way. We define the vector of coordinates associated to the King plot for the transitions  $i$  and 1 as  $\lambda_{i1} = (\lambda_{i1}^1, \dots, \lambda_{i1}^{n-2})^\top$  and introduce the  $(n \times (n-2))$ -matrix  $\Omega = (\Lambda^1, \dots, \Lambda^{n-2})$  of vectors that span the nonlinearity space. Eq. (23) can then be generalised using  $\sum_{\ell=1}^{n-2} \lambda_{i1}^\ell \Lambda^\ell = \Omega \lambda_{i1}$ .

Using the Moore-Penrose pseudoinverse of  $\Omega$  to solve for the coordinates  $\lambda_{i1}^\ell$ , one obtains the generalisation of Eq. (27)

$$\frac{\lambda_{i1}^{\ell'}}{\lambda_{i1}^\ell} = \frac{[(\Omega^\top \Omega)^{-1} \Omega^\top \tilde{\eta}]^{(\ell')}}{[(\Omega^\top \Omega)^{-1} \Omega^\top \tilde{\eta}]^{(\ell)}}, \quad \ell' \neq \ell, \quad (27)$$

where  $[\cdot]^{(\ell)}$  is to be understood as the  $\ell^{\text{th}}$  entry of  $[\cdot]$ . Since  $\ell, \ell'$  run from 1 to  $n-2$ , and  $\ell' \neq \ell$ , this method can resolve  $n-3$  sources of King plot nonlinearity, i.e. one in the case of 4 isotope pairs.

Note that while the slope of the predicted nonlinearity in the King plane is transition-independent, the coordinates  $\lambda_{i1}^\ell$  are not and their magnitude

$$\|\lambda_{i1}\| = \sqrt{\sum_\ell (\lambda_{i1}^\ell)^2} = |H_{i1}| \sqrt{\tilde{\eta}^\top \Omega (\Omega^\top \Omega)^{-2} \Omega^\top \tilde{\eta}}, \quad (28)$$




FIG. 3. Schematic illustration of the Nonlinearity Decomposition Plot in the minimal case of 4 isotope pairs. The two orange points with uncertainty ellipses represent the nonlinearities observed in the King plots constructed from the isotope shift data for the transition pairs (1, 2) and (1, 3). The purple and teal lines show the predicted slopes of the new physics term and a higher-order SM contribution to the isotope shift, respectively.

or

$$\begin{aligned} \lambda_{i1}^+ &= H_i \tilde{\eta} \cdot \frac{\Lambda^+(\Lambda^- \cdot \Lambda^-) - \Lambda^-(\Lambda^- \cdot \Lambda^+)}{(\Lambda^+ \cdot \Lambda^+)(\Lambda^- \cdot \Lambda^-) - (\Lambda^- \cdot \Lambda^+)^2}, \\ \lambda_{i1}^- &= H_i \tilde{\eta} \cdot \frac{\Lambda^-(\Lambda^+ \cdot \Lambda^+) - \Lambda^+(\Lambda^- \cdot \Lambda^+)}{(\Lambda^+ \cdot \Lambda^+)(\Lambda^- \cdot \Lambda^-) - (\Lambda^- \cdot \Lambda^+)^2} \end{aligned} \quad (29)$$

in the case of 4 isotope pairs, can be used to extract the electronic coefficients  $H_{i1}$  from data.

#### Generalised King Plots

In the recent Yb [47–50, 72, 73] and Ca [43] King plots, the leading nonlinearity is not compatible with the new physics term and so-called *Generalised King Plots* [53] were employed to set bounds on  $\alpha_{\text{NP}}$ . These generalise Eq. (10) and (15) to higher dimensions, i.e. they use additional measurements to eliminate higher-order nuclear contributions to the isotope shift equations.

For concreteness, let us assume the isotope shift equations take the form of Eq. (22): As discussed in the case of the *Minimal King Plot*, isotope shift data for two transitions,  $i = 1, 2$ , allow us to eliminate the charge radius variance,  $\delta\langle r^2 \rangle$ , resulting in

$$\tilde{\nu}_2^a = K_{21} + F_{21} \tilde{\nu}_1^a + H_{21} \tilde{\eta}^a + \dots + \frac{\alpha_{\text{NP}}}{\alpha_{\text{EM}}} X_{21} \tilde{\gamma}^a, \quad (30)$$

where  $H_{21} \equiv H_2 - F_{21} H_1$ .

If isotope shifts are measured in a third transition  $i = 3$ , they can be used to eliminate the higher-order SM

	KP	NMKP	GKP	NMGKP	kifit
	Eqs. (10)	(15)	(32)	(33)	Sec. III
Isotope pairs	3	3	$3 \leq n$	$3 \leq n$	$3 \leq n$
Transitions	2	3	$n - 1$	$n$	$2 \leq m$
Spurions	-	-	$n - 3$	$n - 3$	-

TABLE I. Summary of the number of transitions and isotope pairs that are required by the different algebraic methods presented in Section II and by the King plot fit **kifit** introduced in Section III. (Note that  $m$  and  $n$  are independent integers.) For the GKP and the NMGKP we also give the number of higher-order nuclear parameters (spurions) that are eliminated while setting a bound on the coupling  $\alpha_{\text{NP}}$ . KP and NMKP are the minimal cases of GKP and NMGKP, respectively.

term  $\tilde{\eta}^a$ , leading to

$$\tilde{\nu}_3^a = K_{321} + F_{321}\tilde{\nu}_1^a + H_{321}\tilde{\nu}_2^a + \dots + \frac{\alpha_{\text{NP}}}{\alpha_{\text{EM}}}X_{321}\tilde{\gamma}^a, \quad (31)$$

with  $H_{321} \equiv H_{31}/H_{21}$  and  $P_{321} = P_{31} - H_{321}P_{21}$ , with  $P \in \{K, F, X\}$ . Assuming the higher-order SM terms represented by the ellipsis are negligible, the isotope shift data can be used to set bounds on the new physics term (see fit in Ref. [49]). Note, however, that any direct extraction of  $\alpha_{\text{NP}}$  from Eq. (31) will be affected by the large theoretical uncertainties associated with the prediction for the electronic coefficient  $X_{321}$ , corresponding to a difference differences of the  $X$  coefficients introduced in Eq. (7).

One of the main strengths of the *Generalised King Plot* formulas is that they directly use the  $X$  coefficients introduced in Eq. (7) rather than relying on differences thereof. Assuming isotope shifts for  $m$  transitions are measured in  $n = m + 1$  isotope pairs, solving the system of  $m \times n$  isotope shift equations for  $\alpha_{\text{NP}}$  leads to the *Generalised King Plot (GKP)* formula [53]<sup>10</sup>:

$$\frac{\alpha_{\text{NP}}}{\alpha_{\text{EM}}} = \frac{(n-2)! \det(\tilde{\nu}_1, \dots, \tilde{\nu}_{n-1}, \mathbf{1})}{\varepsilon_{i_1 \dots i_{n-1}} \det(X_{i_1} \tilde{\gamma}, \tilde{\nu}_{i_2}, \dots, \tilde{\nu}_{i_{n-1}}, \mathbf{1})}, \quad (32)$$

As briefly suggested in Ref. [53], if higher-order SM contributions are present and the uncertainties on the nuclear masses are a limiting factor, it can be advantageous to combine the benefits of the GKP and NMKP

(Eq. (15)) formulas, obtaining a *No-Mass Generalised King Plot (NMGKP)* formula:

$$\frac{\alpha_{\text{NP}}}{\alpha_{\text{EM}}} = \frac{(n-1)! \det(\nu_1, \nu_2, \dots, \nu_n)}{\varepsilon_{i_1, i_2, \dots, i_n} \det(X_{i_1} \gamma, \nu_{i_2}, \dots, \nu_{i_n})}. \quad (33)$$

### Subtracting SM Nonlinearities

For lighter systems such as Ca, where the mass shift dominates over the field shift, the next-to-leading order SM contribution to Eq. (3) might be the second-order mass shift [52],  $\nu_i^a|_{\text{MS}(2)} = K_i^{(2)} \mu^{a(2)}$ , where  $\mu^{a(2)} \equiv 1/(m^A)^2 - 1/(m^{A'})^2$  is the difference of the squared inverse nuclear masses and  $K_i^{(2)}$  is an electronic coefficient. Since in this case the nuclear parameters (namely, the isotope masses) can be determined experimentally at high precision, the second-order mass shift can be kept explicit in the isotope shift equations, such that

$$\tilde{\nu}_2^a = K_{21} + F_{21}\tilde{\nu}_1^a + \frac{\alpha_{\text{NP}}}{\alpha_{\text{EM}}}X_{21}\tilde{\gamma}^a + K_{21}^{(2)}\tilde{\mu}^{a(2)}. \quad (34)$$

If the frequency shifts  $\nu_i^a$ ,  $i = 1, 2$ , are measured in four isotope pairs, Eq. (34) can then be solved for  $\alpha_{\text{NP}}$ , yielding

$$\frac{\alpha_{\text{NP}}}{\alpha_{\text{EM}}} = \frac{\det(\tilde{\nu}_1, \tilde{\nu}_2, \mathbf{1}, \tilde{\mu}^{(2)})}{\varepsilon_{ij} \det(X_i \tilde{\gamma}, \tilde{\nu}_j, \mathbf{1}, \tilde{\mu}^{(2)})}, \quad (35)$$

where all bold symbols are 4-vectors in isotope-pair space. In this case one higher-order effect can be taken into account without adding an additional transition. We will refer to this equation as the *Nuclear Input King Plot (NIKP)* formula.

Alternatively, if the the electronic coefficients  $K_i^{(2)}$  can be predicted with sufficient accuracy (see e.g. Ref. [43, 52], where a 10% uncertainty is assigned to the prediction of  $K_i^{(2)}$ ), the second-order mass shift can be subtracted from the measured isotope shift [43], yielding an isotope shift equation which resembles Eq. (3):

$$(\tilde{\nu}_2^a - K_{21}^{(2)}\tilde{\mu}^{a(2)}) = K_{21} + F_{21}\tilde{\nu}_1^a + \alpha_{\text{NP}}X_{21}\tilde{\gamma}^a. \quad (36)$$

Now  $\alpha_{\text{NP}}$  can be obtained using Eq. (10) with the modification  $\tilde{\nu}_2^a \rightarrow (\tilde{\nu}_2^a)' = \tilde{\nu}_2^a - K_{21}^{(2)}\tilde{\mu}^{a(2)}$ . Since  $K_i^{(2)}$  depends only on the electronic structure, it induces correlations between the objects  $\{(\tilde{\nu}_2^a)'\}_{a=1}^n$ , which in turn reduce the impact of the uncertainty on the coefficients  $K_i^{(2)}$  on  $\sigma[\alpha_{\text{NP}}]$  [43].

The strategy of subtracting higher-order SM contributions until relations such as Eq. (36) arise, could be promising to bridge the gap between light elements such as hydrogen, deuterium or helium [75–78], where spectroscopic bounds can be derived from a direct comparison of predictions and measurements, and heavy elements such as Yb, where corrections to the charge distribution, such as nuclear deformation, are sizeable but hard to predict from first principles [50].

<sup>10</sup> In the **kifit** code this formula is referred to as the “dimension  $n$  Generalised King Plot”, where  $n$  is the number of isotope pairs. Using frequency-normalised quantities, i.e.  $\tilde{\tau}^a \equiv \tau^a/\nu_1^a$ ,  $\tau \in \{\nu_i, \mu, \gamma\}$ ,  $i = 1, \dots, m$ , Eq. (32) takes the form

$$\frac{\alpha_{\text{NP}}}{\alpha_{\text{EM}}} = \frac{(n-2)! \det(\tilde{\nu}_1, \dots, \tilde{\nu}_{n-1}, \tilde{\mu})}{\varepsilon_{i_1 \dots i_{n-1}} \det(X_{i_1} \tilde{\gamma}, \tilde{\nu}_{i_2}, \dots, \tilde{\nu}_{i_{n-1}}, \tilde{\mu})}.$$



## Combining Data Sets

In Section II, we discussed the algebraic methods summarised in Table I. The advantage of these methods is their simplicity. However, they can only be applied to data sets of the specific dimensions given in Table I. If the available data sets come in different shapes, we are forced to derive bounds using subsets of the data, and how these resulting bounds are best combined is unclear.

In the following section we will follow an alternative approach consisting of a fit to isotope shift data. This strategy is more flexible regarding the dimensions of the input data and in particular enables a fit to isotope shift data from different elements.

## III. THE KING PLOT FIT

The King plot fit presented in this section is inspired by Ref. [36] but significantly extended such that it can handle the contemporary precision of isotope shift data and potentially large hierarchies between the levels of uncertainty of different subsets of the data. In the following, we describe the theoretical framework and the main steps of the algorithm, which is structured into a *build phase*, a *search phase*, an *experiment phase* and a *consolidation phase*, as illustrated in Fig. 6. An implementation of this procedure is provided in the form of the Python package `kifit`, which is publicly available at <https://github.com/QTI-TH/kifit>. More instructions on how to use the package can be found in Appendix III. The data provided by the current version of `kifit` is summarised in Appendix F.

## A. Geometric Construction (Build Phase)

One of the advantages of the King plot fit is that it is more flexible than the algebraic methods described in Section II A (see Table I for an overview) in regard to the number of transitions and isotope pairs that it can combine. As in Section II B,  $m$  denotes the number of transitions and  $n$  the number of isotope pairs considered for a given element.

Depending on the application, it can be advantageous to arrange the same isotope shift data in terms of isotope pairs or transitions: For the construction of the determinants discussed in Section II A, it is beneficial to describe the data in terms of the vectors  $\mathbf{1}$ ,  $\tilde{\nu}_i$  and  $\tilde{\gamma}$ , which are fixed by the transition frequency and mass measurements, as well as the neutron number difference. For the fit, which is performed directly on the level of the King plot and constrains deviations from the King line defined by the electronic coefficients  $F_{i1}$ ,  $K_{i1}$ ,  $i = 2, \dots, m$ , it is more natural to work in transition space.

Fixing  $i = 1$  as the reference transition and assuming the isotope shift data to be linear, we can construct  $(m - 1)$  relations of the form of Eq. (3), and arrange them into




FIG. 4. Illustration of the `kifit` construction in the simplest case of  $m = 2$  transitions and  $n = 3$  isotope pairs. We use the notation  $\vec{O}^a = (\tilde{\nu}_1^a, \tilde{\nu}_2^a|_{\text{lin.}})$ ,  $a = 1, 2, \dots, n$  for the linear prediction (Eq. (37)),  $\vec{Q}^a = (\tilde{\nu}_1^a, \tilde{\nu}_2^a|_{\text{pred.}})$  for the nonlinear prediction including new physics (Eq. (42)) and  $\vec{P}^a = (\tilde{\nu}_1^a, \tilde{\nu}_2^a)$  for the data points. The King line obtained from the initial fit to the isotope shift data is denoted  $\ell^{(0)}$  and described by the fit parameters  $K_{21}^1$  and  $\phi_{21}$ , whereas the parallel line associated to the isotope pair  $a$  is denoted  $\ell^{(a)}$ .

$n$  linear  $m$ -vector equations, one for every isotope pair  $a$  [36]:

$$\tilde{\nu}^a|_{\text{lin.}} \equiv \begin{pmatrix} \tilde{\nu}_1^a \\ \tilde{\nu}_2^a|_{\text{lin.}} \\ \vdots \\ \tilde{\nu}_m^a|_{\text{lin.}} \end{pmatrix} = \begin{pmatrix} 0 \\ K_{21} \\ \vdots \\ K_{m1} \end{pmatrix} + \tilde{\nu}_1^a \begin{pmatrix} 1 \\ F_{21} \\ \vdots \\ F_{m1} \end{pmatrix} \equiv \mathcal{K} + \tilde{\nu}_1^a \mathcal{F}. \quad (37)$$

Here we introduced the  $m$ -dimensional electronic coefficient vectors  $\mathcal{K}$  and  $\mathcal{F}$ . Note that the isotope shift vectors introduced in Eq. (5) and used in Sections II A and II B were vectors in isotope pair space, meaning that we had one for each transition, whereas the vectors that appear in Eq. (37) are vectors in transition space, meaning that we have one for each isotope pair.

Fig. 4 is a 2-dimensional illustration of Eq. (37) for the case of  $m = 2$  transitions and  $n = 3$  isotope pairs: By construction, the points  $\vec{O}^a = (\tilde{\nu}_1^a, \tilde{\nu}_2^a|_{\text{lin.}})$ ,  $a = 1, 2, 3$  lie on a *King line*  $\ell^{(0)}$  with intercept  $K_{21}$  and slope  $F_{21}$ . In the `kifit` code, the vector  $\mathcal{F}$  of electronic field shift coefficients is expressed in terms of the inclination angles  $\phi_{j1}$  of the King lines in the  $(\tilde{\nu}_1, \tilde{\nu}_j)$  plane. This is reasonable, since only its orientation matters. The unit vector  $\hat{e}_{\mathcal{F}}$ , directed along the King line  $\ell^{(0)}$  (orange vector in Fig. 4




FIG. 5. Illustration of a 3 dimensional King plot with a King line  $\ell^{(0)}$  (black), fixed by the fit parameters  $\phi_{j1}$  (see Eq. (38)) and  $K_{j1}^\perp$ ,  $j = 2, 3$  (see Eq. (40)) and oriented in the direction of the unit vector  $\hat{e}_{\mathcal{F}}$ . The lines  $\ell_{j1}$ ,  $j = 2, 3$ , represent the King lines of the 2-dimensional King plots constructed for the transition pairs (1, 2) and (1, 3), respectively.

and Fig. 5), takes the form:

$$\hat{e}_{\mathcal{F}} \equiv \frac{\mathcal{F}}{\|\mathcal{F}\|} = \frac{1}{\sqrt{1 + \sum_{j=2}^m \tan^2 \phi_{j1}}} \begin{pmatrix} 1 \\ \tan \phi_{21} \\ \vdots \\ \tan \phi_{m1} \end{pmatrix}. \quad (38)$$

We can define an angle  $\phi$  which satisfies

$$\cos \phi = \frac{1}{\sqrt{1 + \sum_{j=2}^m \tan^2 \phi_{j1}}} \quad (39)$$

and corresponds to the angle between the direction  $\hat{e}_1 = (1, 0, \dots, 0)$  fixed by the reference transition 1 and the King line. In the case of 2 transitions,  $\phi = \phi_{21}$  (see also Fig. 4).

Instead of taking the  $m - 1$  intercepts  $\{K_{j1}\}_{j=2}^m$  to be fit parameters, we introduce

$$\mathcal{K}^\perp \equiv \begin{pmatrix} K_{11}^\perp \\ K_{21}^\perp \\ \vdots \\ K_{m1}^\perp \end{pmatrix} = \begin{pmatrix} -\sum_{j=2}^m \sin \phi_{j1} K_{j1} \\ \cos \phi_{21} K_{21} \\ \vdots \\ \cos \phi_{m1} K_{m1} \end{pmatrix}, \quad (40)$$

which is orthogonal to  $\hat{e}_{\mathcal{F}}$ , and fit to the the  $m - 1$  projected intercepts  $\{K_{j1}^\perp\}_{j=2}^m$ . This parametrisation is crucial for the numerical stability of the fit in the limit where  $\hat{e}_{\mathcal{F}}$  and  $\mathcal{K}$  approach collinearity and  $K_{j1}^\perp \ll K_{j1}$ .

The initial guesses  $\{\langle K_{j1}^\perp \rangle, \langle \phi_{j1} \rangle\}_{j=2}^m$  for the  $2(m - 1)$  fit parameters and their covariance matrix  $\Sigma_{\mathcal{K}\phi}^{(j1)}$  are determined by a set of  $(m - 1)$  linear fits. Since King plots are constructed from pairs of isotope shifts, all of which have uncertainties, we perform orthogonal distance regression (ODR) [79] that is implemented in the `scipy.odr` method of the Scipy package [80].

Once the background (i.e. the linear King plot) is fixed, we can proceed with the parametrisation of the signal: the new physics term introduced in Eq. (9) leads to

isotope-pair dependent shifts of the data points in transition space (i.e. in the King plot):

$$\delta^a = \frac{\alpha_{\text{NP}}}{\alpha_{\text{EM}}} \tilde{\gamma}^a \mathcal{X}, \quad \mathcal{X} = (0, X_{21}, \dots, X_{m1})^\top. \quad (41)$$

Here,  $X_{j1} \equiv X_j - F_{j1} X_1 = X_j - \tan \phi_{j1} X_1$ ,  $j = 2, \dots, m$ , are the electronic coefficients that are constructed from the fit parameter  $\phi_{j1}$  and the coefficients  $X_i$ ,  $i = 1, \dots, m$  determined by means of atomic structure calculations (see Section II A and Appendix A). The predictions  $\tilde{\nu}^a|_{\text{pred.}}$  incorporating the linear isotope shift behaviour plus the shifts predicted by the new physics term are simply given by

$$\tilde{\nu}^a|_{\text{pred.}} = \tilde{\nu}^a|_{\text{lin.}} + \delta^a, \quad a = 1, \dots, n. \quad (42)$$

For each isotope pair  $a$ , Eq. (42) defines a parallel line  $\ell^{(a)}$  to the King line  $\ell^{(0)}$ . In Fig. 4,  $\vec{Q}^a = (\tilde{\nu}_1^a, \tilde{\nu}_2^a|_{\text{pred.}})$ ,  $a = 1, 2, 3$ , and the vectors  $\delta^a$  with their associated lines  $\ell^{(a)}$  are depicted in purple.

Since it is impossible to distinguish a constant shift of all points  $\tilde{\nu}_i^a$ ,  $1 < i \leq m$ , from a variation of the mass shift coefficients  $K_{i1}$ , we only include the variations  $\sigma[\delta^a]$  with respect to the mean new physics-induced shift in our new physics predictions. The elements of  $\sigma[\delta^a]$  can be estimated as

$$\sigma[\delta_j^a] = \frac{\alpha_{\text{NP}}}{\alpha_{\text{EM}}} (\tilde{\gamma}^a - \langle \tilde{\gamma} \rangle^j) X_{j1}, \quad (43)$$

where

$$\langle \tilde{\gamma} \rangle^j = \frac{\sum_{a=1}^n \tilde{\gamma}^a \sigma[\tilde{\nu}_j^a]}{\sum_{b=1}^n \sigma[\tilde{\nu}_j^b]}, \quad j = 2, \dots, m \quad (44)$$

is the average value of the new physics nuclear parameter  $\tilde{\gamma}^a$ , weighted by the uncertainties on the mass-normalised isotope shifts. (Note that by construction  $\sigma[\delta_1^a] = 0 \forall a$ , since  $X_{11} = 0$ , see Eq. (41)). In this way the constant shift induced by the quantities  $\langle \tilde{\gamma} \rangle^i$  is effectively absorbed in the fit parameters  $\{K_{j1}^\perp\}_{j=2}^m$ , leading to a corrected mass shift vector

$$\mathcal{K}' = \mathcal{K} + \langle \tilde{\gamma} \rangle, \quad (45)$$

with  $\langle \tilde{\gamma} \rangle = (\langle \tilde{\gamma} \rangle^1, \dots, \langle \tilde{\gamma} \rangle^n)^\top$ . Using these definitions, we obtain for the isotope shift

$$\tilde{\nu}^a|_{\text{pred.}} = \mathcal{K}' + \tilde{\nu}_1^a \mathcal{F} + \sigma[\delta^a], \quad a = 1, \dots, n. \quad (46)$$

For simplicity, we will omit the prime in  $\mathcal{K}'$  in the following.

Note that in general Eq. (44) induces a slight asymmetry under exchange of transitions. In order to cross-check the parametrisation, we implement in the `kifit` code a transition-independent version of  $\sigma[\delta^a]$ , with

$$\langle \tilde{\gamma} \rangle \equiv \frac{1}{n} \sum_{a=1}^n \tilde{\gamma}^a = \langle \tilde{\gamma} \rangle^i, \quad \forall i. \quad (47)$$

The invariance of the `kifit` construction under exchange of transitions is discussed in more detail in Appendix D 3.

## Construction of the Log-Likelihood

The aim of the King plot fit is to minimize the distances of the experimental data points  $\tilde{\nu}^a$  (denoted  $\bar{P}^a$  in Fig. 4) from the parallel lines  $\ell^{(a)}$ . These distances are given by

$$\mathbf{d}^a = \mathbf{\Delta}^a - (\mathbf{\Delta}^a \cdot \hat{\mathbf{e}}_{\mathcal{F}}) \hat{\mathbf{e}}_{\mathcal{F}}, \quad a = 1, \dots, n. \quad (48)$$

with  $\hat{\mathbf{e}}_{\mathcal{F}}$  as defined in Eq. (38) and the vectors  $\mathbf{\Delta}^a$  that connect the predictions  $\tilde{\nu}^a|_{\text{pred.}}$  with the data points  $\tilde{\nu}^a$ <sup>11</sup>:

$$\mathbf{\Delta}^a \equiv \tilde{\nu}^a - \tilde{\nu}^a|_{\text{pred.}} = \tilde{\nu}^a - (\mathbf{K} + \tilde{\nu}_1^a \mathcal{F} + \sigma[\delta^a]). \quad (49)$$

More explicitly,

$$d_1^a = -\frac{1}{\|\mathcal{F}\|^2} \sum_{k=2}^m \tan(\phi_{k1}) \Delta_{k1}^a, \quad (50)$$

$$d_{j \neq 1}^a = \Delta_{j1}^a + \tan(\phi_{j1}) d_1^a,$$

where  $\|\mathcal{F}\|^2 = 1 + \sum_{j=2}^m \tan^2 \phi_{j1}$  (see Eq. (38)) and

$$\Delta_{j1}^a = \tilde{\nu}_j^a - \left( K_{j1} + \frac{\tan \phi_{j1}}{\|\mathcal{F}\|} \tilde{\nu}_1^a + \sigma[\delta_{j1}^a] \right). \quad (51)$$

Assuming that the Euclidean norms  $\|\mathbf{d}^a\| = \sqrt{\sum_{i=1}^m (d_i^a)^2}$  approximately follow a multivariate normal distribution, we define the negative log-likelihood

$$-\log \mathcal{L} \propto \frac{1}{2} \sum_{a=1}^n \sum_{b=1}^n \left[ \log \Sigma^{ab} + \|\mathbf{d}^a\| (\Sigma_d^{ab})^{-1} \|\mathbf{d}^b\| \right], \quad (52)$$

where  $\Sigma_d^{ab} \equiv \text{Cov}(\|\mathbf{d}^a\|, \|\mathbf{d}^b\|)$  denotes the covariance of the norms  $\|\mathbf{d}^a\|$  and  $\|\mathbf{d}^b\|$ . In vector notation,

$$-\log \mathcal{L} \propto \frac{1}{2} [\log \det \mathbf{\Sigma}_d + \mathbf{d} \mathbf{\Sigma}_d^{-1} \mathbf{d}], \quad (53)$$

where  $\mathbf{d} \equiv (\|\mathbf{d}^1\|, \dots, \|\mathbf{d}^n\|)^\top$  and  $\mathbf{\Sigma}_d$  is the associated covariance matrix. The latter can be estimated in different ways. For an order-of-magnitude estimate, linear error propagation is sufficient. Assuming the frequency measurements are independent,

$$\begin{aligned} \Sigma_d^{ab} &= \sum_{i=1}^m \sum_{c=1}^n \left( \frac{\partial \|\mathbf{d}^a\|}{\partial \nu_i^c} \sigma[\nu_i^c] \right)^2 \delta^{ab} \\ &+ \sum_A \frac{\partial \|\mathbf{d}^a\|}{\partial m^A} \sigma[m^A]^2 \frac{\partial \|\mathbf{d}^b\|}{\partial m^A} + \sum_{A'} \frac{\partial \|\mathbf{d}^a\|}{\partial m^{A'}} \sigma[m^{A'}]^2 \frac{\partial \|\mathbf{d}^b\|}{\partial m^{A'}} \\ &+ \sum_{j=2}^m \left( \nabla_{\mathbf{K}\phi}^{(j1)} \|\mathbf{d}^a\| \right)^\top \Sigma_{\mathbf{K}\phi}^{(j1)} \left( \nabla_{\mathbf{K}\phi}^{(j1)} \|\mathbf{d}^b\| \right), \end{aligned} \quad (54)$$

<sup>11</sup> Here we are assuming that we only ever compare one data point  $\tilde{\nu}^a$  with a given prediction  $\tilde{\nu}^a|_{\text{pred.}}$ . In presence of  $s$  measurements of the same isotope shift  $\nu_i^a$ ,  $s$  different copies of the element  $\Delta_i^a$  would need to be taken into account. In the current version of the `kifit` code, we assume these complementary measurements to be independent, such that they can be combined in a similar way to separate elements (see Eq. (57)).

where the small indices run over isotope pairs and the capital ones run over isotopes. Here we introduced the notation  $\nabla_{\mathbf{K}\phi}^{j1} \equiv (\partial_{K_{j1}}, \partial_{\phi_{j1}})^\top$  and the  $2 \times 2$  covariance matrix  $\Sigma_{\mathbf{K}\phi}^{(j1)}$  of the fit parameters  $K_{j1}^\perp$  and  $\phi_{j1}$ , which is determined by performing ODR on the data points  $\{(\tilde{\nu}_1^a, \tilde{\nu}_j^a)\}_{a=1}^n$ , as explained in the previous paragraph.

Rather than linear error propagation, the `kifit` code estimates the covariance matrix  $\Sigma_d$  by means of a simple Monte Carlo approach, in which the input parameters  $m^A$ ,  $m^{A'}$  and  $\nu_i^a$ ,  $a = 1, \dots, n$  are sampled from Gaussian distributions fixed by the experimental central values and uncertainties (see Eq. (14)), whereas the fit parameters  $\{K_{j1}^\perp, \phi_{j1}\}_{j=2}^m$  are sampled from the Gaussian distribution fixed by the best fit results  $\{\langle K_{j1}^\perp \rangle, \langle \phi_{j1} \rangle\}_{j=2}^m$  and the covariance matrix  $\Sigma_{\mathbf{K}\phi}^{(j1)}$ :

$$(K_{j1}^\perp, \phi_{j1}) \sim \mathcal{N}(\langle K_{j1}^\perp \rangle, \langle \phi_{j1} \rangle, \Sigma_{\mathbf{K}\phi}^{(j1)}). \quad (55)$$

For better numerical stability, the Cholesky decomposition [81]  $\Sigma_d = \mathbf{L} \mathbf{L}^\top$  into two uniquely determined, lower triangular matrices  $\mathbf{L}$ , is performed and Eq. (53) is reformulated as

$$-\log \mathcal{L} \propto \frac{1}{2} \left[ 2 \sum_{a=1}^n \log(L_{aa}) \mathbf{x} + \mathbf{x}^\top \mathbf{x} \right], \quad (56)$$

where  $\mathbf{x} = \mathbf{L}^{-1} \mathbf{d}$  and  $L_{aa}$  are the diagonal elements of  $\mathbf{L}$ .

The numerical stability can be further improved by employing a regularised version of the covariance matrix,  $\Sigma_d^{(\lambda)} = \Sigma_d + \lambda \mathbf{I}_n$ , where  $\mathbf{I}_n$  is the identity matrix and  $\lambda \ll 1$  is a regulator with the default value  $\lambda = 0$ . We explicitly checked that the spectral decomposition and the Cholesky decomposition lead to comparable results, that these are stable and that the associated numerical uncertainties are negligible.

Since the Fisher information  $I(\alpha_{\text{NP}}) = -\mathbb{E} \left[ \frac{\partial^2 \log \mathcal{L}}{\partial \alpha_{\text{NP}}^2} \right]$ , i.e. the curvature of the log-likelihood with respect to  $\alpha_{\text{NP}}$ , is invariant under linear rescaling of  $\mathbf{d}$ , so are the confidence intervals for  $\alpha_{\text{NP}}$ . This was checked explicitly in the `kifit` code by varying the normalisation of  $\mathbf{d}$ .

Having defined the log-likelihood for one set of independent isotope shifts, the generalisation to a combined log-likelihood is straightforward: Assuming zero correlation between the data sets, the total log-likelihood corresponds to the direct sum of the log-likelihoods  $\log \mathcal{L}^{(E)}$ , associated to the chemical elements (or independent data sets)  $E = 1, 2, \dots$ :

$$-\log \mathcal{L}(\alpha_{\text{NP}}) = - \sum_E \log \mathcal{L}^{(E)}(\alpha_{\text{NP}}). \quad (57)$$

For a fixed value of  $\alpha_{\text{NP}}$  and a set of input and fit parameter samples (see Eqs. (14), (55)), the code evaluates this sum and computes the  $\Delta\chi^2$  values

$$\Delta\chi^2 \equiv 2(x - Q[x, p]), \quad (58)$$

where  $x = -\log \mathcal{L}(\alpha_{\text{NP}})$  and  $Q[x, p]$  denotes the  $p^{\text{th}}$  percentile of the negative log-likelihood values of the full set of  $\alpha_{\text{NP}}$  samples. The value of  $p$  is a hyperparameter that can be fixed by the user (`min_percentile`, in `kifit`). For  $p = 0$ , the minimum of the negative log-likelihood,  $\min(-\log \mathcal{L}(\alpha_{\text{NP}}))$ , is computed. The use of small values of  $p$  can reduce the sensitivity to numerical outliers but too large  $p$ -values will lead to an excessive downward shift of the  $\Delta\chi^2$  values and thus to an overly conservative confidence interval.

In the next section, we will illustrate how the window of favoured  $\alpha_{\text{NP}}$  values is determined in the `kifit` code, while in Sec. III C we will discuss how the confidence intervals and their uncertainties are estimated.

The general structure of the `kifit` code is illustrated in Fig. 6.

### B. Determining the Search Window (Search Phase)

The purpose of the *search phase* is to determine a favoured range of  $\alpha_{\text{NP}}$  values. These will serve as a pool for the Monte Carlo sampling that `kifit` uses to estimate the confidence interval. In the code, two different search options are implemented:

- The generally faster `detloggrid` option constructs a logarithmic grid of  $\alpha_{\text{NP}}$  values, with limits fixed by

$$\begin{aligned} &[\alpha_{\text{NP}}]_{\min}^{\text{alg}} - \text{loggrid\_frac}, \\ &\alpha_{\text{NP}}]_{\max}^{\text{alg}} + \text{loggrid\_frac}, \end{aligned} \quad (59)$$

where  $\alpha_{\text{NP}}]_{\min}^{\text{alg}}$  and  $\alpha_{\text{NP}}]_{\max}^{\text{alg}}$  are the minimum and maximum values found by the algebraic methods (see Eqs. (10), (15), (32) and (33)), for all possible combinations of the experimental input data, while `loggrid_frac` is a user-defined additional number of orders of magnitude used to enlarge the  $\alpha_{\text{NP}}$  scan region.

- The `globaloggrid` option performs an even more agnostic search on a logarithmic grid between  $\alpha_{\text{NP}} = 10^{-15}$  and a maximal exponent which increases with  $m_\phi$ .

As described in the previous section, evaluating the log-likelihood for a given value of  $\alpha_{\text{NP}}$  involves sampling both the input and the fit parameters.

The number of  $\alpha_{\text{NP}}$  samples used in the search phase, and the number of input parameters and fit parameters  $\{K_{j1}^\perp, \phi_{j1}\}_{j=2}^m$  per  $\alpha_{\text{NP}}$  sample can be set by the user. The search window  $[\alpha_\downarrow, \alpha_\uparrow]$  is then fixed by the minimum and maximum  $\alpha_{\text{NP}}$  values,  $\alpha_\downarrow$  and  $\alpha_\uparrow$ , whose  $\Delta\chi^2$  values lie below a  $\Delta\chi^2$  value that is sufficiently large to capture all potentially interesting regions in  $\alpha_{\text{NP}}$  space (see Fig. 7).

Although several orders of magnitude have to be scanned and the Monte Carlo approach only gives reliable results if sufficient statistics are available, we find

that both search strategies work and give comparable results (see Fig. 11).

### C. Estimating the Confidence Interval (Experiment Phase)

Once the *search window*  $[\alpha_\downarrow, \alpha_\uparrow]$  is fixed,  $N_{\text{exp}}$  *experiments* are performed with the aim of determining a reliable confidence interval for  $\alpha_{\text{NP}}$ . In each experiment:

1. A user-defined number  $N_\alpha$  of  $\alpha_{\text{NP}}$  samples are drawn from the normal distribution

$$\alpha_{\text{NP}} \sim \mathcal{N}(\alpha_*, \max(|\alpha_* - \alpha_\downarrow|, |\alpha_\uparrow - \alpha_*|)),$$

where  $\alpha_*$  is the  $\alpha_{\text{NP}}$  value within the search window with the lowest negative log-likelihood.

2. For each of the  $\alpha_{\text{NP}}$  samples, the input and fit parameters are sampled as described in Eq. (14) and (55).
3. Values of  $\Delta\chi^2$  (Eq. (58)) are computed using the samples determined in the previous steps. We obtain  $N_\alpha N_{\text{exp}}$  points in the  $(\alpha_{\text{NP}}, \Delta\chi^2)$ -plane (see Fig. 8).
4. For each experiment  $e$ , a confidence interval  $[\alpha_\downarrow^e, \alpha_\uparrow^e]$  is determined by finding the minimum and maximum  $\alpha_{\text{NP}}$  samples whose  $\Delta\chi^2$  values are below the critical  $\Delta\chi^2$  value associated to  $2m + 1$  degrees of freedom and the number  $N$  of  $\sigma$  ( $N = 2$  being the default value, see red horizontal lines in Fig. 8).
5. For each experiment  $e$ , a *best fit value*  $\alpha_*^e$  is determined, defined as the  $\alpha_{\text{NP}}$  value with the smallest associated negative log-likelihood.

The number of experiments,  $N_{\text{exp}}$ , the number of input and fit parameter samples, the number  $N_\alpha$  of  $\alpha_{\text{NP}}$  samples and the number  $N$  of  $\sigma$  are hyper-parameters that can be fixed by the user. For a stability analysis, see Appendix D 1.

Finally, the results of the  $N_{\text{exp}}$  *experiments* are collected and the final estimation of the confidence interval for  $\alpha_{\text{NP}}$  is determined in the *consolidation phase*.

### D. Producing Results (Consolidation Phase)

Since numerical simulations are used in `kifit` to estimate the new physics bounds, we need a robust strategy to compute both the estimation and its uncertainty. To do so, we apply a technique inspired by the *blocking average method* (BAM), which is commonly used to compute statistically independent estimations starting from a simulated set of possibly correlated data [82]. To facilitate




FIG. 6. Illustration of the *kifit* algorithm used to determine a confidence intervals for  $\alpha_{\text{NP}}$ . The algorithm is composed of a *build phase*, a *search phase*, and *experiment phase* and a *consolidation phase*. See Section III for a description of the individual steps.

the following discussion, we will refer to the illustration presented in Fig. 9.

The BAM involves executing the simulation experiment  $N_{\text{exp}}$  times, thus collecting  $N_{\text{exp}}$  raw estimations. These are then divided into  $B$  blocks of size  $N_b$  (purple subsets in the top right of Fig. 9) and each block is used to produce a single estimation of the target variable. To mitigate eventual correlations among simulated data, it is important to choose the size of the blocks such that the estimation computed from each block can be assumed to be independent from the ones of the other blocks. This is relevant because, if those estimators are independent, their average value and uncertainty stabilises for increasing number of blocks, according to the central limit theorem.

In our case, the raw estimations correspond to confidence intervals of the form  $[\alpha_{\downarrow}^e, \alpha_{\uparrow}^e]$ , where the superscript refers to the result of the experiment  $e$ ,  $1 \leq e \leq$

$N_{\text{exp}}$ . In Ref. [82], and more in general in the context of numerical simulations, the estimation for each block is defined as the average value of the data in the block. We use instead a more conservative approach, with the intention of estimating bounds in the least aggressive way possible. Taking the estimation of the lower bound as an example, we define an estimator as the minimum of values collected in the block  $b$ :

$$\alpha_{\downarrow}^{(b)} \equiv \min_{e \in b} \alpha_{\downarrow}^e. \quad (60)$$

A similar procedure can be used to compute the upper bound, considering instead the maximum value in the block. Since the  $N_{\text{exp}}$  experiments are considered to be independent (contrary to the time-series simulation in Ref. [82]), we can also assume the estimators in Eq. (60) to be statistically independent.

The estimation of the lower bound and its uncertainty






FIG. 7. Output of the *search phase* for a short run of  $N_\alpha = 500$   $\alpha_{\text{NP}}$  samples and only  $N_{\text{exp}} = 10$  experiments. The blue scatter shows the  $(\alpha_{\text{NP}}, \Delta\chi^2(\alpha_{\text{NP}}))$  pairs obtained through the procedure described in Section III B. The upper and lower limits of the search window are indicated by vertical orange lines, the black dashed lines mark the thresholds between which the  $\alpha_{\text{NP}}$ -scaling is linear.

based on  $B$  blocks are computed as follows:

$$\langle \alpha_\downarrow \rangle_B \equiv \frac{1}{B} \sum_{b=1}^B \alpha_\downarrow^{(b)}, \quad (61)$$

$$\sigma[\alpha_\downarrow]_B \equiv \sqrt{\frac{1}{B-1} \sum_{b=1}^B (\alpha_\downarrow^{(b)} - \langle \alpha_\downarrow \rangle_B)^2}, \quad (62)$$

where  $\langle \alpha_\downarrow \rangle_b$  in the second equation corresponds to the estimator defined in the first equation, computed iteratively for any  $b \leq B$ . A similar approach is used to compute the final estimation and uncertainty for the upper bound. The evolution of  $\langle \alpha_\downarrow \rangle_B$  and  $\sigma[\alpha_\downarrow]_B$  for increasing number of blocks is shown in Fig. 10. Ideally, the central value stabilises and the uncertainty decreases for increasing  $B$ . The number of blocks  $B$  is computed from the number of experiments and the block size, both of which can be set by the user. The ideal values for  $N_{\text{exp}}$  and the block size are ones which ensure the stability of the code. See Appendix D 1 for a more detailed discussion.

The final confidence intervals for  $\alpha_{\text{NP}}$  provided by **kifit** take all  $B$  blocks into consideration. We apply the following conservative definition:

$$[\alpha_{\text{NP}}^{\text{LB}}, \alpha_{\text{NP}}^{\text{UB}}] = [\langle \alpha_\downarrow \rangle_B - N \sigma[\alpha_\downarrow]_B, \langle \alpha_\uparrow \rangle_B + N \sigma[\alpha_\uparrow]_B], \quad (63)$$

where  $N$  corresponds the number of  $\sigma$  of the confidence interval. In Fig. 8, the uncertainties on the  $2\sigma$  confidence interval are shown as darker teal bands.

Besides the confidence interval, the **kifit** code provides a *best*  $\alpha_{\text{NP}}$  value (orange star in the upper plot of




FIG. 8. Output of the *experiment phase* for a short run of  $N_\alpha = 500$   $\alpha_{\text{NP}}$  samples and only  $N_{\text{exp}} = 10$  experiments. The teal scatter shows the  $(\alpha_{\text{NP}}, \Delta\chi^2(\alpha_{\text{NP}}))$  pairs obtained through the procedure described in Section III C. Some of the  $\Delta\chi^2$  values are negative due to the choice of  $p = 1$ , (see Eq. (58)). The confidence interval, also shown in teal, is defined as the region in which there are  $\Delta\chi^2(\alpha_{\text{NP}})$  values below the “critical”  $\Delta\chi^2$  value  $\Delta\chi^2_{\text{crit}}$ , indicated by the horizontal red line. The dark teal vertical bars show the uncertainties on the confidence interval. These are estimated by means of the *blocking method* (see Section III D). The best fit  $\alpha_{\text{NP}}$  value is represented by an orange star, its uncertainty being smaller than its width. The lower plot is an enlargement of the confidence interval in the upper plot. The best  $\alpha_{\text{NP}}$  values of the  $N_{\text{exp}}$  experiments are shown in orange.

Fig. 8). This corresponds to the median of the best  $\alpha_{\text{NP}}$  values  $\{\alpha_*^e\}_{e=1}^{N_{\text{exp}}}$  (marked by orange dots in the lower plot in Fig. 8) found in the *experiment phase* (Sec. III C). The uncertainty on the *best*  $\alpha_{\text{NP}}$  value is estimated by the standard deviation of the same set of values  $\{\alpha_*^e\}_{e=1}^{N_{\text{exp}}}$ .

Exclusion plots such as the one shown in Fig. 11 are obtained by repeating the procedure described in Sections III A–III D for multiple values of the mediator mass




FIG. 9. Illustration of the blocking method applied to the estimation of the lower bound (for upper bound:  $\alpha_{\downarrow} \rightarrow \alpha_{\uparrow}$ ). The  $N_{\text{exp}}$  simulated experiments are divided into  $b = 1, \dots, B$  blocks, for each of which the minimum lower bound on  $\alpha_{\text{NP}}$ ,  $\alpha_{\downarrow}^{(b)}$ , is computed. The estimations  $\{\alpha_{\downarrow}^{(b)}\}_{b=1}^B$  are then used to compute an iterative average, the final estimation for the lower bound being  $\langle \alpha_{\downarrow} \rangle_B$ .

$m_{\phi}$ , i.e. for different values of the  $X$  coefficients (see e.g. Eq. (8)).

Fig. 11 compares the  $2\sigma$  bounds obtained by the `kifit` algorithm (orange) to those obtained using the algebraic methods for subsets of the isotope shift data. The results using the Minimal King Plot formula (Eq. (10)) are shown in blue, the No-Mass King Plot results (Eq. (15)) in green and the results using the projection method (Eq. (20)) with  $n = 3$  in purple. For each of these methods, the most stringent limits on  $\alpha_{\text{NP}}$  are connected by lines of matching colour. The horizontal black dashed lines indicate the thresholds between which the plot scale is linear.

As in Fig. 8, the orange stars and bars represent the best fit points and their  $1\sigma$  uncertainties, as obtained by `kifit`. Note that at high-mass values the `kifit` sampling becomes more challenging: the points drawn to evaluate the likelihood are more widely spread around the curve that they trace, and consequently the confidence intervals produced by the experiments exhibit greater variance than at low masses. To mitigate this effect, it is advisable to improve the Monte Carlo sampling, for example by increasing the number of sampled points, using variance reduction techniques, or adopting adaptive strategies in which computational effort grows in proportion to the mass under analysis. In this example we have kept the simulation parameters fixed, so what we observe at high masses is, unsurprisingly, larger error bars. Finally, since we employed a very conservative technique to define the bounds estimates, it is also reasonable to see the bounds widen at high masses.




FIG. 10. Blocking method applied to the estimation of the upper bound on  $\alpha_{\text{NP}}/\alpha_{\text{EM}}$  after executing the procedure described in Sec. III D. These results have been collected from  $N_{\text{exp}} = 500$  experiments divided into  $B = 50$  blocks.

### E. Comparison to a Previous King Plot Fit

As already mentioned in the introduction, the geometric construction used by `kifit` (see Section III A) is inspired by the King plot fit proposed in Ref. [36], but there are some notable differences in the definition of the log-likelihood and the fit parameters. These differences will be discussed in this section.

The fit in Ref. [36] was developed for the case of similar uncertainties across all transitions and isotope pairs (which was the case for the isotope shift data available at that time) and under the (strong) assumptions of uncorrelated uncertainties. In our notation, the  $\chi^2$  function proposed in Ref. [36], takes the form

$$\chi_a^2 = \sum_{i=1}^m \left( \frac{\hat{d}_i^a}{\sigma[\tilde{\nu}_i^a]} \right)^2, \quad \hat{d}_i^a = \tilde{\nu}_i^a - \left( K_{i1} + \frac{\alpha_{\text{NP}}}{\alpha_{\text{EM}}} \tilde{\gamma}^a X_{i1} \right). \quad (64)$$

In order to illustrate the main difference between this definition of  $\chi^2$  and Eq. (58), let us consider the 2-dimensional setup illustrated in Fig. 4, which is characterised by one inclination angle  $\phi_{21}$ . In this case the elements  $\hat{d}_i^a$  take the particularly simple form

$$\hat{d}_1^a = \|\mathbf{d}^a\| \sin \phi_{21}, \quad \hat{d}_2^a = \|\mathbf{d}^a\| \cos \phi_{21}, \quad (65)$$

such that Eq. (64) reads

$$\chi_a^2 = \|\mathbf{d}^a\|^2 \left\{ \left( \frac{\sin \phi_{21}}{\sigma[\tilde{\nu}_1^a]} \right)^2 + \left( \frac{\cos \phi_{21}}{\sigma[\tilde{\nu}_2^a]} \right)^2 \right\}, \quad (66)$$

where  $\|\mathbf{d}^a\|^2 = \|\mathbf{\Delta}^a\| \cos \phi_{21}$ .

Eq. (66) explicitly shows how separately normalising the components of  $\mathbf{d}^a$  by the uncertainties  $\sigma[\tilde{\nu}_i^a]$  distorts the fit: If the data set under consideration has




FIG. 11. Output of the `kikit` code for the `Ca-WT-Aarhus-PTB 2024` dataset, showing the  $2\sigma$  upper and lower bounds on the new physics coupling  $\alpha_{\text{NP}}/\alpha_{\text{EM}}$  as a function of the mass  $m_\phi$  of the new boson with orange connected round markers. The estimation is repeated for the two different *search phase* approaches (see Sec. III B): Left `detloggrid`, adopting the algebraic-induced logarithmic grid and right `globalogrid`, using the global logarithmic grid. The orange stars and bars indicate the best fit points and the associated uncertainties. The bound obtained via the `kikit` algorithm (orange) is compared with the algebraic results: minimal KP (Eq. (10)) in blue, NMKP (Eq. (15)) in green and projection method (Eq. (20)) with  $n = 3$  in purple. The solutions for the different permutations are illustrated by single markers, whereas the envelopes of the most stringent limits are indicated with lines of the same colour. Since the three algebraic methods lead to very similar results, only the purple markers are visible in most cases. For better readability, the regions excluded by any of the plotted bounds are shaded grey. The horizontal black dashed lines indicate the thresholds between which the plot scale is linear.

a direction with significantly smaller uncertainties, e.g.  $\sigma[\tilde{\nu}_1^a] \gg \sigma[\tilde{\nu}_2^a]$ , the minimisation of Eq. (66) will lead to a rotation of the King line, even if this does not reduce the overall distance of the points to the fitted line. Instead, `kikit` minimizes  $\|\mathbf{d}^a\| (\Sigma_d^{ab})^{-1} \|\mathbf{d}^b\|$ , where  $\Sigma_d^{ab}$  is an estimate of the covariance of  $\|\mathbf{d}^a\|$  and  $\|\mathbf{d}^b\|$ . In this way all directions are treated equally, while correlations between the uncertainties are taken into account.

Note also that Eq. (66) does not take into account the fact that a King plot fit cannot capture new physics-induced constant shifts of all data points (see discussion around Eqs. (43)-(45)) and will therefore tend to lead to overly aggressive bounds on  $\alpha_{\text{NP}}$ .

Further improvements with respect to the fit proposed in Ref. [36] are: the estimation of the uncertainties on the best fit point and the confidence interval by means of the *blocking average method* (see Sections III C and III D); the development of an extensible code framework that allows to combine the isotope shift data of several elements in a single fit; an extensive set of numerical tests that ensure the internal consistency of the code and compare `kikit` results in Python with results obtained with Mathematica.

In the following section we provide a comparison of the `kikit` results with results obtained by means of the algebraic methods discussed in Section II.

#### IV. COMPARISON OF FIT AND ALGEBRAIC METHODS

In Section II we reviewed the algebraic methods to extract bounds on  $\alpha_{\text{NP}}$  from King plots, whereas in Section III we presented the King plot fit `kikit`. Although both methods are based on the King plot formalism, they adopt different strategies when it comes to setting bounds on  $\alpha_{\text{NP}}$ . This becomes most evident by stating the questions that either method addresses:

##### Algebraic methods

1.  $\langle \alpha_{\text{NP}} \rangle$ : Given a data set of dimension as specified in Tab. I, which value of  $\alpha_{\text{NP}}$  is required to perfectly describe the experimental central values of the data points in the King plot?
2.  $\sigma[\alpha_{\text{NP}}]$ : How large is the impact of the experimental uncertainties on the value of  $\alpha_{\text{NP}}$ ?

##### King plot fit

1.  $\{K_{j1}^\perp, \phi_{j1}\}_{j=2}^m$ : What is the best linear fit, given the central values and uncertainties on the data points in the King plot (mass-normalised isotope shifts)?
2. Given the central values and uncertainties on the fit parameters and on the data points, what is the likelihood (Eq. (52)) associated to the set of sample  $\alpha_{\text{NP}}$  values?

3.  $[\alpha_{\text{NP}}^{\text{LB}}, \alpha_{\text{NP}}^{\text{UB}}]$ : What are the minimum and maximum  $\alpha_{\text{NP}}$  values whose  $\Delta\chi^2$  values are below the critical  $\Delta\chi^2$  value associated to  $2m + 1$  degrees of freedom and  $N\sigma$ ?

In the rest of the section, we will contrast these approaches in more detail, discuss their respective advantages and disadvantages, present our results and suggest when to apply which method.

### A. Blind Directions

Since both approaches are based on the King plot formalism, they inherit the blind directions of the King plot: As discussed in Section II A, in particular in Eqs. (11) and (12), the King plot method is only sensitive to effects that have a component orthogonal to the mass shift or the field shift, i.e. that lead to a “misalignment” both in transition space and in isotope pair space. In a standard King plot in transition space such a misalignment manifests itself in a deviation from the King line (Eq. (5)). This is the effect that the King plot fit looks for. Equivalently, it can be viewed as a component orthogonal to the plane of King linearity fixed by the basis vectors  $\boldsymbol{\mu}$  and  $\boldsymbol{\delta}\langle\mathbf{r}^2\rangle$  (or equivalently  $\mathbf{1}$  and  $\tilde{\mathbf{v}}_1$ , as in Fig. 2) in isotope pair space. This is the effect exploited by the algebraic methods and the nonlinearity decomposition (Eq. (23)).

The blind directions of the King plot method and their manifestations in the algebraic methods and on the King plot fit `kifit` are summarised in Table II: Using the notation of Eq. (12), they can be described by vanishing determinants of the matrix of electronic or nuclear factors or, in `kifit`, either by an alignment of the electronic coefficients  $X_j/X_1$  and  $F_{j1} = F_j/F_1$  leading to a vanishing coefficient  $X_{j1} \rightarrow 0$ , or by a global shift of the King line, which cannot be distinguished from a redefinition of the electronic mass shift coefficient  $\mathcal{K}$  (see Fig. 4).

Another common feature of the fit and the algebraic methods which is inherited from the King plot, is the insensitivity to the experimental uncertainties parallel to the King line: Whereas this is true by construction for the King plot fit (see Fig. 4), it is less obvious in the case of the algebraic methods. In Appendix C we check explicitly that the effect of the isotope shift uncertainties parallel to the King line has a negligible impact on  $\sigma[\alpha_{\text{NP}}]$ .

### B. Geometric Construction & Form of Data Sets

The most obvious difference lies in the geometric constructions. The algebraic methods compare volumes constructed from a combination of data points and predictions (see Fig. 1), providing one constraint on  $\alpha_{\text{NP}}$  per data set. It is for this reason that the data sets used by the algebraic methods must have a fixed size (see Table I).

Algebraic Methods	<code>kifit</code>
$\det(\mathcal{N}) \rightarrow 0$	$\tilde{\gamma}^a \rightarrow \langle\tilde{\gamma}\rangle \quad \forall a$
$\det(\mathcal{M}) \rightarrow 0$	$X_{j1} \rightarrow 0 \quad \forall j$

TABLE II. Blind directions of the King plot method and how they manifest themselves in the algebraic methods and in the fit. The matrices  $\mathcal{M}$  and  $\mathcal{N}$  are defined in Eq. (12) and (16) for the cases of the simple King plot and the No-Mass King Plot, respectively, whereas  $X_{j1}$  and  $\langle\tilde{\gamma}\rangle$  are defined in Eqs. (41) and (43), respectively.

The fit, on the other hand, minimizes the distances  $\|\mathbf{d}^a\|$  of the data points  $\tilde{\mathbf{v}}^a$  from the set of King lines  $\{\ell^{(a)}\}_{a=1}^n$ , which are determined by a combination of the best linear fit to data,  $\ell^{(0)}$ , and the predicted nonlinearity induced by new physics (see Fig. 4). Since the fit is based on a log-likelihood, which sums over the data points (Eq. (52)), it can handle data sets of dimension  $(n, m)$ , with  $3 \leq n$  and  $2 \leq m$ , but where  $n$  and  $m$  are independent (see Table I), and also combine multiple data sets, enabling global fits to data. This makes the fit particularly valuable for the search for new physics, which is expected to couple in the same way to electrons and neutrons, irrespective of the element.

#### Size of Data Sets

Whereas the algebraic methods work best for small data sets, the fit relies on the presence of sufficiently large and diverse data sets. When this is not the case, there is a risk of overfitting. Overfitting happens when a statistical model is too expressive for a given dataset. The optimal fit parameters are obtained by learning the statistical fluctuations or any other source of error. For example, using a data set with two points only, we would obtain a perfect interpolation and could define a margin for new physics in the form of a finite value for  $\sigma[\alpha_{\text{NP}}]$ , but at the same time, we would be performing maximum over-fitting: All data is used to fix the fit parameters  $\{K_{j1}^\perp, \phi_{j1}\}_{j=2}^m$  and the bound on the remaining fit parameter  $\alpha_{\text{NP}}$  is meaningless. Indeed, in order to fit  $2(m-1) + 1 = 2m - 1$  degrees of freedom with  $n$  data points, each of which contributes an additional scalar constraint (distance  $\|\mathbf{d}^a\|$ ),  $n \geq 2m - 1$  data points are required (see also Table I).

Data sets with just 2 transitions and 2 isotope pairs are not considered in King plots for precisely this reason. However, since the statistical uncertainty associated to the spread of the points is expected to scale as  $1/\sqrt{n}$ , the fit, which minimises the distances from the data points to the King line, will significantly underestimate the bounds on  $|\alpha_{\text{NP}}|$  in the limit of sparse data.

In Appendix E, we generate mock data starting from a set of linear relations. This allows us to study the impact

Data Set	$(\alpha_{\text{NP}} \pm \sigma[\alpha_{\text{NP}}]) _{\text{KP}}^{(1)}$	$(\alpha_{\text{NP}} \pm \sigma[\alpha_{\text{NP}}]) _{\text{KP}}^{\text{MC}}$	$[\alpha_{\text{NP}}^{\text{LB}} \pm \sigma[\alpha_{\text{NP}}]_B, \alpha_{\text{NP}}^{\text{UB}} \pm \sigma[\alpha_{\text{NP}}]_B]$
Ca3pointTEST	$(1.36 \pm 4.84) \times 10^{-10}$	$(1.36 \pm 4.81) \times 10^{-10}$	$[-3.51 \pm 0.16, 3.59 \pm 0.13] \times 10^{-11}$
Ca4pointTEST	$(-0.14 \pm 4.50) \times 10^{-11}$	$(-0.14 \pm 4.50) \times 10^{-11}$	$[-3.44 \pm 0.15, 3.29 \pm 0.09] \times 10^{-11}$
Ca10pointTEST	$(0.19 \pm 1.35) \times 10^{-10}$	$(0.19 \pm 1.31) \times 10^{-10}$	$[0.32 \pm 0.07, 1.65 \pm 0.08] \times 10^{-10}$
Ca_PTB_2015	$(0.65 \pm 1.10) \times 10^{-8}$	$(0.65 \pm 1.10) \times 10^{-8}$	$[-7.04 \pm 0.20, 8.04 \pm 0.50] \times 10^{-10}$
Camn	$(1.36 \pm 1.12) \times 10^{-9}$	$(1.36 \pm 1.1) \times 10^{-9}$	$[-0.91 \pm 0.03, 1.0 \pm 0.05] \times 10^{-10}$
Ca24min	$(1.36 \pm 4.70) \times 10^{-10}$	$(1.36 \pm 4.74) \times 10^{-10}$	$[-2.66 \pm 0.13, 2.55 \pm 0.07] \times 10^{-11}$
Ca_WT_Aarhus_PTB_2024	$(1.36 \pm 4.70) \times 10^{-10}$	$(1.36 \pm 4.72) \times 10^{-10}$	$[-1.99 \pm 0.04, 1.45 \pm 0.05] \times 10^{-9}$

TABLE III. Estimates of the  $1\sigma$  confidence interval for  $\alpha_{\text{NP}}$  for  $m_\phi = 1$  eV. The first column shows the central value of  $\alpha_{\text{NP}}$  (Eq. (10)) and its uncertainty obtained using linear error propagation (Eq. (C11)). The uncertainties in the second column were estimated using a Monte Carlo approach (see Eq. (14) to Eq. (10)) with 2000 samples. Both for column 1 and column 2, those subsets of the data were employed that lead to the most stringent bounds on  $\alpha_{\text{NP}}$ . The *kifit* results are listed in the last column. These were obtained using a search phase on a *detlogrid* with *logrid\_frac*=2, 500  $\alpha_{\text{NP}}$  samples and 200 input and fit parameter samples per  $\alpha_{\text{NP}}$  sample. The experiment phase consisted of 150 experiments, with 1000  $\alpha_{\text{NP}}$  samples each and 500 input and fit parameter samples per  $\alpha_{\text{NP}}$  sample. The experiments were divided into blocks of size 25.




FIG. 12. Comparison of the fit results with the algebraic results (KP: Eq. (10), NMKP: Eq. (15), PROJ: Eq. (20)) for the minimal dataset (3 isotope pairs, 2 transitions) Ca24min. For better readability, the regions excluded by any of the plotted bounds are shaded grey, even if these are considered too aggressive in the case of Ca24min (see text for discussion). The black dashed lines indicate the thresholds between which the scaling in the plot is linear.

of data sparsity on the fit results in more detail.

#### Numerical Comparisons

To explicitly compare results obtained using the algebraic methods and the fit, we list in Table III the values of  $\sigma[\alpha_{\text{NP}}]$  estimated by means of linear error prop-




FIG. 13. Fit results for two data sets of dimensions  $(n, m) = (3, 2)$  (Ca24min, Ca\_PTB\_2015) and their combination of dimension  $(3, 4)$  (Ca\_WT\_Aarhus\_PTB\_2024). The most stringent algebraic bounds (KP: Eq. (10)) for Ca\_WT\_Aarhus\_PTB\_2024 correspond to the algebraic bounds for Ca24min and are shown in blue. Whereas the difference between the fit results and the algebraic results are sizeable for the  $(3, 2)$ -dimensional data sets, the agreement is better for Ca\_WT\_Aarhus\_PTB\_2024.

agation through Eq. (10) ( $\sigma[\alpha_{\text{NP}}]|_{\text{KP}}^{(1)}$ , first column) and the Monte Carlo approach outlined in Section II A, in particular in Eqs. (13) and (14) ( $\sigma[\alpha_{\text{NP}}]|_{\text{KP}}^{\text{MC}}$ , second column), as well as the  $1\sigma$  confidence interval determined in *kifit* by the BAM, Eq. (63) (last column). We observe good agreement between the linear uncertainty propagation and the Monte Carlo approach. In contrast, the *kifit* results show some deviations. Notably,






FIG. 14. Fit results for a Ca data set, an Yb data set and the combination thereof. The (most stringent) algebraic results for the Ca data set are also shown.

for all minimal data sets (2 transitions and 3 isotope pairs), the fit results are roughly an order of magnitude stronger. However, we notice that the discrepancy is lifted for larger datasets, such as **Ca4pointTEST** and **Ca10pointTEST** (each with 2 transitions and 4 or 10 isotope pairs) and **Ca\_WT\_Aarhus\_PTB\_2024** (4 transitions and 3 isotope pairs). For a description of these data sets and the relevant references, see Appendix F.

This behaviour can also be observed in the exclusion plot in Fig. 12, which shows the bounds on  $\alpha_{NP}$  for the minimal data set **Ca24min**. Here the fit bounds (orange) appear to be nearly two orders of magnitude more stringent than the bounds obtained using the minimal algebraic method (Eq. (10), blue) and the projection method (Eq. (20), purple). Fig. 13 shows the same fit result as Fig. 12, but complemented with those using the minimal data set **Ca\_PTB\_2015**, and **Ca\_WT\_Aarhus\_PTB\_2024**, which combines **Ca\_PTB\_2015** and **Ca24min** to a data set for 3 isotope pairs and 4 transitions. The fit and the algebraic results using **Ca\_WT\_Aarhus\_PTB\_2024** show significantly better agreement.

A similar situation can be observed in Fig. 14, which combines **Ca\_WT\_Aarhus\_PTB\_2024** with **Yb\_Kyoto\_MIT\_GSI\_MPIK\_2024**, a set of Yb isotope shifts for 4 isotope pairs and 5 transitions. The fit result for Yb must be interpreted with caution, since it is known to contain SM nonlinearities, which are, however, not taken into account by the current version of **kifit**. In Appendix E we discuss the impact of data sparsity in more detail.

In Figs. 13 and 14, the fit appears to lose sensitivity more quickly than the algebraic methods at the high-mass end. We trace this behaviour back to the **Ca\_PTB\_2015** data set, which starts to lose sensitivity comparatively rapidly at around  $m_\phi = 10^{-6}$  eV. Since the algebraic bound in Fig. 13 shows the envelope of the bounds, and the algebraic bounds for **Ca24min** turn out

to be more stringent, the behaviour of the **Ca\_PTB\_2015** bound is not visible in the algebraic results plotted in Figs. 13 and 14. Nonetheless, the robustness of the fit in the high-mass region could be improved by ameliorating the Monte Carlo sampling discussed in Section IIID and by increasing the resolution in  $m_\phi$ . For completeness, we performed a series of higher-resolution simulations executing the algebraic KP method on the **Ca\_PTB\_2015** dataset (equivalent to the blue curve in Fig. 13, but considering eighty mass values in the same range) and we were able to resolve similar spikes to the one shown in the **kifit** bound in Fig. 13.

## V. CONCLUSIONS AND OUTLOOK

Over the past few years, several precision isotope shift measurements have been performed, spurring the development of a wide range of King plot-inspired methods to search for new physics and study nuclear structure. In this work, we systematically compare the available analysis methods, complementing them with the No-Mass Generalised King Plot and the Nuclear Input King Plot formulas, and taking a first step towards a global analysis of isotope shift data with our novel fit framework **kifit**.

The key strengths of **kifit** originate from its construction based on a log-likelihood: While the algebraic methods require data sets with  $n = m$  or  $n = m + 1$  isotope pairs for  $m$  transitions, the fit is flexible with respect to  $n$  and  $m$ . It thus avoids the omission of data or cherry-picking the subset that yields the most stringent bound on new physics. Under the assumption of uncorrelated measurements, it further allows to combine isotope shifts from different elements.

This is particularly valuable for the search for new bosons whose couplings to atomic nuclei do not introduce element-dependent parameters. The Standard Model extension consisting of a new boson coupling linearly to neutrons and electrons, which is commonly investigated using King plots, is particularly predictive. **kifit** can, for the first time, test this model across elements, setting *global* bounds on the product of the new boson's couplings to neutrons and to electrons, for new boson masses spanning eight orders of magnitude.

The fit does, however, rely on the existence of sufficiently large data sets, since it tends to underestimate the margin for new physics when data is sparse. For this reason, measuring isotope shifts and nuclear masses in larger arrays of isotopes would not only provide new opportunities to resolve higher-order nuclear effects, but also allow more stable and reliable bounds on new physics to be established. Example systems with a large number of even stable isotopes are Zn (5 stable isotopes of which 4 are even, see Ref. [58] for recent isotope shift data), Sn [83] (10 stable isotopes of which 7 are even, plus one trace radioisotope with  $\tau_{1/2} \sim 10^5$  y), Ba (6 stable isotopes of which 4 are even, plus one even isotope with a half-life  $\tau_{1/2} \sim 10^{21}$  y, measured so far with 100-400 MHz

uncertainty [84]) and Cd (6 stable isotopes of which 5 are even, plus one even isotope with  $\tau_{1/2} \sim 10^{19}$  y; Ref. [85] reached a resolution of few MHz for isotope shifts in two transitions), plus the hitherto unexploited metastable isotopes in Ca and Yb, amongst others.

Recently, several additional isotope shift measurements were completed: Ref. [86] achieved a precision below 10 Hz in the 431 nm transition in neutral Yb; Ref. [87] reduced the uncertainty of isotope shifts in the 369 nm and 935 nm transitions in  $\text{Yb}^+$  to the sub-MHz level; Ref [88] measured isotope shifts in four metastable state transitions in  $\text{Yb}^+$  at 30 MHz and sub-MHz level. The isotope shift in the fine structure splitting of  $P$  states in  $\text{Kr}^+$  was measured at sub-MHz precision in Ref. [89]. Isotope shifts measured at sub-10 MHz precision in  $\text{Th}^{3+}$  were reported in Ref. [90]. Ref. [91] determined isotope shifts in 3 isotope pairs in Rydberg transitions of Sr (including the odd isotope  $^{87}\text{Sr}$ ) at a sub-MHz precision. Ref. [92] extended the King plot analysis to isotope shifts in diatomic molecules. These developments add to the growing treasure of isotope shift data and pave the path towards high-precision isotope shifts in a multitude of elements and even molecules.

Whereas the current version of the `kifit` code is limited to King plots that are linear within uncertainties, King plot nonlinearities could be included in future versions, e.g. by subtracting them as in Ref. [43], provided accurate predictions or complementary experimental input are available. The framework provided by the `kifit` code might even be applied to obtain insights into higher-order nuclear effects in a data-driven way, as was suggested in Ref. [50].

Note that all King plot searches for new physics, including `kifit`, rely on predictions for the electronic coefficients that enter the new physics term, while the analysis of nonlinear King plots using `kifit`-like setups would additionally require predictions for higher-order SM terms. For this reason, it may be beneficial to focus on elements whose dominant King nonlinearities are higher-order mass shifts, in which case the nuclear contributions can be determined from experimental data and the electronic coefficients can be predicted with comparatively good precision [43, 52]. In systems where the second-order mass shift is the only resolvable King nonlinearity, the Nuclear Input King Plot formula presented in this work can be employed to extract the new physics coupling from isotope shift data for just two transitions and four isotope pairs.

In conclusion, we encourage experimentalists to measure isotope shifts across additional transitions

and isotopes, even if the initial precision is lower than that of the best current measurements. Additional data points can help eliminate higher-order effects in the isotope shift equations and stabilise the bounds on new physics. They also provide new insights into nuclear structure and can thus be used to refine nuclear models. If new isotope shifts are measured before the corresponding isotope masses, the No-Mass King Plot and its generalised version facilitate a first analysis of the new data and indicate the required precision of the mass measurements to achieve competitive sensitivity to new physics and the nuclear shape. Ultimately, it is the combination of new measurements, advancements in atomic and nuclear structure theory and global (King plot) analyses that will enhance the sensitivity to new physics and to nuclear effects.

## ACKNOWLEDGMENTS

We thank Julian Berengut for providing the  $X$  coefficients of Yb that he calculated using the open-source atomic structure code `AMBiT` [59], and for allowing us to make them publicly available in `kifit`. FK would like to thank Peter Stangl for fruitful discussions during the early stages of this project. We thank our collaborators in the  $\text{Yb}^+$  [50] and  $\text{Ca}^+/\text{Ca}^{14+}$  [43] King plot analyses for the insightful discussions. EF, FK and JR acknowledge funding by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany’s Excellence Strategy – EXC-2123 QuantumFrontiers – 390837967. AM is partially funded by the DFG via the project B10 of the Collaborative Research Cluster 1227 (DQ-mat) – Project-ID 274200144 and also acknowledges support by the EXC-2123 QuantumFrontiers. EF was funded by CERN in the initial phase of this work and thanks CERN-TH and the CERN QTI for the stimulating environment. FK thanks CERN-TH for its hospitality during the early stages of this project and for support via the CERN-TH visitor program. AM thanks the Graduate Academy of Leibniz University Hannover for financial support of an extended research stay at CERN during the early stages of the project and CERN-TH for its hospitality. MR is supported by CERN through the CERN Quantum Technology Initiative (QTI). This work has been partially funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) - 491245950. This collaboration came about thanks to the CERN QTI Forum.

---

[1] M. S. Safronova, D. Budker, D. DeMille, Derek F. Jackson Kimball, A. Derevianko, and C. W. Clark, “Search for New Physics with Atoms and Molecules,” *Rev. Mod. Phys.* **90**, 025008 (2018), arXiv:1710.01833 [physics.atom-ph].

[2] M. G. Kozlov, M. S. Safronova, J. R. Crespo López-Urrutia, and P. O. Schmidt, “Highly charged ions: Optical clocks and applications in fundamental physics,” *Rev. Mod. Phys.* **90**, 045005 (2018), arXiv:1803.06532 [physics.atom-ph].

- [3] D. Antypas *et al.*, “New Horizons: Scalar and Vector Ultraviolet Dark Matter,” (2022), [arXiv:2203.14915 \[hep-ex\]](#).
- [4] T.W. Hänsch and A.L. Schawlow, “Cooling of gases by laser radiation,” *Optics Communications* **13**, 68–69 (1975).
- [5] D.J. Wineland and H. Dehmelt, “Proposed  $10^{14} \Delta\nu < \nu$  Laser Fluorescence Spectroscopy on  $\text{Ti}^+$  Mono-Ion Oscillator,” *Bulletin of the American Physical Society* **20**, 637 (1975).
- [6] Stig Stenholm, “The semiclassical theory of laser cooling,” *Rev. Mod. Phys.* **58**, 699–739 (1986).
- [7] Harold Metcalf and Peter van der Straten, “Cooling and trapping of neutral atoms,” *Physics Reports* **244**, 203–286 (1994).
- [8] Theodor W. Hänsch, “Nobel lecture: Passion for precision,” *Rev. Mod. Phys.* **78**, 1297–1309 (2006).
- [9] John L. Hall, “Nobel lecture: Defining and measuring optical frequencies,” *Rev. Mod. Phys.* **78**, 1279–1295 (2006).
- [10] A. Ashkin, “Acceleration and trapping of particles by radiation pressure,” *Phys. Rev. Lett.* **24**, 156–159 (1970).
- [11] A. Ashkin, J. M. Dziedzic, J. E. Bjorkholm, and Steven Chu, “Observation of a single-beam gradient force optical trap for dielectric particles,” *Opt. Lett.* **11**, 288 (1986).
- [12] Steven Chu, J. E. Bjorkholm, A. Ashkin, and A. Cable, “Experimental Observation of Optically Trapped Atoms,” *Phys. Rev. Lett.* **57**, 314–317 (1986).
- [13] V. S. Letokhov, “Narrowing of the Doppler Width in a Standing Wave,” *Soviet Journal of Experimental and Theoretical Physics Letters* **7**, 272 (1968).
- [14] V I Balykin, V S Letokhov, and V G Minogin, “Laser control of the motion of neutral atoms and optical atomic traps,” *Physica Scripta* **1988**, 119 (1988).
- [15] F. Mauri and E. Arimondo, “Two dimension selective coherent population trapping controlled by a phase shift,” *Physical Review Letters* **54** (1992), [10.1007/bf00325388](#).
- [16] Rudolf Grimm, Matthias Weidemüller, and Yurii B. Ovchinnikov, “Optical dipole traps for neutral atoms,” (1999), [arXiv:physics/9902072 \[physics.atom-ph\]](#).
- [17] P. O. Schmidt, T. Rosenband, C. Langer, W. M. Itano, J. C. Bergquist, and D. J. Wineland, “Spectroscopy using quantum logic,” *Science* **309**, 749–752 (2005).
- [18] D. J. Heinzen and D. J. Wineland, “Quantum-limited cooling and detection of radio-frequency oscillations by laser-cooled ions,” *Phys. Rev. A* **42**, 2977–2994 (1990).
- [19] J. I. Cirac and P. Zoller, “Quantum computations with cold trapped ions,” *Phys. Rev. Lett.* **74**, 4091–4094 (1995).
- [20] D. J. Wineland, C. Monroe, W. M. Itano, D. Leibfried, B. E. King, and D. M. Meekhof, “Experimental issues in coherent quantum state manipulation of trapped atomic ions,” *J. Res. Natl. Inst. Stand. Tech.* **103**, 259 (1998), [arXiv:quant-ph/9710025](#).
- [21] Alexander Aepli, Kyungtae Kim, William Warfield, Marianna S. Safronova, and Jun Ye, “Clock with  $8 \times 10^{-19}$  Systematic Uncertainty,” *Phys. Rev. Lett.* **133**, 023401 (2024).
- [22] H. N. Hausser, J. Keller, T. Nordmann, N. M. Bhatt, J. Kiethe, H. Liu, I. M. Richter, M. von Boehn, J. Rahm, S. Weyers, E. Benkler, B. Lipphardt, S. Dörscher, K. Stahl, J. Klose, C. Lisdat, M. Filzinger, N. Huntemann, E. Peik, and T. E. Mehlstäubler, “ $^{115}\text{In}^+ - ^{172}\text{Yb}^+$  Coulomb Crystal Clock with  $2.5 \times 10^{-18}$  Systematic Uncertainty,” *Phys. Rev. Lett.* **134**, 023201 (2025).
- [23] W. H. King, “Comments on the article “Peculiarities of the isotope shift in the samarium spectrum”,” *J. Opt. Soc. Am.* **53**, 638 (1963).
- [24] W. H. King, *Isotope shifts in atomic spectra* (Springer Science & Business Media, 1984).
- [25] I. Angeli and K.P. Marinova, “Table of experimental nuclear ground state charge radii: An update,” *Atomic Data and Nuclear Data Tables* **99**, 69–95 (2013).
- [26] Florian Gebert, Yong Wan, Fabian Wolf, Christopher N. Angstmann, Julian C. Berengut, and Piet O. Schmidt, “Precision Isotope Shift Measurements in Calcium Ions Using Quantum Logic Detection Schemes,” *Phys. Rev. Lett.* **115**, 053003 (2015), [arXiv:1504.03139 \[physics.atom-ph\]](#).
- [27] C Shi, F Gebert, C Gorges, S Kaufmann, W Nörtershäuser, BK Sahoo, A Surzhykov, VA Yerokhin, JC Berengut, F Wolf, *et al.*, “Unexpectedly large difference of the electron density at the nucleus in the  $4p^2\text{P}_{1/2,3/2}$  fine-structure doublet of  $\text{Ca}^+$ ,” *Applied Physics B*, 1–19 (2018), [arXiv:1608.07572 \[physics.atom-ph\]](#).
- [28] Cédric Delaunay, Roe Ozeri, Gilad Perez, and Yotam Soreq, “Probing Atomic Higgs-like Forces at the Precision Frontier,” *Phys. Rev. D* **96**, 093001 (2017), [arXiv:1601.05087 \[hep-ph\]](#).
- [29] Julian C. Berengut *et al.*, “Probing New Long-Range Interactions by Isotope Shift Spectroscopy,” *Phys. Rev. Lett.* **120**, 091801 (2018), [arXiv:1704.05068 \[hep-ph\]](#).
- [30] Michael Bordag, Galina Leonidovna Klimchitskaya, Umar Mohideen, and Vladimir Mikhaylovich Mostepanenko, *Advances in the Casimir Effect* (Oxford University Press, 2009).
- [31] Rouven Essig, Philip Schuster, and Natalia Toro, “Probing Dark Forces and Light Hidden Sectors at Low-Energy  $e^+e^-$  Colliders,” *Phys. Rev. D* **80**, 015003 (2009), [arXiv:0903.3941 \[hep-ph\]](#).
- [32] Rouven Essig, Philip Schuster, Natalia Toro, and Bogdan Wojtsekhowski, “An Electron Fixed Target Experiment to Search for a New Vector Boson  $A'$  Decaying to  $e^+e^-$ ,” *JHEP* **02**, 009 (2011), [arXiv:1001.2557 \[hep-ph\]](#).
- [33] Brian Batell, Maxim Pospelov, and Adam Ritz, “Exploring Portals to a Hidden Sector Through Fixed Targets,” *Phys. Rev. D* **80**, 095024 (2009), [arXiv:0906.5614 \[hep-ph\]](#).
- [34] J. P. Lees *et al.* (BaBar), “Search for  $B \rightarrow K^{(*)}\nu\bar{\nu}$  and invisible quarkonium decays,” *Phys. Rev. D* **87**, 112005 (2013), [arXiv:1303.7465 \[hep-ex\]](#).
- [35] A. V. Artamonov *et al.* (E949), “New Measurement of the  $K^+ \rightarrow \pi^+\nu\bar{\nu}$  Branching Ratio,” *Phys. Rev. Lett.* **101**, 191802 (2008), [arXiv:0808.2459 \[hep-ex\]](#).
- [36] Claudia Frugiuele, Elina Fuchs, Gilad Perez, and Matthias Schlaffer, “Constraining New Physics Models with Isotope Shift Spectroscopy,” *Phys. Rev. D* **96**, 015011 (2017), [arXiv:1602.04822 \[hep-ph\]](#).
- [37] Simon Knapen, Tongyan Lin, and Kathryn M. Zurek, “Light Dark Matter: Models and Constraints,” *Phys. Rev. D* **96**, 115021 (2017), [arXiv:1709.07882 \[hep-ph\]](#).
- [38] A. Kramida, “Isotope shifts in neutral and singly-



- ionized calcium,” *Atomic Data and Nuclear Data Tables* **133–134**, 101322 (2020).
- [39] David Röser, Lukas Möller, Hans Keffler, and Simon Stellmer, “Isotope-shift measurement of the 423-nm transition in neutral Ca,” *Phys. Rev. A* **110**, 032809 (2024), arXiv:2406.06806 [physics.atom-ph].
- [40] Felix W. Knollmann, Ashay N. Patel, and S. Charles Doret, “Part-per-billion measurement of the  $4^2S_{1/2} \rightarrow 3^2D_{5/2}$  electric quadrupole transition isotope shifts between  $^{42,44,48}\text{Ca}^+$  and  $^{40}\text{Ca}^+$ ,” *Phys. Rev. A* **100**, 022514 (2019), [Erratum: *Phys. Rev. A* **107**, 069902 (2023)], arXiv:1906.04105 [physics.atom-ph].
- [41] Cyrille Solaro, Steffen Meyer, Karin Fisher, Julian C. Berengut, Elina Fuchs, and Michael Drewsen, “Improved isotope-shift-based bounds on bosons beyond the Standard Model through measurements of the  $^2D_{3/2}-^2D_{5/2}$  interval in  $\text{Ca}^+$ ,” *Phys. Rev. Lett.* **125**, 123003 (2020), [Erratum: *Phys. Rev. Lett.* **127**, 029901 (2021)], arXiv:2005.00529 [physics.atom-ph].
- [42] Timothy T. Chang, Bless Bah Awazi, Julian C. Berengut, Elina Fuchs, and S. Charles Doret, “Systematic-free limit on new light scalar bosons via isotope-shift spectroscopy in  $\text{Ca}^+$ ,” *Phys. Rev. A* **110**, L030801 (2024), arXiv:2311.17337 [physics.atom-ph].
- [43] Alexander Wilzewski *et al.*, “Nonlinear calcium King plot constrains new bosons and nuclear properties,” (2024), arXiv:2412.10277 [physics.atom-ph].
- [44] D. L. Clark, M. E. Cage, D. A. Lewis, and G. W. Greenlees, “Optical isotopic shifts and hyperfine splittings for Yb,” *Phys. Rev. A* **20**, 239–253 (1979).
- [45] C. J. Bowers, D. Budker, S. J. Freedman, G. Gwinner, J. E. Stalnaker, and D. DeMille, “Experimental investigation of the  $6s^{21}S_0 \rightarrow 5d6s^3D_{1,2}$  forbidden transitions in atomic ytterbium,” *Phys. Rev. A* **59**, 3513–3526 (1999).
- [46] Michaela Kleinert, M. E. Gold Dahl, and Scott Bergeson, “Measurement of the  $\text{Yb I } ^1S_0-^1P_1$  transition frequency at 399 nm using an optical frequency comb,” *Phys. Rev. A* **94**, 052511 (2016).
- [47] N. L. Figueroa, J. C. Berengut, V. A. Dzuba, V. V. Flambaum, D. Budker, and D. Antypas, “Precision Determination of Isotope Shifts in Ytterbium and Implications for New Physics,” *Phys. Rev. Lett.* **128**, 073001 (2022).
- [48] Ian Counts, Joonseok Hur, Diana P. L. Aude Craik, Honggi Jeon, Calvin Leung, Julian C. Berengut, Amy Geddes, Akio Kawasaki, Wonho Jhe, and Vladan Vuletić, “Evidence for Nonlinear Isotope Shift in  $\text{Yb}^+$  Search for New Boson,” *Phys. Rev. Lett.* **125**, 123002 (2020), arXiv:2004.11383 [physics.atom-ph].
- [49] Joonseok Hur *et al.*, “Evidence of Two-Source King Plot Nonlinearity in Spectroscopic Search for New Boson,” *Phys. Rev. Lett.* **128**, 163201 (2022), arXiv:2201.03578 [physics.atom-ph].
- [50] Menno Door *et al.*, “Probing New Bosons and Nuclear Structure with Ytterbium Isotope Shifts,” *Phys. Rev. Lett.* **134**, 063002 (2025), arXiv:2403.07792 [physics.atom-ph].
- [51] Saleh O. Allehabi, V. A. Dzuba, V. V. Flambaum, and A. V. Afanasjev, “Nuclear deformation as a source of the nonlinearity of the King plot in the  $\text{Yb}^+$  ion,” *Phys. Rev. A* **103**, L030801 (2021).
- [52] Anna V. Viatkina, Vladimir A. Yerokhin, and Andrey Surzhykov, “Calculation of isotope shifts and King-plot nonlinearities in  $\text{Ca}^+$ ,” *Phys. Rev. A* **108**, 022802 (2023), arXiv:2306.06990 [physics.atom-ph].
- [53] Julian C. Berengut, Cédric Delaunay, Amy Geddes, and Yotam Soreq, “Generalized King linearity and new physics searches with isotope shifts,” *Phys. Rev. Res.* **2**, 043444 (2020), arXiv:2005.06144 [hep-ph].
- [54] Julian C. Berengut and Cédric Delaunay, “Precision isotope-shift spectroscopy for new physics searches and nuclear insights,” *Nature Rev. Phys.* **7**, 119–125 (2025).
- [55] CWP Palmer and DN Stacey, “Theory of anomalous isotope shifts in samarium,” *Journal of Physics B: Atomic and Molecular Physics* **15**, 997 (1982).
- [56] Julian C. Berengut and Natalia S. Oreshkina, “Second-order hyperfine structure in strontium and impact on new physics searches using isotope shift spectroscopy,” (2024), arXiv:2409.01530 [physics.atom-ph].
- [57] Simon Hofsäss, J. Eduardo Padilla-Castillo, Sid C. Wright, Sebastian Kray, Russell Thomas, Boris G. Sartakov, Ben Ohayon, Gerard Meijer, and Stefan Truppe, “High-resolution isotope-shift spectroscopy of  $\text{Cd I}$ ,” *Phys. Rev. Res.* **5**, 013043 (2023), arXiv:2210.11425 [physics.atom-ph].
- [58] David Röser, J. Eduardo Padilla-Castillo, Ben Ohayon, Russell Thomas, Stefan Truppe, Gerard Meijer, Simon Stellmer, and Sid C. Wright, “Hyperfine structure and isotope shifts of the  $(4s^2)^1S_0 \rightarrow (4s4p)^1P_1$  transition in atomic zinc,” *Phys. Rev. A* **109**, 012806 (2024).
- [59] E.V. Kahl and J.C. Berengut, “ambit: A programme for high-precision relativistic atomic structure calculations,” *Computer Physics Communications* **238**, 232–243 (2019).
- [60] G. Breit, “Theory of Isotope Shift,” *Rev. Mod. Phys.* **30**, 507–516 (1958).
- [61] W. H. King, “Comments on the Article “Peculiarities of the Isotope Shift in the Samarium Spectrum,”,” *J. Opt. Soc. Am.* **53**, 638–639 (1963).
- [62] D N Stacey, “Isotope shifts and nuclear charge distributions,” *Reports on Progress in Physics* **29**, 171 (1966).
- [63] K. Heilig and A. Steudel, “Changes in mean-square nuclear charge radii from optical isotope shifts,” *Atomic Data and Nuclear Data Tables* **14**, 613–638 (1974), nuclear Charge and Moment Distributions.
- [64] CWP Palmer, “Reformulation of the theory of the mass shift,” *Journal of Physics B: Atomic and Molecular Physics* **20**, 5987 (1987).
- [65] S A Blundell, P E G Baird, C W P Palmer, D N Stacey, and G K Woodgate, “A reformulation of the theory of field isotope shift in atoms,” *Journal of Physics B: Atomic and Molecular Physics* **20**, 3663 (1987).
- [66] Meng Wang, W. J. Huang, F. G. Kondev, G. Audi, and S. Naimi, “The AME 2020 atomic mass evaluation (II). Tables, graphs and references,” *Chin. Phys. C* **45**, 030003 (2021).
- [67] A. Steudel J. E. Hansen and H. Walther, “Isotopieverschiebung der natürlichen geraden und ungeraden Sm- und Nd-Isotope,” *Z. Physik* **203**, 296–329 (1967).
- [68] C W P Palmer, P E G Baird, S A Blundell, J R Brandenberger, C J Foot, D N Stacey, and G K Woodgate, “Laser spectroscopy of calcium isotopes,” *Journal of Physics B: Atomic and Molecular Physics* **17**, 2197 (1984).
- [69] H. E. Haber, Gordon L. Kane, and T. Sterling, “The Fermion Mass Scale and Possible Effects of Higgs Bosons

- on Experimental Observables,” *Nucl. Phys. B* **161**, 493–532 (1979).
- [70] A R Striganov, V A Katulin, and V V Eliseev, “Peculiarities of the isotopic shift in the samarium spectrum,” *Optics and Spectroscopy (U.S.S.R.) (English Translation)* **2** (1962).
- [71] D.N. Stacey, “A note on the interpretation of isotope shifts,” *Physics Letters* **20**, 644–645 (1966).
- [72] Koki Ono, Yugo Saito, Taiki Ishiyama, Toshiya Higomoto, Tetsushi Takano, Yosuke Takasu, Yasuhiro Yamamoto, Minoru Tanaka, and Yoshiro Takahashi, “Observation of nonlinearity of generalized king plot in the search for new boson,” *Phys. Rev. X* **12**, 021033 (2022).
- [73] Akio Kawasaki, Takumi Kobayashi, Akiko Nishiyama, Takehiko Tanabe, and Masami Yasuda, “Isotope shift analysis with the  $4f^{14}6s^2\ ^1S_0 - 4f^{13}5d6s^2(J = 2)$  transition in ytterbium,” (2024), [arXiv:2402.13541 \[physics.atom-ph\]](https://arxiv.org/abs/2402.13541).
- [74] B. Ohayon, S. Hofsäss, J. E. Padilla-Castillo, S. C. Wright, G. Meijer, S. Truppe, K. Gibble, and B. K. Sahoo, “Isotope shifts in cadmium as a sensitive probe for physics beyond the standard model,” *New J. Phys.* **24**, 123040 (2022), [arXiv:2208.13599 \[physics.atom-ph\]](https://arxiv.org/abs/2208.13599).
- [75] Cédric Delaunay, Claudia Frugiuele, Elina Fuchs, and Yotam Soreq, “Probing new spin-independent interactions through precision spectroscopy in atoms with few electrons,” *Phys. Rev. D* **96**, 115002 (2017), [arXiv:1709.02817 \[hep-ph\]](https://arxiv.org/abs/1709.02817).
- [76] Matthew P. A. Jones, Robert M. Potvliege, and Michael Spannowsky, “Probing new physics using Rydberg states of atomic hydrogen,” *Phys. Rev. Res.* **2**, 013244 (2020), [arXiv:1909.09194 \[hep-ph\]](https://arxiv.org/abs/1909.09194).
- [77] Robert M. Potvliege, Adair Nicolson, Matthew P. A. Jones, and Michael Spannowsky, “Deuterium spectroscopy for enhanced bounds on physics beyond the standard model,” *Phys. Rev. A* **108**, 052825 (2023), [arXiv:2309.03732 \[hep-ph\]](https://arxiv.org/abs/2309.03732).
- [78] R. M. Potvliege, “Spectroscopy of light atoms and bounds on physics beyond the standard model,” (2024), [arXiv:2412.16315 \[physics.atom-ph\]](https://arxiv.org/abs/2412.16315).
- [79] P. T. Boggs and J. E. Rogers, “Orthogonal Distance Regression: Statistical analysis of measurement error models and applications: proceedings of the AMS-IMS-SIAM joint summer research conference held June 10–16, 1989,” *Contemporary Mathematics* **112**, 186 (1990).
- [80] Pauli Virtanen, Ralf Gommers, Travis E. Oliphant, Matt Haberland, Tyler Reddy, David Cournapeau, Evgeni Burovski, Pearu Peterson, Warren Weckesser, Jonathan Bright, Stéfan J. van der Walt, Matthew Brett, Joshua Wilson, K. Jarrod Millman, Nikolay Mayоров, Andrew R. J. Nelson, Eric Jones, Robert Kern, Eric Larson, C J Carey, İlhan Polat, Yu Feng, Eric W. Moore, Jake VanderPlas, Denis Laxalde, Josef Perktold, Robert Cimrman, Ian Henriksen, E. A. Quintero, Charles R. Harris, Anne M. Archibald, Antônio H. Ribeiro, Fabian Pedregosa, Paul van Mulbregt, and SciPy 1.0 Contributors, “SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python,” *Nature Methods* **17**, 261–272 (2020).
- [81] Mohsen Pourahmadi, “Cholesky Decompositions and Estimation of A Covariance Matrix: Orthogonality of Variance–Correlation Parameters,” *Biometrika* **94**, 1006–1013 (2007), <https://academic.oup.com/biomet/article-pdf/94/4/1006/681653/asm073.pdf>.
- [82] Henrik Flyvbjerg and H.G. Petersen, “Error estimates on averages of correlated data,” *The Journal of Chemical Physics* **91** (1989), 10.1063/1.457480.
- [83] A T Goble, J D Silver, and D N Stacey, “Isotope shifts in the atomic spectrum of tin:  $^{112}\text{Sn}$ ,  $^{114}\text{Sn}$  and  $^{115}\text{Sn}$ ,” *Journal of Physics B: Atomic and Molecular Physics* **7**, 26 (1974).
- [84] Jungwoo Choi, Eunhwi Lee, Dahyun Yum, Kyungwon An, and Junki Kim, “Direct measurement of isotope shifts in the barium  $6s^2\ ^1S_0 - 5d6p\ ^3D_1$  transition,” *Phys. Rev. A* **110**, 032812 (2024).
- [85] Simon Hofsäss, J. Eduardo Padilla-Castillo, Sid C. Wright, Sebastian Kray, Russell Thomas, Boris G. Sartakov, Ben Ohayon, Gerard Meijer, and Stefan Truppe, “High-resolution isotope-shift spectroscopy of Cd I,” *Phys. Rev. Res.* **5**, 013043 (2023).
- [86] Taiki Ishiyama, Koki Ono, Hokuto Kawase, Tetsushi Takano, Reiji Asano, Ayaki Sunaga, Yasuhiro Yamamoto, Minoru Tanaka, and Yoshiro Takahashi, “Orders-of-magnitude improved precision spectroscopy of an inner-shell orbital clock transition in neutral ytterbium,” (2025), [arXiv:2505.04154 \[physics.atom-ph\]](https://arxiv.org/abs/2505.04154).
- [87] Jize Han, Ying Zheng, Yanmei Yu, Jiguang Li, Zhiguo Huang, Jingwei Wen, Ling Qian, and Lijun Wang, “Enhanced accuracy of isotope-shift measurements of  $\text{Yb}^+$  369- and 935-nm transitions via sympathetic cooling,” *Phys. Rev. A* **111**, 033101 (2025).
- [88] N. A. Diepeveen, C. Robalo Pereira, M. Mazzanti, Z. E. D. Ackerman, L. P. H. Gallagher, T. Timmerman, R. Gerritsma, and R. X. Schüssler, “Single-ion spectroscopy of four metastable-state clear-out transitions in  $\text{Yb}^+$ : Isotope shifts and hyperfine structure,” *Phys. Rev. A* **110**, 042809 (2024).
- [89] Toshiaki Ando, Kana Yamada, Atsushi Iwasaki, and Kaoru Yamanouchi, “Isotope shift of fine structure of  $\text{Kr}^+$  and hyperfine structure of  $^{83}\text{Kr}^+$  by strong-field ultrahigh-resolution Fourier-transform spectroscopy,” *Phys. Rev. Res.* **7**, L022025 (2025).
- [90] G. Zitzer, J. Tiedau, Ch. E. Düllmann, M. V. Okhapkin, and E. Peik, “Laser spectroscopy on the hyperfine structure and isotope shift of sympathetically cooled  $^{229}\text{Th}^{3+}$  ions,” *Phys. Rev. A* **111**, L050802 (2025).
- [91] Chao Zhang, Ryohei Terabayashi, and Shuichi Hasegawa, “Investigation of isotope shifts and Stark shifts in the strontium Rydberg state via  $5s^2\ ^1S_0 \rightarrow 5s5p\ ^1P_1^0 \rightarrow 5p_{1/2}5p_{1/2} \rightarrow 4d_{3/2}nl_j (n^* = 39.4)$ ,” *Spectrochimica Acta Part B: Atomic Spectroscopy* **225**, 107118 (2025).
- [92] Michail Athanasakis-Kaklamanakis, Shane G. Wilkins, Alexander A. Breier, and Gerda Neyens, “King-Plot Analysis of Isotope Shifts in Simple Diatomic Molecules,” *Phys. Rev. X* **13**, 011015 (2023).
- [93] V. A. Dzuba, V. V. Flambaum, and M. G. Kozlov, “Combination of the many-body perturbation theory with the configuration-interaction method,” *Phys. Rev. A* **54**, 3948–3959 (1996).
- [94] M.G. Kozlov, S.G. Porsev, M.S. Safronova, and I.I. Tupitsyn, “CI-MBPT: A package of programs for relativistic atomic calculations based on a method combining configuration interaction and many-body perturbation theory,” *Computer Physics Communications* **195**, 199–213 (2015).
- [95] J. C. Berengut, “Particle-hole configuration interaction



- and many-body perturbation theory: Application to  $\text{Hg}^+$ ,” *Phys. Rev. A* **94**, 012502 (2016).
- [96] F. Torretti, A. Windberger, A. Ryabtsev, S. Dobrodey, H. Bekker, W. Ubachs, R. Hoekstra, E. V. Kahl, J. C. Berengut, J. R. Crespo López-Urrutia, and O. O. Versolato, “Optical spectroscopy of complex open- $4d$ -shell ions  $\text{Sn}^{7+} - \text{Sn}^{10+}$ ,” *Phys. Rev. A* **95**, 042503 (2017).
  - [97] Jack Sugar and Charles Corliss, “Atomic energy levels of the iron-period elements: potassium through nickel,” *Journal of Physical and Chemical Reference Data* **14**, Supplement No. 2 (1985).
  - [98] Felix W. Knollmann, Ashay N. Patel, and S. Charles Doret, “Part-per-billion measurement of the  $4^2S_{1/2} \rightarrow 3^2D_{5/2}$  electric-quadrupole-transition isotope shifts between  $^{42,44,48}\text{Ca}^+$  and  $^{40}\text{Ca}^+$ ,” *Phys. Rev. A* **100**, 022514 (2019).
  - [99] Felix W. Knollmann, Ashay N. Patel, and S. Charles Doret, “Erratum: Part-per-billion measurement of the  $4^2S_{1/2} \rightarrow 3^2D_{5/2}$  electric-quadrupole-transition isotope shifts between  $^{42,44,48}\text{Ca}^+$  and  $^{40}\text{Ca}^+$  [Phys. Rev. A 100, 022514 (2019)],” *Phys. Rev. A* **107**, 069902 (2023).
  - [100] A. Kramida, Yu. Ralchenko, and J. et al. Reader, “Nist atomic spectra database (version 5.11), [online],” (2023), <https://doi.org/10.18434/T4W30F>.
  - [101] FG Kondev and S Naimi, “The AME2016 atomic mass evaluation (I). Evaluation of input data; and adjustment procedures,” *Chinese physics C* **41**, 030002 (2017).
  - [102] Timothy T. Chang, Bless Bah Awazi, Julian C. Berengut, Elina Fuchs, and S. Charles Doret, “Systematic-free limit on new light scalar bosons via isotope shift spectroscopy in  $\text{Ca}^+$ ,” (2024), [arXiv:2311.17337 \[physics.atom-ph\]](https://arxiv.org/abs/2311.17337).
  - [103] and, F. G. Kondev, , and S. and Naimi, “The AME2016 atomic mass evaluation (II). Tables, graphs and references,” *Chinese Physics C* **41**, 030003 (2017).
  - [104] Patrick Müller, Kristian König, Phillip Ingram, Jörg Krämer, and Wilfried Nörtershäuser, “Collinear laser spectroscopy of  $\text{Ca}^+$ : Solving the field-shift puzzle of the  $4s^2S_{1/2} \rightarrow 4p^2P_{1/2,3/2}$  transitions,” *Phys. Rev. Res.* **2**, 043351 (2020).
  - [105] W Alt, M Block, V Schmidt, T Nakamura, P Seibert, X Chu, and G Werth, “Shifts of the  $3d - 4p$  transitions in different isotopes of positive calcium ions,” *Journal of Physics B: Atomic, Molecular and Optical Physics* **30**, L677 (1997).
  - [106] Ronald Fernando Garcia Ruiz, ML Bissell, Klaus Blaum, A Ekström, N Frömmgen, G Hagen, M Hammen, K Hebeler, JD Holt, GR Jansen, *et al.*, “Unexpectedly large charge radii of neutron-rich calcium isotopes,” *Nature Physics* **12**, 594–598 (2016), [arXiv:1602.07906 \[nucl-ex\]](https://arxiv.org/abs/1602.07906).
  - [107] Taiki Ishiyama, Koki Ono, Tetsushi Takano, Ayaki Sunaga, and Yoshiro Takahashi, “Observation of an inner-shell orbital clock transition in neutral ytterbium atoms,” *Phys. Rev. Lett.* **130**, 153402 (2023).
  - [108] Ann-Marie Mårtensson-Pendrill, David S. Gough, and Peter Hannaford, “Isotope shifts and hyperfine structure in the 369.4-nm  $6s-6p_{1/2}$  resonance line of singly ionized ytterbium,” *Phys. Rev. A* **49**, 3351–3365 (1994).
  - [109] R. W. Berends and L. Maleki, “Hyperfine structure and isotope shifts of transitions in neutral and singly ionized ytterbium,” *J. Opt. Soc. Am. B* **9**, 332–338 (1992).

## Appendix A: Calculation of Electronic Coefficients in $\text{Ca}^+$

In this section we present some details of the calculation of the electronic new physics coefficients in  $\text{Ca}^+$ <sup>12</sup>. Similar to Ref. [29], we applied the finite field method. In this approach, the new physics potential

$$V_{\text{NP}}(r, m_\phi) = \lambda \frac{1}{4\pi r} e^{-m_\phi r} \quad (\text{A1})$$

is directly added to the Hamiltonian in the Dirac equation, including a dimensionless scaling parameter  $\lambda$ . Then, the eigenenergies of the considered states are calculated by solving the Dirac equation for different values of the parameter  $\lambda$ . Finally, the  $X$  coefficient can be extracted by taking the numerical derivative of the eigenenergies with respect to  $\lambda$ :

$$X_i = \left. \frac{\partial E_i(\lambda)}{\partial \lambda} \right|_{\lambda=0}. \quad (\text{A2})$$

In a similar manner, the FS coefficients  $F_i$  are evaluated by calculating the energies for different nuclear radii and taking the numerical derivative of the eigen-energies with respect to  $\delta\langle r^2 \rangle$ :

$$F_i = \left. \frac{\partial E_i(\lambda)}{\partial \delta\langle r^2 \rangle} \right|_{\delta\langle r^2 \rangle=0}. \quad (\text{A3})$$

To obtain the eigenenergies, electronic structure calculations are performed using the AMBiT code, which is based on a combination of configuration interaction (CI) and many-body perturbation theory (MBPT) (see e.g. Refs. [93–96]) and is thoroughly described in Ref. [59]. The specifics of the calculations for  $\text{Ca}^+$  are overviewed in the appendices of Ref. [43].

The results for the  $X$  and  $F$  coefficients are summarised in Tab. IV and Fig. 15 and are in good agreement with reference values from [52].

As discussed in the main text in the high-mass regime the ratio of the  $X$  coefficients approaches the ratio of field shift coefficients, leading to the loss of sensitivity to new physics. Since even a small discrepancy between the experimental and theoretical values of the field shift ratio  $F_j/F_i$  can significantly impact the high-mass behavior of the bounds on new physics, the  $X$  coefficients are rescaled so that in the high-mass limit the ratio  $X_j/X_i$  approaches the experimental field shift ratio  $F_j^{\text{exp}}/F_i^{\text{exp}}$ , which is obtained from a linear fit to King plot measurements.

Note that the conventions for the new physics coefficients (referred to as  $D$  coefficients e.g. in Refs. [48, 49])

<sup>12</sup> The  $X$  coefficients for Yb used in this paper were calculated in Ref. [47–49] and obtained from J. Berengut via personal communication.

vary in the literature. For instance, one can include all prefactors of the Yukawa potential of Eq. (A1) in the calculation of the new physics coefficients. As a result, when utilising these coefficients to derive constraints on new physics, the obtained bounds are expressed in terms of  $y_e y_n$ . However, in other conventions a factor of  $1/(4\pi\alpha_{\text{EM}})$  is factored out from the  $X$  coefficient, resulting in bounds being expressed in terms of  $y_e y_n/(4\pi\alpha_{\text{EM}}) = \frac{\alpha_{\text{NP}}}{\alpha_{\text{EM}}}$ . In this work, the  $X$  coefficients are always presented in the latter convention.




FIG. 15. Dependence of the electronic new physics coefficients of the  $\nu_{729}$  and  $\nu_{397}$  transitions in  $\text{Ca}^+$  on the mediator mass  $m_\phi$ . The coefficient of the  $\nu_{732}$  transition is nearly identical to that of  $\nu_{729}$  and is therefore not displayed in the figure.

## Appendix B: Sensitivity to New Physics

To gain an understanding of how the choice of transitions can affect the bounds on the new physics coupling, we compare bounds obtained by applying Eqs. (10) and (13) to subsets of isotope shift measurements in  $\text{Ca}^+$ . We fix  $\nu_{729}$ :  $3p^6 4s^2 S_{1/2} \rightarrow 3p^6 3d^2 D_{5/2}$  as the reference transition  $\nu_1$  and combine it with either of the following transitions:

- $\nu_{397}$ :  $3p^6 4s^2 S_{1/2} \rightarrow 3p^6 4p^2 P_{1/2}$
- $\nu_{732}$ :  $3p^6 4s^2 S_{1/2} \rightarrow 3p^6 3d^2 D_{3/2}$
- $\nu_{\text{DD}}$ :  $3p^6 3d^2 D_{3/2} \rightarrow 3p^6 3d^2 D_{5/2}$

where the subscripts correspond to the transition wavelength in nm, while DD is associated to fine-structure splitting. A comprehensive list of available isotope shift data in Ca can be found in Tab. IX.




FIG. 16. Sensitivity of different transition pairs to new physics. The reference transition is fixed to  $\nu_1 = \nu_{729}$ , the second transitions are as specified in the legend. The inverse of the electronic dependence of  $V_{\text{pred}}$  (see Eq. (12)) as a function of the mass  $m_\phi$  of the new boson, suggesting the combination  $(\nu_{729}, \nu_{397})$  is more sensitive than the other two, which are equally sensitive.

### Identifying King Plot Candidates

The shape of  $V_{\text{pred}}$  can hint at the best transitions for new physics searches: a combination of electronic coefficients that maximises this volume suggests a high sensitivity of the associated transitions to  $\alpha_{\text{NP}}$ . In Tab. IV we report our calculations of the electronic coefficients of the transitions listed above and in Fig. 16 we plot the inverse of the electronic dependence of  $V_{\text{pred}}$ :

$$\frac{1}{V_{\text{pred}}} \propto \frac{1}{\det(\mathcal{M}(m_\phi))} = \frac{1}{F_1 X_2(m_\phi) - F_2 X_1(m_\phi)}, \quad (\text{B1})$$

which determines the  $m_\phi$ -dependence of the bounds on  $\alpha_{\text{NP}}$  (see Eqs. (10) and (12)). The  $m_\phi$ -independent low-mass behaviour, as well as the loss of sensitivity at high mediator masses  $m_\phi$ , are clearly visible. Moreover, we observe peaks, which are the consequence of accidental cancellations of  $\det(\mathcal{M}(m_\phi))$  at specific values of  $m_\phi$ .

We identify the combination of  $\nu_{729}$  and  $\nu_{397}$  as the pair of transitions with the largest value for  $\det(\mathcal{M}(m_\phi))$ , suggesting a high sensitivity to new physics. This can be understood as a consequence of the complementarity of the electronic levels involved in these transitions. In contrast, the combinations of  $\nu_{729}$  with  $\nu_{732}$  or  $\nu_{\text{DD}}$  yield a smaller value for the electronic part of  $V_{\text{pred}}$ . Indeed, the similarity of the  $D_{3/2}$  and  $D_{5/2}$  states results in similar electronic coefficients leading to a higher degree of cancellation in  $F_1 X_2(m_\phi) - F_2 X_1(m_\phi)$ .

### The Perfectly Linear King Plot

Next, we consider the idealised case of a perfectly linear King plot ( $V_{\text{dat}} = 0$ ). To this end, we generate perfectly linear mock values for the isotope shifts, i.e. we use Eq. (1) to calculate linear isotope shift data based on experimental values for the nuclear radii [25] and masses [66], as well as on theory calculations of the SM electronic coefficients (reported in Tab. IV).

Clearly, in this case  $\langle\alpha_{\text{NP}}\rangle = 0$  and the bounds are fully determined by  $\sigma[\alpha_{\text{NP}}]$ . We can gain some insight into the behaviour of  $\sigma[\alpha_{\text{NP}}]$  by performing linear error propagation on Eq. (10):

$$\sigma[\alpha_{\text{NP}}] = \sqrt{\left(\frac{\sigma[V_{\text{dat}}]}{V_{\text{pred}}}\right)^2 + \left(\frac{V_{\text{dat}}}{V_{\text{pred}}^2}\sigma[V_{\text{pred}}]\right)^2}. \quad (\text{B2})$$

with  $V_{\text{dat}}$  and  $V_{\text{pred}}$  as defined in Eq. (10). In linear King plots, only the first term is relevant and  $\sigma[\alpha_{\text{NP}}]$  is dominated by experimental uncertainties. What determines the sensitivity to new physics is thus the interplay of  $V_{\text{pred}}$  and  $\sigma[V_{\text{dat}}]$ . Since the latter includes terms of the form  $\tilde{\nu}_i^a \sigma[\tilde{\nu}_j^b]$ , more constraining bounds can be expected from comparatively small but precisely measured isotope shifts. We assign realistic uncertainties to our mock transitions (20 Hz for  $\nu_{729}$ ,  $\nu_{\text{DD}}$  and  $\nu_{732}$  [41], 80 kHz for  $\nu_{397}$  [26]) and compare the projections for the  $2\sigma$  upper bounds in Fig. 17. Here, we limit the analysis to  $|\frac{\alpha_{\text{NP}}}{\alpha_{\text{EM}}}|$  for simplicity. Although the combination involving  $\nu_{397}$  appeared to be the most promising based on the behaviour of its electronic coefficients, the higher experimental uncertainty limits its sensitivity to new physics, relative to the other combinations.

### The Linear King Plot

Having identified the combination of the  $\nu_{729}$  and  $\nu_{\text{DD}}$  transitions to generate the most stringent bounds to new physics, we use real data from Refs. [41, 98, 99], which produce a linear King plot within the experimental uncertainties ( $\sigma[V_{\text{dat}}]/V_{\text{dat}} \lesssim 1$ ). In Fig. 18, we compare the resulting bounds with those from perfectly linear mock data.

Since real data predict a non-zero central value for the new physics coupling, the bounds are shifted away from zero. This effect is most pronounced in the low-mass limit, where the King plot method is most sensitive to new physics. Moreover, the second term in Eq. (B2) is now non-zero. This has a noticeable impact if the uncertainties on the electronic new physics coefficients  $X$ , which we take to be 10% of their computed value (see Appendix A), are introduced: Since this term scales with  $1/V_{\text{pred}}^2$  and  $V_{\text{pred}}$  becomes small for large values of  $m_\phi$ , uncertainties on the  $X$  coefficients lead to a loss of sensitivity in the high mass region. This highlights the




FIG. 17. Predicted  $2\sigma$  upper bounds on  $|\alpha_{\text{NP}}/\alpha_{\text{EM}}|$  for linear mock data with uncertainties as reported in the legend. Although the combination involving  $\nu_{397}$  appears to be the most promising based on the behaviour of  $|1/\det(\mathcal{M})|$ , the bounds on  $|\alpha_{\text{NP}}/\alpha_{\text{EM}}|$  obtained for the other two combinations turn out to be more stringent due to smaller experimental uncertainties.  $\nu_{\text{DD}}$  wins against  $\nu_{732}$  thanks to the smaller isotope shift.




FIG. 18. The bounds from the combination of  $\nu_{\text{DD}}$  and  $\nu_{729}$ . The black curve shows the results for perfectly linear mock data while the orange dashed and teal dotted curves represent bounds derived from the experimental data reported in Refs. [41, 98], with and without including uncertainties on the electronic new physics coefficient  $X_i$ .

importance of considering the uncertainty of  $X$  and suggests that more precise  $X$  coefficient calculations could improve the bounds, particularly in the high-mass range.

Transition ( $i$ )	$\Delta E$ [eV]	$F_i$ [eV/fm <sup>2</sup> ]	$K_i$ [eV·u]	$X_i(m_\phi = 1 \text{ keV})$ [eV]	$X_i(m_\phi = 10^4 \text{ keV})$ [eV]
$\nu_{397}: 3p^6 4s^2 S_{1/2} \rightarrow 3p^6 4p^2 P_{1/2}$	3.16 3.12 [97]	$-1.19 \times 10^{-6}$ $-1.18 \times 10^{-6}$ [52]	$-1.59 \times 10^{-3}$ [52]	1.11	$1.29 \times 10^{-4}$
$\nu_{732}: 3p^6 4s^2 S_{1/2} \rightarrow 3p^6 3d^2 D_{3/2}$	1.677 1.692 [97]	$-1.559 \times 10^{-6}$ $-1.567 \times 10^{-6}$ [52]	$-1.017 \times 10^{-2}$ [52]	-2.95	$1.71 \times 10^{-4}$
$\nu_{729}: 3p^6 4s^2 S_{1/2} \rightarrow 3p^6 3d^2 D_{5/2}$	1.685 1.700 [97]	$-1.557 \times 10^{-6}$ $-1.565 \times 10^{-6}$ [52]	$-1.012 \times 10^{-2}$ [52]	-2.93	$1.70 \times 10^{-4}$

TABLE IV. Transition energies ( $\Delta E$ ) and electronic field shift ( $F_i$ ), mass shift ( $K_i$ ) and new physics ( $X_i$ ) coefficients for three transitions in  $\text{Ca}^+$ . Values of  $\Delta E$ ,  $F_i$  and  $X_i$  were calculated using the CI + MBPT method implemented in **AMBiT** [59] (see also Appendix A). The latter are reported for two different values of the mediator mass  $m_\phi$ . The results are in good agreement with the transition energies given in Ref. [97] and the electronic coefficients of Ref. [52]. The values for  $K_i$  are taken from Ref. [52]. Values for  $\nu_{\text{DSIS}}$  can be obtained from the difference of  $\nu_{729}$  and  $\nu_{732}$ . In Ref. [52], the uncertainties on  $F_i$  and  $K_i$  are estimated to be 5% and 10%, respectively. For  $X_i$ , we estimate the uncertainty to be about 10%.

### Appendix C: Projection of Uncertainties

To better understand how the uncertainties on the measured isotope shifts and atomic masses propagate to the bounds on the new physics coupling, we can project the experimental uncertainties onto the directions parallel and perpendicular to the King line and evaluate their impact on the uncertainty on  $\alpha_{\text{NP}}$ :

$$\begin{aligned}\Sigma_{\parallel}^{\alpha} &= (\nabla_{\tilde{\nu}} \alpha_{\text{NP}})^{\top} \mathbf{P}_{\parallel} \Sigma_{\tilde{\nu}} \mathbf{P}_{\parallel} \nabla_{\tilde{\nu}} \alpha_{\text{NP}} \equiv \left( \sigma[\alpha_{\text{NP}}]_{\text{KP}, \parallel}^{(1)} \right)^2 \\ \Sigma_{\perp}^{\alpha} &= (\nabla_{\tilde{\nu}} \alpha_{\text{NP}})^{\top} \mathbf{P}_{\perp} \Sigma_{\tilde{\nu}} \mathbf{P}_{\perp} \nabla_{\tilde{\nu}} \alpha_{\text{NP}} \equiv \left( \sigma[\alpha_{\text{NP}}]_{\text{KP}, \perp}^{(1)} \right)^2 \\ \Sigma_{\parallel\perp}^{\alpha} &= (\nabla_{\tilde{\nu}} \alpha_{\text{NP}})^{\top} \mathbf{P}_{\parallel} \Sigma_{\tilde{\nu}} \mathbf{P}_{\perp} \nabla_{\tilde{\nu}} \alpha_{\text{NP}} = (\Sigma_{\perp\parallel}^{\alpha})^{\top}, \quad (\text{C1})\end{aligned}$$

Here the superscript (1) indicates that we are applying linear error propagation, whereas

$$\nabla_{\tilde{\nu}} \alpha_{\text{NP}} = \begin{pmatrix} \frac{\partial \alpha_{\text{NP}}}{\partial \tilde{\nu}_1^1} & \cdots & \frac{\partial \alpha_{\text{NP}}}{\partial \tilde{\nu}_1^n} \\ \vdots & \ddots & \vdots \\ \frac{\partial \alpha_{\text{NP}}}{\partial \tilde{\nu}_1^n} & \cdots & \frac{\partial \alpha_{\text{NP}}}{\partial \tilde{\nu}_m^n} \end{pmatrix} \quad (\text{C2})$$

corresponds to the gradient of  $\alpha_{\text{NP}}$  with respect to the  $n \times m$  mass-normalised isotope shifts  $\{\tilde{\nu}_i^a\}_{1 \leq a \leq n}^{1 \leq i \leq m}$  (points in the King plot), and

$$\Sigma_{\tilde{\nu}} = \begin{pmatrix} \Sigma_{\tilde{\nu}}^{11} & \cdots & \Sigma_{\tilde{\nu}}^{1n} \\ \vdots & \ddots & \vdots \\ \Sigma_{\tilde{\nu}}^{n1} & \cdots & \Sigma_{\tilde{\nu}}^{nn} \end{pmatrix} \quad (\text{C3})$$

denotes the  $(m \times m \times n \times n)$ -dimensional covariance matrix of the set of  $n$  mass-normalised isotope shift vectors  $\{\tilde{\nu}^a\}_{a=1}^n$ . The entries of  $\Sigma_{\tilde{\nu}}$  are  $(m \times m)$ -dimensional covariance matrices associated to  $\tilde{\nu}^a = (\tilde{\nu}_1^a, \dots, \tilde{\nu}_m^a)$  and  $\tilde{\nu}^b = (\tilde{\nu}_1^b, \dots, \tilde{\nu}_m^b)$ :

$$\Sigma_{\tilde{\nu}}^{ab} = \begin{pmatrix} \text{Cov}(\tilde{\nu}_1^a, \tilde{\nu}_1^b) & \cdots & \text{Cov}(\tilde{\nu}_1^a, \tilde{\nu}_m^b) \\ \vdots & \ddots & \vdots \\ \text{Cov}(\tilde{\nu}_m^a, \tilde{\nu}_m^b) & \cdots & \text{Cov}(\tilde{\nu}_m^a, \tilde{\nu}_m^b) \end{pmatrix}, \quad (\text{C4})$$

where

$$\begin{aligned}\text{Cov}(\tilde{\nu}_i^a, \tilde{\nu}_j^b) &= \sum_{k=1}^m \sum_{c=1}^n \left( \frac{\partial \tilde{\nu}_i^a}{\partial \nu_k^c} \sigma[\nu_k^c]^2 \frac{\partial \tilde{\nu}_j^b}{\partial \nu_k^c} \right. \\ &\quad \left. + \frac{\partial \tilde{\nu}_i^a}{\partial m^c} \sigma[m^c]^2 \frac{\partial \tilde{\nu}_j^b}{\partial m^c} + \frac{\partial \tilde{\nu}_i^a}{\partial m^{c'}} \sigma[m^{c'}]^2 \frac{\partial \tilde{\nu}_j^b}{\partial m^{c'}} \right), \quad (\text{C5})\end{aligned}$$

assuming the isotope shift and mass measurements are independent. The objects  $\mathbf{P}_{\parallel}$  and  $\mathbf{P}_{\perp}$  in Eq. (C1) are  $n \times n \times m \times m$ -dimensional projectors onto the parallel and perpendicular directions,

$$\mathbf{P}_{\parallel} = \mathbb{I}_n \otimes \mathbf{P}_{\parallel} = \text{diag}(\mathbf{P}_{\parallel}, \dots, \mathbf{P}_{\parallel})_n \quad (\text{C6})$$

$$\mathbf{P}_{\perp} = \mathbb{I}_n \otimes \mathbf{P}_{\perp} = \text{diag}(\mathbf{P}_{\perp}, \dots, \mathbf{P}_{\perp})_n. \quad (\text{C7})$$

Here  $\otimes$  denotes the tensor product and  $\mathbf{P}_{\parallel}$  and  $\mathbf{P}_{\perp}$  are the  $m \times m$ -dimensional projectors

$$\mathbf{P}_{\parallel} = \hat{\mathbf{e}}_{\mathcal{F}} \hat{\mathbf{e}}_{\mathcal{F}}^{\top}, \quad \mathbf{P}_{\perp} = \mathbb{I}_m - \hat{\mathbf{e}}_{\mathcal{F}} \hat{\mathbf{e}}_{\mathcal{F}}^{\top}, \quad (\text{C8})$$

$$\mathbf{P}_{\parallel}^2 = \mathbf{P}_{\parallel}, \quad \mathbf{P}_{\parallel}^{\top} = \mathbf{P}_{\parallel}, \quad \mathbf{P}_{\perp}^2 = \mathbf{P}_{\perp}, \quad \mathbf{P}_{\perp}^{\top} = \mathbf{P}_{\perp}, \quad (\text{C9})$$

where  $\hat{\mathbf{e}}_{\mathcal{F}}$  is the unit vector along the King line (see Eq. (38)) and  $\mathbb{1}_m$  is the  $(m \times m)$ -dimensional identity matrix.

Finally, the impact of the uncertainties perpendicular to the King line on the estimation of  $\sigma[\alpha_{\text{NP}}]$  can be evaluated by computing

$$\sqrt{\frac{\Sigma_{\perp}^{\alpha}}{\Sigma^{\alpha}}} \equiv \frac{\sigma[\alpha_{\text{NP}}]_{\text{KP},\perp}^{(1)}}{\sigma[\alpha_{\text{NP}}]_{\text{KP}}^{(1)}}, \quad (\text{C10})$$

where  $\sigma[\alpha_{\text{NP}}]_{\text{KP}}^{(1)}$  is the full uncertainty without projectors:

$$\sigma[\alpha_{\text{NP}}]_{\text{KP}}^{(1)} \equiv \sqrt{\Sigma_{\alpha}} \equiv \sqrt{(\nabla_{\hat{\nu}} \alpha_{\text{NP}})^{\top} \Sigma_{\hat{\nu}} \nabla_{\hat{\nu}} \alpha_{\text{NP}}}. \quad (\text{C11})$$

For all elements in Table III we checked that  $\sigma[\alpha_{\text{NP}}]_{\text{KP},\perp}^{(1)} / \sigma[\alpha_{\text{NP}}]_{\text{KP}}^{(1)} \approx 1$  within numerical uncertainties.

## Appendix D: The kifit Package

**kifit** is a prototype code and is published strictly for reasons of transparency and reproducibility. In this section, we briefly explain the structure of the package and give an introductory example of how to use **kifit**. We also summarise the **pytest**s implemented in the code.

### 1. Structure of the kifit Package

The source code of **kifit** can be divided into three main modules: **build**, **tools** and **run**, which are represented with yellow boxes in the diagram of Fig. 19.

The operative module is **run**, where a **Runner** class makes use of **build** and **tools** to process data and perform either the algebraic methods and the King plot fit.




FIG. 19. Schematic structure of kifit 0.1.0.

To launch the **kifit** scripts, a **kifit\_data** folder is required, in which experimental data have to be provided in the same format as in the example which can be found

in the code repository and as described below. The execution of the code can be customised by defining a set of initial parameters in an appropriate configuration file (see **README.md** file of our repository for more detailed information).

The module **build** constructs one instance of the class **kifit.Elem** for each experimental data set collected in a **kifit.ElemCollection**. The module **tools** contains two sets of functions which compute the bounds on  $\alpha_{\text{NP}}$ , either using the fit (**fitools**) or the algebraic methods (**detools**). Finally, the script **plot** is a compilation of plot scripts that can be used to visualise the fit and determinant results.

### 2. How to Use kifit

#### Data Preparation

First, the input data needs to be organised in subfolders of the **kifit\_data** subfolder and with names corresponding to the element identification (e.g. **elem**):

```

1 kifit/
2 |-- src/
3 |   |-- kifit/
4 |   |   |-- ...
5 |   |   |-- user_elements.py
6 |   |-- kifit_data/
7 |   |   |-- elem/
8 |   |   |   |-- binding_energies_elem.dat
9 |   |   |   |-- isotopes_elem.dat
10 |   |   |   |-- nu_elem.dat
11 |   |   |   |-- sig_nu_elem.dat
12 |   |   |   |-- sig_Xcoeffs_elem.dat
13 |   |   |   |-- Xcoeffs_elem.dat
14 |   |-- tests/

```

The **kifit** datafiles inside the **elem** subfolder are to be named as follows:

**type\_elem\_ISinstitutions\_massinstitution\_year.dat**

where **type** refers to the type of data saved in the file (for example data files, see the code repository):

- **binding\_energies**: electron binding energies, extracted from NIST [100]
- **isotopes**: isotope numbers  $A, A'$ , masses  $(m^A, m^{A'})$ , mass uncertainties  $(\sigma[m^A], \sigma[m^{A'}])$ .
- **nu**: isotope shift measurements  $(\nu_i^a)$ ; (columns: transitions, rows: isotope pairs).
- **sig nu**: uncertainties on the isotope shift measurements  $(\sigma[\nu_i^a])$ .
- **Xcoeffs** and **sig\_Xcoeffs**: electronic coefficients  $(X_i)$  and their uncertainties  $(\sigma[X_i])$  for the transitions listed in **nu** datafiles, for different values of



the new boson masses  $m_\phi$  (each line corresponding to a separate value of  $m_\phi$ ). In the current version of kikit, the `sigXcoeffs` files are not employed.

`ISinstitutions` and `massinstitution` list the institutions where the isotope shift measurements contained in `nu` and the isotope masses measurements contained in `isotopes` datafiles were respectively carried out, while `year` reports the year of the most recent isotope shift measurements collected in the folder. If no `massinstitution` is specified, isotope masses measurement are taken from either AME2020 [101] or AME2016 [66], as reported in the heading of the `isotopes` datafiles. Similarly, the heading of `ISinstitutions` datafiles reports, in shorthand notation, the publications from which we extracted the isotope shift measurements, as well as for the electronic transition. Table V provides a list of the data folders available within kikit, the shorthand notation adopted to refer to the institutions and the “dimension” of the dataset, i.e. number of transitions ( $m$ ) and isotope pairs ( $n$ ).

In order for the element `elem` to become a “valid” kikit element, its identification (e.g. `elem`) needs to be added to the list of elements collected in `kikit/src/user.elements.py`.

#### Setting up the Simulation Parameters

The kikit run can be customised through various hyperparameters. These can be parsed by kikit directly via the command line, or provided in the form of a JSON file. An example kikit configuration could be the following:

```
1 # config.json
2 {
3     # Here we list the elements whose data is to
4     # be combined:
5     "element_list": ["elem", "extra_elem"],
6     # When more than one element is listed above
7     # , use globalgrid:
8     "search_mode": "globalgrid",
9     # Experiments and block sizes for blocking
10    "num_exp": 100,
11    "block_size": 20,
12    # Sampled points in the MC
13    "num_alphasamples_search": 500,
14    "num_elemsamples_per_alphasample_search":
15    500,
16    "num_elemsamples_exp": 500,
17    "num_alphasamples_exp": 500,
18    "min_percentile": 2,
19    # The target mass indices
20    "x0_fit": [0],
21    "x0_det": [0],
22    "gkp_dims": [3],
23    # Algebraic methods' params
24    "nmgkp_dims": [3],
25    "proj_dims": [3],
26    "num_det_samples": 5000,
27    "num_sigmas": 2,
28 }
```

More parameters can be added to the configuration file, and we recommend referring to the official documentation for a more detailed explanation.

#### Running the Algorithm

Once the input files are organised and the kikit configuration is defined, we can write a python script to compute the estimation of the new physics bounds. As shown in Fig. 19, we use a `Runner` object (`Runner.config`) to share information among the kikit tools.

```
1 from kikit.run import Runner
2 from kikit.config import RunParams
3
4 # Set the kikit parameters given in the
5 # configuration file
6 # (Leave the arguments empty if parsing from
7 # command line)
8 params = RunParams(configuration_file="config.
9 json")
10
11 # initialize the runner
12 runner = Runner(params)
13
14 # Run
15 runner.run()
16
17 # If needed, generate plots
18 runner.generate_all_alphaNP_ll_plots()
```

#### Collecting results

The output of kikit is stored in the `kikit/results` folder, which can contain two sub-folders: `output_data`, containing the numerical results in the form of JSON files, and `plots`, containing the final plots plus the real-time plots, if requested by the user. Once a kikit run is completed and the numerical results are collected, the user can exploit the same configuration file as above to process data and generate more plots, using the functions provided in `plot.py`. (Simply replace `runner.run()` with a command such as `runner.generate_all_alphaNP_ll_plots()`.)

### 3. Validation of the Algorithm

We performed a series of validation simulations to highlight the robustness to the procedure sketched above, and to gain a better understanding of the optimal hyperparameters required to run a kikit experiment. Table VI lists the benchmark values, while the bar plots in Fig. 20 illustrate the stability of the code as we vary one of the hyperparameters after the other.

The sensitivity of the code to the choice of the reference transition was investigated explicitly. Although the

kifit folder	Isotope shifts	Masses	$n \times m$
Ca_PTB_2015	Gebert2015 [26]	[101]	$3 \times 2$
Ca_WT_Aarhus_2020	Knollmann2019[98]+Solaro2020 [41]	[101]	$4 \times 2$
Ca_WT_Aarhus_2024	Solaro2020 [41], Knollmann2023 [99], Chang2024 [102]	[101]	$4 \times 2$
Ca_WT_Aarhus_PTB_2020	Knollmann2019 [98], Solaro2020 [41], Gebert2015 [26]	[101]	$3 \times 4$
Ca_WT_Aarhus_PTB_2024	Knollmann2023 [99], Chang2024 [102], Gebert2015 [26]	[101]	$3 \times 4$
Ca24min	Knollmann2023 [99], Chang2024 [102]	[66]	$3 \times 2$
Camin	Knollmann2019 [98], Solaro2020 [41]	[66]	$3 \times 2$
Ca_testdata	Knollmann2019 [98], Solaro2020 [41], Gebert2015 [26]	[66]	$3 \times 4$
Ca3pointsTEST	* Knollmann2023 [99], Chang2024 [102]	[66]	$3 \times 2$
Ca4pointsTEST	* Knollmann2023 [99], Chang2024 [102]	[66]	$4 \times 2$
Ca10pointsTEST	mock data	[66]	$10 \times 2$
Yb_Kyoto_MIT_GSI_2022	Ono2022 [72], Counts2020 [48], Hur2022 [49], Figueroa2022 [47]	[66]	$4 \times 5$
Yb_Kyoto_MIT_GSI_PTB_2024	Ono2022 [72], Door2024 [50], Counts2020 [48], Figueroa2022 [47]	[66]	$4 \times 5$
Yb_Kyoto_MIT_GSI_PTB_MPIK_2024	Ono2022 [72], Door2024 [50], Counts2020 [48], Figueroa2022 [47]	[50]	$4 \times 5$
Yb_PTB_MPIK_2024	Door2024 [50]	[50]	$4 \times 2$

TABLE V. `kifit` data folders with the corresponding references (2<sup>nd</sup> & 3<sup>rd</sup> columns) and dimensions (number of transitions  $m$  and isotope pairs  $n$ , 4<sup>th</sup> column). The datafiles marked with an asterisk use artificially modified uncertainties for the isotope shifts and were used as test datafiles. The mock data for `Ca10pointsTEST` was generated using Eq. (1), the electronic coefficients from Tab. IV and the nuclear masses and charge radii from Refs. [25, 66], plus small artificial King nonlinearities.

Hyperparameter	Value
<code>num_elemsaples_per_alphasample_search</code>	500
<code>num_exp</code>	200
<code>block_size</code>	20
<code>min_percentile</code>	5
<code>num_alphasamples_exp</code>	500
<code>num_elemsamples_exp</code>	500

TABLE VI. Default values of the `kifit` hyperparameters set in our validation test. In Fig. 20 these values are varied one by one to investigate their impact on the `kifit` estimation.

reference transition seems to play a special role in the construction of the King plot fit (see e.g. Eqs. (37), (41) or Fig. 4), the relations between the reference transition and the other transitions are symmetric and

$$|d_{ji}^a| = |d_{ij}^a|, \quad (\text{D1})$$

where we denoted  $d_j^a \equiv d_{ji}^a$  (see Eq. (48)), i.e. we explicitly mentioned the reference transition  $i$ .

This property is particularly easy to observe in the 2-

dimensional case:

$$\begin{aligned}
|d_{21}^a| &= |\delta_{21}^a \cos \phi_{21}| \\
&= \left| \frac{\alpha_{\text{NP}}}{\alpha_{\text{EM}}} (X_2 - \tan \phi_{21} X_1) (\tilde{\gamma}^a - \langle \tilde{\gamma} \rangle^2) \cos \phi_{21} \right| \\
&\stackrel{(*)}{=} \left| \frac{\alpha_{\text{NP}}}{\alpha_{\text{EM}}} (\tan \phi_{12} X_2 - X_1) (\tilde{\gamma}^a - \langle \tilde{\gamma} \rangle^1) \cos \phi_{12} \right| \\
&= |\delta_{12}^a \cos \phi_{12}| = |d_{12}^a|, \quad (\text{D2})
\end{aligned}$$

where in (\*) we assumed that  $\langle \tilde{\gamma} \rangle^2 = \langle \tilde{\gamma} \rangle^1$ , which is the case if we employ the definition given in Eq. (47).

The symmetry of the log-likelihood construction can be observed in Fig. 21, which shows the outputs of one of our numerical tests of `kifit` (can be found in the `kifit/src/tests` folder). Both were produced using the same input data (central values and uncertainties of the mass and frequency measurements) and  $\alpha_{\text{NP}}$  samples, but the upper plot in Fig. 21 shows the  $\Delta\chi^2$  values computed using the manifestly symmetric version of  $\sigma[\delta_{ii}^a]$  (Eq. (47)), while the definition given in Eq. (44) was used in the making of the lower plot in Fig. 21. The different colours represent different sets of samples of the input parameters (transition frequencies and masses, see Eq. (14)) and fit parameters  $K_{21}^\perp$  and  $\phi_{21}$  (see Eq. (55)): The label “Ca  $ji$ , Ca  $lk$  samples”, means that the element Ca with the set of transitions  $k, l$ , where  $k$  is the reference transition, was used to generate samples of the input parameters and fit parameters, and that the  $\Delta\chi^2$  function associated to the element Ca  $ji$ , where  $i$  is the reference transition, was evaluated. As can be seen in




FIG. 20. Estimated  $\alpha_{NP}/\alpha_{EM}$  values (central dots) and bounds (bars) obtained executing `kifit`. The subplots show the impact of varying one of the relevant hyperparameters at a time.

Fig. 21, the manifestly symmetric version of  $\langle \tilde{\gamma} \rangle$  given in Eq. (47) leads to a  $\Delta\chi^2$  that is invariant under the exchange of transitions ( $12 \rightarrow 21$ ). The  $\Delta\chi^2$  values vary slightly between different sets of samples (Ca 21 samples vs. Ca 12 samples), but this is precisely the spread which is captured in the *experiment phase* (Section III C) and by the *blocking method* described in Section III D. This spread is present between all four sets of results when using the definition for  $\langle \tilde{\gamma} \rangle$  given in Eq. (44) (Fig. 21).

A number of complementary tests were directly implemented as `pytests` in the `kifit` code. A non-exhaustive list of the tests is given in the following. For details, please directly check the `kifit/src/tests` folder.

#### `test_build`

- Loading elements, checking dimensions of input data.
- Comparison of results of ODR and linear regression in linear fit to King plot data.

- Numerical cross-checks of `kifit` construction against results obtained with `Mathematica`.
- Numerical cross-checks of implementation of algebraic methods in `kifit` against results obtained with `Mathematica`.

#### `test_detools`

- Numerical cross-checks of uncertainties computed for algebraic methods in `kifit` against results obtained with `Mathematica`.
- Plot distribution of  $\alpha_{NP}$  values.

#### `test_fitools`

- Numerical cross-checks of `kifit` construction for single point (sample) in parameter space against results obtained with `Mathematica`. The procedure




FIG. 21. Test of the symmetry of the `kikit` code with respect to exchanges of the transitions. `Ca12` and `Ca21` correspond to two test data sets that differ only in the order of the transitions, and consequently in which of the two transitions is used as a reference transition (in `Caij`,  $i, j \in \{1, 2\}$ ,  $i$  is considered to be the reference transition). `Caij samples` denote the input parameter and fit parameter samples obtained for the data set `Caij`. Based on these samples, the  $\Delta\chi^2$  values are computed for each  $\alpha_{NP}$  sample, once using the `kikit.Elem` setup provided by the element `Caij` and once by adapting the samples to the conventions of the “dual” element `Caji` and using the setup provided by `Caji`. The left (right) figure shows the  $(\alpha_{NP}/\alpha_{EM}, \Delta\chi^2)$  values obtained adopting the symmetric (asymmetric) definition of  $\langle\tilde{\gamma}\rangle$  (Eq. (47) vs Eq. (44)).

is repeated for different values of the fit parameters  $\{K_{j1}^\perp, \phi_{j1}\}_{j=2}^m$ , for  $\frac{\alpha_{NP}}{\alpha_{EM}} = 0$  and for  $\frac{\alpha_{NP}}{\alpha_{EM}} = 10^{-11}$ .

- Cross-check of  $\|d\|$ , the covariance matrix  $\Sigma_d$ , the log-likelihood and various intermediate steps against results obtained with `Mathematica`. The procedure is repeated for different values of the fit parameters  $\{K_{j1}^\perp, \phi_{j1}\}_{j=2}^m$ , with and without sampling of the fit parameters, for different values of  $\alpha_{NP}$  and for different sample numbers.
- Cross-check of  $\Sigma_d$  and the log-likelihood for  $N_s = 10^2, 10^3, 10^4, 10^5$  samples. Check of the condition number of the covariance matrices and, by computing the spectral difference, Frobenius norm difference and the Kullback-Leibler divergence, comparison to the covariance matrices determined in `Mathematica`.  
For  $\alpha_{NP} = 0$  and  $\alpha_{NP} = 10^{-11}$ , comparison of the distribution of log-likelihood values for the  $N_s$  samples obtained using `kikit` and `Mathematica`.
- Check of the numerical accuracy of the inversion of the covariance matrix  $\Sigma_d$ .
- Tests of the symmetry of the `kikit` code with respect to exchange of the transitions or the choice of the reference transition (see discussion above), varying only the input parameters or varying both the input parameters and the fit parameters.
- Comparison of fit and algebraic methods for the test data set `Ca24min_mod`.

- Test of the numerical impact of the regulation of the covariance matrix  $\Sigma_d^{(\lambda)} = \Sigma_d + \lambda \mathbf{I}_n$ .

- Test of `kikit` run procedure

Finally, `test_cache_update` tests whether the `kikit`-internal cache is working and updated as expected.

## Appendix E: The Impact of Data Sparsity on the Fit

In this section we investigate how many data points (isotope pairs) are needed in a given number of dimensions (transitions) to ensure that the initial ODR is not over-fitting the data. We stress that the purpose of this analysis is to simulate the scaling of the relevant metrics in our estimation procedure with respect to the dimensionality of the King line ( $m$ ) and the number of data points ( $n$ ). The analysis is performed on mock data constructed from  $N_s$  samples of  $n$  “data points”  $\{(x^{(a,s)}, y_1^{(a,s)}, \dots, y_m^{(a,s)})\}_{a=1}^n\}_{s=1}^{N_s}$  which are normally distributed around a linear relation  $y_j = b_j x + c_j$ ,  $j = 1, \dots, m$  with variance  $\sigma_n$ . The uncertainties on the mock data points are modelled using Gaussian random variables with variance  $\sigma$ .

Concretely, for a given number  $n$  of points in  $m$  dimensions, we generate the parameters

$$b_j, c_j \sim \mathcal{N}(0, 10), \quad j = 1, \dots, m \quad (\text{E1})$$

which fix the linear relation  $y_j = b_j x + c_j$  and then generate  $N_s = 500$  samples ( $s = 1, \dots, N_s$ ) from the distri-




FIG. 22. Illustration of the goodness a fit to  $y_j = b_j x + c_j$ ,  $j = 1, \dots, m$  as a function of the number  $n$  of mock data points  $(x, y_1, \dots, y_m)$ .  $\zeta^{(n)} \equiv \langle \zeta^{(n,s)} \rangle_{(s)}$  with  $\zeta^{(n,s)}$  as defined in Eq. (E3) corresponds to the average relative uncertainty on the  $2(n-1)$  fit parameters  $\{b_j, c_j\}_{j=2}^m$  used to fix the line in  $m$ -dimensional space. We show the results for the minimal case of  $m = 2$  and for  $m = 6$ . For each number  $n$  of data points, 500 mock data sets with Gaussian uncertainties  $\sigma = \sigma_n = 10^{-10}$  were generated, leading to relative uncertainties  $\sigma_x/x$ ,  $\sigma_y/y$  of the same order. In black we show the curve defined by  $f(n) = \theta_1/\sqrt{n} - \theta_2 + \theta_3$ , with  $(\theta_1, \theta_2, \theta_3) \approx (1.5 \times 10^{-9}, 1.7, -3 \times 10^{-10})$ .




FIG. 23.  $\delta^{(n,m)}$  (see Eq. (E6)) as a function of the number  $n$  of the mock data points. In black we show the curve defined by  $f(n) = \theta_1/\sqrt{n} - \theta_2 + \theta_3$ , with  $(\theta_1, \theta_2, \theta_3) \approx (-0.6, 1.6, 1)$ .  $\delta^{(n,m)}$  can be interpreted as the average distance of a point  $a$  in an  $n$ -dimensional data set to the corresponding best-fit line.

butions

$$\begin{aligned} x^{(a,s)} &\sim \mathcal{N}(x^{(a)}, \sigma_n), \quad a = 1, \dots, n \\ y_j^{(a,s)} &\sim \mathcal{N}(b_j x^{(a)} + c_j, \sigma_n), \quad j = 1, \dots, m \\ \sigma[x^{(a,s)}], \sigma[y^{(a,s)}] &\sim \mathcal{N}(0, \sigma), \end{aligned} \quad (\text{E2})$$

where we take  $\sigma_n = \sigma = 10^{-10}$ . For each sample  $\{(x^{(a,s)}, y_1^{(a,s)}, \dots, y_m^{(a,s)})\}_{a=1}^n$  of the  $n$  data points in  $m$  dimensions,  $2m$  fit parameters  $\{(b_j^{(n,s)}, c_j^{(n,s)})\}_{j=1}^m$  are determined by means of orthogonal distance regression.




FIG. 24.  $\delta^{(n,m)}$  (see Eq. (E6)) as a function of the number  $m$  of dimensions, for selected sample sizes  $n$ .  $\delta^{(n,m)}$  can be interpreted as the average distance of a point  $a$  in an  $n$ -dimensional data set to the corresponding best-fit line.

As a first check, we compute the combined fractional differences of the fit parameter samples with respect to the original slopes and intercepts  $\{(b_j, c_j)\}_{j=1}^m$ ,

$$\zeta^{(n,s)} = \sqrt{\sum_{j=2}^m \left( \frac{(b_j^{(n,s)} - b_j)^2}{b_j^2} + \frac{(c_j^{(n,s)} - c_j)^2}{c_j^2} \right)}. \quad (\text{E3})$$

The averages and standard deviations over the samples,  $\zeta^{(n)} \equiv \langle \zeta^{(n,s)} \rangle_{(s)}$  and  $\sigma[\zeta^{(n,s)}]_{(s)}$  can be viewed as a measure of the error in the fit parameters due to the finite sample size  $n$ . They are plotted in Fig. 22 as a function of the number of points  $n$ . As expected,  $\zeta^{(n)}$  approximately follows a relation of the form  $f(n) = \theta_1/\sqrt{n} - \theta_2 + \theta_3$ , which is motivated by the expected asymptotic behaviour of Gaussian uncertainties, by the fact that a linear fit to one point is ill-defined and by the finite number of  $n$  values considered for the curve fit. The explicit parameters obtained when fitting the first 15 data points for  $m = 2$  to this curve are  $(\theta_1, \theta_2, \theta_3) \approx (1.5 \times 10^{-9}, 1.7, -3 \times 10^{-10})$ .

Since relevant quantity for the log-likelihood in Eq. (52) is the distance of the points from the line, normalised by the respective uncertainties, we apply a definition very similar to that used in `kifit` (see Eqs. (48),



(38), (49)):

$$\delta^{(a,m,s)} = \frac{1}{\sigma\sqrt{m}} \|\Delta^{(a,m,s)} - (\Delta^{(a,m,s)} \cdot \hat{\mathbf{e}}_{\mathbf{f}}) \hat{\mathbf{e}}_{\mathbf{f}}\| \quad (\text{E4})$$

with

$$\Delta^{(a,m,s)} = \begin{pmatrix} 0 \\ y_1^{(a,s)} - (b_1^{(a,s)} x^{(a,s)} + c_1^{(a,s)}) \\ \vdots \\ y_m^{(a,s)} - (b_m^{(a,s)} x^{(a,s)} + c_m^{(a,s)}) \end{pmatrix} \quad (\text{E5})$$

and  $\hat{\mathbf{e}}_{\mathbf{f}} = (1, b_1^{(a,s)}, \dots, b_m^{(a,s)})$ , and a normalisation  $\sigma\sqrt{m}$  that captures the asymptotic behaviour of the uncertainties on the sum. We then define

$$\begin{aligned} \delta^{(n,m)} &\equiv \left\langle \frac{1}{n} \sum_{a=1}^n \delta^{(a,m,s)} \right\rangle_{(s)}, \\ \sigma[\delta^{(n,m)}] &\equiv \sigma \left[ \frac{1}{n} \sum_{a=1}^n \delta^{(a,m,s)} \right]_{(s)}, \end{aligned} \quad (\text{E6})$$

where  $\langle \cdot \rangle_{(s)}$  and  $\sigma[\cdot]_{(s)}$  denote the average and the standard deviation over the samples ( $s$ ). In Fig. 23,  $\delta^{(n,m)}$  and  $\sigma[\delta^{(n,m)}]$  are plotted as a function of the number of

points  $n$  and for different dimensions  $m$ .  $\delta^{(n,m)}$  can be viewed as the average distance, normalised by the experimental uncertainties, of a point  $a$  in a data set of size  $n$  to the best-fit line.  $\delta^{(n,m)}$  increases with  $n$  because the uncertainty on the distance to the line receives an additional contribution from the increasing spread of the data points, which for the mock data employed here has variance  $\sigma_n$  (see Eq. (E2)). This residual  $n$ -dependence is again captured by a function of the form  $f(n) = \theta_1/\sqrt{n-\theta_2} + \theta_3$ , this time with  $(\theta_1, \theta_2, \theta_3) \approx (-0.6, 1.6, 1)$  for the first 15 points in  $m = 2$  dimensions. Fig. 23 shows that increasing the number of data points from 3 by a few can significantly improve the reliability of the fit results.

The  $m$ -dependence of  $\delta^{(n,m)}$  is less pronounced, as can be observed in Fig. 24. Nonetheless, measurements of additional transitions can improve the heterogeneity of the data and thus the reliability of the bounds on  $\alpha_{\text{NP}}$ .

## Appendix F: Isotope Shift and Atomic/Nuclear Mass Data

In Tables IX and X, we collect the available isotope shift measurements for Ca and Yb. The most recent measurements for the isotope masses can be found in Tables VII and VIII. We mark with an asterisk the values that are included in the `kifit` folders.

Isotope A	$m_A$ [amu] [103]	$m_A$ [amu] [66]	$m_A/m_{40}$ [43]
40	39.962590866(22)	39.962590851(22)	1
42	41.95861783(16)	41.95861778(16)	1.049 961 066 498(15)
44	43.9554815(3)	43.9554815(3)	1.099 943 105 797(15)
46	45.9536880(24)	45.9536877(24)	1.149 958 773 895(30)
48	47.95252290(10)	47.952522654(19)	1.199 990 087 090(40)

TABLE VII. Atomic masses of Ca isotopes from Refs. [66, 103]. The third column shows the ratios of bare nuclear masses to the mass of isotope  $A = 40$ , as reported in Ref. [43].

Isotope A	$m_A$ [amu] [103]	$m_A$ [amu] [66]	$m_A/m_{172}$ [50]
168	167.9338891(13)	167.93389130(10)	0.976715921749(4)
170	169.934767246(11)	169.934767243(11)	0.988355799258(4)
172	171.936386659(15)	171.936386654(15)	1
174	173.938867548(12)	173.938867546(12)	1.011649212140(4)
176	175.942574709(16)	175.942574706(16)	1.023305557965(4)

TABLE VIII. Atomic masses of Yb isotopes from Refs. [66, 103]. The third column shows ratios of bare nuclear masses to the mass of isotope  $A = 172$ , as reported in Ref. [50].

$(A, A')$	$\text{Ca}^+: 4^2S_{1/2} \rightarrow 3^2D_{5/2}$ [729 nm]	$\text{Ca}^+: 3^2D_{3/2} \rightarrow 3^2D_{5/2}$ DD	$\text{Ca}^+: 4s^2S_{1/2} \rightarrow 4p^2P_{1/2}$ [397 nm]
(40, 42)	*2 771 872 467.6(7.6) [98] 2 771 873 000(2000) [41] 2 771 872 430.217(27) [43]	*-3 519 896(24) [41] *-3 519 910(9.7) [102] -3 519 944.6(60) [43]	*425 706 000(94 000) [26] 425 490 000(150 000) [104]
(40, 44)	*5 340 887 394.6(7.8) [98] 5 340 888 000(2000) [41] 5 340 887 395.288(38) [43]	*-6 792 470(22) [41] *-6 792 440(6) [102] -6 792 440.1(59) [43]	*849 534 000(74 000) [26] 849 000 000(140 000) [104]
(40, 46)	*7 768 401 000(2000) [41] 7 768 401 432.916(63) [43] 9 990 382 525.0(4.9) [99]	*-9 901 524(21) [41] *-9 901 520(2828.43) [102] -9 901 524(21) [43] -12 746 610(27) [41]	*1 297 610 000(340 000) [104] 1 705 389 000(60 000) [26] 1 705 460 000(140 000) [104]
(40, 48)	9 990 383 000(2000) [41] 9 990 382 526.834(55) [43]	*-12 746 600(7.5) [102] -12 746 588.2(57) [43]	
$(A, A')$	$\text{Ca}^+: 4^2S_{1/2} \rightarrow 3^2D_{3/2}$ [732 nm]	$\text{Ca}^+: 4^2S_{1/2} \rightarrow 4p^2P_{3/2}$ [393 nm]	$\text{Ca}^+: 3d^2D_{3/2} \rightarrow 4p^2P_{1/2}$ [866 nm]
(40, 42)	2 775 392 374.8(6.0) [102]	426 040 000(150 000) [104] 425 932 000 (71 000) [27]	*-2 349 974 000(99 000) [26] 2 366 000 000(59 000 000) [105] 2 352 100 000(2 100 000) [38]
(40, 44)	5 347 679 835.4(5.9) [102]	850 090 000(140 000) [104] 850 231 000(65 000) [27]	*-4 498 883 000(80 000) [26] 4 509 000 000(24 000 000) [105] 4 499 300 000(2 300 000) [38]
(40, 46)		1 299 070 000(580 000) [104] 1 301 000 000(3 600 000) [106]	
(40, 48)	10 003 129 115.1(5.7) [102]	1 707 580 000(160 000) [104] 1 707 945 000(67 000) [27]	*-8 297 769 000(81 000) [26] 8 296 700 000(3 200 000) [38]
$(A, A')$	$\text{Ca}^+: ^2D_{5/2} \rightarrow ^2P_{3/2}$ [854 nm]	$\text{Ca}^+: ^2D_{3/2} \rightarrow ^2P_{3/2}$ [850 nm]	$\text{Ca}^{14+}: ^3P_0 \rightarrow ^3P_1$ [570 nm]
(40, 42)	2 272 000 000(94 000 000) [105] 2 347 600 000(3 900 000) [38]	2 359 000 000(64 000 000) [105] 2 351 450 000(700 000) [38]	539 088 421.24(12) [43]
(40, 44)	4 510 000 000(19 000 000) [105] 4 489 800 000(3 600 000) [38]	4 538 000 000(27 000 000) [105] 4 497 270 000(900 000) [38]	1 030 447 731.64(11) [43]
(40, 46)	6 470 100 000(2 400 000) [38]		1 481 135 946.74(14) [43]
(40, 48)	8 277 900 000(5 000 000) [38]	8 295 060 000(1 100 000) [38]	1 894 297 294.53(14) [43]

TABLE IX. Isotope shifts of  $\text{Ca}^+$  and  $\text{Ca}^{14+}$ , expressed in Hz. Values included in `kifit` are marked with an asterisk. Notice that Ref. [102] reports measurements of the  $4^2S_{1/2} \rightarrow 3^2D_{3/2}$  transition, and the corresponding DD-transition values are derived therein from their measurement of  $4^2S_{1/2} \rightarrow 3^2D_{3/2}$  and the  $4^2S_{1/2} \rightarrow 3^2D_{5/2}$  reported in Ref.s [41, 98, 99]. Similarly, the DD-transition values reported in Ref. [43] are derived from their measurement of  $4^2S_{1/2} \rightarrow 3^2D_{5/2}$  and the  $4^2S_{1/2} \rightarrow 3^2D_{3/2}$  reported in Ref.[102].

$(A, A')$	$(\alpha) \text{ Yb}^+ : ^2S_{1/2} \rightarrow ^2D_{5/2}$ [411 nm]	$(\beta) \text{ Yb}^+ : ^2S_{1/2} \rightarrow ^2D_{3/2}$ [436 nm]	$(\gamma) \text{ Yb}^+ : ^2S_{1/2} \rightarrow ^2F_{7/2}$ [467 nm]
(168, 170)	*2 179 098 930(210) [48] *2 179 098 868.0(5.3) [50]	*2 212 391 850(370) [48]	*-4 438 160 300(500) [49] *-4 438 159 671.1(15.7) [50]
(170, 172)	*2 044 854 780(340) [48] *2 044 851 281.0(4.9) [50]	*2 076 421 580(390) [48]	*-4 149 190 380(450) [49] *-4 149 190 501.1(15.7) [50]
(172, 174)	*1 583 068 420(360) [48] *1 583 064 149.3(4.8) [50]	*1 609 181 470(220) [48]	*-3 132 321 600(500) [49] *-3 132 320 458.1(15.7) [50]
(174, 176)	*1 509 055 290(280) [48] *1 509 053 195.8(4.7) [50]	*1 534 144 060(240) [48]	*-2 976 391 600(480) [49] *-2 976 392 045.3(15.7) [50]
(168, 172)			-8 587 352 000(470) [49]
(170, 174)	3 627 922950(500) [48]	3 685 601950(330) [48]	-7 281 511 880(450) [49]
(172, 176)			-6 108 712 930(440) [49]
$(A, A')$	$(\epsilon) \text{ Yb} : ^1S_0 \rightarrow ^1D_2$ [361 nm]	$(\delta) \text{ Yb} : ^1S_0 \rightarrow ^3P_0$ [578 nm]	$\text{Yb} : 4f^{14}6s^{21}S_0 \longleftrightarrow 4f^{13}5d^6s^2$ [431 nm]
(168, 170)	*1 781 785 360(710) [47]	*1 358 484 476.2(2.2) <sub>tot</sub> [72]	-1 753 930 000(3000) <sub>stat</sub> [107] -1 753 952 000(26 000) [73]
(170, 172)	*1 672 021 510(300) [47]	*1 275 772 006(2.8) <sub>stat</sub> [72]	-1 630 050 000(3000) <sub>stat</sub> [107] -1 630 028 000(26 000) [73]
(172, 174)	*1 294 454 440(240) [47]	*992 714 586.6(2.1) <sub>tot</sub> [72]	-1 180 614 000(2000) <sub>stat</sub> [107] -1 180 616 000(25 000) [73]
(174, 176)	*1 233 942 190(310) [47]	*946 921 774.9(2.9) <sub>tot</sub> [72]	-1 115 766 000(6000) <sub>stat</sub> [107] -1 115 787 000(24 000) [73]
(168, 172)	3 453 805 270(83) [47]		
(172, 176)	2 528 396 500(34) [47]		
$(A, A')$	$\text{Yb}^+ : 6^2S_{1/2} \rightarrow 6^2P_{1/2}^o$ [369 nm]	$\text{Yb}^+ : 6^2S_{1/2} \rightarrow 6^2P_{3/2}^o$ [329 nm]	$\text{Yb} : 6^1S_0 \rightarrow 6^3P_1^o$ [556 nm]
(168, 170)			-1 368 630 000(500000) [44]
(170, 172)	-1 623 300(800) [108]	-1 459 000(21000) [109]	-1 286 470(500) [44]
(172, 174)	-1 275 300(700) [108]	-1 154 000(11000) [109]	-1 000 280(500) [44]
(172, 176)	-2 492 800(1000) [108]	-2 259 000(13000) [109]	-1 955 040(500) [44]
(174, 176)			-954 760 000(5000000) [44]
$(A, A')$	$\text{Yb}^+ : ^1S_0 \rightarrow ^1P_1$ [399 nm]		
(174, 168)	1 888 800 000(110 000) [46]		
(174, 172)	1 190 360 000(490 000) [46]		
(174, 172)	-250 780 000(330 000) [46]		
(174, 176)	-508 890 000(090 000) [46]		

TABLE X. Isotope shifts for Yb and  $\text{Yb}^+$ , expressed in Hz. Values included in **kifit** are marked with an asterisk. Isotopes used in this work are organised in pairs  $(A, A')$  with  $A' = A + 2$ . The data for the other isotope pairs can be used for cross-checks.