CERN-EP-2025-169
2025/08/22

CMS-HIG-24-009

Search for a Higgs boson produced in association with a charm quark and decaying to a W boson pair in proton-proton collisions at $\sqrt{s} = 13$ TeV

The CMS Collaboration*

Abstract

This paper presents a search for a Higgs boson produced in association with a charm quark (cH) which allows to probe the Higgs-charm Yukawa coupling strength modifier κ_c . Higgs boson decays to a pair of W bosons are considered, where one W boson decays to an electron and a neutrino, and the other W boson decays to a muon and a neutrino. The data, corresponding to an integrated luminosity of 138 fb^{-1} , were collected between 2016 and 2018 with the CMS detector at the LHC at a center-of-mass energy of $\sqrt{s} = 13$ TeV. Upper limits at the 95% confidence level (CL) are set on the ratio of the measured yield to the standard model expectation for cH production. The observed (expected) upper limit is 1065 (506). When combined with the previous search for cH in the diphoton decay channel of the Higgs boson, the limits are interpreted as observed (expected) constraints at 95% CL on the value of κ_c , $|\kappa_c| < 47$ (51).

Submitted to the Journal of High Energy Physics

1 Introduction

Over a decade has passed since the ATLAS and CMS Collaborations at the CERN LHC discovered the Higgs (H) boson [1–3]. With more data and improved analysis techniques, precise measurements of the properties of the H boson have become possible, along with its rare decay channels. The couplings of the H boson to other standard model (SM) particles are among its most important properties [4, 5]. The coupling strength modifiers, κ , are introduced to parameterize potential deviations from the SM predictions in the H boson interactions with other particles. The couplings to bosons (Z, W, and γ) are well determined, based on precise measurements from the $H \rightarrow ZZ \rightarrow 4\ell$, $H \rightarrow WW$, and $H \rightarrow \gamma\gamma$ decay channels [4, 5]. Among the Yukawa couplings, only those to third-generation fermions have been firmly established. The coupling modifiers κ_b and κ_τ have been constrained through measurements of the $H \rightarrow b\bar{b}$ [6–8] and $H \rightarrow \tau\tau$ [9, 10] processes, while κ_t has been constrained through a global fit [4, 5]. The coupling to muons has been probed in the $H \rightarrow \mu\mu$ channel [11, 12], reaching evidence at the level of 3 standard deviations [4].

The direct constraints on κ_c have been obtained through searches for $H \rightarrow c\bar{c}$ decays. These constraints have been derived in several production processes, including gluon fusion (ggH) [13] and associated production with a vector boson (VH) [14, 15]. With the advanced analysis techniques and recent improvements in charm jet identification methods (charm tagging) and calibrations [16, 17], the direct measurements of the decay provide observed (expected) constraints of $|\kappa_c| < 5.5$ (3.4) at 95% confidence level (CL) [15]. In addition to direct searches, κ_c can also be constrained indirectly through differential measurements of the transverse momentum (p_T) of the H boson across multiple decay channels [14, 18]. An alternative method to probe the Higgs-charm coupling is via H boson production in association with charm quarks (cH) [19, 20]. This channel probes κ_c via radiation of a H boson of a charm quark. Charm quarks appear abundant in the initial state together with a gluon. Leading-order (LO) Feynman diagrams contributing to the $pp \rightarrow cH$ process are shown in Fig. 1. Despite a small cross section of this channel, its advantage is a possibility to explore simpler event signatures compared to those involving c jets from the H boson decay. Recently, the CMS Collaboration reported on this production mechanism in the diphoton decay channel using the proton-proton (pp) data set at $\sqrt{s} = 13$ TeV collected between 2016 and 2018 [21]. An observed (expected) limit $|\kappa_c| < 38.1$ (72.5) at 95% CL has been obtained. Additionally, a search for the H boson with a charm quark in the final state was released by the ATLAS Collaboration [22]. The ATLAS analysis did not measure the cH component that is sensitive to κ_c . Hence, no constraint on $|\kappa_c|$ was reported.




Figure 1: Leading-order Feynman diagrams that contribute to the $pp \rightarrow cH$ process. The red dots indicate vertices where the Higgs-charm coupling modifier κ_c is involved.

In this paper, a search for a H boson produced in association with a charm quark is presented, where the H boson decays to two W bosons, one decaying to an electron-neutrino pair and the other to a muon-neutrino pair. This final state is chosen to suppress background from Z+jets production, which has a significantly larger cross section and predominantly affects the same-flavor dilepton final states (ee and $\mu\mu$). Tabulated results are provided in the HEPData record for this analysis [23].

2 The CMS detector

The central feature of the CMS apparatus is a superconducting solenoid of 6 m internal diameter, providing a magnetic field of 3.8 T. Within the solenoid volume are a silicon pixel and strip tracker, a lead tungstate crystal electromagnetic calorimeter (ECAL), and a brass and scintillator hadron calorimeter (HCAL), each composed of a barrel and two endcap sections. Forward calorimeters extend the pseudorapidity (η) coverage provided by the barrel and endcap detectors. Muons are reconstructed in gas-ionization detectors embedded in the steel flux-return yoke outside the solenoid. More detailed descriptions of the CMS detector, together with a definition of the coordinate system used and the relevant kinematic variables, can be found in Refs. [24, 25].

Events of interest are selected using a two-tiered trigger system. The first level (L1), composed of custom hardware processors, uses information from the calorimeters and muon detectors to select events at a rate of around 100 kHz within a fixed latency of $4\mu\text{s}$ [26]. The second level, known as the high-level trigger, consists of a farm of processors running a version of the full event reconstruction software optimized for fast processing and reduces the event rate to around 1 kHz before data storage [27, 28]. During the 2016 and 2017 data-taking, a gradual shift in the timing of the inputs of the ECAL L1 trigger caused a specific trigger inefficiency. Correction factors were computed from data and applied to the simulated events, to mitigate this effect.

3 Data sets and simulated samples

The data sets used in the analysis were recorded by the CMS detector in the 2016, 2017, and 2018 data-taking periods, corresponding to an integrated luminosity of 36.3, 41.5, and 59.8 fb^{-1} , respectively [29–31]. The 2016 data set is split into two separate periods, due to a substantial change in the tracker and muon system configurations in the middle of the data-taking, with the corresponding integrated luminosities of 19.5 and 16.8 fb^{-1} , respectively. Events are selected by a combination of triggers requiring the presence of either one isolated muon with $p_T > 24$ (27) GeV for 2016 and 2018 (2017), or one isolated electron with p_T of 27, 35, and 32 GeV for 2016, 2017, and 2018, respectively, or the presence of both ($e\mu$), with values of $p_T > 23$ (12) GeV for the lepton with the highest (second-highest) p_T .

Processes for H boson production in association with a c or b quark, cH and bH, are generated with MADGRAPH5_aMC@NLO 2.6 [32, 33] at next-to-leading order (NLO). Dedicated studies have been performed to assess the impact of heavy-quark masses in cH and bH processes. In the 4-flavor scheme (4FS) [34], the calculations are performed with massless c quarks, while in the 3-flavor scheme (3FS) massive c quarks are assumed. A similar approach is applied to b quarks: in the 5-flavor scheme (5FS) massless b quarks are used, whereas in the 4FS their mass is taken into account. These two schemes are designed to model heavy quarks in different kinematic regimes, considering whether the mass of the quark plays a role of hard scattering scale in this process. The massless quark assumption simplifies the calculations significantly




Figure 2: Diagram illustrating H boson production in association with a charm quark in the absence of charm Yukawa coupling. The crossed vertex represents the top quark loop.

compared to including their masses, but this approach introduces divergences at low energies. To mitigate this effect, dedicated events are generated in order to estimate the uncertainties arising from the kinematic differences. This analysis assumes massless c or b quarks, using the 4FS for cH and the 5FS for bH . These calculations are combined with parton shower (PS) processes using FxFx jet merging [35] at NLO. The cross sections are found to be $\sigma_{cH} = 90$ fb and $\sigma_{bH} = 660$ fb. The backward evolution recovers the information of the initial state c and b quarks [34]. The same method was used in our previous $cH(H \rightarrow \gamma\gamma)$ search [21].

Other H boson production processes, including ggH , vector boson fusion (VBF), VH , and $t\bar{t}H$, are simulated with POWHEG v2 [36–42]. All H boson production processes other than cH are considered as background in this analysis and denoted as H-bkg. The cH events from diagrams without charm Yukawa coupling, as shown in Fig. 2, are considered as part of the H-bkg background. The final state kinematics of the processes in Figs. 1 and 2 are very similar. Therefore, a multivariate discriminator based on a combination of kinematic variables is used to distinguish the processes represented by these diagrams, as discussed in Section 5. In addition, the modeling of H-bkg processes involving heavy-flavor jets poses significant theoretical challenges. To account for these difficulties, a 50% uncertainty is assigned based on calculations for the $ggH+b\bar{b}$ process [43].

Other background processes, such as top quark-antiquark ($t\bar{t}$), single top quark, and diboson (WW, WZ, ZZ) production are produced with POWHEG v2 at NLO. Production of Z bosons in association with jets (Z+jets) is simulated by MADGRAPH5_aMC@NLO 2.6 at NLO. For the W boson production in association with jets (W+jets), a combination of simulated events of LO and NLO from different generators (MADGRAPH5_aMC@NLO 2.6 and SHERPA 2.2 [44]) are used to obtain sufficiently large samples. The contributions of W+jets and Z+jets are denoted inclusively as V+jets in this paper.

Signal and background processes are simulated using various event generators. For all the simulated events, PYTHIA 8.240 [45] simulates underlying event (UE), PS, and hadronization. The simulations employ the NNPDF 3.1 [46] parton distribution functions at next-to-next-to-leading order (NNLO), along with the CP5 underlying event tune [47]. The simulated events include additional pp interactions within the same or nearby bunch crossings (pileup), which are corrected to match the distribution observed in data. The CMS detector response is modeled with GEANT4 [48].

4 Event reconstruction and selection

The primary vertex is taken to be the vertex corresponding to the hardest scattering in the event, evaluated using tracking information alone, as described in Section 9.4.1 of Ref. [49]. Charged hadrons originating from other vertices (pileup vertices) are removed from the analysis.

A particle-flow (PF) algorithm [50] aims to reconstruct and identify each individual particle in an event (PF candidate), with an optimized combination of information from the various elements of the CMS detector.

Electrons are measured in $|\eta| < 2.5$ with a gap at $1.44 < |\eta| < 1.57$. The energy of electrons is determined from a combination of the track momentum at the main interaction vertex, the corresponding ECAL cluster energy, and the energy sum of all bremsstrahlung photons attached to the track. The electron momentum is estimated by combining the energy measurement in the ECAL with the momentum measurement in the tracker. The momentum resolution for electrons with $p_T \approx 45$ GeV from $Z \rightarrow ee$ decays ranges from 1.6 to 5%. It is generally better in the barrel region than in the endcaps, and also depends on the bremsstrahlung energy emitted by the electron as it traverses the material in front of the ECAL [51, 52]. Muons are measured in $|\eta| < 2.4$, with detection planes made using three technologies: drift tubes, cathode strip chambers, and resistive-plate chambers. The momentum of muons is obtained from the corresponding track curvature. Matching muons to tracks measured in the silicon tracker results in a relative p_T resolution, for muons with p_T up to 100 GeV, of 1% in the barrel and 3% in the endcaps. The p_T resolution in the barrel is better than 7% for muons with p_T up to 1 TeV [53]. The selected electrons and muons are required to pass tight identification criteria, ensuring a low misidentification rate with lepton efficiencies of approximately 70% [51, 53]. To suppress leptons originating from b and c hadron decays, while retaining those from W boson decays, isolation from jet activity is required within a cone of radius $\Delta R = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2} = 0.3$ (0.4) for electrons (muons) [53], where ϕ is the azimuthal angle in radians. The isolation is defined as the scalar p_T sum of the PF candidates within the cone divided by the lepton p_T .

Events are selected if they contain exactly one electron and one muon, with the leading lepton ($\ell 1$) $p_T > 25$ GeV and subleading lepton ($\ell 2$) $p_T > 15$ GeV. A higher p_T threshold of 35 (30) GeV is applied to the leading electron (muon) if an event is triggered only by a single-lepton trigger. Additionally, due to higher online p_T thresholds, electrons must have $p_T > 38$ GeV in the 2017 data set.

For each event, hadronic jets are reconstructed by clustering PF candidates using the infrared- and collinear-safe anti- k_T algorithm [54, 55] with a distance parameter of 0.4. Jet momentum is determined as the vectorial sum of all particle momenta in the jet, and is found from simulation to be, on average, within 5 to 10% of the true momentum over the whole p_T spectrum and detector acceptance. Pileup can contribute additional tracks and calorimetric energy deposits to the jet momentum. To mitigate this effect, charged particles identified to be originating from pileup vertices are discarded and an offset correction is applied to correct for remaining contributions [56]. Jet energy corrections are derived from simulation to bring the measured response of jets to that of particle-level jets on average. In situ measurements of the momentum balance in dijet, photon+jet, Z+jet, and multijet events are used to account for any residual differences in the jet energy scale between data and simulation [57]. The jet energy resolution amounts typically to 15–20% at 30 GeV, 10% at 100 GeV, and 5% at 1 TeV [57]. Additional selection criteria are applied to each jet to remove jets potentially dominated by anomalous contributions from various subdetector components or reconstruction failures.

Jets originating from charm quarks, c jets, are identified using the DEEPJET algorithm [16, 58,

59]. This is a deep neural network algorithm which utilizes several input features, including secondary-vertex properties, track-based variables, and PF jet constituents. Discrimination between c jets and light jets (jets originating from u, d, and s quarks or gluons) is achieved via the probability ratio defined as $CvsL = p(c)/[p(c) + p(udsg)]$, where the $p(c)$ and $p(udsg)$ represent the classifier output for c and light jets, respectively. Similarly, discrimination between c jets and b jets is based on the probability ratio defined as $CvsB = p(c)/[p(c) + p(b)]$. These two discriminant ratios are the charm jet discriminants, which are used to select c jet candidates with thresholds of $CvsL > 0.12$ and $CvsB > 0.5$. These selections provide a charm jet tagging efficiency of 43% while the b (light) jet mistag rate is 27% (5%). This allows discrimination between cH and bH processes. The c jet with the highest $CvsL$ score is considered the leading c jet candidate. Events with at least one jet that passes the above charm jet selection criteria and that satisfies $p_T > 20$ GeV, $|\eta| < 2.4$, and $\Delta R(\ell, j) > 0.4$ (with $\ell = e, \mu$) are selected. The shapes of c jet discriminants in the simulated samples are corrected by flavor-dependent simulation-to-data efficiency scale factors [17].

The missing transverse momentum ($\vec{p}_T^{\text{miss}}$) is computed as the negative vector sum of the transverse momenta of all the PF candidates in an event, and its magnitude is denoted as p_T^{miss} [60]. The $\vec{p}_T^{\text{miss}}$ is modified to account for corrections to the energy scale of the reconstructed jets in the event. Only events passing $p_T^{\text{miss}} > 20$ GeV are selected. Anomalous high- p_T^{miss} events can arise due to a variety of reconstruction failures, detector malfunctions, or non-collision backgrounds. Such events are rejected by event filters that are designed to identify greater than 85% of the spurious high- p_T^{miss} events with a mistagging rate less than 0.1% [60].

In this paper, only the $WW \rightarrow e\nu\mu\nu + c$ final state is considered. The $WW \rightarrow e\nu e\nu$ and $\mu\nu\mu\nu$ decay channels are not considered due to the large Z+jet background contribution. Therefore, the presence of an electron and a muon with opposite charges and $\Delta R(e, \mu) > 0.4$ is required. The invariant mass (the p_T) of the dilepton pair must be greater than 12 (30) GeV to remove low mass resonances. Furthermore, the event selection criteria employ the transverse mass defined as

$$m_T^{\text{obj}} = \sqrt{2p_T^{\text{obj}} p_T^{\text{miss}} [1 - \cos \Delta\phi(\text{obj}, \vec{p}_T^{\text{miss}})]}, \quad (1)$$

where the p_T^{obj} represents the transverse momentum of an object (a lepton, or dilepton system), and $\Delta\phi$ is the difference in ϕ between the object and $\vec{p}_T^{\text{miss}}$. A requirement on the transverse mass with the subleading lepton, $m_T^{\ell\ell} > 30$ GeV, is applied to suppress backgrounds not originating from W boson decays.

5 Analysis strategy

Events are categorized into two signal regions (SRs) and two control regions (CRs). The c jet multiplicity requirement, $N_{c,j} = 1$ and $N_{c,j} > 1$, defines two SRs with different sensitivities. In each SR, events must satisfy $m_T^{\ell\ell} > 60$ GeV and $m_{\ell\ell} < 72$ GeV to maximize the signal efficiencies and to suppress the background contributions. Two CRs are defined, denoted as *high- $m_{\ell\ell}$* and *top*. The *high- $m_{\ell\ell}$* CR follows the SR selection criteria, except for the dilepton invariant mass requirement, which is inverted to maintain a similar background composition as in the SRs. The *top* CR, in addition to the inverted $m_{\ell\ell}$ selection, requires at least two c jet candidates and $m_T^{\ell\ell} < 60$ GeV to enrich the contribution of $t\bar{t}$ events. Table 1 summarizes the definitions of the SRs and CRs. The dominant background process is $t\bar{t}$, which contributes approximately 90% to the total background.

A boosted decision tree (BDT) is used to construct two binary classifiers for distinguishing sig-

Table 1: Definitions of the SRs and CRs. Two SRs are defined according to c jet multiplicity in selected events. Events with dilepton invariant mass above 72 GeV are considered as CRs. Events with an inverted transverse masses requirement that have at least two c jets are considered in *top* CR and the remaining events are tagged as *high- $m_{\ell\ell}$* CR.

Selection		$N_{c-j} = 1$ SR	$N_{c-j} > 1$ SR	High $m_{\ell\ell}$ CR	Top CR
c jet multiplicity	$N_{c-j} = 1$	=1	>1	≥ 1	≥ 2
Dilepton invariant mass (GeV)	$m_{\ell\ell}$	<72	<72	>72	>72
Transverse mass for the subleading lepton (GeV)	$m_T^{\ell 2}$	>30	>30	>30	>30
Transverse mass for the dilepton (GeV)	$m_T^{\ell\ell}$	>60	>60	>60	<60




Figure 3: The area-normalized distributions of the two BDT classifiers: D_{bkg} (left) and D_{H-bkg} (right). The D_{bkg} classifier effectively separates the non H boson background ($t\bar{t}$, single top, diboson, and V+jets, denoted as other background) contributions from the cH production, but it is not sufficient for suppressing the H-bkg (ggH, VBF, VH, $t\bar{t}H$, and bH) contributions. The dedicated BDT classifier, D_{H-bkg} , is capable of distinguishing between other H boson production processes and cH production.

nal from background events. The first classifier, D_{bkg} , is designed to disentangle the cH process from the main background contributions ($t\bar{t}$, single top quark, diboson, and V+jets). The second classifier, D_{H-bkg} , is trained to separate the cH signal from other H boson processes. Both BDTs use the same variables for training, including the p_T , the angular separations of the leptons, p_T^{miss} and the c jet candidate, as well as the $CvsL$ and $CvsB$ values of the c jet candidate. The full list of input variables to the BDT training are summarized in Table 2. The classifiers are implemented using the XGBOOST [61] package with the gradient boosting method. The training data set includes 20 k events for the cH signal, 40 k for H-bkg, and 100 k for the other background processes. These data sets are split equally between training and validation. Events are weighted according to their corresponding cross sections, and a class-balancing weight is applied to the signal to ensure balanced training. The BDT trainings are optimized separately for each data-taking era to account for differences in detector conditions. The performance of D_{bkg} and D_{H-bkg} is evaluated using the area under the ROC curve, which ranges from 93% to 98% for D_{bkg} and from 73% to 78% for D_{H-bkg} . The shapes of the two BDT classifiers are shown in Fig. 3.

Furthermore, H-bkg events are classified into three categories based on the generator-level

Table 2: Summary of input variables used in the BDT trainings.

Variable	Description
$p_T^{\ell\ell}$	Transverse momentum of dilepton system ($e\mu$)
$p_T^{\ell 1}$	Transverse momentum of leading lepton (sorted by p_T)
$p_T^{\ell 2}$	Transverse momentum of subleading lepton (sorted by p_T)
p_T^c	Transverse momentum of c jet candidate
p_T^{miss}	Missing transverse momentum
$m_{\ell\ell}$	Mass of dilepton system ($e\mu$)
$m_T^{\ell 1}$	Transverse mass of leading lepton and $\vec{p}_T^{\text{miss}}$
$m_T^{\ell 2}$	Transverse mass of subleading lepton and $\vec{p}_T^{\text{miss}}$
$\Delta R(\ell\ell, \ell 1)$	ΔR separation between dilepton system and leading lepton
$\Delta R(\ell\ell, \ell 2)$	ΔR separation between dilepton system and subleading lepton
$\Delta R(\ell\ell, c)$	ΔR separation between dilepton system and c jet
$\Delta\phi(\ell\ell + \vec{p}_T^{\text{miss}}, c)$	$\Delta\phi$ separation of dilepton+ $\vec{p}_T^{\text{miss}}$ system and c jet
$\Delta\phi(\ell 1, \vec{p}_T^{\text{miss}})$	$\Delta\phi$ separation of leading lepton and $\vec{p}_T^{\text{miss}}$
$\Delta\phi(\ell 2, \vec{p}_T^{\text{miss}})$	$\Delta\phi$ separation of subleading lepton and $\vec{p}_T^{\text{miss}}$
DEEPJET $CvsL$	Discriminant separating light and c jets
DEEPJET $CvsB$	Discriminant separating b and c jets
nSV	Number of secondary vertices in the event

flavor of the additional jet: H+c-bkg, H+b-bkg, and H+udsg-bkg, where the latter two terms are combined and denoted as H+notc-bkg. The generator-level flavor of the jet is determined by ghost matching [62]. Due to the presence of c jets in the H+c-bkg component, its D_{bkg} distribution has a similar shape as the cH process as shown in Fig. 4 (left). However, its $D_{\text{H-bkg}}$ shape is different. The splitting of H-bkg into three components depending on the flavor of the associated jets is used in the signal extraction described later.

Two BDT classifiers, D_{bkg} and $D_{\text{H-bkg}}$, are combined to construct the distributions used for signal extraction. The k -means [63] clustering algorithm is used to define the bin boundaries of the 2D BDT distribution, in order to preserve the information of the 2D BDT output while keeping the statistical uncertainties sufficiently small. The k -means clustering algorithm partitions data into clusters with approximately equal variance by minimizing the distance between each point and its cluster centroid. Each resulting cluster is collapsed into a single bin, preserving its approximate location in the original 2D phase space. These bins are then reordered into a one-dimensional (1D) distribution, sorted by ascending BDT score according to their 2D position ($D_{\text{H-bkg}}$ ascending first, D_{bkg} ascending second). The number of bins is optimized separately for each data-taking period and SR category to maximize sensitivity. For the $N_{\text{c-j}} = 1$ category, 40 bins are used for the 2016 and 2017 data sets, and 50 bins for 2018 data set, due to its larger size. For the $N_{\text{c-j}} > 1$ category, 30 bins are used across all years.

The background contributions after event selection are estimated from both the CRs and SRs in data and simulated events. The normalizations for single top quark, diboson, and V+jets processes are obtained from simulated events, while the dominant background ($t\bar{t}$) is deter-




Figure 4: Distributions of D_{bkg} (left) and $D_{\text{H-bkg}}$ (right) for different H boson production processes. The event counts correspond to the expected yields for 138 fb^{-1} . The splitting of H-bkg into three components is based on the flavor of the additional jet associated with the H boson. The yield of the bH process is scaled by a factor of 10, and cH by a factor of 100.

mined from data by a simultaneous fit in the SRs and CRs with unconstrained normalization factors. These normalization factors are determined separately for each data-taking period and c jet multiplicity category. The $t\bar{t}$ contribution in the $N_{\text{c-j}} = 1$ SR is constrained by the normalization factor in the *high- $m_{\ell\ell}$* CR, whereas its contribution in $N_{\text{c-j}} > 1$ SR is constrained from the *top* CR due to the different charm jet multiplicity. In the fit, only the total event yield in the *top* CR is used, while the binned $D_{\text{H-bkg}}$ distribution is used in the *high- $m_{\ell\ell}$* CR. In the SRs, the clustered 1D distributions from k -means binning are used. The bH and H+notc-bkg contributions are obtained from the simulation. Two scenarios are considered for the treatment of the H+c-bkg contribution. In the first scenario, the H+c-bkg contribution is set to the simulation expectation, within its uncertainties. In the second scenario, the H+c-bkg contribution is extracted simultaneously with cH signal, thus constrained from a fit to the data.

6 Statistical procedure and systematic uncertainties

The statistical interpretation is performed using the CMS statistical analysis tool COMBINE [64], which is based on the ROOFIT [65] and ROOSTATS [66] frameworks. Signal extraction is performed by a binned maximum likelihood fit to data in all SRs and CRs, using binned templates described in Section 5.

Systematic uncertainties affecting the results are categorized into theoretical and experimental. Furthermore, they are divided into two groups: uncertainties that only modify the normalization of the processes, and shape uncertainties that change the form of the fitted distributions. Some shape uncertainties, particularly in background templates, are poorly modeled due to the finite sample size of simulated events. To mitigate this, the relative deviations are smoothed using the LOWESS algorithm [67, 68], which preserves local structure. Smoothing is applied only to adjacent bins with correlated fluctuations in the 2D phase space.

The theoretical uncertainties are described in the following. For cH and bH processes, the FS uncertainty arises from differences in matrix-element calculations that treat the heavy quark in the proton PDF as either massive or massless. For bH, the uncertainty is derived from the

difference between the 5FS and 4FS; for cH, it comes from the difference between 4FS and 3FS. These are treated as shape uncertainties and can reach up to 30%, making them the dominant theoretical uncertainty in this analysis. A 50% normalization uncertainty is assigned to the modeling of heavy-flavor contributions in other H boson production processes (H-bkg), based on the theoretical uncertainty of the $ggH+b\bar{b}$ process [43]. Production cross section uncertainties are treated as normalization uncertainties and are taken from theoretical predictions. A 1% uncertainty is assigned to the single top quark production cross section [69–71], 3% for V+jets processes [72–76], 4% for diboson production [77–84], and 1–3% for H boson production and decay [85], depending on the production processes. The UE and PS uncertainties, stemming from generator tuning and modeling choices, are included as shape uncertainties and typically amount to 3%. The PDF uncertainties are evaluated by summing the 100 eigenvectors of the NNPDF set, contributing normalization uncertainties between 1 and 3%, and up to 6% for the cH process. Renormalization and factorization scale uncertainties, resulting from independent variations of these two scales by factors of 0.5 and 2, are included as shape uncertainties with a typical size of 1%. Lastly, $t\bar{t}$ p_T mismodeling is incorporated as a shape uncertainty, based on reweighting the top quark p_T distribution to match NNLO predictions [86–89], with effects particularly significant in signal-enriched regions, reaching up to 3%. The theoretical uncertainties are considered fully correlated across data-taking periods.

Experimental uncertainties are also assessed. Statistical uncertainties from limited simulated sample sizes are accounted for using the Barlow–Beeston-lite method [90], assigning one parameter per bin across all processes, with typical magnitudes between 5 and 10%. Jet energy scale and resolution uncertainties, propagated to jets and p_T^{miss} , are included as shape uncertainties and can reach up to 5%, with enhancements in signal-enriched regions. Jet identification uncertainties are treated as shape variations, ranging between 1 and 2% depending on the data-taking year. Charm jet identification uncertainties, derived from corrections to the c jet discriminator distributions using control samples [17], are included as shape uncertainties and range from 3 to 7%, with notable impact in signal-enriched regions. The integrated luminosity uncertainty ranges from 1.2 to 2.5% across the 2016–2018 data-taking period [29–31], with an overall uncertainty of 1.6%, and is treated as a normalization uncertainty. Lepton trigger and identification efficiencies, including uncertainties from trigger, reconstruction, identification, and isolation corrections, are also treated as normalization uncertainties and are in the range between 1 and 3%. A 1% normalization uncertainty is assigned to account for L1 trigger inefficiency during 2016–2017. Pileup correction uncertainties are evaluated by varying the total inelastic cross section by 4.6% [91], and are treated as shape uncertainties. The statistical uncertainties of simulated events are treated as uncorrelated among all data-taking periods. Experimental uncertainties from c jet identification, jet energy scale, jet energy resolution, and integrated luminosity are treated as partially correlated between data-taking years, while all other uncertainties are considered to be uncorrelated.

7 Results

The results are evaluated in two scenarios. In the first scenario, the contribution from H-bkg is fixed to its SM prediction. This corresponds to a one parameter of interest (1POI) fit, with the cH signal strength modifier μ_{cH} which is defined as the ratio of measured yield to the one predicted by the SM. In the second scenario, the normalization of the charm jet associated H boson production, $\mu_{\text{H+c-bkg}}$, is allowed to vary in the fit, resulting in a two parameter of interest (2POI) fit, with μ_{cH} and $\mu_{\text{H+c-bkg}}$ as parameters of interest. The fit is performed simultaneously in SRs and CRs using data collected between 2016 and 2018. Figure 5 displays the observed and expected distributions of data and simulated events after performing the fit in the 1POI

scenario.

The observed (expected) upper limit on the cH signal strength at 95% CL is 1065 (506) times the SM value in the 1POI fit. In the 2POI fit, the corresponding upper limit is 1804 (1097) times the SM prediction. The measured signal strength for the cH process in the 1POI fit is found to be $\mu_{\text{cH}} = 390^{+380}_{-270}$, which is compatible with the SM within 1.5 standard deviations. The results from different data-taking periods are summarized in Fig. 6. A strong correlation of -87% is observed between H+c-bkg and cH productions in the 2POI fit, leading to a weaker constraint of $\mu_{\text{cH}} = 230^{+1000}_{-640}$, as shown in Fig. 7. Similarly, the bH production is highly correlated to cH, with a size of -89% observed in the fit. The sensitivity of this analysis is primarily limited by statistical uncertainties in data, with nonnegligible contributions from theoretical uncertainties, such as the FS choice of cH production and heavy-flavor modeling in ggH production. In view of the high-luminosity LHC, it can be expected that the theoretical uncertainty would be the dominant uncertainty of this analysis. The impact of each uncertainty on the measurement of μ_{cH} is summarized in Table 3.

The resulting upper limit on μ_{cH} is translated into a constraint on κ_c , using the flat direction assumption [20, 92]. Under this assumption, the modifier of all other H couplings κ_H changes according to κ_c , such as to keep the signal strengths of other H boson processes at unity. The observed (expected) interval at 95% CL is constrained to $|\kappa_c| < 211$ (95).

A combined analysis of the 1POI fit with the previous search in the diphoton decay channel [21] is performed. Common experimental and theoretical uncertainties related to H boson production cross sections are treated as correlated. The combination improves the expected upper limit on μ_{cH} at 95% CL to 281, while the observed limit is 257, as shown in Fig. 8. This leads to constraints on the observed (expected) value of $|\kappa_c| < 47$ (51) at 95% CL.

Table 3: Relative impact (in percent) of individual uncertainty sources on the measurement of μ_{cH} expressed as the change in the one standard deviation when each source is fixed. Statistical uncertainties are derived by fixing all constrained nuisance parameters.

Uncertainty source	1POI	2POI
Theoretical		
cH modeling	22.1	10.9
H-bkg	21.4	10.0
Other backgrounds	4.5	3.5
Experimental		
MC statistical (bin by bin)	6.2	9.1
Pileup correction	4.3	1.2
Lepton efficiencies	4.6	3.2
Jet energy scale and resolution	6.0	7.7
Charm jet identification	5.9	4.1
$t\bar{t}$ normalization	5.6	2.0
Statistical uncertainty	60.0	77.1




Figure 5: Observed and expected distributions of one-parameter-of-interest scenario after performing the fit in signal and control regions. The upper left plot shows the distribution of events as a function of $\log(S/B)$ in the $N_{c-j} = 1$ signal region, while the upper right shows the $N_{c-j} > 1$ signal region, where S and B are the signal and background yields, respectively. The lower left plot shows the distribution of D_{bkg} used in the $high\text{-}m_{\ell\ell}$ control region, while the normalization factor of the top CR is shown in the lower right. Within each plot, the upper panel provides the number of events from simulation and data in logarithmic scale, the middle panel displays the fractional contribution of each background component to the total expected yield in each bin, and the lower panel provides the ratio of data to simulated background events. The red lines represent background plus signal with $\mu_{cH} = 390$ divided by background. The hashed area shows the sum of systematic and statistical uncertainties.




Figure 6: Upper limits of μ_{CH} at 95% CL for each data-taking period, and the combination of the periods. The light blue (red) bars show the 1 (2) standard deviation expected result of the 1POI fit with fixing the H+c-bkg contribution to the SM prediction. The green (yellow) bars show the 1 (2) standard deviation expected result of the 2POI fit with floating H+c-bkg contribution. The blue circles represent the median of the expected limit, while the black circles represent the observed limit.




Figure 7: Two-dimensional likelihood contour of μ_{cH} and $\mu_{H+c-bkg}$. The color scale represents twice the negative log likelihood difference with respect to the best fit point. The observed 95% CL (dashed) and 68% CL (solid) contours are shown in black lines, and the best fit point as a black cross. The SM expectation is marked by a red diamond. The kink in the contour arises from a local minimum in the likelihood, driven by the shape uncertainty associated with the flavor scheme used in the cH process modeling.




Figure 8: Upper limits of μ_{CH} at 95% CL of the combined analysis with the previous result in the diphoton channel [21] using the 1POI fit. The light blue (red) bars show the 1 (2) standard deviation range of the expected results. The blue circles represent the median of the expected limit, while the black circles represent the observed limit.

8 Summary

The first search for the associated production of a charm quark and a Higgs boson (cH) with the Higgs boson decays to a pair of W bosons has been presented. Decays of W bosons to $e\nu\mu\nu$ are considered. The search is based on the data collected from 2016 to 2018 with the CMS detector at the LHC, corresponding to an integrated luminosity of 138 fb^{-1} . The observed (expected) upper limit at 95% confidence level (CL) on the ratio of the measured production cross section with respect to the value expected from the standard model (SM) for cH production μ_{cH} is set at 1065 (506) times the SM prediction. This search provides an alternative probe of the Yukawa coupling between the Higgs boson and charm quark. When combined with the previous search for cH in the diphoton decay channel, the limits are interpreted as observed (expected) constraints on the Yukawa coupling modifier of the Higgs boson to the charm quark, yielding $|\kappa_c| < 47$ (51) at 95% CL.

Acknowledgments

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centers and personnel of the Worldwide LHC Computing Grid and other centers for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector, and the supporting computing infrastructure provided by the following funding agencies: SC (Armenia), BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MoST, and NSFC (China); MINCIENCIAS (Colombia); MSES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); ERC PRG, TARISTU24-TK10 and MoER TK202 (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); SRNSF (Georgia); BMFTR, DFG, and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIT and NRF (Republic of Korea); MES (Latvia); LMTLT (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MES, NSC, and NAWA (Poland); FCT (Portugal); MESTD (Serbia); MICIU/AEI and PCTI (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHESI and NSTDA (Thailand); TUBITAK and TENMAK (Türkiye); NASU (Ukraine); STFC (United Kingdom); DOE and NSF (USA).

Individuals have received support from the Marie-Curie program and the European Research Council and Horizon 2020 Grant, contract Nos. 675440, 724704, 752730, 758316, 765710, 824093, 101115353, 101002207, 101001205, and COST Action CA16108 (European Union); the Leventis Foundation; the Alfred P. Sloan Foundation; the Alexander von Humboldt Foundation; the Science Committee, project no. 22rl-037 (Armenia); the Fonds pour la Formation à la Recherche dans l'Industrie et dans l'Agriculture (FRIA-Belgium); the Beijing Municipal Science & Technology Commission, No. Z191100007219010, the Fundamental Research Funds for the Central Universities, the Ministry of Science and Technology of China under Grant No. 2023YFA1605804, and the Natural Science Foundation of China under Grant No. 12061141002 (China); the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Shota Rustaveli National Science Foundation, grant FR-22-985 (Georgia); the Deutsche Forschungsgemeinschaft (DFG), among others, under Germany's Excellence Strategy – EXC 2121 “Quantum Universe” – 390833306, and under project number 400140256 - GRK2497; the Hellenic

Foundation for Research and Innovation (HFRI), Project Number 2288 (Greece); the Hungarian Academy of Sciences, the New National Excellence Program - ÚNKP, the NKFIH research grants K 131991, K 133046, K 138136, K 143460, K 143477, K 146913, K 146914, K 147048, 2020-2.2.1-ED-2021-00181, TKP2021-NKTA-64, and 2021-4.1.2-NEMZ.KI-2024-00036 (Hungary); the Council of Science and Industrial Research, India; ICSC – National Research Center for High Performance Computing, Big Data and Quantum Computing, FAIR – Future Artificial Intelligence Research, and CUP I53D23001070006 (Mission 4 Component 1), funded by the NextGenerationEU program (Italy); the Latvian Council of Science; the Ministry of Education and Science, project no. 2022/WK/14, and the National Science Center, contracts Opus 2021/41/B/ST2/01369, 2021/43/B/ST2/01552, 2023/49/B/ST2/03273, and the NAWA contract BPN/PPO/2021/1/00011 (Poland); the Fundação para a Ciência e a Tecnologia, grant CEECIND/01334/2018 (Portugal); the National Priorities Research Program by Qatar National Research Fund; MICIU/AEI/10.13039/501100011033, ERDF/EU, “European Union NextGenerationEU/PRTR”, and Programa Severo Ochoa del Principado de Asturias (Spain); the Chulalongkorn Academic into Its 2nd Century Project Advancement Project, and the National Science, Research and Innovation Fund via the Program Management Unit for Human Resources & Institutional Development, Research and Innovation, grant B39G680009 (Thailand); the Kavli Foundation; the Nvidia Corporation; the SuperMicro Corporation; the Welch Foundation, contract C-1845; and the Weston Havens Foundation (USA).

References

- [1] CMS Collaboration, “Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC”, *Phys. Lett. B* **716** (2012) 30, doi:10.1016/j.physletb.2012.08.021, arXiv:1207.7235.
- [2] ATLAS Collaboration, “Observation of a new particle in the search for the standard model higgs boson with the ATLAS detector at the LHC”, *Phys. Lett. B* **716** (2012) 1, doi:10.1016/j.physletb.2012.08.020, arXiv:1207.7214.
- [3] CMS Collaboration, “Observation of a new boson with mass near 125 GeV in pp collisions at $\sqrt{s} = 7$ and 8 TeV”, *JHEP* **06** (2013) 081, doi:10.1007/JHEP06(2013)081, arXiv:1303.4571.
- [4] CMS Collaboration, “A portrait of the Higgs boson by the CMS experiment ten years after the discovery”, *Nature* **607** (2022) 60, doi:10.1038/s41586-022-04892-x, arXiv:2207.00043. [Erratum: *Nature* 623 (2023)].
- [5] ATLAS Collaboration, “A detailed map of Higgs boson interactions by the ATLAS experiment ten years after the discovery”, *Nature* **607** (2022) 52, doi:10.1038/s41586-022-04893-w, arXiv:2207.00092. [Erratum: *Nature* 612, E24 (2022)].
- [6] CMS Collaboration, “Measurement of simplified template cross sections of the Higgs boson produced in association with W or Z bosons in the $H \rightarrow b\bar{b}$ decay channel in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *Phys. Rev. D* **109** (2024) 092011, doi:10.1103/PhysRevD.109.092011, arXiv:2312.07562.
- [7] ATLAS Collaboration, “Measurements of WH and ZH production in the $H \rightarrow b\bar{b}$ decay channel in pp collisions at 13 TeV with the ATLAS detector”, *Eur. Phys. J. C* **81** (2021) 178, doi:10.1140/epjc/s10052-020-08677-2, arXiv:2007.02873.

- [8] ATLAS Collaboration, “Measurements of WH and ZH production with Higgs boson decays into bottom quarks and direct constraints on the charm Yukawa coupling in 13 TeV pp collisions with the ATLAS detector”, *JHEP* **04** (2025) 075, doi:10.1007/JHEP04(2025)075, arXiv:2410.19611.
- [9] ATLAS Collaboration, “Measurements of Higgs boson production cross-sections in the $H \rightarrow \tau^+\tau^-$ decay channel in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **08** (2022) 175, doi:10.1007/JHEP08(2022)175, arXiv:2201.08269.
- [10] CMS Collaboration, “Measurements of Higgs boson production in the decay channel with a pair of τ leptons in proton–proton collisions at $\sqrt{s} = 13$ TeV”, *Eur. Phys. J. C* **83** (2023) 562, doi:10.1140/epjc/s10052-023-11452-8, arXiv:2204.12957.
- [11] CMS Collaboration, “Evidence for Higgs boson decay to a pair of muons”, *JHEP* **01** (2021) 148, doi:10.1007/JHEP01(2021)148, arXiv:2009.04363.
- [12] ATLAS Collaboration, “A search for the dimuon decay of the standard model Higgs boson with the ATLAS detector”, *Phys. Lett. B* **812** (2021) 135980, doi:10.1016/j.physletb.2020.135980, arXiv:2007.07830.
- [13] CMS Collaboration, “Search for Higgs boson and observation of Z boson through their decay into a charm quark-antiquark pair in boosted topologies in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *Phys. Rev. Lett.* **131** (2023) 041801, doi:10.1103/PhysRevLett.131.041801, arXiv:2211.14181.
- [14] ATLAS Collaboration, “Direct constraint on the Higgs-charm coupling from a search for Higgs boson decays into charm quarks with the ATLAS detector”, *Eur. Phys. J. C* **82** (2022) 717, doi:10.1140/epjc/s10052-022-10588-3, arXiv:2201.11428.
- [15] CMS Collaboration, “Search for Higgs boson decay to a charm quark-antiquark pair in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *Phys. Rev. Lett.* **131** (2023) 061801, doi:10.1103/PhysRevLett.131.061801, arXiv:2205.05550.
- [16] E. Bols et al., “Jet flavour classification using DeepJet”, *JINST* **15** (2020) P12012, doi:10.1088/1748-0221/15/12/P12012, arXiv:2008.10519.
- [17] CMS Collaboration, “A new calibration method for charm jet identification validated with proton-proton collision events at $\sqrt{s} = 13$ TeV”, *JINST* **17** (2022) P03014, doi:10.1088/1748-0221/17/03/P03014, arXiv:2111.03027.
- [18] CMS Collaboration, “Combination and interpretation of differential Higgs boson production cross sections in proton-proton collisions at $\sqrt{s} = 13$ TeV”, 2025. arXiv:2504.13081. Submitted to *JHEP*, doi:10.48550/arXiv.2504.13081.
- [19] N. M. Coyle, C. E. M. Wagner, and V. Wei, “Bounding the charm Yukawa coupling”, *Phys. Rev. D* **100** (2019) 073013, doi:10.1103/PhysRevD.100.073013, arXiv:1905.09360.
- [20] I. Brivio, F. Goertz, and G. Isidori, “Probing the charm quark Yukawa coupling in Higgs+Charm production”, *Phys. Rev. Lett.* **115** (2015) 211801, doi:10.1103/PhysRevLett.115.211801, arXiv:1507.02916.
- [21] CMS Collaboration, “Search for the associated production of a Higgs boson with a charm quark in the diphoton decay channel in pp collisions at $\sqrt{s} = 13$ TeV”, 2025. arXiv:2503.08797. Submitted to *JHEP*, doi:10.48550/arXiv.2503.08797.

-
- [22] ATLAS Collaboration, “Search for the associated production of charm quarks and a Higgs boson decaying into a photon pair with the ATLAS detector”, *JHEP* **02** (2025) 045, doi:10.1007/JHEP02(2025)045, arXiv:2407.15550.
- [23] HEPData record for this analysis, 2025. doi:10.17182/hepdata.123456.
- [24] CMS Collaboration, “The CMS experiment at the CERN LHC”, *JINST* **3** (2008) S08004, doi:10.1088/1748-0221/3/08/S08004.
- [25] CMS Collaboration, “Development of the CMS detector for the CERN LHC Run 3”, *JINST* **19** (2024) P05064, doi:10.1088/1748-0221/19/05/P05064, arXiv:2309.05466.
- [26] CMS Collaboration, “Performance of the CMS Level-1 trigger in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JINST* **15** (2020) P10017, doi:10.1088/1748-0221/15/10/P10017, arXiv:2006.10165.
- [27] CMS Collaboration, “The CMS trigger system”, *JINST* **12** (2017) P01020, doi:10.1088/1748-0221/12/01/P01020, arXiv:1609.02366.
- [28] CMS Collaboration, “Performance of the CMS high-level trigger during LHC Run 2”, *JINST* **19** (2024) P11021, doi:10.1088/1748-0221/19/11/P11021, arXiv:2410.17038.
- [29] CMS Collaboration, “Precision luminosity measurement in proton-proton collisions at $\sqrt{s} = 13$ TeV in 2015 and 2016 at CMS”, *Eur. Phys. J. C* **81** (2021) 800, doi:10.1140/epjc/s10052-021-09538-2, arXiv:2104.01927.
- [30] CMS Collaboration, “CMS luminosity measurement for the 2017 data-taking period at $\sqrt{s} = 13$ TeV”, CMS Physics Analysis Summary CMS-PAS-LUM-17-004, 2018.
- [31] CMS Collaboration, “CMS luminosity measurement for the 2018 data-taking period at $\sqrt{s} = 13$ TeV”, CMS Physics Analysis Summary CMS-PAS-LUM-18-002, 2019.
- [32] J. Alwall et al., “The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations”, *JHEP* **07** (2014) 079, doi:10.1007/JHEP07(2014)079, arXiv:1405.0301.
- [33] R. Frederix et al., “The automation of next-to-leading order electroweak calculations”, *JHEP* **07** (2018) 185, doi:10.1007/JHEP11(2021)085, arXiv:1804.10017. [Erratum: doi:10.1007/JHEP11(2021)085].
- [34] M. Wiesemann et al., “Higgs production in association with bottom quarks”, *JHEP* **02** (2015) 132, doi:10.1007/JHEP02(2015)132, arXiv:1409.5301.
- [35] R. Frederix and S. Frixione, “Merging meets matching in MC@NLO”, *JHEP* **12** (2012) 061, doi:10.1007/JHEP12(2012)061, arXiv:1209.6215.
- [36] P. Nason, “A new method for combining NLO QCD with shower Monte Carlo algorithms”, *JHEP* **11** (2004) 040, doi:10.1088/1126-6708/2004/11/040, arXiv:hep-ph/0409146.
- [37] P. Nason and C. Oleari, “NLO Higgs boson production via vector-boson fusion matched with shower in POWHEG”, *JHEP* **02** (2010) 037, doi:10.1007/JHEP02(2010)037, arXiv:0911.5299.

- [38] S. Frixione, P. Nason, and C. Oleari, “Matching NLO QCD computations with parton shower simulations: the POWHEG method”, *JHEP* **11** (2007) 070, doi:10.1088/1126-6708/2007/11/070, arXiv:0709.2092.
- [39] S. Alioli, P. Nason, C. Oleari, and E. Re, “A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX”, *JHEP* **06** (2010) 043, doi:10.1007/JHEP06(2010)043, arXiv:1002.2581.
- [40] E. Bagnaschi, G. Degrandi, P. Slavich, and A. Vicini, “Higgs production via gluon fusion in the POWHEG approach in the SM and in the MSSM”, *JHEP* **02** (2012) 088, doi:10.1007/JHEP02(2012)088, arXiv:1111.2854.
- [41] G. Luisoni, P. Nason, C. Oleari, and F. Tramontano, “ $HW^\pm/HZ+0$ and 1 jet at NLO with the POWHEG BOX interfaced to GoSam and their merging within MiNLO”, *JHEP* **10** (2013) 083, doi:10.1007/JHEP10(2013)083, arXiv:1306.2542.
- [42] H. B. Hartanto, B. Jager, L. Reina, and D. Wackeroth, “Higgs boson production in association with top quarks in the POWHEG BOX”, *Phys. Rev. D* **91** (2015) 094003, doi:10.1103/PhysRevD.91.094003, arXiv:1501.04498.
- [43] S. Manzoni et al., “Taming a leading theoretical uncertainty in HH measurements via accurate simulations for bbH production”, *JHEP* **09** (2023) 179, doi:10.1007/JHEP09(2023)179, arXiv:2307.09992.
- [44] Sherpa Collaboration, “Event generation with Sherpa 2.2”, *SciPost Phys.* **7** (2019) 034, doi:10.21468/SciPostPhys.7.3.034, arXiv:1905.09127.
- [45] T. Sjöstrand et al., “An introduction to PYTHIA 8.2”, *Comput. Phys. Commun.* **191** (2015) 159, doi:10.1016/j.cpc.2015.01.024, arXiv:1410.3012.
- [46] NNPDF Collaboration, “Parton distributions from high-precision collider data”, *Eur. Phys. J. C* **77** (2017) 663, doi:10.1140/epjc/s10052-017-5199-5, arXiv:1706.00428.
- [47] CMS Collaboration, “Extraction and validation of a new set of CMS PYTHIA8 tunes from underlying-event measurements”, *Eur. Phys. J. C* **80** (2020) 4, doi:10.1140/epjc/s10052-019-7499-4, arXiv:1903.12179.
- [48] GEANT4 Collaboration, “GEANT4—a simulation toolkit”, *Nucl. Instrum. Meth. A* **506** (2003) 250, doi:10.1016/S0168-9002(03)01368-8.
- [49] CMS Collaboration, “Technical proposal for the Phase-II upgrade of the Compact Muon Solenoid”, CMS Technical Proposal CERN-LHCC-2015-010, CMS-TDR-15-02, 2015.
- [50] CMS Collaboration, “Particle-flow reconstruction and global event description with the CMS detector”, *JINST* **12** (2017) P10003, doi:10.1088/1748-0221/12/10/P10003, arXiv:1706.04965.
- [51] CMS Collaboration, “Electron and photon reconstruction and identification with the CMS experiment at the CERN LHC”, *JINST* **16** (2021) P05014, doi:10.1088/1748-0221/16/05/P05014, arXiv:2012.06888.
- [52] CMS Collaboration, “ECAL 2016 refined calibration and Run2 summary plots”, CMS Detector Performance Note CMS-DP-2020-021, 2020.

-
- [53] CMS Collaboration, “Performance of the CMS muon detector and muon reconstruction with proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JINST* **13** (2018) P06015, doi:10.1088/1748-0221/13/06/P06015, arXiv:1804.04528.
- [54] M. Cacciari, G. P. Salam, and G. Soyez, “The anti- k_T jet clustering algorithm”, *JHEP* **04** (2008) 063, doi:10.1088/1126-6708/2008/04/063, arXiv:0802.1189.
- [55] M. Cacciari, G. P. Salam, and G. Soyez, “FastJet user manual”, *Eur. Phys. J. C* **72** (2012) 1896, doi:10.1140/epjc/s10052-012-1896-2, arXiv:1111.6097.
- [56] CMS Collaboration, “Pileup mitigation at CMS in 13 TeV data”, *JINST* **15** (2020) P09018, doi:10.1088/1748-0221/15/09/P09018, arXiv:2003.00503.
- [57] CMS Collaboration, “Jet energy scale and resolution in the CMS experiment in pp collisions at 8 TeV”, *JINST* **12** (2017) P02014, doi:10.1088/1748-0221/12/02/P02014, arXiv:1607.03663.
- [58] CMS Collaboration, “Performance of the DeepJet b tagging algorithm using 41.9/fb of data from proton-proton collisions at 13 TeV with Phase 1 CMS detector”, CMS Detector Performance Note CMS-DP-2018-058, 2018.
- [59] CMS Collaboration, “Performance summary of AK4 jet charm tagging with the CMS Run2 legacy dataset”, CMS Detector Performance Note CMS-DP-2023-006, 2023.
- [60] CMS Collaboration, “Performance of missing transverse momentum reconstruction in proton-proton collisions at $\sqrt{s} = 13$ TeV using the CMS detector”, *JINST* **14** (2019) P07004, doi:10.1088/1748-0221/14/07/P07004, arXiv:1903.06078.
- [61] T. Chen and C. Guestrin, “XGBoost: A scalable tree boosting system”, in *Proc. 22nd ACM SIGKDD Int. Conf. on Knowledge Discovery and Data Mining*, KDD ’16, p. 785. ACM, 2016. doi:10.1145/2939672.2939785.
- [62] M. Cacciari and G. P. Salam, “Pileup subtraction using jet areas”, *Phys. Lett. B* **659** (2008) 119, doi:10.1016/j.physletb.2007.09.077, arXiv:0707.1378.
- [63] D. Sculley, “Web-scale k-means clustering”, in *Proc. 19th Int. Conf. on World Wide Web*, WWW ’10, p. 1177. Association for Computing Machinery, 2010. doi:10.1145/1772690.1772862.
- [64] CMS Collaboration, “The CMS statistical analysis and combination tool: Combine”, *Comput. Softw. Big Sci.* **8** (2024) 19, doi:10.1007/s41781-024-00121-4, arXiv:2404.06614.
- [65] W. Verkerke and D. Kirkby, “The ROOFIT toolkit for data modeling”, in *Proc. 13th Int. Conf. on Computing in High Energy and Nuclear Physics (CHEP 2003): La Jolla CA, United States, March 24–28*. 2003. arXiv:physics/0306116.
- [66] L. Moneta et al., “The RooStats project”, *PoS ACAT2010* (2010) 057, doi:10.22323/1.093.0057, arXiv:1009.1003.
- [67] W. S. Cleveland, “Robust locally weighted regression and smoothing scatterplots”, *J. Am. Stat. Assoc.* **74** (1979) 829, doi:10.1080/01621459.1979.10481038.
- [68] W. S. Cleveland and S. J. Devlin, “Locally weighted regression: An approach to regression analysis by local fitting”, *J. Am. Stat. Assoc.* **83** (1988) 596, doi:10.1080/01621459.1988.10478639.

- [69] J. Campbell, T. Neumann, and Z. Sullivan, “Single-top-quark production in the t -channel at NNLO”, *JHEP* **02** (2021) 040, doi:10.1007/JHEP02(2021)040, arXiv:2012.01574.
- [70] PDF4LHC Working Group Collaboration, “The PDF4LHC21 combination of global PDF fits for the LHC Run III”, *J. Phys. G* **49** (2022) 080501, doi:10.1088/1361-6471/ac7216, arXiv:2203.05506.
- [71] N. Kidonakis and N. Yamanaka, “Higher-order corrections for tW production at high-energy hadron colliders”, *JHEP* **05** (2021) 278, doi:10.1007/JHEP05(2021)278, arXiv:2102.11300.
- [72] K. Melnikov and F. Petriello, “Electroweak gauge boson production at hadron colliders through $\mathcal{O}(\alpha_s^2)$ ”, *Phys. Rev. D* **74** (2006) 114017, doi:10.1103/PhysRevD.74.114017, arXiv:hep-ph/0609070.
- [73] S. Catani et al., “Vector boson production at hadron colliders: a fully exclusive QCD calculation at NNLO”, *Phys. Rev. Lett.* **103** (2009) 082001, doi:10.1103/PhysRevLett.103.082001, arXiv:0903.2120.
- [74] J. Baglio, C. Duhr, B. Mistlberger, and R. Szafron, “Inclusive production cross sections at $N^3\text{LO}$ ”, *JHEP* **12** (2022) 066, doi:10.1007/JHEP12(2022)066, arXiv:2209.06138.
- [75] C. Anastasiou, L. J. Dixon, K. Melnikov, and F. Petriello, “High precision QCD at hadron colliders: Electroweak gauge boson rapidity distributions at NNLO”, *Phys. Rev. D* **69** (2004) 094008, doi:10.1103/PhysRevD.69.094008, arXiv:hep-ph/0312266.
- [76] S. Dittmaier, A. Huss, and C. Schwinn, “Mixed QCD-electroweak $\mathcal{O}(\alpha_s\alpha)$ corrections to Drell-Yan processes in the resonance region: pole approximation and non-factorizable corrections”, *Nucl. Phys. B* **885** (2014) 318, doi:10.1016/j.nuclphysb.2014.05.027, arXiv:1403.3216.
- [77] J. M. Campbell and R. K. Ellis, “MCFM for the Tevatron and the LHC”, *Nucl. Phys. B Proc. Suppl.* **205-206** (2010) 10, doi:10.1016/j.nuclphysbps.2010.08.011, arXiv:1007.3492.
- [78] Caola, Fabrizio and Melnikov, Kirill and Rötsch, Raoul and Tancredi, Lorenzo, “QCD corrections to W^+W^- production through gluon fusion”, *Phys. Lett. B* **754** (2016) 275, doi:10.1016/j.physletb.2016.01.046, arXiv:1511.08617.
- [79] T. Melia, P. Nason, R. Röntsch, and G. Zanderighi, “ W^+W^- , WZ and ZZ production in the POWHEG BOX”, *JHEP* **11** (2011) 078, doi:10.1007/JHEP11(2011)078, arXiv:1107.5051.
- [80] J. M. Campbell, R. K. Ellis, and C. Williams, “Vector boson pair production at the LHC”, *JHEP* **07** (2011) 018, doi:10.1007/JHEP07(2011)018, arXiv:1105.0020.
- [81] F. Cascioli et al., “ ZZ production at hadron colliders in NNLO QCD”, *Phys. Lett. B* **735** (2014) 311, doi:10.1016/j.physletb.2014.06.056, arXiv:1405.2219.
- [82] M. Grazzini, S. Kallweit, and D. Rathlev, “ ZZ production at the LHC: fiducial cross sections and distributions in NNLO QCD”, *Phys. Lett. B* **750** (2015) 407, doi:10.1016/j.physletb.2015.09.055, arXiv:1507.06257.

-
- [83] S. Kallweit and M. Wiesemann, “ZZ production at the LHC: NNLO predictions for $2\ell 2\nu$ and 4ℓ signatures”, *Phys. Lett. B* **786** (2018) 382, doi:10.1016/j.physletb.2018.10.016, arXiv:1806.05941.
- [84] J. M. Lindert et al., “Precise predictions for V+jets dark matter backgrounds”, *Eur. Phys. J. C* **77** (2017) 829, doi:10.1140/epjc/s10052-017-5389-1, arXiv:1705.04664.
- [85] LHC Higgs Cross Section Working Group, “Handbook of LHC Higgs cross sections: 4. deciphering the nature of the Higgs sector”, 2016. arXiv:1610.07922. doi:10.23731/CYRM-2017-002.
- [86] N. Kidonakis, “NNLL threshold resummation for top-pair and single-top production”, *Phys. Part. Nucl.* **45** (2014) 714, doi:10.1134/S1063779614040091, arXiv:1210.7813.
- [87] S. Catani et al., “Top-quark pair production at the LHC: Fully differential QCD predictions at NNLO”, *JHEP* **07** (2019) 100, doi:10.1007/JHEP07(2019)100, arXiv:1906.06535.
- [88] M. Czakon, D. Heymes, and A. Mitov, “High-precision differential predictions for top-quark pairs at the LHC”, *Phys. Rev. Lett.* **116** (2016) 082003, doi:10.1103/PhysRevLett.116.082003, arXiv:1511.00549.
- [89] M. Czakon et al., “Top-pair production at the LHC through NNLO QCD and NLO EW”, *JHEP* **10** (2017) 186, doi:10.1007/JHEP10(2017)186, arXiv:1705.04105.
- [90] J. S. Conway, “Incorporating nuisance parameters in likelihoods for multisource spectra”, in *PHYSTAT 2011*, p. 115. 2011. arXiv:1103.0354. doi:10.5170/CERN-2011-006.115.
- [91] CMS Collaboration, “Measurement of the inelastic proton-proton cross section at $\sqrt{s} = 13$ TeV”, *JHEP* **07** (2018) 161, doi:10.1007/JHEP07(2018)161, arXiv:1802.02613.
- [92] J. A. Aguilar-Saavedra, J. M. Cano, and J. M. No, “More light on Higgs flavor at the LHC: Higgs boson couplings to light quarks through $h+\gamma$ production”, *Phys. Rev. D* **103** (2021) 095023, doi:10.1103/PhysRevD.103.095023, arXiv:2008.12538.

A The CMS Collaboration

Yerevan Physics Institute, Yerevan, Armenia

A. Hayrapetyan, V. Makarenko , A. Tumasyan¹ 

Institut für Hochenergiephysik, Vienna, Austria

W. Adam , J.W. Andrejkovic, L. Benato , T. Bergauer , M. Dragicevic , C. Giordano, P.S. Hussain , M. Jeitler² , N. Krammer , A. Li , D. Liko , M. Matthewman, I. Mikulec , J. Schieck² , R. Schöffbeck² , D. Schwarz , M. Shooshtari, M. Sonawane , W. Waltenberger , C.-E. Wulz² 

Universiteit Antwerpen, Antwerpen, Belgium

T. Janssen , H. Kwon , D. Ocampo Henao , T. Van Laer , P. Van Mechelen 

Vrije Universiteit Brussel, Brussel, Belgium

J. Bierkens , N. Breugelmans, J. D'Hondt , S. Dansana , A. De Moor , M. Delcourt , F. Heyen, Y. Hong , P. Kashko , S. Lowette , I. Makarenko , D. Müller , J. Song , S. Tavernier , M. Tytgat³ , G.P. Van Onsem , S. Van Putte , D. Vannerom 

Université Libre de Bruxelles, Bruxelles, Belgium

B. Bilin , B. Clerbaux , A.K. Das, I. De Bruyn , G. De Lentdecker , H. Evard , L. Favart , P. Gianneios , A. Khalilzadeh, F.A. Khan , A. Malara , M.A. Shahzad, L. Thomas , M. Vanden Bemden , C. Vander Velde , P. Vanlaer , F. Zhang 

Ghent University, Ghent, Belgium

M. De Coen , D. Dobur , G. Gokbulut , J. Knolle , D. Marckx , K. Skovpen , A.M. Tomaru, N. Van Den Bossche , J. van der Linden , J. Vandenbroeck , L. Wezenbeek 

Université Catholique de Louvain, Louvain-la-Neuve, Belgium

S. Bein , A. Benecke , A. Bethani , G. Bruno , A. Cappati , J. De Favereau De Jeneret , C. Delaere , A. Giammanco , A.O. Guzel , V. Lemaitre, J. Lidrych , P. Malek , P. Mastrapasqua , S. Turkcapar 

Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro, Brazil

G.A. Alves , M. Barroso Ferreira Filho , E. Coelho , C. Hensel , T. Menezes De Oliveira , C. Mora Herrera⁴ , P. Rebello Teles , M. Soeiro , E.J. Tonelli Manganote⁵ , A. Vilela Pereira⁴ 

Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

W.L. Aldá Júnior , H. Brandao Malbouisson , W. Carvalho , J. Chinellato⁶ , M. Costa Reis , E.M. Da Costa , G.G. Da Silva⁷ , D. De Jesus Damiao , S. Fonseca De Souza , R. Gomes De Souza , S. S. Jesus , T. Laux Kuhn⁷ , M. Macedo , K. Mota Amarilo , L. Mundim , H. Nogima , J.P. Pinheiro , A. Santoro , A. Sznajder , M. Thiel , F. Torres Da Silva De Araujo⁸ 

Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo, Brazil

C.A. Bernardes⁷ , F. Damas , T.R. Fernandez Perez Tomei , E.M. Gregores , B. Lopes Da Costa , I. Maietto Silverio , P.G. Mercadante , S.F. Novaes , B. Orzari , Sandra S. Padula , V. Scheurer

Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

A. Aleksandrov , G. Antchev , P. Danev, R. Hadjiiska , P. Iaydjiev , M. Shopova , G. Sultanov 

University of Sofia, Sofia, Bulgaria

A. Dimitrov , L. Litov , B. Pavlov , P. Petkov , A. Petrov 

Instituto De Alta Investigación, Universidad de Tarapacá, Casilla 7 D, Arica, Chile

S. Keshri , D. Laroze , S. Thakur 

Universidad Tecnica Federico Santa Maria, Valparaiso, Chile

W. Brooks 

Beihang University, Beijing, China

T. Cheng , T. Javaid , L. Wang , L. Yuan 

Department of Physics, Tsinghua University, Beijing, China

Z. Hu , Z. Liang, J. Liu, X. Wang 

Institute of High Energy Physics, Beijing, China

G.M. Chen⁹ , H.S. Chen⁹ , M. Chen⁹ , Y. Chen , Q. Hou , X. Hou, F. Iemmi , C.H. Jiang, A. Kapoor¹⁰ , H. Liao , G. Liu , Z.-A. Liu¹¹ , J.N. Song¹¹, S. Song, J. Tao , C. Wang⁹, J. Wang , H. Zhang , J. Zhao 

State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

A. Agapitos , Y. Ban , A. Carvalho Antunes De Oliveira , S. Deng , B. Guo, Q. Guo, C. Jiang , A. Levin , C. Li , Q. Li , Y. Mao, S. Qian, S.J. Qian , X. Qin, C. Quaranta , X. Sun , D. Wang , J. Wang, H. Yang, M. Zhang, Y. Zhao, C. Zhou 

State Key Laboratory of Nuclear Physics and Technology, Institute of Quantum Matter, South China Normal University, Guangzhou, China

S. Yang 

Sun Yat-Sen University, Guangzhou, China

Z. You 

University of Science and Technology of China, Hefei, China

K. Jaffel , N. Lu 

Nanjing Normal University, Nanjing, China

G. Bauer^{12,13}, Z. Cui¹³, B. Li¹⁴, H. Wang , K. Yi¹⁵ , J. Zhang 

Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) - Fudan University, Shanghai, China

Y. Li

Zhejiang University, Hangzhou, Zhejiang, China

Z. Lin , C. Lu , M. Xiao¹⁶ 

Universidad de Los Andes, Bogota, Colombia

C. Avila , D.A. Barbosa Trujillo , A. Cabrera , C. Florez , J. Fraga , J.A. Reyes Vega

Universidad de Antioquia, Medellin, Colombia

C. Rendón , M. Rodriguez , A.A. Ruales Barbosa , J.D. Ruiz Alvarez 

University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia

N. Godinovic , D. Lelas , A. Sculac 

University of Split, Faculty of Science, Split, Croatia

M. Kovac , A. Petkovic , T. Sculac 

Institute Rudjer Boskovic, Zagreb, Croatia

P. Bargassa , V. Brigljevic , B.K. Chitroda , D. Ferencek , K. Jakovcic, A. Starodumov , T. Susa 

University of Cyprus, Nicosia, Cyprus

A. Attikis , K. Christoforou , A. Hadjiagapiou, C. Leonidou , C. Nicolaou, L. Paizanos , F. Ptochos , P.A. Razis , H. Rykaczewski, H. Saka , A. Stepenov 

Charles University, Prague, Czech Republic

M. Finger[†] , M. Finger Jr. 

Escuela Politecnica Nacional, Quito, Ecuador

E. Ayala 

Universidad San Francisco de Quito, Quito, Ecuador

E. Carrera Jarrin 

Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt

S. Elgammal¹⁷, A. Ellithi Kamel¹⁸ 

Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum, Egypt

M.A. Mahmoud , Y. Mohammed 

National Institute of Chemical Physics and Biophysics, Tallinn, Estonia

K. Ehataht , M. Kadastik, T. Lange , C. Nielsen , J. Pata , M. Raidal , N. Seeba , L. Tani 

Department of Physics, University of Helsinki, Helsinki, Finland

A. Milieva , K. Osterberg , M. Voutilainen 

Helsinki Institute of Physics, Helsinki, Finland

N. Bin Norjoharuddeen , E. Brücken , F. Garcia , P. Inkaew , K.T.S. Kallonen , R. Kumar Verma , T. Lampén , K. Lassila-Perini , B. Lehtela , S. Lehti , T. Lindén , N.R. Mancilla Xinto , M. Myllymäki , M.m. Rantanen , S. Saariokari , N.T. Toikka , J. Tuominiemi 

Lappeenranta-Lahti University of Technology, Lappeenranta, Finland

H. Kirschenmann , P. Luukka , H. Petrow 

IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France

M. Besancon , F. Couderc , M. Dejardin , D. Denegri, P. Devouge, J.L. Faure , F. Ferri , P. Gagne, S. Ganjour , P. Gras , G. Hamel de Monchenault , M. Kumar , V. Lohezic , J. Malcles , F. Orlandi , L. Portales , S. Ronchi , M.Ö. Sahin , A. Savoy-Navarro¹⁹ , P. Simkina , M. Titov , M. Tornago 

Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France

R. Amella Ranz , F. Beaudette , G. Boldrini , P. Busson , C. Charlot , M. Chiusi , T.D. Cuisset , O. Davignon , A. De Wit , T. Debnath , I.T. Ehle , S. Ghosh , A. Gilbert , R. Granier de Cassagnac , L. Kalipoliti , M. Manoni , M. Nguyen , S. Obraztsov , C. Ochando , R. Salerno , J.B. Sauvan , Y. Sirois , G. Sokmen, L. Urda Gómez , A. Zabi , A. Zghiche 

Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France

J.-L. Agram²⁰ , J. Andrea , D. Bloch , J.-M. Brom , E.C. Chabert , C. Collard 

G. Coulon, S. Falke , U. Goerlach , R. Haeberle , A.-C. Le Bihan , M. Meena , O. Poncet , G. Saha , P. Vaucelle 

Centre de Calcul de l'Institut National de Physique Nucleaire et de Physique des Particules, CNRS/IN2P3, Villeurbanne, France

A. Di Florio 

Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France

D. Amram, S. Beauceron , B. Blancon , G. Boudoul , N. Chanon , D. Contardo , P. Depasse , H. El Mamouni, J. Fay , S. Gascon , M. Gouzevitch , C. Greenberg , G. Grenier , B. Ille , E. Jourdain, I.B. Laktineh, M. Lethuillier , B. Massoteau, L. Mirabito, A. Purohit , M. Vander Donckt , J. Xiao 

Georgian Technical University, Tbilisi, Georgia

G. Adamov, I. Lomidze , Z. Tsamalaidze²¹ 

RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany

V. Botta , S. Consuegra Rodríguez , L. Feld , K. Klein , M. Lipinski , D. Meuser , P. Nattland , V. Oppenländer, A. Pauls , D. Pérez Adán , N. Röwert , M. Teroerde 

RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

C. Daumann, S. Diekmann , A. Dodonova , N. Eich , D. Eliseev , F. Engelke , J. Erdmann , M. Erdmann , B. Fischer , T. Hebbeker , K. Hoepfner , F. Ivone , A. Jung , N. Kumar , M.y. Lee , F. Mausolf , M. Merschmeyer , A. Meyer , F. Nowotny, A. Pozdnyakov , W. Redjeb , H. Reithler , U. Sarkar , V. Sarkisovi , A. Schmidt , C. Seth, A. Sharma , J.L. Spah , V. Vaulin, S. Zaleski

RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany

M.R. Beckers , C. Dziwok , G. Flügge , N. Hoeflich , T. Kress , A. Nowack , O. Pooth , A. Stahl , A. Zotz 

Deutsches Elektronen-Synchrotron, Hamburg, Germany

H. Aarup Petersen , A. Abel, M. Aldaya Martin , J. Alimena , S. Amoroso, Y. An , I. Andreev , J. Bach , S. Baxter , M. Bayatmakou , H. Becerril Gonzalez , O. Behnke , A. Belvedere , F. Blekman²² , K. Borrás²³ , A. Campbell , S. Chatterjee , L.X. Coll Saravia , G. Eckerlin, D. Eckstein , E. Gallo²² , A. Geiser , V. Guglielmi , M. Guthoff , A. Hinzmann , L. Jeppe , M. Kasemann , C. Kleinwort , R. Kogler , M. Komm , D. Krücker , W. Lange, D. Leyva Pernia , K.-Y. Lin , K. Lipka²⁴ , W. Lohmann²⁵ , J. Malvaso , R. Mankel , I.-A. Melzer-Pellmann , M. Mendizabal Morentin , A.B. Meyer , G. Milella , K. Moral Figueroa , A. Mussgiller , L.P. Nair , J. Niedziela , A. Nürnberg , J. Park , E. Ranken , A. Raspereza , D. Rastorguev , L. Rygaard, M. Scham^{26,23} , S. Schnake²³ , P. Schütze , C. Schwanenberger²² , D. Selivanova , K. Sharke , M. Shchedrolosiev , D. Stafford , M. Torkian, F. Vazzoler , A. Ventura Barroso , R. Walsh , D. Wang , Q. Wang , K. Wichmann, L. Wiens²³ , C. Wissing , Y. Yang , S. Zakharov, A. Zimmermann Castro Santos 

University of Hamburg, Hamburg, Germany

A.R. Alves Andrade , M. Antonello , S. Bollweg, M. Bonanomi , K. El Morabit , Y. Fischer , M. Frahm, E. Garutti , A. Grohsjean , A.A. Guvenli , J. Haller , D. Hundhausen, G. Kasieczka , P. Keicher , R. Klanner , W. Korcar , T. Kramer , C.c. Kuo, F. Labe , J. Lange , A. Lobanov , L. Moureaux , A. Nigamova , K. Nikolopoulos , A. Paasch , K.J. Pena Rodriguez , N. Prouvost, T. Quadfasel , B. Raciti , M. Rieger , D. Savoie , P. Schleper , M. Schröder , J. Schwandt 

M. Sommerhalder , H. Stadie , G. Steinbrück , R. Ward , B. Wiederspan, M. Wolf 

Karlsruher Institut fuer Technologie, Karlsruhe, Germany

S. Brommer , E. Butz , Y.M. Chen , T. Chwalek , A. Dierlamm , G.G. Dincer , U. Elicabuk, N. Faltermann , M. Giffels , A. Gottmann , F. Hartmann²⁷ , M. Horzela , F. Hummer , U. Husemann , J. Kieseler , M. Klute , R. Kunnilan Muhammed Rafeek, O. Lavoryk , J.M. Lawhorn , A. Lintuluoto , S. Maier , M. Mormile , Th. Müller , E. Pfeffer , M. Presilla , G. Quast , K. Rabbertz , B. Regnery , R. Schmieder, N. Shadskiy , I. Shvetsov , H.J. Simonis , L. Sowa , L. Stockmeier, K. Tauqeer, M. Toms , B. Topko , N. Trevisani , C. Verstege , T. Voigtländer , R.F. Von Cube , J. Von Den Driesch, M. Wassmer , R. Wolf , W.D. Zeuner , X. Zuo 

Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece

G. Anagnostou , G. Daskalakis , A. Kyriakis 

National and Kapodistrian University of Athens, Athens, Greece

G. Melachroinos, Z. Painesis , I. Paraskevas , N. Saoulidou , K. Theofilatos , E. Tziaferi , E. Tzovara , K. Vellidis , I. Zisopoulos 

National Technical University of Athens, Athens, Greece

T. Chatzistavrou, G. Karapostoli , K. Kousouris , E. Siamarkou, G. Tsipolitis 

University of Ioánnina, Ioánnina, Greece

I. Bestintzanos, I. Evangelou , C. Foudas, P. Katsoulis, P. Kokkas , P.G. Kosmoglou Kioseoglou , N. Manthos , I. Papadopoulos , J. Strologas 

HUN-REN Wigner Research Centre for Physics, Budapest, Hungary

D. Druzhkin , C. Hajdu , D. Horvath^{28,29} , K. Márton, A.J. Rádl³⁰ , F. Sikler , V. Veszpremi 

MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

M. Csanád , K. Farkas , A. Fehérkuti³¹ , M.M.A. Gadallah³² , Á. Kadlecik , M. León Coello , G. Pásztor , G.I. Veres 

Faculty of Informatics, University of Debrecen, Debrecen, Hungary

B. Ujvari , G. Zilizi 

HUN-REN ATOMKI - Institute of Nuclear Research, Debrecen, Hungary

G. Bencze, S. Czellar, J. Molnar, Z. Szillasi

Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary

T. Csorgo³¹ , F. Nemes³¹ , T. Novak , I. Szanyi³³ 

Panjab University, Chandigarh, India

S. Bansal , S.B. Beri, V. Bhatnagar , G. Chaudhary , S. Chauhan , N. Dhingra³⁴ , A. Kaur , A. Kaur , H. Kaur , M. Kaur , S. Kumar , T. Sheokand, J.B. Singh , A. Singla 

University of Delhi, Delhi, India

A. Bhardwaj , A. Chhetri , B.C. Choudhary , A. Kumar , A. Kumar , M. Naimuddin , S. Phor , K. Ranjan , M.K. Saini

University of Hyderabad, Hyderabad, India

S. Acharya³⁵ , B. Gomber , B. Sahu³⁵ 

Indian Institute of Technology Kanpur, Kanpur, IndiaS. Mukherjee **Saha Institute of Nuclear Physics, HBNI, Kolkata, India**S. Baradia , S. Bhattacharya , S. Das Gupta, S. Dutta , S. Dutta, S. Sarkar**Indian Institute of Technology Madras, Madras, India**M.M. Ameen , P.K. Behera , S. Chatterjee , G. Dash , A. Dattamunsi, P. Jana , P. Kalbhor , S. Kamble , J.R. Komaragiri³⁶ , T. Mishra , P.R. Pujahari , A.K. Sikdar , R.K. Singh , P. Verma , S. Verma , A. Vijay **IISER Mohali, India, Mohali, India**

B.K. Sirasva

Tata Institute of Fundamental Research-A, Mumbai, IndiaL. Bhatt, S. Dugad , G.B. Mohanty , M. Shelake , P. Suryadevara**Tata Institute of Fundamental Research-B, Mumbai, India**A. Bala , S. Banerjee , S. Barman³⁷ , R.M. Chatterjee, M. Guchait , Sh. Jain , A. Jaiswal, B.M. Joshi , S. Kumar , M. Maity³⁷, G. Majumder , K. Mazumdar , S. Parolia , R. Saxena , A. Thachayath **National Institute of Science Education and Research, An OCC of Homi Bhabha National Institute, Bhubaneswar, Odisha, India**S. Bahinipati³⁸ , D. Maity³⁹ , P. Mal , K. Naskar³⁹ , A. Nayak³⁹ , S. Nayak, K. Pal , R. Raturi, P. Sadangi, S.K. Swain , S. Varghese³⁹ , D. Vats³⁹ **Indian Institute of Science Education and Research (IISER), Pune, India**A. Alpana , S. Dube , P. Hazarika , B. Kansal , A. Laha , R. Sharma , S. Sharma , K.Y. Vaish **Indian Institute of Technology Hyderabad, Telangana, India**S. Ghosh **Isfahan University of Technology, Isfahan, Iran**H. Bakhshiansohi⁴⁰ , A. Jafari⁴¹ , V. Sedighzadeh Dalavi , M. Zeinali⁴² **Institute for Research in Fundamental Sciences (IPM), Tehran, Iran**S. Bashiri , S. Chenarani⁴³ , S.M. Etesami , Y. Hosseini , M. Khakzad , E. Khazaie , M. Mohammadi Najafabadi , S. Tizchang⁴⁴ **University College Dublin, Dublin, Ireland**M. Felcini , M. Grunewald **INFN Sezione di Bari^a, Università di Bari^b, Politecnico di Bari^c, Bari, Italy**M. Abbrescia^{a,b} , M. Barbieri^{a,b}, M. Buonsante^{a,b} , A. Colaleo^{a,b} , D. Creanza^{a,c} , N. De Filippis^{a,c} , M. De Palma^{a,b} , W. Elmetenawee^{a,b,45} , N. Ferrara^{a,c} , L. Fiore^a , L. Longo^a , M. Louka^{a,b} , G. Maggi^{a,c} , M. Maggi^a , I. Margjeka^a , V. Mastrapasqua^{a,b} , S. My^{a,b} , F. Nenna^{a,b} , S. Nuzzo^{a,b} , A. Pellicchia^{a,b} , A. Pompili^{a,b} , G. Pugliese^{a,c} , R. Radogna^{a,b} , D. Ramos^a , A. Ranieri^a , L. Silvestris^a , F.M. Simone^{a,c} , Ü. Sözbilir^a , A. Stamerra^{a,b} , D. Troiano^{a,b} , R. Venditti^{a,b} , P. Verwilligen^a , A. Zaza^{a,b} **INFN Sezione di Bologna^a, Università di Bologna^b, Bologna, Italy**G. Abbiendi^a , C. Battilana^{a,b} , D. Bonacorsi^{a,b} , P. Capiluppi^{a,b} , F.R. Cavallo^a , M. Cuffiani^{a,b} , G.M. Dallavalle^a , T. Diotallevi^{a,b} , F. Fabbri^a , A. Fanfani^{a,b} 

D. Fasanella^a , P. Giacomelli^a , C. Grandi^a , S. Lo Meo^{a,46} , M. Lorusso^{a,b} , L. Lunerti^a , S. Marcellini^a , G. Masetti^a , F.L. Navarria^{a,b} , G. Paggi^{a,b} , A. Perrotta^a , F. Primavera^{a,b} , A.M. Rossi^{a,b} , S. Rossi Tisbeni^{a,b} , T. Rovelli^{a,b} , G.P. Siroli^{a,b} 

INFN Sezione di Catania^a, Università di Catania^b, Catania, Italy

S. Costa^{a,b,47} , A. Di Mattia^a , A. Lapertosa^a , R. Potenza^{a,b} , A. Tricomi^{a,b,47} 

INFN Sezione di Firenze^a, Università di Firenze^b, Firenze, Italy

J. Altork^{a,b} , P. Assiouras^a , G. Barbagli^a , G. Bardelli^a , M. Bartolini^{a,b} , A. Calandri^{a,b} , B. Camaiani^{a,b} , A. Cassese^a , R. Ceccarelli^a , V. Ciulli^{a,b} , C. Civinini^a , R. D'Alessandro^{a,b} , L. Damenti^{a,b} , E. Focardi^{a,b} , T. Kello^a , G. Latino^{a,b} , P. Lenzi^{a,b} , M. Lizzo^a , M. Meschini^a , S. Paoletti^a , A. Papanastassiou^{a,b} , G. Sguazzoni^a , L. Viliani^a 

INFN Laboratori Nazionali di Frascati, Frascati, Italy

L. Benussi , S. Bianco , S. Meola⁴⁸ , D. Piccolo 

INFN Sezione di Genova^a, Università di Genova^b, Genova, Italy

M. Alves Gallo Pereira^a , F. Ferro^a , E. Robutti^a , S. Tosi^{a,b} 

INFN Sezione di Milano-Bicocca^a, Università di Milano-Bicocca^b, Milano, Italy

A. Benaglia^a , F. Brivio^a , V. Camagni^{a,b} , F. Cetorelli^{a,b} , F. De Guio^{a,b} , M.E. Dinardo^{a,b} , P. Dini^a , S. Gennai^a , R. Gerosa^{a,b} , A. Ghezzi^{a,b} , P. Govoni^{a,b} , L. Guzzi^a , M.R. Kim^a , G. Lavizzari^{a,b} , M.T. Lucchini^{a,b} , M. Malberti^a , S. Malvezzi^a , A. Massironi^a , D. Menasce^a , L. Moroni^a , M. Paganoni^{a,b} , S. Palluotto^{a,b} , D. Pedrini^a , A. Perego^{a,b} , G. Pizzati^{a,b} , S. Ragazzi^{a,b} , T. Tabarelli de Fatis^{a,b} 

INFN Sezione di Napoli^a, Università di Napoli 'Federico II'^b, Napoli, Italy; Università della Basilicata^c, Potenza, Italy; Scuola Superiore Meridionale (SSM)^d, Napoli, Italy

S. Buontempo^a , C. Di Fraia^{a,b} , F. Fabozzi^{a,c} , L. Favilla^{a,d} , A.O.M. Iorio^{a,b} , L. Lista^{a,b,49} , P. Paolucci^{a,27} , B. Rossi^a 

INFN Sezione di Padova^a, Università di Padova^b, Padova, Italy; Università degli Studi di Cagliari^c, Cagliari, Italy

P. Azzi^a , N. Bacchetta^{a,50} , D. Bisello^{a,b} , P. Bortignon^{a,c} , G. Bortolato^{a,b} , A.C.M. Bulla^{a,c} , R. Carlin^{a,b} , T. Dorigo^{a,51} , F. Gasparini^{a,b} , S. Giorgetti^a , E. Lusiani^a , M. Margoni^{a,b} , G. Maron^{a,52} , A.T. Meneguzzo^{a,b} , J. Pazzini^{a,b} , P. Ronchese^{a,b} , R. Rossin^{a,b} , F. Simonetto^{a,b} , M. Tosi^{a,b} , A. Triossi^{a,b} , S. Ventura^a , M. Zanetti^{a,b} , P. Zotto^{a,b} , A. Zucchetta^{a,b} , G. Zumerle^{a,b} 

INFN Sezione di Pavia^a, Università di Pavia^b, Pavia, Italy

A. Braghieri^a , S. Calzaferri^a , P. Montagna^{a,b} , M. Pelliccioni^a , V. Re^a , C. Riccardi^{a,b} , P. Salvini^a , I. Vai^{a,b} , P. Vitulo^{a,b} 

INFN Sezione di Perugia^a, Università di Perugia^b, Perugia, Italy

S. Ajmal^{a,b} , M.E. Ascioti^{a,b} , G.M. Bilei^a , C. Carrivale^{a,b} , D. Ciangottini^{a,b} , L. Della Penna^{a,b} , L. Fanò^{a,b} , V. Mariani^{a,b} , M. Menichelli^a , F. Moscatelli^{a,53} , A. Rossi^{a,b} , A. Santocchia^{a,b} , D. Spiga^a , T. Tedeschi^{a,b} 

INFN Sezione di Pisa^a, Università di Pisa^b, Scuola Normale Superiore di Pisa^c, Pisa, Italy; Università di Siena^d, Siena, Italy

C. Aimè^{a,b} , C.A. Alexe^{a,c} , P. Asenov^{a,b} , P. Azzurri^a , G. Bagliesi^a , L. Bianchini^{a,b} 

T. Boccali^a , E. Bossini^a , D. Bruschini^{a,c} , L. Calligaris^{a,b} , R. Castaldi^a , F. Cattafesta^{a,c} , M.A. Ciocci^{a,d} , M. Cipriani^{a,b} , R. Dell'Orso^a , S. Donato^{a,b} , R. Forti^{a,b} , A. Giassi^a , F. Ligabue^{a,c} , A.C. Marini^{a,b} , D. Matos Figueiredo^a , A. Messineo^{a,b} , S. Mishra^a , V.K. Muraleedharan Nair Bindhu^{a,b} , S. Nandan^a , F. Palla^a , M. Riggirello^{a,c} , A. Rizzi^{a,b} , G. Rolandi^{a,c} , S. Roy Chowdhury^{a,54} , T. Sarkar^a , A. Scribano^a , P. Solanki^{a,b} , P. Spagnolo^a , F. Tenchini^{a,b} , R. Tenchini^a , G. Tonelli^{a,b} , N. Turini^{a,d} , F. Vaselli^{a,c} , A. Venturi^a , P.G. Verdini^a 

INFN Sezione di Roma^a, Sapienza Università di Roma^b, Roma, Italy

P. Akrap^{a,b} , C. Basile^{a,b} , S.C. Behera^a , F. Cavallari^a , L. Cunqueiro Mendez^{a,b} , F. De Ruggi^{a,b} , D. Del Re^{a,b} , E. Di Marco^a , M. Diemoz^a , F. Errico^a , L. Frosina^{a,b} , R. Gargiulo^{a,b} , B. Harikrishnan^{a,b} , F. Lombardi^{a,b} , E. Longo^{a,b} , L. Martikainen^{a,b} , J. Mijuskovic^{a,b} , G. Organtini^{a,b} , N. Palmeri^{a,b} , R. Paramatti^{a,b} , S. Rahatlou^{a,b} , C. Rovelli^a , F. Santanastasio^{a,b} , L. Soffi^a , V. Vladimirov^{a,b} 

INFN Sezione di Torino^a, Università di Torino^b, Torino, Italy; Università del Piemonte Orientale^c, Novara, Italy

N. Amapane^{a,b} , R. Arcidiacono^{a,c} , S. Argiro^{a,b} , M. Arneodo^{a,c} , N. Bartosik^{a,c} , R. Bellan^{a,b} , A. Bellora^{a,b} , C. Biino^a , C. Borca^{a,b} , N. Cartiglia^a , M. Costa^{a,b} , R. Covarelli^{a,b} , N. Demaria^a , M. Ferrero^a , L. Finco^a , M. Grippio^{a,b} , B. Kiani^{a,b} , L. Lanteri^{a,b} , F. Legger^a , F. Luongo^{a,b} , C. Mariotti^a , S. Maselli^a , G. Mazza^a , A. Mecca^{a,b} , L. Menzio^{a,b} , P. Meridiani^a , E. Migliore^{a,b} , M. Monteno^a , M.M. Obertino^{a,b} , G. Ortona^a , L. Pacher^{a,b} , N. Pastrone^a , M. Ruspa^{a,c} , F. Siviero^{a,b} , V. Sola^{a,b} , A. Solano^{a,b} , C. Tarricone^{a,b} , D. Trocino^a , G. Umoret^{a,b} , R. White^{a,b} 

INFN Sezione di Trieste^a, Università di Trieste^b, Trieste, Italy

J. Babbar^{a,b} , S. Belforte^a , V. Candelise^{a,b} , M. Casarsa^a , F. Cossutti^a , K. De Leo^a , G. Della Ricca^{a,b} , R. Delli Gatti^{a,b} 

Kyungpook National University, Daegu, Korea

S. Dogra , J. Hong , J. Kim, T. Kim , D. Lee, H. Lee , J. Lee, S.W. Lee , C.S. Moon , Y.D. Oh , S. Sekmen , B. Tae, Y.C. Yang 

Department of Mathematics and Physics - GWNu, Gangneung, Korea

M.S. Kim 

Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea

G. Bak , P. Gwak , H. Kim , D.H. Moon , J. Seo 

Hanyang University, Seoul, Korea

E. Asilar , F. Carnevali , J. Choi⁵⁵ , T.J. Kim , Y. Ryou

Korea University, Seoul, Korea

S. Ha , S. Han, B. Hong , J. Kim , K. Lee, K.S. Lee , S. Lee , J. Yoo 

Kyung Hee University, Department of Physics, Seoul, Korea

J. Goh , J. Shin , S. Yang 

Sejong University, Seoul, Korea

Y. Kang , H. S. Kim , Y. Kim , S. Lee

Seoul National University, Seoul, Korea

J. Almond, J.H. Bhyun, J. Choi , J. Choi, W. Jun , H. Kim , J. Kim , T. Kim, Y. Kim,

Y.W. Kim , S. Ko , H. Lee , J. Lee , J. Lee , B.H. Oh , S.B. Oh , J. Shin , U.K. Yang, I. Yoon 

University of Seoul, Seoul, Korea

W. Jang , D.Y. Kang, D. Kim , S. Kim , B. Ko, J.S.H. Lee , Y. Lee , I.C. Park , Y. Roh, I.J. Watson 

Yonsei University, Department of Physics, Seoul, Korea

G. Cho, K. Hwang , B. Kim , S. Kim, K. Lee , H.D. Yoo 

Sungkyunkwan University, Suwon, Korea

M. Choi , Y. Lee , I. Yu 

College of Engineering and Technology, American University of the Middle East (AUM), Dasman, Kuwait

T. Beyrouthy , Y. Gharbia 

Kuwait University - College of Science - Department of Physics, Safat, Kuwait

F. Alazemi 

Riga Technical University, Riga, Latvia

K. Dreimanis , O.M. Eberlins , A. Gaile , C. Munoz Diaz , D. Osite , G. Pikurs , R. Plese , A. Potrebko , M. Seidel , D. Sidiropoulos Kontos 

University of Latvia (LU), Riga, Latvia

N.R. Strautnieks 

Vilnius University, Vilnius, Lithuania

M. Ambrozys , A. Juodagalvis , S. Nargelas , A. Rinkevicius , G. Tamulaitis 

National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia

I. Yusuff⁵⁶ , Z. Zolkapli

Universidad de Sonora (UNISON), Hermosillo, Mexico

J.F. Benitez , A. Castaneda Hernandez , A. Cota Rodriguez , L.E. Cuevas Picos, H.A. Encinas Acosta, L.G. Gallegos Maríñez, J.A. Murillo Quijada , L. Valencia Palomo 

Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico

G. Ayala , H. Castilla-Valdez , H. Crotte Ledesma , R. Lopez-Fernandez , J. Mejia Guisao , R. Reyes-Almanza , A. Sánchez Hernández 

Universidad Iberoamericana, Mexico City, Mexico

C. Oropeza Barrera , D.L. Ramirez Guadarrama, M. Ramírez García 

Benemerita Universidad Autonoma de Puebla, Puebla, Mexico

I. Bautista , F.E. Neri Huerta , I. Pedraza , H.A. Salazar Ibarguen , C. Uribe Estrada 

University of Montenegro, Podgorica, Montenegro

I. Bubanja , N. Raicevic 

University of Canterbury, Christchurch, New Zealand

P.H. Butler 

National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan

A. Ahmad , M.I. Asghar , A. Awais , M.I.M. Awan, W.A. Khan 

AGH University of Krakow, Krakow, Poland

V. Avati, L. Forthomme , L. Grzanka , M. Malawski , K. Piotrkowski 

National Centre for Nuclear Research, Swierk, Poland

M. Bluj , M. Górski , M. Kazana , M. Szleper , P. Zalewski 

Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland

K. Bunkowski , K. Doroba , A. Kalinowski , M. Konecki , J. Krolikowski , A. Muhammad 

Warsaw University of Technology, Warsaw, Poland

P. Fokow , K. Pozniak , W. Zabolotny 

Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal

M. Araujo , D. Bastos , C. Beirão Da Cruz E Silva , A. Boletti , M. Bozzo , T. Camporesi , G. Da Molin , M. Gallinaro , J. Hollar , N. Leonardo , G.B. Marozzo , A. Petrilli , M. Pisano , J. Seixas , J. Varela , J.W. Wulff 

Faculty of Physics, University of Belgrade, Belgrade, Serbia

P. Adzic , L. Markovic , P. Milenovic , V. Milosevic 

VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia

D. Devetak , M. Dordevic , J. Milosevic , L. Nadderd , V. Rekovic , M. Stojanovic 

Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain

M. Alcalde Martinez , J. Alcaraz Maestre , Cristina F. Bedoya , J.A. Brochero Cifuentes , Oliver M. Carretero , M. Cepeda , M. Cerrada , N. Colino , B. De La Cruz , A. Delgado Peris , A. Escalante Del Valle , D. Fernández Del Val , J.P. Fernández Ramos , J. Flix , M.C. Fouz , M. Gonzalez Hernandez , O. Gonzalez Lopez , S. Goy Lopez , J.M. Hernandez , M.I. Josa , J. Llorente Merino , C. Martin Perez , E. Martin Viscasillas , D. Moran , C. M. Morcillo Perez , Á. Navarro Tobar , R. Paz Herrera , C. Perez Dengra , A. Pérez-Calero Yzquierdo , J. Puerta Pelayo , I. Redondo , J. Vazquez Escobar

Universidad Autónoma de Madrid, Madrid, Spain

J.F. de Trocóniz 

Universidad de Oviedo, Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Oviedo, Spain

B. Alvarez Gonzalez , J. Ayllon Torresano , A. Cardini , J. Cuevas , J. Del Riego Badas , D. Estrada Acevedo , J. Fernandez Menendez , S. Folgueras , I. Gonzalez Caballero , P. Leguina , M. Obeso Menendez , E. Palencia Cortezon , J. Prado Pico , A. Soto Rodríguez , C. Vico Villalba , P. Vischia

Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain

S. Blanco Fernández , I.J. Cabrillo , A. Calderon , J. Duarte Campderros , M. Fernandez , G. Gomez , C. Lasasa García , R. Lopez Ruiz , C. Martinez Rivero , P. Martinez Ruiz del Arbol , F. Matorras , P. Matorras Cuevas , E. Navarrete Ramos , J. Piedra Gomez , C. Quintana San Emeterio , L. Scodellaro , I. Vila , R. Vilar Cortabitarte , J.M. Vizan Garcia

University of Colombo, Colombo, Sri Lanka

B. Kailasapathy⁵⁷ , D.D.C. Wickramarathna 

University of Ruhuna, Department of Physics, Matara, Sri Lanka

W.G.D. Dharmaratna⁵⁸ , K. Liyanage , N. Perera 

CERN, European Organization for Nuclear Research, Geneva, Switzerland

D. Abbaneo , C. Amendola , R. Ardino , E. Auffray , J. Baechler, D. Barney , J. Bendavid , M. Bianco , A. Bocci , L. Borgonovi , C. Botta , A. Bragagnolo , C.E. Brown , C. Caillol , G. Cerminara , P. Connor , D. d'Enterria , A. Dabrowski , A. David , A. De Roeck , M.M. Defranchis , M. Deile , M. Dobson , P.J. Fernández Manteca , B.A. Fontana Santos Alves , W. Funk , A. Gaddi, S. Giani, D. Gigi, K. Gill , F. Glege , M. Glowacki, A. Gruber , J. Hegeman , J.K. Heikkilä , R. Hofsaess , B. Huber , T. James , P. Janot , O. Kaluzinska , O. Karacheban²⁵ , G. Karathanasis , S. Laurila , P. Lecoq , C. Lourenço , A.-M. Lyon , M. Magherini , L. Malgeri , M. Mannelli , A. Mehta , F. Meijers , J.A. Merlin, S. Mersi , E. Meschi , M. Migliorini , F. Monti , F. Moortgat , M. Mulders , M. Musich , I. Neutelings , S. Orfanelli, F. Pantaleo , M. Pari , G. Petrucciani , A. Pfeiffer , M. Pierini , M. Pitt , H. Qu , D. Rabady , A. Reimers , B. Ribeiro Lopes , F. Riti , P. Rosado , M. Rovere , H. Sakulin , R. Salvatico , S. Sanchez Cruz , S. Scarfi , M. Selvaggi , A. Sharma , K. Shchelina , P. Silva , P. Sphicas⁵⁹ , A.G. Stahl Leiton , A. Steen , S. Summers , D. Treille , P. Tropea , E. Vernazza , J. Wanczyk⁶⁰ , J. Wang, S. Wuchterl , M. Zarucki , P. Zehetner , P. Zejdl , G. Zevi Della Porta 

PSI Center for Neutron and Muon Sciences, Villigen, Switzerland

T. Bevilacqua⁶¹ , L. Caminada⁶¹ , W. Erdmann , R. Horisberger , Q. Ingram , H.C. Kaestli , D. Kotlinski , C. Lange , U. Langenegger , L. Noehte⁶¹ , T. Rohe , A. Samalan 

ETH Zurich - Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland

T.K. Aarrestad , M. Backhaus , G. Bonomelli , C. Cazzaniga , K. Datta , P. De Bryas Dexmiers D'Archiacchiac⁶⁰ , A. De Cosa , G. Dissertori , M. Dittmar, M. Donegà , F. Eble , K. Gedia , F. Glessgen , C. Grab , N. Härringer , T.G. Harte , W. Lustermann , M. Malucchi , R.A. Manzoni , M. Marchegiani , L. Marchese , A. Mascellani⁶⁰ , F. Nessi-Tedaldi , F. Pauss , V. Perovic , B. Ristic , R. Seidita , J. Steggemann⁶⁰ , A. Tarabini , D. Valsecchi , R. Wallny 

Universität Zürich, Zurich, Switzerland

C. Amsler⁶² , P. Bärttschi , F. Bilandzija , M.F. Canelli , G. Celotto , K. Cormier , M. Huwiler , W. Jin , A. Jofrehei , B. Kilminster , T.H. Kwok , S. Leontsinis , V. Lukashenko , A. Macchiolo , F. Meng , M. Missiroli , J. Motta , P. Robmann, M. Senger , E. Shokr , F. Stäger , R. Tramontano , P. Viscone 

National Central University, Chung-Li, Taiwan

D. Bhowmik, C.M. Kuo, P.K. Rout , S. Taj , P.C. Tiwari³⁶ 

National Taiwan University (NTU), Taipei, Taiwan

L. Ceard, K.F. Chen , Z.g. Chen, A. De Iorio , W.-S. Hou , T.h. Hsu, Y.w. Kao, S. Karmakar , G. Kole , Y.y. Li , R.-S. Lu , E. Paganis , X.f. Su , J. Thomas-Wilsker , L.s. Tsai, D. Tsionou, H.y. Wu , E. Yazgan 

High Energy Physics Research Unit, Department of Physics, Faculty of Science, Chulalongkorn University, Bangkok, Thailand

C. Asawatangkuldee , N. Srimanobhas 

Tunis El Manar University, Tunis, Tunisia

Y. Maghrbi 

Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey

D. Agyel , F. Dolek , I. Dumanoglu⁶³ , Y. Guler⁶⁴ , E. Gurpinar Guler⁶⁴ , C. Isik , O. Kara , A. Kayis Topaksu , Y. Komurcu , G. Onengut , K. Ozdemir⁶⁵ , B. Tali⁶⁶ , U.G. Tok , E. Uslan , I.S. Zorbakir 

Middle East Technical University, Physics Department, Ankara, Turkey

M. Yalvac⁶⁷ 

Bogazici University, Istanbul, Turkey

B. Akgun , I.O. Atakisi⁶⁸ , E. Gülmez , M. Kaya⁶⁹ , O. Kaya⁷⁰ , M.A. Sarkisla⁷¹, S. Tekten⁷² 

Istanbul Technical University, Istanbul, Turkey

A. Cakir , K. Cankocak^{63,73} , S. Sen⁷⁴ 

Istanbul University, Istanbul, Turkey

B. Hacisahinoglu , I. Hos⁷⁵ , B. Kaynak , S. Ozkorucuklu , O. Potok , H. Sert , C. Simsek , C. Zorbilmez 

Yildiz Technical University, Istanbul, Turkey

S. Cerci , C. Dozen⁷⁶ , B. Isildak⁷⁷ , E. Simsek , D. Sunar Cerci , T. Yetkin⁷⁶ 

Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkiv, Ukraine

A. Boyaryntsev , O. Dadazhanova, B. Grynyov 

National Science Centre, Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine

L. Levchuk 

University of Bristol, Bristol, United Kingdom

J.J. Brooke , A. Bundock , F. Bury , E. Clement , D. Cussans , D. Dharmender, H. Flacher , J. Goldstein , H.F. Heath , M.-L. Holmberg , L. Kreczko , S. Paramesvaran , L. Robertshaw, M.S. Sanjrani⁴⁰, J. Segal, V.J. Smith 

Rutherford Appleton Laboratory, Didcot, United Kingdom

A.H. Ball, K.W. Bell , A. Belyaev⁷⁸ , C. Brew , R.M. Brown , D.J.A. Cockerill , A. Elliot , K.V. Ellis, J. Gajownik , K. Harder , S. Harper , J. Linacre , K. Manolopoulos, M. Moallemi , D.M. Newbold , E. Olaiya , D. Petyt , T. Reis , A.R. Sahasransu , G. Salvi , T. Schuh, C.H. Shepherd-Themistocleous , I.R. Tomalin , K.C. Whalen , T. Williams 

Imperial College, London, United Kingdom

I. Andreou , R. Bainbridge , P. Bloch , O. Buchmuller, C.A. Carrillo Montoya , D. Colling , J.S. Dancu, I. Das , P. Dauncey , G. Davies , M. Della Negra , S. Fayer, G. Fedi , G. Hall , H.R. Hoorani , A. Howard, G. Iles , C.R. Knight , P. Krueper , J. Langford , K.H. Law , J. León Holgado , E. Leutgeb , L. Lyons , A.-M. Magnan , B. Maier , S. Mallios, A. Mastronikolis , M. Mieskolainen , J. Nash⁷⁹ , M. Pesaresi , P.B. Pradeep , B.C. Radburn-Smith , A. Richards, A. Rose , L. Russell , K. Savva , C. Seez , R. Shukla , A. Tapper , K. Uchida , G.P. Uttley , T. Virdee²⁷ , M. Vojinovic , N. Wardle , D. Winterbottom 

Brunel University, Uxbridge, United Kingdom

J.E. Cole , A. Khan, P. Kyberd , I.D. Reid 

Baylor University, Waco, Texas, USA

S. Abdullin , A. Brinkerhoff , E. Collins , M.R. Darwish , J. Dittmann 

K. Hatakeyama , V. Hegde , J. Hiltbrand , B. McMaster , J. Samudio , S. Sawant ,
C. Sutantawibul , J. Wilson 

Bethel University, St. Paul, Minnesota, USA

J.M. Hogan 

Catholic University of America, Washington, DC, USA

R. Bartek , A. Dominguez , S. Raj , A.E. Simsek , S.S. Yu 

The University of Alabama, Tuscaloosa, Alabama, USA

B. Bam , A. Buchot Perraguin , S. Campbell, R. Chudasama , S.I. Cooper , C. Crovella ,
G. Fidalgo , S.V. Gleyzer , A. Khukhunaishvili , K. Matchev , E. Pearson, C.U. Perez ,
P. Rumerio⁸⁰ , E. Usai , R. Yi 

Boston University, Boston, Massachusetts, USA

S. Cholak , G. De Castro, Z. Demiragli , C. Erice , C. Fangmeier , C. Fernandez Madrazo , E. Fontanesi , J. Fulcher , F. Golf , S. Jeon , J. O'Cain, I. Reed ,
J. Rohlf , K. Salyer , D. Sperka , D. Spitzbart , I. Suarez , A. Tsatsos , E. Wurtz,
A.G. Zecchinelli 

Brown University, Providence, Rhode Island, USA

G. Barone , G. Benelli , D. Cutts , S. Ellis , L. Gouskos , M. Hadley , U. Heintz ,
K.W. Ho , T. Kwon , L. Lambrecht , G. Landsberg , K.T. Lau , J. Luo , S. Mondal ,
J. Roloff, T. Russell , S. Sagir⁸¹ , X. Shen , M. Stamenkovic , N. Venkatasubramanian 

University of California, Davis, Davis, California, USA

S. Abbott , B. Barton , R. Breedon , H. Cai , M. Calderon De La Barca Sanchez ,
E. Cannaert, M. Chertok , M. Citron , J. Conway , P.T. Cox , R. Erbacher , O. Kukral ,
G. Mocellin , S. Ostrom , I. Salazar Segovia, J.S. Tafoya Vargas , W. Wei , S. Yoo 

University of California, Los Angeles, California, USA

K. Adamidis, M. Bachtis , D. Campos, R. Cousins , A. Datta , G. Flores Avila ,
J. Hauser , M. Ignatenko , M.A. Iqbal , T. Lam , Y.f. Lo , E. Manca ,
A. Nunez Del Prado , D. Saltzberg , V. Valuev 

University of California, Riverside, Riverside, California, USA

R. Clare , J.W. Gary , G. Hanson 

University of California, San Diego, La Jolla, California, USA

A. Aportela , A. Arora , J.G. Branson , S. Cittolin , S. Cooperstein , B. D'Anzi ,
D. Diaz , J. Duarte , L. Giannini , Y. Gu, J. Guiang , V. Krutelyov , R. Lee ,
J. Letts , H. Li, M. Masciovecchio , F. Mokhtar , S. Mukherjee , M. Pieri , D. Primosch,
M. Quinnan , V. Sharma , M. Tadel , E. Vourliotis , F. Würthwein , A. Yagil , Z. Zhao

University of California, Santa Barbara - Department of Physics, Santa Barbara, California, USA

A. Barzdukas , L. Brennan , C. Campagnari , S. Carron Montero⁸² , K. Downham ,
C. Grieco , M.M. Hussain, J. Incandela , M.W.K. Lai, A.J. Li , P. Masterson ,
J. Richman , S.N. Santpur , U. Sarica , R. Schmitz , F. Setti , J. Sheplock , D. Stuart ,
T.Á. Vámi , X. Yan , D. Zhang 

California Institute of Technology, Pasadena, California, USA

A. Albert , S. Bhattacharya , A. Bornheim , O. Cerri, R. Kansal , J. Mao , H.B. Newman ,
G. Reales Gutiérrez, T. Sievert, M. Spiropulu , J.R. Vlimant , R.A. Wynne , S. Xie 

Carnegie Mellon University, Pittsburgh, Pennsylvania, USA

J. Alison , S. An , M. Cremonesi, V. Dutta , E.Y. Ertorer , T. Ferguson , T.A. Gómez Espinosa , A. Harilal , A. Kallil Tharayil, M. Kanemura, C. Liu , P. Meiring , T. Mudholkar , S. Murthy , P. Palit , K. Park , M. Paulini , A. Roberts , A. Sanchez , W. Terrill 

University of Colorado Boulder, Boulder, Colorado, USA

J.P. Cumalat , W.T. Ford , A. Hart , S. Kwan , J. Pearkes , C. Savard , N. Schonbeck , K. Stenson , K.A. Ulmer , S.R. Wagner , N. Zipper , D. Zuolo 

Cornell University, Ithaca, New York, USA

J. Alexander , X. Chen , J. Dickinson , A. Duquette, J. Fan , X. Fan , J. Grassi , S. Hogan , P. Kotamnives , J. Monroy , G. Niendorf , M. Oshiro , J.R. Patterson , A. Ryd , J. Thom , P. Wittich , R. Zou , L. Zygala 

Fermi National Accelerator Laboratory, Batavia, Illinois, USA

M. Albrow , M. Alyari , O. Amram , G. Apollinari , A. Apresyan , L.A.T. Bauerdick , D. Berry , J. Berryhill , P.C. Bhat , K. Burkett , J.N. Butler , A. Canepa , G.B. Cerati , H.W.K. Cheung , F. Chlebana , C. Cosby , G. Cummings , I. Dutta , V.D. Elvira , J. Freeman , A. Gandrakota , Z. Gecse , L. Gray , D. Green, A. Grummer , S. Grünendahl , D. Guerrero , O. Gutsche , R.M. Harris , T.C. Herwig , J. Hirschauer , V. Innocente , B. Jayatilaka , S. Jindariani , M. Johnson , U. Joshi , T. Klijsma , B. Klima , K.H.M. Kwok , S. Lammel , C. Lee , D. Lincoln , R. Lipton , T. Liu , K. Maeshima , D. Mason , P. McBride , P. Merkel , S. Mrenna , S. Nahn , J. Ngadiuba , D. Noonan , S. Norberg, V. Papadimitriou , N. Pastika , K. Pedro , C. Pena⁸³ , C.E. Perez Lara , F. Ravera , A. Reinsvold Hall⁸⁴ , L. Ristori , M. Safdari , E. Sexton-Kennedy , N. Smith , A. Soha , L. Spiegel , S. Stoynev , J. Strait , L. Taylor , S. Tkaczyk , N.V. Tran , L. Uplegger , E.W. Vaandering , C. Wang , I. Zoi 

University of Florida, Gainesville, Florida, USA

C. Aruta , P. Avery , D. Bourilkov , P. Chang , V. Cherepanov , R.D. Field, C. Huh , E. Koenig , M. Kolosova , J. Konigsberg , A. Korytov , N. Menendez , G. Mitselmakher , K. Mohrman , A. Muthirakalayil Madhu , N. Rawal , S. Rosenzweig , V. Sulimov , Y. Takahashi , J. Wang 

Florida State University, Tallahassee, Florida, USA

T. Adams , A. Al Kadhimi , A. Askew , S. Bower , R. Goff, R. Hashmi , A. Hassani , R.S. Kim , T. Kolberg , G. Martinez , M. Mazza , H. Prosper , P.R. Prova, R. Yohay 

Florida Institute of Technology, Melbourne, Florida, USA

B. Alsufyani , S. Butalla , S. Das , M. Hohlmann , M. Lavinsky, E. Yanes

University of Illinois Chicago, Chicago, Illinois, USA

M.R. Adams , N. Barnett, A. Baty , C. Bennett , R. Cavanaugh , R. Escobar Franco , O. Evdokimov , C.E. Gerber , H. Gupta , M. Hawksworth, A. Hingrajiya, D.J. Hofman , J.h. Lee , C. Mills , S. Nanda , G. Nigmatkulov , B. Ozek , T. Phan, D. Pilipovic , R. Pradhan , E. Prifti, P. Roy, T. Roy , N. Singh, M.B. Tonjes , N. Varelas , M.A. Wadud , J. Yoo 

The University of Iowa, Iowa City, Iowa, USA

M. Alhusseini , D. Blend , K. Dilsiz⁸⁵ , O.K. Köseyan , A. Mestvirishvili⁸⁶ , O. Neogi, H. Ogul⁸⁷ , Y. Onel , A. Penzo , C. Snyder, E. Tiras⁸⁸ 

Johns Hopkins University, Baltimore, Maryland, USA

B. Blumenfeld , J. Davis , A.V. Gritsan , L. Kang , S. Kyriacou , P. Maksimovic , M. Roguljic , S. Sekhar , M.V. Srivastav , M. Swartz 

The University of Kansas, Lawrence, Kansas, USA

A. Abreu , L.F. Alcerro Alcerro , J. Anguiano , S. Arteaga Escatel , P. Baringer , A. Bean , R. Bhattacharya , Z. Flowers , D. Grove , J. King , G. Krintiras , M. Lazarovits , C. Le Mahieu , J. Marquez , M. Murray , M. Nickel , S. Popescu⁸⁹ , C. Rogan , C. Royon , S. Rudrabhatla , S. Sanders , C. Smith , G. Wilson 

Kansas State University, Manhattan, Kansas, USA

B. Allmond , N. Islam , A. Ivanov , K. Kaadze , Y. Maravin , J. Natoli , G.G. Reddy , D. Roy , G. Sorrentino 

University of Maryland, College Park, Maryland, USA

A. Baden , A. Belloni , J. Bistany-riebman , S.C. Eno , N.J. Hadley , S. Jabeen , R.G. Kellogg , T. Koeth , B. Kronheim , S. Lascio , P. Major , A.C. Mignerey , C. Palmer , C. Papageorgakis , M.M. Paranjpe , E. Popova⁹⁰ , A. Shevelev , L. Zhang 

Massachusetts Institute of Technology, Cambridge, Massachusetts, USA

C. Baldenegro Barrera , H. Bossi , S. Bright-Thonney , I.A. Cali , Y.c. Chen , P.c. Chou , M. D'Alfonso , J. Eysermans , C. Freer , G. Gomez-Ceballos , M. Goncharov , G. Grosso , P. Harris , D. Hoang , G.M. Innocenti , D. Kovalskyi , J. Krupa , L. Lavezzo , Y.-J. Lee , K. Long , C. McGinn , A. Novak , M.I. Park , C. Paus , C. Reissel , C. Roland , G. Roland , S. Rothman , T.a. Sheng , G.S.F. Stephans , D. Walter , Z. Wang , B. Wyslouch , T. J. Yang 

University of Minnesota, Minneapolis, Minnesota, USA

B. Crossman , W.J. Jackson , C. Kapsiak , M. Krohn , D. Mahon , J. Mans , B. Marzocchi , R. Rusack , O. Sancar , R. Saradhy , N. Strobbe 

University of Nebraska-Lincoln, Lincoln, Nebraska, USA

K. Bloom , D.R. Claes , G. Haza , J. Hossain , C. Joo , I. Kravchenko , A. Rohilla , J.E. Siado , W. Tabb , A. Vagnerini , A. Wightman , F. Yan 

State University of New York at Buffalo, Buffalo, New York, USA

H. Bandyopadhyay , L. Hay , H.w. Hsia , I. Iashvili , A. Kalogeropoulos , A. Kharchilava , A. Mandal , M. Morris , D. Nguyen , S. Rappoccio , H. Rejeb Sfar , A. Williams , P. Young , D. Yu 

Northeastern University, Boston, Massachusetts, USA

G. Alverson , E. Barberis , J. Bonilla , B. Bylsma , M. Campana , J. Dervan , Y. Haddad , Y. Han , I. Israr , A. Krishna , M. Lu , N. Manganelli , R. McCarthy , D.M. Morse , T. Orimoto , L. Skinnari , C.S. Thoreson , E. Tsai , D. Wood 

Northwestern University, Evanston, Illinois, USA

S. Dittmer , K.A. Hahn , Y. Liu , M. McGinnis , Y. Miao , D.G. Monk , M.H. Schmitt , A. Taliencio , M. Velasco , J. Wang 

University of Notre Dame, Notre Dame, Indiana, USA

G. Agarwal , R. Band , R. Bucci , S. Castells , A. Das , A. Ehnis , R. Goldouzian , M. Hildreth , K. Hurtado Anampa , T. Ivanov , C. Jessop , A. Karneyeu , K. Lannon , J. Lawrence , N. Loukas , L. Lutton , J. Mariano , N. Marinelli , I. Mcalister , T. McCauley , C. Mcgrady , C. Moore , Y. Musienko²¹ , H. Nelson , M. Osherson , A. Piccinelli , R. Ruchti , A. Townsend , Y. Wan , M. Wayne , H. Yockey 

The Ohio State University, Columbus, Ohio, USA

A. Basnet , M. Carrigan , R. De Los Santos , L.S. Durkin , C. Hill , M. Joyce ,
M. Nunez Ornelas , D.A. Wenzl , B.L. Winer , B. R. Yates 

Princeton University, Princeton, New Jersey, USA

H. Bouchamaoui , G. Dezoort , P. Elmer , A. Frankenthal , M. Galli , B. Greenberg ,
N. Haubrich , K. Kennedy , G. Kopp , Y. Lai , D. Lange , A. Loeliger , D. Marlow ,
I. Ojalvo , J. Olsen , F. Simpson , D. Stickland , C. Tully 

University of Puerto Rico, Mayaguez, Puerto Rico, USA

S. Malik , R. Sharma 

Purdue University, West Lafayette, Indiana, USA

S. Chandra , R. Chawla , A. Gu , L. Gutay , M. Jones , A.W. Jung , D. Kondratyev ,
M. Liu , G. Negro , N. Neumeister , G. Paspalaki , S. Piperov , N.R. Saha ,
J.F. Schulte , F. Wang , A. Wildridge , W. Xie , Y. Yao , Y. Zhong 

Purdue University Northwest, Hammond, Indiana, USA

N. Parashar , A. Pathak , E. Shumka 

Rice University, Houston, Texas, USA

D. Acosta , A. Agrawal , C. Arbour , T. Carnahan , P. Das , K.M. Ecklund , S. Freed,
P. Gardner, F.J.M. Geurts , T. Huang , I. Krommydas , N. Lewis, W. Li , J. Lin ,
O. Miguel Colin , B.P. Padley , R. Redjimi , J. Rotter , M. Wulansatiti , E. Yigitbasi ,
Y. Zhang 

University of Rochester, Rochester, New York, USA

O. Bessidskaia Bylund, A. Bodek , P. de Barbaro[†] , R. Demina , A. Garcia-Bellido ,
H.S. Hare , O. Hindrichs , N. Parmar , P. Parygin⁹⁰ , H. Seo , R. Taus 

Rutgers, The State University of New Jersey, Piscataway, New Jersey, USA

B. Chiarito, J.P. Chou , S.V. Clark , S. Donnelly, D. Gadkari , Y. Gershtein ,
E. Halkiadakis , C. Houghton , D. Jaroslawski , A. Kobert , S. Konstantinou ,
I. Laflotte , A. Lath , J. Martins , M. Perez Prada , B. Rand , J. Reichert , P. Saha ,
S. Salur , S. Schnetzer, S. Somalwar , R. Stone , S.A. Thayil , S. Thomas, J. Vora 

University of Tennessee, Knoxville, Tennessee, USA

D. Ally , A.G. Delannoy , S. Fiorendi , J. Harris, T. Holmes , A.R. Kanuganti ,
N. Karunarathna , J. Lawless, L. Lee , E. Nibigira , B. Skipworth, S. Spanier 

Texas A&M University, College Station, Texas, USA

D. Aebi , M. Ahmad , T. Akhter , K. Androsov , A. Bolshov, O. Bouhali⁹¹ ,
A. Cagnotta , V. D'Amante , R. Eusebi , P. Flanagan , J. Gilmore , Y. Guo, T. Kamon ,
S. Luo , R. Mueller , A. Safonov 

Texas Tech University, Lubbock, Texas, USA

N. Akchurin , J. Damgov , Y. Feng , N. Gogate , Y. Kazhykarim, K. Lamichhane ,
S.W. Lee , C. Madrid , A. Mankel , T. Peltola , I. Volobouev 

Vanderbilt University, Nashville, Tennessee, USA

E. Appelt , Y. Chen , S. Greene, A. Gurrola , W. Johns , R. Kunnawalkam Elayavalli ,
A. Melo , D. Rathjens , F. Romeo , P. Sheldon , S. Tuo , J. Velkovska , J. Viinikainen ,
J. Zhang

University of Virginia, Charlottesville, Virginia, USA

B. Cardwell , H. Chung , B. Cox , J. Hakala , G. Hamilton Ilha Machado, R. Hirosky , M. Jose, A. Ledovskoy , C. Mantilla , C. Neu , C. Ramón Álvarez , Z. Wu

Wayne State University, Detroit, Michigan, USA

S. Bhattacharya , P.E. Karchin 

University of Wisconsin - Madison, Madison, Wisconsin, USA

A. Aravind , S. Banerjee , K. Black , T. Bose , E. Chavez , S. Dasu , P. Everaerts , C. Galloni, H. He , M. Herndon , A. Herve , C.K. Koraka , S. Lomte , R. Loveless , A. Mallampalli , A. Mohammadi , S. Mondal, T. Nelson, G. Parida , L. Pétré , D. Pinna , A. Savin, V. Shang , V. Sharma , W.H. Smith , D. Teague, H.F. Tsoi , W. Vetens , A. Warden 

Authors affiliated with an international laboratory covered by a cooperation agreement with CERN

S. Afanasiev , V. Alexakhin , Yu. Andreev , T. Aushev , D. Budkouski , R. Chistov , M. Danilov , T. Dimova , A. Ershov , S. Gninenko , I. Gorbunov , A. Gribushin , A. Kamenev , V. Karjavine , M. Kirsanov , V. Klyukhin , O. Kodolova⁹² , V. Koronkov , I. Korsakov, A. Kozyrev , N. Krasnikov , A. Lanev , A. Malakhov , V. Matveev , A. Nikitenko^{93,92} , V. Palichik , V. Perelygin , S. Petrushanko , S. Polikarpov , O. Radchenko , M. Savina , V. Shalaev , S. Shmatov , S. Shulha , Y. Skovpen , K. Slizhevskiy, V. Smirnov , O. Teryaev , I. Tlisova , A. Toropin , N. Voytishin , A. Zarubin , I. Zhizhin 

Authors affiliated with an institute formerly covered by a cooperation agreement with CERN

E. Boos , V. Bunichev , M. Dubinin⁸³ , V. Savrin , A. Snigirev , L. Dudko , K. Ivanov , V. Kim²¹ , V. Murzin , V. Oreshkin , D. Sosnov 

†: Deceased

¹Also at Yerevan State University, Yerevan, Armenia

²Also at TU Wien, Vienna, Austria

³Also at Ghent University, Ghent, Belgium

⁴Also at Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

⁵Also at FACAMP - Faculdades de Campinas, Sao Paulo, Brazil

⁶Also at Universidade Estadual de Campinas, Campinas, Brazil

⁷Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil

⁸Also at The University of the State of Amazonas, Manaus, Brazil

⁹Also at University of Chinese Academy of Sciences, Beijing, China

¹⁰Also at China Center of Advanced Science and Technology, Beijing, China

¹¹Also at University of Chinese Academy of Sciences, Beijing, China

¹²Also at School of Physics, Zhengzhou University, Zhengzhou, China

¹³Now at Henan Normal University, Xinxiang, China

¹⁴Also at University of Shanghai for Science and Technology, Shanghai, China

¹⁵Now at The University of Iowa, Iowa City, Iowa, USA

¹⁶Also at Center for High Energy Physics, Peking University, Beijing, China

¹⁷Now at British University in Egypt, Cairo, Egypt

¹⁸Now at Cairo University, Cairo, Egypt

¹⁹Also at Purdue University, West Lafayette, Indiana, USA

²⁰Also at Université de Haute Alsace, Mulhouse, France

²¹Also at an institute formerly covered by a cooperation agreement with CERN

²²Also at University of Hamburg, Hamburg, Germany

²³Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

- ²⁴Also at Bergische University Wuppertal (BUW), Wuppertal, Germany
- ²⁵Also at Brandenburg University of Technology, Cottbus, Germany
- ²⁶Also at Forschungszentrum Jülich, Juelich, Germany
- ²⁷Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland
- ²⁸Also at HUN-REN ATOMKI - Institute of Nuclear Research, Debrecen, Hungary
- ²⁹Now at Universitatea Babes-Bolyai - Facultatea de Fizica, Cluj-Napoca, Romania
- ³⁰Also at MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary
- ³¹Also at HUN-REN Wigner Research Centre for Physics, Budapest, Hungary
- ³²Also at Physics Department, Faculty of Science, Assiut University, Assiut, Egypt
- ³³Also at The University of Kansas, Lawrence, Kansas, USA
- ³⁴Also at Punjab Agricultural University, Ludhiana, India
- ³⁵Also at University of Hyderabad, Hyderabad, India
- ³⁶Also at Indian Institute of Science (IISc), Bangalore, India
- ³⁷Also at University of Visva-Bharati, Santiniketan, India
- ³⁸Also at IIT Bhubaneswar, Bhubaneswar, India
- ³⁹Also at Institute of Physics, Bhubaneswar, India
- ⁴⁰Also at Deutsches Elektronen-Synchrotron, Hamburg, Germany
- ⁴¹Also at Isfahan University of Technology, Isfahan, Iran
- ⁴²Also at Sharif University of Technology, Tehran, Iran
- ⁴³Also at Department of Physics, University of Science and Technology of Mazandaran, Behshahr, Iran
- ⁴⁴Also at Department of Physics, Faculty of Science, Arak University, ARAK, Iran
- ⁴⁵Also at Helwan University, Cairo, Egypt
- ⁴⁶Also at Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy
- ⁴⁷Also at Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy
- ⁴⁸Also at Università degli Studi Guglielmo Marconi, Roma, Italy
- ⁴⁹Also at Scuola Superiore Meridionale, Università di Napoli 'Federico II', Napoli, Italy
- ⁵⁰Also at Fermi National Accelerator Laboratory, Batavia, Illinois, USA
- ⁵¹Also at Lulea University of Technology, Lulea, Sweden
- ⁵²Also at Laboratori Nazionali di Legnaro dell'INFN, Legnaro, Italy
- ⁵³Also at Consiglio Nazionale delle Ricerche - Istituto Officina dei Materiali, Perugia, Italy
- ⁵⁴Also at UPES - University of Petroleum and Energy Studies, Dehradun, India
- ⁵⁵Also at Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France
- ⁵⁶Also at Department of Applied Physics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia
- ⁵⁷Also at Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka
- ⁵⁸Also at Saegis Campus, Nugegoda, Sri Lanka
- ⁵⁹Also at National and Kapodistrian University of Athens, Athens, Greece
- ⁶⁰Also at Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland
- ⁶¹Also at Universität Zürich, Zurich, Switzerland
- ⁶²Also at Stefan Meyer Institute for Subatomic Physics, Vienna, Austria
- ⁶³Also at Near East University, Research Center of Experimental Health Science, Mersin, Turkey
- ⁶⁴Also at Konya Technical University, Konya, Turkey
- ⁶⁵Also at Izmir Bakircay University, Izmir, Turkey
- ⁶⁶Also at Adiyaman University, Adiyaman, Turkey
- ⁶⁷Also at Bozok Universitetesi Rektörlüğü, Yozgat, Turkey

-
- ⁶⁸Also at Istanbul Sabahattin Zaim University, Istanbul, Turkey
- ⁶⁹Also at Marmara University, Istanbul, Turkey
- ⁷⁰Also at Milli Savunma University, Istanbul, Turkey
- ⁷¹Also at Informatics and Information Security Research Center, Gebze/Kocaeli, Turkey
- ⁷²Also at Kafkas University, Kars, Turkey
- ⁷³Now at Istanbul Okan University, Istanbul, Turkey
- ⁷⁴Also at Hacettepe University, Ankara, Turkey
- ⁷⁵Also at Istanbul University - Cerrahpasa, Faculty of Engineering, Istanbul, Turkey
- ⁷⁶Also at Istinye University, Istanbul, Turkey
- ⁷⁷Also at Yildiz Technical University, Istanbul, Turkey
- ⁷⁸Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom
- ⁷⁹Also at Monash University, Faculty of Science, Clayton, Australia
- ⁸⁰Also at Università di Torino, Torino, Italy
- ⁸¹Also at Karamanoğlu Mehmetbey University, Karaman, Turkey
- ⁸²Also at California Lutheran University,, Thousand Oaks, California, USA
- ⁸³Also at California Institute of Technology, Pasadena, California, USA
- ⁸⁴Also at United States Naval Academy, Annapolis, Maryland, USA
- ⁸⁵Also at Bingöl University, Bingöl, Turkey
- ⁸⁶Also at Georgian Technical University, Tbilisi, Georgia
- ⁸⁷Also at Sinop University, Sinop, Turkey
- ⁸⁸Also at Erciyes University, Kayseri, Turkey
- ⁸⁹Also at Horia Hulubei National Institute of Physics and Nuclear Engineering (IFIN-HH), Bucharest, Romania
- ⁹⁰Now at another institute formerly covered by a cooperation agreement with CERN
- ⁹¹Also at Hamad Bin Khalifa University (HBKU), Doha, Qatar
- ⁹²Also at Yerevan Physics Institute, Yerevan, Armenia
- ⁹³Also at Imperial College, London, United Kingdom