



Search for a new pseudoscalar decaying into a pair of bottom and antibottom quarks in top-associated production in $\sqrt{s} = 13$ TeV proton–proton collisions with the ATLAS detector

ATLAS Collaboration*

CERN, 1211 Geneva 23, Switzerland

Received: 24 March 2025 / Accepted: 10 July 2025
 © The Author(s) 2025

Abstract A search for a pseudoscalar a produced in association with a top-quark pair, or in association with a single top quark plus a W boson, with the pseudoscalar decaying into b -quarks ($a \rightarrow b\bar{b}$), is performed using the full Run 2 data sample using a dileptonic decay mode signature. The search covers pseudoscalar boson masses between 12 and 100 GeV and involves both the kinematic regime where the decay products of the pseudoscalar are reconstructed as two standard b -tagged small-radius jets, or merged into a large-radius jet due to its Lorentz boost. No significant excess relative to expectations is observed. Assuming a branching ratio $\text{BR}(a \rightarrow b\bar{b}) = 100\%$, the range of pseudoscalar masses between 50 and 80 GeV is excluded at 95% confidence level for a coupling of the pseudoscalar to the top quark of 0.5, while a coupling of 1.0 is excluded at 95% confidence level for the masses considered, with the coupling defined as the strength modifier of the Standard Model Yukawa coupling.

Contents

1	Introduction	
2	ATLAS detector	
3	Data and simulated event samples	
4	Object definition	
4.1	Physics objects	
4.2	Corrections to physics objects	
5	Event selection	
5.1	Preselection	
5.2	Signal and control regions	
6	Background estimate	
7	Analysis strategy	
7.1	Event reconstruction BDTs	
7.2	Signal-versus-background discriminating neural networks	
8	Statistical treatment	

* e-mail: atlas.publications@cern.ch

9	Systematic uncertainties	
9.1	Experimental uncertainties	
9.2	Modelling uncertainties	
10	Results	
11	Conclusions	
	References	

1 Introduction

Since the discovery of the Higgs boson [1,2], there is an ongoing effort at the Large Hadron Collider (LHC) [3] to measure its properties and search for new physics. The Higgs boson was discovered by observing few decay modes [1,2] consistent with the Standard Model (SM) predictions. Part of the interest is about the nature of the discovered particle and whether it is the single boson predicted by the SM or, alternatively, part of an extended Higgs sector as suggested by models such as the two-Higgs-doublet model (2HDM) [4,5], which can be embedded into supersymmetric models [6–11]. It predicts a total of five bosons: a light (h) and a heavy (H) CP-even Higgs boson, with the light one corresponding to the observed Higgs boson; two charged Higgs bosons (H^+ and H^-); and a CP-odd particle (a), also referred to as pseudoscalar. The additional scalar/pseudoscalar states of these models may also provide a portal into dark matter, serving as a mediator between the SM and dark matter sector [12,13]. A pseudoscalar a is also predicted in axion models [14].

This analysis performs a search for a light pseudoscalar with a mass smaller than the SM Higgs boson, produced in association either with a top-quark pair or a single top quark and a W boson, where the pseudoscalar decays into a bottom-antibottom quark pair, as proposed in [15]. It is based on a simplified model with the following Yukawa lagrangian:

$$\mathcal{L} = -\frac{g_t y_t}{\sqrt{2}} a \bar{t}(i\gamma^5)t - \frac{g_b y_b}{\sqrt{2}} a \bar{b}(i\gamma^5)b,$$

where $y_j/\sqrt{2} = m_j/v$ is the SM Yukawa coupling of particle j to the pseudoscalar a and g_j is the coupling modifier, with $j = t$ or b . Figure 1 shows two example Feynman diagrams for this process. This decay channel is favoured by many models for the range of explored pseudoscalar masses, $m_a < m_h$, although the branching ratios of the pseudoscalar depend on the specific model parameters. This is the first search for this process, exploring the couplings of the pseudoscalar to bottom quarks. Previously, the ATLAS and CMS Collaborations performed similar searches of $t\bar{t}a$ associated production exploiting leptonic decays of the pseudoscalar. The CMS Collaboration studied the decay of a pseudoscalar to the three families of leptons [16], while the ATLAS Collaboration studied the pseudoscalar coupling to muons [17]. None of these searches found significant excesses, but these decay channels are typically disfavoured compared with the $b\bar{b}$ decays when assuming a Yukawa coupling.

The search is performed in the dileptonic decay channel, with both top and antitop quarks (or both W bosons) decaying leptonically. Despite the reduced branching ratio of this decay channel, the reduced jet multiplicity of the final state and the precisely measured kinematics of the two leptons allow for a more efficient identification of the b -jets originating from the top and antitop quark decays than its semileptonic (with only one W boson from either the top or antitop quarks decaying leptonically) or fully hadronic (with the two W bosons decaying hadronically) counterparts.

For masses of the pseudoscalar below ~ 30 GeV, the $b\bar{b}$ pair has a large Lorentz-boost and is thus reconstructed as a single large-radius jet. On the other hand, for higher masses, the jets are well separated at detector level. The analysis is designed to exploit both kinematic regimes to have good signal sensitivity, making use of multiple signal regions, reconstructed objects, and machine learning techniques.

2 ATLAS detector

The ATLAS detector [18] at the LHC covers nearly the entire solid angle around the collision point.¹ It consists of an inner tracking detector surrounded by a thin superconducting solenoid, electromagnetic and hadronic calorimeters, and a muon spectrometer incorporating three large superconducting air-core toroidal magnetic systems.

¹ ATLAS uses a right-handed coordinate system with its origin at the nominal interaction point (IP) in the centre of the detector and the z -axis along the beam pipe. The x -axis points from the IP to the centre of the LHC ring, and the y -axis points upwards. Polar coordinates (r, ϕ) are used in the transverse plane, ϕ being the azimuthal angle around the z -axis. The pseudorapidity is defined in terms of the polar angle θ as $\eta = -\ln \tan(\theta/2)$ and is equal to the rapidity $y = \frac{1}{2} \ln \left(\frac{E+p_z c}{E-p_z c} \right)$ in the relativistic limit. Angular distance is measured in units of $\Delta R \equiv \sqrt{(\Delta y)^2 + (\Delta \phi)^2}$.

The inner-detector system (ID) is immersed in a 2 T axial magnetic field and provides charged-particle tracking in the range of $|\eta| < 2.5$. The high-granularity silicon pixel detector covers the vertex region and typically provides four measurements per track, the first hit generally being in the insertable B-layer installed before Run 2 [19,20]. It is followed by the SemiConductor Tracker, which usually provides eight measurements per track. These silicon detectors are complemented by the transition radiation tracker (TRT), which enables radially extended track reconstruction up to $|\eta| = 2.0$. The TRT also provides electron identification information based on the fraction of hits (typically 30 in total) above a higher energy-deposit threshold corresponding to transition radiation.

The calorimeter system covers the pseudorapidity range $|\eta| < 4.9$. Within the region $|\eta| < 3.2$, electromagnetic calorimetry is provided by barrel and endcap high-granularity lead/liquid-argon (LAr) calorimeters, with an additional thin LAr presampler covering $|\eta| < 1.8$ to correct for energy loss in material upstream of the calorimeters. Hadronic calorimetry is provided by the steel/scintillator-tile calorimeter, segmented into three barrel structures within $|\eta| < 1.7$, and two copper/LAr hadronic endcap calorimeters. The solid angle coverage is completed with forward copper/LAr and tungsten/LAr calorimeter modules optimised for electromagnetic and hadronic energy measurements respectively.

The muon spectrometer comprises separate trigger and high-precision tracking chambers measuring the deflection of muons in a magnetic field generated by the superconducting air-core toroidal magnets. The field integral of the toroids ranges between 2.0 and 6.0 Tm across most of the detector. Three layers of precision chambers, each consisting of layers of monitored drift tubes, cover the region $|\eta| < 2.7$, complemented by cathode-strip chambers in the forward region, where the background is highest. The muon trigger system covers the range $|\eta| < 2.4$ with resistive-plate chambers in the barrel, and thin-gap chambers in the endcap regions.

The luminosity is measured mainly by the LUCID-2 [21] detector that records Cherenkov light produced in the quartz windows of photomultipliers located close to the beam pipe.

Events are selected by the first-level trigger system implemented in custom hardware, followed by selections made by algorithms implemented in software in the high-level trigger [22]. The first-level trigger accepts events from the 40 MHz bunch crossings at a rate below 100 kHz, which the high-level trigger further reduces to record complete events to disk at about 1 kHz.

A software suite [23] is used in data simulation, in the reconstruction and analysis of real and simulated data, in detector operations, and in the trigger and data acquisition systems of the experiment.




Fig. 1 Feynman diagrams for (a) $t\bar{t}$ - and (b) tW -associated production of a pseudoscalar particle a that decays into a pair of b -quarks

3 Data and simulated event samples

This search is based on proton–proton (pp) collision data at a centre-of-mass energy of 13 TeV collected with the ATLAS detector at the LHC from 2015 to 2018, referred to as the Run 2 data sample in the following. After applying the data quality requirements that ensure that all subdetectors were operational, the integrated luminosity of the data sample is $140.1 \pm 1.2 \text{ fb}^{-1}$ [24].

The signal consists in the production of a light pseudoscalar a in association with a top-quark pair, $t\bar{t}a$, or in association with a single top quark and a W boson, $tW a$. The main background in this search is $t\bar{t}$ production in association with jets, followed by smaller contributions from single top quark, $t\bar{t}H$, $t\bar{t}V$, V +jets, diboson and other rare processes involving the production of a top quark. The analysis only considers the decay of the pseudoscalar to a $b\bar{b}$ pair. Decays of the pseudoscalar to other final states like a $\tau\bar{\tau}$ pair or a $c\bar{c}$ pair are not considered, as these are suppressed both by the Yukawa coupling for the masses considered (roughly a factor 5 and 20, respectively) and by the large b -jet multiplicity required in the analysis signal regions.

All signal and background samples are simulated using various Monte Carlo (MC) matrix-element (ME) generators interfaced with different algorithms for the parton shower, hadronisation, and underlying event. The effect from multiple pp interactions originating from the same or neighbouring bunch crossings, usually referred to as pile-up, is simulated by overlaying the simulated hard-scattering event with inelastic pp collisions simulated using PYTHIA 8.1 [25] and the A3 set of tuned parameters (tune) [26]. A reweighting is applied to the simulated samples such that they match the pile-up conditions in data. For the detector simulation two different approaches are used. The full ATLAS detector simulation (FS) is based on GEANT 4 [27], while the “fast” detector simulation (AF2) uses a parameterisation of the calorimeter response [28]. Most background samples are produced with

FS while signal samples are produced with AF2. Both MC and data are processed using the same reconstruction and analysis software.

The $t\bar{t}a$ signal samples are simulated with MADGRAPH5_AMC@NLO 2.3.3 [29] generator at next-to-leading order (NLO) in the strong coupling constant α_s . A simplified model based on the decoupling limit of the 2HDM+a type II is used. Additionally, the subdominant $tW a$ signal samples are simulated with the same generator at leading-order (LO). For notational simplicity, in the following $t\bar{t}a$ refers to the production of the pseudoscalar a in association with either a top-quark pair or a single top quark and a W boson. Samples are simulated for the following values of the pseudoscalar mass m_a : 12, 16, 20, 25, 30, 40, 50, 60, 80 and 100 GeV. Additional $t\bar{t}a$ signal samples for 20 and 60 GeV are also simulated with FS to check that no significant differences between the two detector simulations are observed.

The production of top-quark pairs with additional jets represents the main background source, especially the production of $t\bar{t}$ plus heavy flavour ($t\bar{t}$ + HF): $t\bar{t}$ + b -jets and $t\bar{t}$ + c -jets. For the modelling of $t\bar{t}$ + b -jets events, four flavour-scheme samples (4FS), with massive b -quarks, are simulated using the POWHEGBOX-Res framework at NLO [30] and the NNPDF3.1nnlo parton distribution function (PDF) set is used. For the modelling of $t\bar{t}$ + c -jets and $t\bar{t}$ +light-jets events, five flavour-scheme (5FS) samples with massless b -quarks are simulated using POWHEGBOX-v2 [31–34], also at NLO. To avoid double-counting of events, neither the $t\bar{t}$ + b -jets events from the 5FS samples nor the $t\bar{t}$ + c -jets nor the $t\bar{t}$ +light-jets events from the 4FS samples are used. The simulated $t\bar{t}$ events are categorised based on the number of additional jets matched to b - or c -hadrons with transverse momentum p_T larger than 5 GeV within $\Delta R < 0.3$ of the jet axis.

To model the production of single-top-quark events, which mostly contribute through the tW -channel, and the produc-

tion of $t\bar{t}H$, the same POWHEGBOX-v2 [31–34] settings, as used in the $t\bar{t}$ +light/ c -jets production, are used. The production of $t\bar{t}Z$ and $t\bar{t}W$ events is modelled using the MADGRAPH5_AMC@NLO2.3.3 [29] generator at NLO. For all samples listed above, the NNPDF3.0nlo [35] PDF sets are used and a top-quark mass of $m_{\text{top}} = 172.5$ GeV is set. The events are interfaced with PYTHIA8.230 [36] using the A14 tune [37] and the NNPDF2.3lo set of PDFs [38] for the parton shower and hadronisation modelling.

The production of tZq and tWZ events is performed using MADGRAPH5_aMC@NLO v2.3.3, at LO and NLO in QCD respectively, in the 4FS with the CTEQ6L1 PDF set [39] and using PYTHIA 8.212 for the parton shower.

Finally, the production of V +jets and diboson samples (VV) is simulated with different versions of the SHERPA [40] generator and the simulated events are matched with the SHERPA parton shower [41] using the MEPS@NLO prescription [42–45] with the set of tuned parameters developed by the SHERPA authors. All samples and their basic generation parameters are summarised in Table 1.

For all samples, except those generated with SHERPA, the decays of b - and c -hadrons are simulated using the EVTGEN programme [46].

4 Object definition

In this section, the reconstruction and definition of the physics objects are described, together with the additional corrections applied to each.

4.1 Physics objects

Electron candidates are reconstructed from energy deposits (clusters) in the electromagnetic calorimeter associated with reconstructed tracks in the inner detector. Candidates are selected with $p_T > 10$ GeV and $|\eta| < 2.47$, excluding the calorimeter transition region $1.37 < |\eta| < 1.52$. Electrons satisfy the `TightLH` [58] likelihood-based identification criterion and are required to match the `PromptLeptonTagger` working point `PLVLoose` [59]. They are further required to have $|z_0 \sin \theta| < 0.5$ mm and a d_0 significance $|\frac{d_0}{\sigma(d_0)}| < 5$, where the transverse impact parameter (d_0) is calculated relative to the beamline, and the longitudinal impact parameter (z_0) as the longitudinal distance between the primary vertex and the point where d_0 is measured.

Muon candidates are reconstructed from track segments in the various layers of the muon spectrometer and matched with tracks from the inner detector. The final muon candidates are refitted using the complete track information from both detector systems and must have $p_T > 10$ GeV and $|\eta| < 2.5$. Muons are required to satisfy the `Medium` quality requirements and match the `PromptLeptonTagger` working

point `PLVLoose` [60]. Further requirements are $|\frac{d_0}{\sigma(d_0)}| < 3$, and $|z_0 \sin \theta| < 0.5$ mm.

Small- R jet candidates are reconstructed by clustering particle flow objects [61] using the anti- k_t algorithm [62,63] with a radius parameter of $R = 0.4$ and a four-momentum recombination scheme. The energy of the jet is corrected to the particle level by the application of a jet energy scale calibration derived from $\sqrt{s} = 13$ TeV data and simulation [64]. Baseline jets are required to have $p_T > 20$ GeV and $|\eta| < 2.5$. For pile-up rejection, jets with $p_T \in [20, 60]$ GeV and $|\eta| < 2.4$ are required to have a jet vertex tagger weight [65] larger than 0.5.

The b -tagging is the identification of jets that originate from the decay of b -hadrons using dedicated algorithms. A deep neural-network, called `DL1r` [66–69] is used. Small- R jets with a `DL1r` score above a certain threshold are defined as b -tagged jets. The pseudo-continuous (PC) b -tagging working point is used: each jet is classified with an integer from one to five depending on how many calibrated b -tagging working points (WP) the jet fulfils. The four calibrated `DL1r` WPs are 85%, 77%, 70% and 60%, each corresponding to the approximate average b -tagging efficiency in an inclusive $t\bar{t}$ MC sample. Jets not satisfying any WP are assigned a value of one, and this value is increased by one for every WP that they fulfil. The sum of the PC b -tagging over all jets in an event is defined as `sumPCBTag`. Jets satisfying the 70% WP are referred to as b -jets, while jets satisfying the 85% WP, but not the 70% WP, are classified as loose- b -jets.

Large- R jet candidates are formed by reclustering the small- R jets and tracks with a larger radius parameter of $R = 0.8$ using the anti- k_t algorithm [62,63]. The larger radius for track association allows more tracks from the targeted double b -hadron decays to be associated with the reclustered jet. The tracks in and around the small- R jet associated with the reclustered jet through ghost association [70,71] are selected with a loose track selection [72]. In this procedure, the p_T of the tracks is set to infinitesimal values, such that the “ghost” tracks can then be reclustered with the constituents of the reclustered jets with the appropriate radius parameter. Since the p_T of the tracks is infinitesimally small, they do not influence the reconstruction of the jet, allowing the use of additional tracks that leak outside the small- R jets.

To resolve the substructure within a large- R jet originating from a boosted $X \rightarrow b\bar{b}$ decay, which the small- R jet reconstruction fails to completely capture, additional information is extracted from the large- R jet by reconstructing track-subjets inside the large- R jet. The track-subjets are derived using the tracks that are ghost associated to each large- R jet as inputs to the exclusive- k_T method [73]. The selected tracks for a given jet are clustered using the k_T algorithm with a radius parameter of $R = 0.8$. The clustering stops when there are exactly two track clusters left.

Table 1 Nominal simulated signal and background event samples. The matrix element generator, PDF set, parton-shower (PS) generator and calculation accuracy of the cross section in QCD and EW used for normalisation are shown. MADGRAPH is abbreviated to MG

Process	Matrix element generator	PDF set	PS generator	Normalisation
$t\bar{t}a, a \rightarrow b\bar{b}$	MG_aMC@NLO v2.3.3	NNPDF3.0 NLO	PYTHIA 8.230	–
$tW a, a \rightarrow b\bar{b}$				
$t\bar{t}$ + jets	POWHEGBOX-v2	NNPDF3.0 NLO	PYTHIA 8.230	(NLO+NNLL) _{QCD} [47–53]
$t\bar{t} + b\bar{b}$	POWHEGBOX-Res	NNPDF3.1 NNLO	PYTHIA 8.244	–
Single-top	POWHEGBOX-v2	NNPDF3.0 NLO	PYTHIA 8.230	(NLO+NNLL) _{QCD} [54,55]
$t\bar{t}H$	POWHEGBOX-v2	NNPDF3.0 NLO	PYTHIA 8.230	NLO _{QCD+EW} [56]
$t\bar{t}Z$	MG5_aMC@NLO v2.3.3	NNPDF3.0 NLO	PYTHIA 8.210	NLO _{QCD+EW} [56]
$t\bar{t}W$	MG5_aMC@NLO v2.3.3	NNPDF3.0 NLO	PYTHIA 8.210	NLO _{QCD+EW} [56]
tZq, tWZ	MG5_aMC@NLO v2.3.3	CTEQ6L1	PYTHIA 8.212	–
		NNPDF3.0 NLO		
Z/W + jets	SHERPA 2.2.11	NNPDF3.1 NNLO	SHERPA	NNLO _{QCD} [57]
Diboson	SHERPA 2.2.1	NNPDF3.1 NNLO	SHERPA	–
	SHERPA 2.2.2			

These clusters are used as the track-subjets associated with a given jet. For signal events, each track-subjet should originate from the decay of one b -hadron ideally. The associated large- R jet is required to satisfy $|\eta| < 2.0$ to account for their extended radius and the acceptance of the ID. Furthermore, each track-subjet is required to satisfy $p_T > 5$ GeV where the track-subjet p_T is estimated from the sum of its constituent tracks' four-momenta. The four-momentum of the large- R jet is defined as the sum of the four-momenta of its track-subjets.

In addition, secondary vertices (SV) inside the large- R jets are reconstructed to help the identification of b -hadrons. For this purpose, an algorithm that combines the track-cluster-based low- p_T vertex tagger (TC-LVT) [74] and the multiple-secondary-vertex finder algorithms (MSVF) [75] is used. The TC-LVT algorithm was developed for soft b -hadron tagging and optimised to reconstruct low- p_T b -hadron decays. The clustering algorithm from TC-LVT is used to identify displaced tracks not originating from the primary vertex. The MSVF algorithm is used to identify multiple SVs in the track cluster. The algorithm builds all two-track proto-vertices consistent with displaced tracks that are not compatible with a hadronic material interaction, a photon conversion, or the decay of long-lived light-flavoured hadrons. All displaced tracks reconstructed in the ID are used to build proto-vertices. Proto-vertices define track-to-track relations, since a single track can be associated with more than one proto-vertex. Each set of tracks that are mutually connected to each other forms a secondary vertex. After secondary vertices are formed, tracks not compatible with the vertex are removed, and the ambiguity caused by distant vertices sharing common tracks is resolved. Nearby vertices are also merged by the MSVF algorithm. Finally, reconstructed SVs are required to be ΔR -

matched to a large- R jet. Further details and studies about the large- R jets can be found in Ref. [76].

In contrast to b -tagging, the B -tagging is the identification of pairs of b -jets that are too close to be resolved and identified individually. For this purpose the DeXTer tagger is used. It is a double b -tagger based on a deep sets neural network (NN) architecture designed to do flavour tagging of merged reconstructed jets [77] and uses information of the SVs and jet kinematics. This is done in two transverse momentum ranges: a low p_T range between 20 and 200 GeV and a high p_T one, above 200 GeV. Two working points are defined: the 0–40% tagging interval is referred to as **Tight** WP, and the **Loose** WP is defined by the inclusive 40–60% tagging interval. A sample of Z +jets and $t\bar{t}$ events is used to measure the DeXTer efficiency in data, and to derive B -tagging and b -mistagging rate correction factors for the simulated events. Large- R jets satisfying the **Tight** WP are referred to as B -jets.

The missing transverse momentum E_T^{miss} measures the event momentum imbalance in the transverse plane of the detector. It is defined as the magnitude of the negative vector sum of p_T for all selected and calibrated physics objects in the event, with an extra term added to account for soft energy that is not associated with any of the selected objects. This soft term is calculated from inner detector tracks matched to the primary vertex to make it more resilient to pile-up contamination [78]. The E_T^{miss} computation is based on the momenta of the objects defined previously, and after applying the overlap removal procedure defined in the next section.

4.2 Corrections to physics objects

The overlap removal is the procedure followed to prevent double-counting of objects. First, electrons that share a track with a muon are removed. To prevent double-counting of electron energy deposits as jets, the closest small- R jet within $\Delta R < 0.2$ of a selected electron is removed. If the nearest jet surviving that selection is within $\Delta R = 0.4$ of the electron, the electron is discarded. To reduce the background from muons from heavy-flavour decays inside jets, muon candidates are required to be separated by $\Delta R > 0.4$ from the nearest small- R jet, removing the muon if the jet has at least three associated tracks, and removing the jet otherwise. This avoids an inefficiency for high-energy muons undergoing significant energy loss in the calorimeter.

To avoid double-counting of jets, the jet overlap removal is done as follows. First, every small- R jet is tested to see if it is eligible to be DeXTer-tagged. This requires the small- R jet to have $p_T > 20$ GeV and be isolated, meaning it is the only constituent of its reclustered jet [79] with the anti- k_t algorithm and radius parameter $R = 0.8$. These jets are then tested by the DeXTer tagger: if a small- R jet passes the Loose working point selection, it is defined as a B -tagged large- R jet and removed from the small- R jet list. Otherwise, it is kept as a small- R jet. Thus, this first step gives two jet lists: the DeXTer-tagged large- R jets and the small- R jets, which are either not eligible for DeXTer tagging or fail the tagger. Finally, the jet overlap removal procedure with leptons is repeated for large- R jets using $\Delta R = 0.8$.

The μ -in-jet p_T correction is the procedure of adding the muons reconstructed inside of a b -jet to the four-momentum of the respective b -jet. Around 20% of all b -hadron decays produce a low-momentum or soft muon inside of the resulting jet, but those soft muons are removed in the overlap removal as described above. This correction recovers the original energy of those b -jets and reduces biases from the invariant masses calculated with them. The soft muons used are required to have $p_T > 4$ GeV and $|\eta| < 2.5$, and fulfil the Medium soft muon quality requirement.

In the case of the DeXTer-tagged jet, the μ -in-jet p_T correction is carried out as follows. First, the soft muons are matched to track-subjets within an angular distance of $\Delta R < 0.3$. At most the two highest p_T soft muons are taken into account for each track-subjet, and any muon is only matched once to the closest subjet. At last, the matched muons are added to the four-momentum of the track-subjet.

5 Event selection

Only events recorded with a single-electron [80] or single-muon trigger [81] under stable beam conditions and for which all detector subsystems were operational are considered [82].

Single-lepton triggers with p_T thresholds varying from 20 to 140 GeV, depending on the year, lepton flavour, isolation requirement and luminosity, are combined in a logical OR to increase the overall efficiency. The triggers with the lower p_T thresholds include isolation requirements on the lepton candidate, resulting in inefficiencies at high p_T that are recovered by the triggers with higher p_T thresholds.

5.1 Preselection

Events are required to have exactly two leptons (electrons, muons, or both) with opposite charge, satisfying the criteria defined in Sect. 4. Since single-lepton triggers are used, at least one of the two reconstructed leptons is required to have a $p_T > 27$ GeV and match a lepton with the same flavour reconstructed by the trigger algorithm within ΔR of 0.15. The chosen p_T threshold ensures a fully efficient trigger for the whole Run 2 period. In the ee and $\mu\mu$ channels, the dilepton invariant mass must be above 15 GeV and outside the Z boson mass window 83–99 GeV. Further suppression of the background is achieved by requiring at least three jets (either large- R or small- R) with at least one of which being b -tagged using the DL1r 85% WP. The fraction of signal events in the preselection region is negligible for all masses.

5.2 Signal and control regions

After preselection, the data sample is dominated by background from $t\bar{t}$ events. To take advantage of the high jet and b -object multiplicities of the $t\bar{t}a$ signal process, events are classified into non-overlapping regions based on the total number of B -jets and b -jets. Some of the regions also require the presence of at least one loose (and not tight) b -jet (small- R jets tagged with the DL1r 85% WP but failing to meet the 70% WP). The name of every signal region (SR) or control region (CR) includes the number of B -jets followed by “B” and of b -jets followed by “b”. The names of those regions requiring at least an extra loose b -jet indicate it in their name with “+1bL”. Due to the high b -jet multiplicity of the signal, only regions with at least three b -objects are considered as signal regions (B -jets count as two b -objects). To maximise the signal sensitivity, signal events are classified into two boosted regions (SR 1B1b+1bL and SR 1B2b) and two resolved regions (SR 0B3b and SR 0B4b). The loose b -tagged jet in one of the signal regions is required to suppress $t\bar{t}$ -light events. The complementarity between boosted and resolved regions is illustrated in Fig. 2, which shows the invariant mass of the B -jet in SR 1B2b and the invariant mass of the pair of b -jets with the largest p_T in SR 0B4b for different values of the pseudoscalar mass. Regions containing one B -jet are particularly relevant in the boosted regime ($m_a < 30$ GeV). Regions with no B -jets and a high b -jet multiplicity are more powerful in the resolved regime ($m_a > 30$ GeV).




Fig. 2 Invariant mass of the (a) B -jet in SR 1B2b and of the (b) pair of b -jets with the largest p_T in SR 0B4b for various values of m_a . The distributions are normalised according to the predicted m_a -dependent theoretical cross sections with a Yukawa coupling of the a -boson to the top quark of 0.5

Table 2 Overview of the jet multiplicities considered per region in the fit

Region	Large- R jets	Small- R jets	B -jets	b -jets	Loose b -jets
SR 0B4b	≥ 0	≥ 4	$= 0$	≥ 4	–
SR 0B3b	≥ 0	≥ 3	$= 0$	$= 3$	–
SR 1B2b	≥ 1	≥ 2	$= 1$	≥ 2	–
SR 1B1b+1bL	≥ 1	≥ 2	$= 1$	$= 1$	≥ 1
CR 0B2b+1bL	≥ 0	≥ 3	$= 0$	$= 2$	≥ 1

In addition to the four signal regions described above, a control region is included in the fit to improve the $t\bar{t}+\geq 1c$ normalisation (Sect. 8). The control region (CR 0B2b+1bL), orthogonal to all four signal regions, is composed of events with no B -jets and exactly two b -jets, as well as at least one additional loose (and non-tight) b -jet. Similarly to the SR, the loose b -tagged jet is required to suppress $t\bar{t}$ +light events that would otherwise dominate the control region. Finally, events entering the CR are required to have a sum of the pseudo-continuous b -tagging scores between 12 and 15. The number of b -jets in the 0B3b, 1B1b+1bL and 0B2b+1bL regions is exclusive, while it is inclusive in the 0B4b and 1B2b regions. Table 2 summarises the selections for each region, which are applied in addition to the previously mentioned preselection requirements.

6 Background estimate

Data-driven corrections are derived for the $t\bar{t}$ Monte Carlo simulation, the main background process in this search. These corrections are derived to improve the description

of the rates of $t\bar{t}$ plus heavy flavour jets and the transverse momenta of lepton and jets observed in data, using a method similar to the one developed for other ATLAS searches [83–85]. The corrections are derived in very inclusive control regions where the contamination of signal is predicted to be below 1% for all considered pseudoscalar mass hypotheses. The region of choice satisfies the preselection requirements described in Sect. 5.1, with an extra requirement of at least two b -jets and no B -jets. Additionally, to suppress the Z +jets contribution, only the different lepton flavour ($e\mu$) region is considered.

The first correction targets the production rate of heavy-flavour jets: c -jets and b -jets. It was observed in previous ATLAS and CMS analyses [86–88] that the rate of $t\bar{t}$ +HF events is underestimated in MC simulation. Due to the high b -object multiplicity of the $t\bar{t}a$ signal, these HF events represent a large fraction of the $t\bar{t}$ +jets background in the signal regions, and therefore the MC simulation must be corrected. To have a more accurate flavour composition, an event reweighting procedure is applied based on the sumPCBTag distribution. Figure 3a shows how the $t\bar{t}$ +light production is dominant at low values of sumPCBTag, while the $t\bar{t}$ +HF




Fig. 3 Comparison of the data versus MC distribution of the sum of the pseudo-continuous b -tagging score of all the jets per event (a) before and (b) after applying the normalisation factors extracted from the heavy-flavour correction fit

Table 3 Normalisation factors for the three $t\bar{t}$ +jets HF categories resulting from the likelihood fit performed using the sumPCBTag distribution

HF category	Norm. factor
$t\bar{t}$ + light, tW	0.91 ± 0.03
$t\bar{t} + \geq 1c$	1.58 ± 0.14
$t\bar{t} + \geq 1b$	1.13 ± 0.07

production populates the tail of the distribution. The correction procedure consists in deriving three normalisation factors: one for each component, using a likelihood fit to the sumPCBTag MC distributions compared with data. In this fit, DL1r b -tagging systematic uncertainties (detailed in Sect. 9) are included. Figure 3b shows the improved agreement after the fit. The normalisation factors and their corresponding uncertainties are shown in Table 3.

This procedure also corrects the distribution of the number of jets (inclusive of all jet types) per event, N_{jets} , which was also observed to be mismodelled in the simulation. Figure 4 shows the corresponding distribution, before and after applying the normalisation factors from Table 3. In the following, in all Figures and Tables, the category “Other” includes the following minor background processes: Z +jets, $t\bar{t}W$, tq , tZ , tWZ , WW , ZZ , WZ and W +jets.

The second correction targets the transverse momenta of the jets and leptons originating from the decay of top quark/antiquark, a quantity that was also observed to be mis-

modelled by current $t\bar{t}$ 5FS MC generators. The disagreement between data and MC persists even after applying the $t\bar{t}$ +HF correction. To improve the agreement between data and MC for these variables, a kinematic reweighting factor for the $t\bar{t} + \geq 1c$, $t\bar{t}$ +light, and tW components is derived from the data/MC ratio, after subtracting other background components from the data. These mismodellings are assumed to be independent of the flavour of the extra radiation, and applied equally to $t\bar{t}$ +light, $t\bar{t} + \geq 1c$, and tW .

The event hardness or H_T , which is defined as the scalar sum of the p_T of all the jets and leptons in the event, is largely correlated with the total number of jets in the event, as every additional jet in the event shifts H_T to larger values. Performing the kinematic reweighting directly with H_T would therefore spoil the data/MC agreement achieved after the HF correction to the number of jets distribution shown in Fig. 4. To reduce the N_{jets} dependency, a new variable, $H_T^{\text{red}}(n)$, is defined:

$$H_T^{\text{red}}(n) = H_T - (n - 3)\Delta H_T(n),$$

where n is the number of jets (small- R and large- R jets, with a minimum of three jets) and $\Delta H_T(n)$ is the average offset in H_T caused by the addition of each extra jet to the event. Correction factors are derived over a binned H_T^{red} distribution, and the results are fitted using a continuous hyperbolic function which is later used for the MC reweighting. Figure 5 shows the H_T distribution before and after applying the




Fig. 4 Comparison of the data versus MC N_{jets} distribution (a) before and (b) after applying the $t\bar{t}$ +jets normalisation factors extracted from the heavy flavour correction fit. Only statistical uncertainties are shown

correction. Similar improvements are observed for individual leptons and jets. No significant changes in the number of jets distribution are observed after the kinematic corrections are applied. Following the same procedure, dedicated kinematic reweighting corrections are derived for the alternative $t\bar{t}$ +light, $t\bar{t}$ + $\geq 1c$, and tW samples employed in the evaluation of systematic uncertainties described in Sect. 9. Residual differences between data and the MC are taken into account in the analysis fit by including free-floating individual normalisations for the various $t\bar{t}$ +jets background contributions.

7 Analysis strategy

The analysis uses various machine learning (ML) algorithms to improve the sensitivity to the target signal. First, two different boosted decision trees (BDTs) are trained to identify the jets originating from the decay of the top quark and antiquark and the jets originating from the pseudoscalar decay. Second, a mass-parameterised NN is trained for signal/background discrimination in each of the SRs described in Sect. 5.2. The final fit uses the NN output score distribution in each of the four SRs and the $\text{sumPCBT}_{\text{tag}}$ distribution in the CR. In addition to the signal strength μ , three normalisation factors, corresponding to the main background contributions, are left to freely float in the fit. Figure 6 shows a diagram summarising the ML approach followed in the analysis as well as the CR and SRs used in the final fit. Further details on each step are given in the following.

7.1 Event reconstruction BDTs

Two different BDTs are trained to do partial event reconstruction. One targets the identification of the lepton-jet pairs associated with the top quark/antiquark decays, while the other identifies the pair of jets from the pseudoscalar decay. The two BDTs use the five leading small- R jets, together with the two charged leptons in the case of the top quark/antiquark BDT, and in each case select the pair of jets or the lepton/jet pair most likely to correspond to the pseudoscalar or the top quark/antiquark, respectively. Both BDTs were designed to reconstruct resolved topologies, thus they do not use large- R track-jets. Also, no attempt to reconstruct the two neutrinos from the top quark/antiquark decay is made. The two BDTs are trained using all signal samples inclusively, such that they are generic and valid for all considered masses. During the BDT training process, target labels for each jet are assigned based on a one-to-one matching between reconstructed jets at the detector level and parton-level b -quarks/leptons. Consequently, a reconstructed jet/lepton is assigned as originating from a (anti-)top quark, or pseudoscalar decay candidate based on the aforementioned generator information.

The BDT targeting the top quark/antiquark decay attempts to pair each lepton with its corresponding b -jet, considering each lepton in turn. For this, the BDT receives as input several kinematic variables that depend on the tag lepton/jet pair (jl -pair), such as its invariant mass or transverse momentum, or the separation angle between the lepton and the jet. It also uses information about the lepton and jet candidates them-




Fig. 5 Comparison between data and MC of the H_T distribution (a) before and (b) after correcting it using H_T^{red} . Only statistical uncertainties are shown




Fig. 6 Diagram of the analysis strategy, illustrating the data-driven corrections, jet/lepton and dijet pairings by two BDTs and the NN for signal versus background discrimination. The CR and SRs are used in the final fits to extract the signal strength (μ) as well as normalisation factors for the main backgrounds

selves, such as their pseudorapidity and transverse momenta, or the jet index indicating in decreasing order the hardness of the jets. In addition, the BDT uses information from the auxiliary jl -pair built with the lepton that is not being evaluated, together with information from the top/antitop system formed by the tag and auxiliary jl -pairs, and variables that refer to the full event. In a similar way, the BDT targeting the pseudoscalar decay receives various kinematic variables connected to the pair of two jets, jj -pair, as its mass or transverse momentum, together with information about the jets themselves or about the overall event. The full list of variables used by each BDT is shown in Table 4.

For the training of both BDTs, generator information is used to define the targets (b -quarks and leptons from the decays of the top quark, top antiquark and pseudoscalar a) and to identify the correct permutations at detector level, which are used as signal (or target) during the training, while all wrong permutations are used as background. A mix of signal and $t\bar{t}$ samples is used during the training of both BDTs. In both cases, two sets of BDTs are trained using k-2 fold training to avoid biases. The BDT trained with odd events is applied to even events and vice versa. The training is performed with the TMVA package of ROOT [89].

Following the training of both BDTs, they are applied to data and MC as follows. For the top quark/antiquark BDT, the lepton/jet permutation with the highest BDT score is identified for each lepton as the most likely jl -pair and the selected jet is assigned to the top quark or antitop quark decay depending on the lepton charge. If both leptons are initially assigned to the same jet, only the one with the highest BDT score keeps the assignment, while the other lepton is reassigned to the second most likely jet in terms of BDT score. In a similar way, for the pseudoscalar BDT, the permutation of two jets with the highest BDT score is selected and the two corresponding jets are assigned to the pseudoscalar decay. The selected jl - and jj -pairs are used to define related variables, such as the top-quark or pseudoscalar reconstructed invariant mass or separation angles, that are later used as input by the signal-versus-background discrimination neural networks. Figure 7 shows the prefit data/MC comparison of the reconstructed mass of the lb -pair selected by the top-quark BDT and of the jj -pair selected by the pseudoscalar BDT in SR 0B3b, the signal region with the largest statistics.

7.2 Signal-versus-background discriminating neural networks

As described in Sect. 5.2, events are divided into four signal regions according to their B - and b -jet multiplicity to better separate signal from background. The four signal regions used in the final fit are SR 0B4b, SR 0B3b, SR 1B2b and SR 1B1b+1bL. Independent NNs are trained individually per region to separate signal from background.

To make better use of the MC samples in the training, five different trainings are performed independently per region, where 80% of the events in the region constitute the training sample and the remaining 20% are used as the validation sample. An appropriate distribution of events in the various samples guarantees that no event is used both in the training and the evaluation of the NNs.

Each NN contains two hidden layers with twice as many nodes as the input layer, connected by Rectified Linear Unit (ReLU) activation functions. The final layer is a single node, normalised by a sigmoid function. The dropout for every layer is set to 0.3. To avoid overtraining, early stopping is implemented when the validation loss function does not improve during the last four epochs. The training is done using PyTorch 1.13.1 [90], and each of the NNs combines basic four-momentum information with high-level variables, such as invariant masses or angular distances, as well as relevant variables from the BDTs described in Sect. 7.1. The full list of input variables depends slightly on the region, given the slightly different signal topologies per region. Table 5 shows the overall list of input variables used by the NNs. Some of the most important variables in the NNs are H_T^{jets} , the invariant mass of two small- R jets or the mass and p_T of the large- R jet, among others.



All NNs are mass-parameterised, meaning that they receive the mass hypothesis as input during the evaluation. For the training, background MC samples are randomly assigned a mass from the grid of generated signal samples, while appropriate mass values are assigned to the signal events. Once the NNs are trained, the data scores are evaluated for each value of m_a considered in the analysis. In each SR, the resulting NN score is the distribution used in the profile likelihood fit, as discussed in the next section. Figure 8 shows the prefit distributions of the four NN scores corresponding to the 30 GeV mass hypothesis.

8 Statistical treatment

To test for the presence of a $t\bar{t}a$ signal, for each mass hypothesis, a binned maximum-likelihood fit to the data is performed simultaneously in all SRs and the CR (Sect. 5.2). In each SR, the input to the fit is the corresponding NN distribution described in Sect. 7.2, evaluated at the appropriate mass hypothesis. In the CR, the input to the fit is the sumPCBTtag distribution. The parameter of interest is the signal strength, μ , a multiplicative factor to the cross section of the signal process. In addition to the signal strength μ , the fit includes three additional free parameters that work as scale factors to the normalisation for the three main background components: $k(t\bar{t}+\text{light}, tW)$, $k(t\bar{t}+\geq 1c)$, and $k(t\bar{t}+\geq 1b)$. To estimate the signal strength, a binned likelihood function $\mathcal{L}(\mu, \theta)$ is used,

Table 4 Input variables used in the (a) top quark/antiquark reconstruction BDT and (b) pseudoscalar reconstruction BDT

(a) Top quark/antiquark reconstruction BDT		(b) Pseudoscalar reconstruction BDT	
Object	Variables	Object	Variables
Full event	$N_{\text{jets}}, N_{b\text{-jets}}$	Full event	$N_{\text{jets}}, N_{b\text{-jets}}, \text{sumPCBTag}$
Lepton (tag, aux.)	p_T, η	Jet (1st, 2nd)	$p_T, \eta, \text{PC } b\text{-tag, jet index}$
Jet (tag, aux.)	$p_T, \eta, \text{PC } b\text{-tag, jet index}$	jj pair	$m, p_T, \eta, E, \phi, \Delta R$
jl pair (tag, aux.)	$m, p_T, \eta, \Delta R$		
$t\bar{t}$ pair	$m, p_T, \eta, \Delta R, \Delta\phi$		
jj pair	ΔR		

**Fig. 7** Distribution (a) of the antitop-quark m_{jl} and (b) pseudoscalar m_{jj} selected by the top quark and pseudoscalar BDTs, respectively in SR 0B3b before the fit. The dashed line shows the distribution of the

30 GeV signal normalised to the total or half the total number of events in the region. The band displays the total pre-fit uncertainty

Table 5 List of input variables of the signal-versus-background discrimination NN. The distributions corresponding to both the pair with the maximum p_T and minimum ΔR are included for bb variables. Angular variables with one b or one B use the pair with the minimum ΔR . The m_{bbbb} and m_{bbb} variables correspond to the combination with the maximum scalar sum of p_T

Object	Variables
Full event	$N_{\text{jets}}, H_T^{\text{jets}}, E_T^{\text{miss}}$
BDT $t \rightarrow jl$	Score, $p_T^{jl}, \Delta R_{jl}, \Delta \eta_{jl}, \Delta \phi_{jl}, \text{jet index}$
BDT $a \rightarrow jj$	Score, $p_T^{jj}, \eta_{jj}, m_{jj}, \Delta R_{jj}, \Delta \eta_{jj}, \Delta \phi_{jj}, \text{jet index}$
Leptons	$\Delta R_{ll}, \Delta \eta_{ll}, \Delta \phi_{ll}, \Delta \phi_{E_T^{\text{miss}}, l}, \Delta R_{ll,bb}, \Delta R_{ll,B}, \Delta R_{ll,b}$
Large- R jets	$p_T, \eta, m, \Delta R_{Bb}, \Delta \phi_{E_T^{\text{miss}}, B}$
Small- R jets	$p_T^{bb}, m_{bb}, m_{bbb}, m_{bbbb}, \Delta R_{bb}, \Delta \eta_{bb}, \Delta \phi_{bb}, \Delta \phi_{E_T^{\text{miss}}, b}$
	$p_T, \eta, \text{PC } b\text{-tag}$




Fig. 8 Pre-fit distributions corresponding to the NN output score of (a) SR 0B4b, (b) SR 0B3b, (c) SR 1B2b and (d) SR 1B1b+1bL for the 30 GeV mass hypothesis fit. The dashed line shows the distribution of

signal normalised to the total number of events in each region. The band displays the total pre-fit uncertainty. Arrows appearing in the bottom panels indicate the ratio being outside the displayed range

$$\mathcal{L}(\mu, \theta) = \prod_i \frac{(E[n_i(\mu, \theta)])^{n_i}}{n_i!} e^{-E[n_i(\mu, \theta)]} \prod_{\theta_j \in \theta} \rho(\theta_j | \tilde{\theta}_j).$$

The function is constructed as a product of Poisson probability terms with one Poisson term included for every bin i

of the NN distribution in the analysis regions. The binning of the NN distributions for each signal is chosen to provide a good separation of signal and background while maintaining a stable performance of the fit. The expected number of events, $E[n_i(\mu, \theta)]$, in each bin, n_i , is a function of μ ,

and a set of nuisance parameters, θ . The nuisance parameters encode effects from the normalisation of backgrounds, including the systematic uncertainties and one parameter per bin to model statistical uncertainties in the simulated samples. Unlike the free-floating parameters, all nuisance parameters are constrained by prior distributions, $\rho(\theta|\hat{\theta})$, which follow Gaussian, log-normal, or Poisson distributions centred around their nominal values, $\hat{\theta}$. This procedure allows the reduction of the impact of the uncertainties by taking advantage of the separated populations of signal and background. The best-fit value of the signal strength is obtained by performing a fit to the data under the signal-plus-background hypothesis, maximising $\mathcal{L}(\mu, \theta)$ over μ and θ . To set upper limits on μ , the following test statistic is used:

$$\tilde{q}_\mu = \begin{cases} -2 \ln \frac{\mathcal{L}(\mu, \hat{\theta}(\mu))}{\mathcal{L}(0, \hat{\theta}(0))} & \hat{\mu} < 0, \\ -2 \ln \frac{\mathcal{L}(\mu, \hat{\theta}(\mu))}{\mathcal{L}(\hat{\mu}, \hat{\theta})} & 0 \leq \hat{\mu} \leq \mu, \\ 0 & \hat{\mu} > \mu. \end{cases}$$

The values of the signal strength and nuisance parameters that maximise the likelihood function are represented by $\hat{\mu}$ and $\hat{\theta}$, respectively. For a given value of μ , the values of the nuisance parameters that maximise the likelihood function are represented by $\hat{\theta}(\mu)$. This test statistic measures the compatibility of the observed data with the background-only hypothesis ($\mu = 0$), represented by the p -value, and is estimated by integrating the distribution of $\tilde{q}_0$ based on the asymptotic formula in Ref. [91]. The test statistic is set to zero for $\hat{\mu} > \mu$, as this case indicates that the μ hypothesis is compatible with the observed data and cannot be rejected. Upper limits on μ are derived by using $\tilde{q}_\mu$ in the CL_s method [92, 93].

The systematic uncertainties, including those derived from MC samples, can show fluctuations due to generator weights or statistical variations. To ensure the quality of the templates and the stability of the fit, smoothing algorithms are applied to the histograms before the fit. In addition, systematic uncertainties are pruned to reduce computing time. Only uncertainties with an effect greater than 1% are included in the fit. This is done separately for shape and normalisation effects.

9 Systematic uncertainties

Various sources of systematic uncertainties are considered. Each systematic uncertainty is introduced as a nuisance parameter (NP) in the statistical analysis described in Sect. 8. Section 9.1 describes all experimental uncertainties, related to the luminosity and pile-up or the reconstruction and iden-

tification of jets and leptons. They are applied to all MC samples equally and their effects are treated in a correlated way across all four SRs and the CR in the final fit. The signal and background modelling uncertainties are detailed in Sect. 9.2, and can be different depending on the process. They are implemented as decorrelated between regions, given their different coverage of phase spaces, and decorrelated between signal and background samples in the fit.

9.1 Experimental uncertainties

Luminosity and pile-up modelling. The uncertainty in the integrated luminosity for the full Run 2 data sample is 0.83% [24], obtained using the LUCID-2 detector [21] for the primary luminosity measurements. A variation in the pile-up reweighting of simulated events is included to cover the uncertainty in the ratio of the simulated and measured distribution of inelastic cross sections.

Leptons. Uncertainties associated with leptons are related to the trigger, reconstruction, identification and isolation, as well as the lepton energy or momentum scale and resolution. The reconstruction, identification, and isolation efficiency of electrons and muons, as well as the efficiency of the trigger used to record the events, differ slightly between data and simulation, and is corrected by dedicated scale factors. Efficiency scale factors are measured using tag-and-probe techniques on $Z \rightarrow ll$ data and simulated samples [58, 60], and are applied to the simulation to correct for differences. Additional sources of uncertainty originate from the corrections applied to adjust the lepton momentum scale and resolution in the simulation to match those in data, measured using $Z \rightarrow ll$ and $J/\psi \rightarrow ll$ events [58, 60].

Jets. Uncertainties associated with jets arise from the efficiency of pile-up rejection by the jet vertex tagger (JVT), from the jet energy scale (JES) and resolution (JER), and from the different flavour-tagging algorithms used, DL1r and DeXTer. Scale factors are applied to correct for discrepancies between data and MC for JVT efficiencies, and are estimated by using $Z \rightarrow \mu\mu$ with tag-and-probe techniques [65]. The jet energy scale and its uncertainty are derived by combining information from test-beam data, LHC collision data and simulation [64]. The jet energy resolution is measured in Run 2 data and simulation as a function of jet p_T and rapidity using dijet events.

To correct flavour-tagging efficiencies in simulated samples to match those measured in data, scale factors are derived. They are calculated as a function of p_T for b -jets, c -jets, and light jets separately in dedicated calibration analyses. For b -jet efficiencies, $t\bar{t}$ events in the dilepton topology are used, exploiting the very pure sample of b -jets arising from the decay of the top quarks [67]. For c -jet mistag rates, $t\bar{t}$ events in the single-lepton topology are used, exploiting c -jets from the hadronically decaying W boson [68].

The negative-tag method is used in Z +jets events [69] for light-jets mistag rates. The use of DeXTer introduces additional scale factors to correct for the differences in efficiency between simulated samples and data. The scale factors for DeXTer are derived as a function of p_T for B - and b -jets. The calibration measurements with data are performed using both $t\bar{t}$ and Z +jets events simultaneously to measure B -jet tagging and b -jet mistagging efficiency in data. Nevertheless, the DeXTer uncertainties are provided with conservative error bands, leaving the calibration to be performed in situ in the final fit of the analysis. Further details on the methodology can be found in Refs. [77,77].

Missing transverse momentum. All the described uncertainties in energy scales or resolutions of the reconstructed objects (hard components) are propagated to the missing transverse momentum. Additional uncertainties in the scale and resolution of the soft term are considered, to account for the disagreement between data and MC for the p_T balance between the hard and soft components [78].

Tracks. Systematic uncertainties related to the track selection efficiency are determined by changing the amount of tracker material and the physical models in the GEANT4 simulation [94,95]. Dedicated uncertainties are considered for the track parameters, including the transverse and longitudinal impact parameters and the track sagitta.

Large- R jet mass scale correction. To correct for the mis-modelling in the large- R jet mass, additional mass scale corrections are estimated. The large- R jet mass scale is varied by $\pm 5\%$ and compared with the nominal results.

9.2 Modelling uncertainties

Renormalisation (μ_r) and factorisation (μ_f) scales. Variations in the renormalisation and factorisation scales are used to estimate the uncertainty due to missing higher order corrections. The uncertainties are combined by taking an envelope of all the variations.

Initial-state radiation and final-state radiation modelling. For the ISR, the amount of radiation is increased (decreased) using the Var3cUp (Var3cDown) variation of the A14 tune [37]. For the FSR, the amount of radiation is increased (decreased) varying the coupling of the QCD emission in the final state by a factor of 0.5 (2).

PDF uncertainties. The PDF uncertainties follow the PDF4LHC recommendations [96]. The α_s uncertainty is derived using the same PDF set evaluated with two different α_s values. The uncertainties from the PDF and α_s are added in quadrature.

Parton shower. The uncertainty associated with hadronisation and parton shower is evaluated by comparing samples with different parton shower models. The nominal $t\bar{t}a$ samples simulated using POWHEG+PYTHIA 8 are compared with samples simulated using MADGRAPH5_AMC@NLO




Fig. 9 Expected and observed 95% CL upper limits of $\sigma(t\bar{t}a) \times \text{BR}(a \rightarrow b\bar{b})$ as a function of the a -boson mass. The lines correspond to the signal cross sections calculated using different coupling strengths of the a boson to the top quark assuming a $\text{BR}(a \rightarrow b\bar{b}) = 100\%$

+HERWIG 7 [97]. The comparison is done after normalising both $t\bar{t}a$ samples. The nominal $t\bar{t}$ (4FS and 5FS) POWHEG+PYTHIA 8 samples are compared with samples simulated using POWHEG+HERWIG 7. The tW and $t\bar{t}H$ MADGRAPH5_AMC@NLO+PYTHIA 8 samples are compared with MADGRAPH5_AMC@NLO+HERWIG 7 samples.

Matrix element uncertainties. For the 5FS and 4FS $t\bar{t}$ samples, the uncertainty associated with the matching between the Matrix element calculations and the parton shower is calculated by comparing the nominal POWHEG+PYTHIA 8 sample with an alternative set of samples simulated also in POWHEG+PYTHIA 8 but using the pThard=1 setting. For tW and $t\bar{t}H$, the matrix element uncertainty is evaluated by comparing the nominal POWHEG+PYTHIA 8 samples to those simulated with MADGRAPH5_AMC@NLO+PYTHIA 8. For $t\bar{t}Z$, the nominal samples are compared with an alternative sample simulated using SHERPA 2.2.0, which accounts both for the matrix element and parton shower uncertainties.

POWHEG damping function. In the $t\bar{t}b\bar{b}$ (4FS) samples, the effect of the choice of a damping scale h_{bzd} that controls the resummation of infrared divergences is evaluated by comparing the nominal sample ($h_{\text{bzd}} = 5$) with an alternative sample in which the scale is set to 2 [30].

Initial-state shower recoil. The uncertainty due to the recoil choice of ISR emissions is evaluated by comparing the nominal sample, in which the whole final state recoils the ISR emission, with an alternative one, in which only one final-state parton recoils against the ISR emission [30].




Fig. 10 Post-fit distributions corresponding to the NN output score of (a) SR 0B4b, (b) SR 0B3b, (c) SR 1B2b and (d) SR 1B1b+1bL and to the sum of the pseudo-continuous b -tagging score of (e) CR 0B2b+1bL for the 30 GeV mass hypothesis fit. The dashed lines in the top and ratio

panels show the post-fit distribution of the signal scaled by a factor of 25 and the post-fit ratio of the background over the total prediction, respectively. The band displays the total post-fit uncertainty

Table 6 Post-fit background and signal yields in the four signal regions and in the control region for the $m_a = 30$ GeV hypothesis. The uncertainties in each yield are the total uncertainties of each component after the fit

Sample	SR 0B4b	SR 0B3b	SR 1B2b	SR 1B1b+1bL	CR 0B2b+1bL
Signal 30 GeV	28 ± 14	180 ± 98	71 ± 31	35 ± 16	110 ± 63
$t\bar{t}$ +light	5 ± 2	1400 ± 320	130 ± 24	1100 ± 180	$17,000 \pm 3000$
$t\bar{t}+\geq 1c$	58 ± 21	4700 ± 1200	380 ± 140	740 ± 230	$12,000 \pm 3500$
$t\bar{t}+\geq 1b$	1093 ± 47	9820 ± 700	1758 ± 97	815 ± 70	5510 ± 650
tW	22 ± 12	360 ± 140	46 ± 20	64 ± 17	830 ± 220
$t\bar{t}H$	62 ± 9	222 ± 22	31 ± 4	14 ± 12	136 ± 13
$t\bar{t}Z$	27 ± 6	120 ± 22	15 ± 3	11 ± 2	128 ± 25
Other	14 ± 2	394 ± 35	47 ± 4	78 ± 10	1060 ± 120
Total pred.	1300 ± 35	$17,000 \pm 130$	2500 ± 50	2900 ± 53	$36,000 \pm 190$
Data	1301	17,242	2479	2866	36,350

Table 7 Table of the impact of each group of uncertainties in the fitted cross section for the hypothesis masses of 12, 30 and 80 GeV. The values shown are the average of up and down uncertainties. The fitted

cross section values include the $\text{BR}(t\bar{t} \rightarrow WbWb) \times \text{BR}(W \rightarrow l\nu) \times \text{BR}(W \rightarrow l\nu)$ in addition to the $\text{BR}(a \rightarrow b\bar{b})$

Fitted cross section [fb]	$m_a = 12$ GeV $\hat{\sigma} = 9$	$m_a = 30$ GeV $\hat{\sigma} = 46$	$m_a = 80$ GeV $\hat{\sigma} = -6.1$
Uncertainty source	$\Delta\hat{\sigma}$	$\Delta\hat{\sigma}$	$\Delta\hat{\sigma}$
Data statistics	6.1	11.0	6.0
MC statistics	2.4	4.2	1.8
Luminosity & pile-up	0.1	0.4	0.1
Jet reconstruction	0.5	4.9	1.2
Lepton reconstruction	<0.1	<0.1	<0.1
E_T^{miss} reconstruction	<0.1	0.3	<0.1
Track reconstruction	4.1	1.5	0.1
DL1r	0.4	3.5	1.4
DeXTer	4.2	18	1.1
Modelling signal	1.7	7.5	1.3
Modelling $t\bar{t} + b$	2.7	13	5.5
Modelling $t\bar{t} + c$	0.9	1.8	1.4
Modelling $t\bar{t}$ +light	0.8	2.0	2.2
Modelling tW	0.3	0.7	0.6
Modelling $t\bar{t}H$	0.1	0.3	0.2
Modelling $t\bar{t}Z$	0.1	0.2	1.0
Norm factors	0.7	6.7	4.7
Reweighting	<0.1	<0.1	<0.1
Total systematic uncertainty	8.0	22	7.8
Total uncertainty	10	24	9.7

Interference between $t\bar{t}$ and tW . To account for uncertainties in the interference between $t\bar{t}$ +jets and tW , the nominal tW sample simulated using diagram removal (DR) is compared with another sample simulated using diagram subtraction (DS).

Reweighting uncertainties. To account for the systematic uncertainties associated with the reweighting functions described in Sect. 6, several uncertainties are determined by

the variations of $t\bar{t}$ +light + tW , $t\bar{t}+\geq 1c$ and $t\bar{t}+\geq 1b$ normalisation factors and the variations of the parameters of the H_T^{red} hyperbolic fit. The uncertainties are evaluated after diagonalising the fit correlation matrix and propagating the diagonal variations in a correlated way.

10 Results

The expected and observed upper limits on the inclusive $\sigma(t\bar{t}a) \times \text{BR}(a \rightarrow b\bar{b})$ are shown in Fig. 9 as a function of the a -boson mass, which ranges from 12 to 100 GeV. This result is compared with the predicted cross sections for the signal corresponding to three different values of the coupling of the a -boson to the top quark, defined as a strength modifier to the SM Yukawa coupling. No significant excess is observed: the largest excess corresponds to the 30 GeV mass hypothesis, with a local significance of 2.0 standard deviations. Assuming $\text{BR}(a \rightarrow b\bar{b}) = 100\%$, the mass region between 50 and 80 GeV is excluded for a coupling of the pseudoscalar to the top quark of 0.5, while a coupling of 1.0 is excluded for all masses. Post-fit distributions of the NN output score corresponding to this mass in each of the four signal regions and of the sum of the pseudo-continuous b -tagging score of all jets in the control region are shown in Fig. 10. Table 6 shows the post-fit event yields per signal and background component in each of the signal and control regions for the same mass hypothesis.

Table 7 summarises the impact of the different sources of uncertainties in the fitted signal strength for three different mass hypotheses: 12, 30 and 80 GeV, which are representative of the low, medium and high mass ranges, respectively. Fits to low-mass hypotheses are limited by data statistics, track reconstruction and DeXTer-related uncertainties. Fits to medium-mass hypotheses are dominated by DeXTer-related uncertainties, followed by the modelling of the $t\bar{t}+\geq 1b$ process and data statistics. Finally, fits to high-mass hypotheses are limited mainly by data statistics, the modelling of the $t\bar{t}+\geq 1b$ process and the normalisation of $t\bar{t}+\text{HF}$. In all cases, the uncertainties in the modelling of the signal are subdominant compared with that of $t\bar{t}+\geq 1b$. No large pulls are observed in any of the fits. Including the pre-fit reweighting corrections detailed in Table 3, the final normalisation factors extracted in the fit corresponding to the 30 GeV mass hypothesis are 1.0 ± 0.3 for $t\bar{t}+\text{light}$ and tW , 1.5 ± 0.5 for $t\bar{t}+\geq 1c$ and 1.2 ± 0.2 for $t\bar{t}+\geq 1b$. These results are compatible with the latest ATLAS $t\bar{t}H$ Run 2 analysis [98].

11 Conclusions

A search for a pseudoscalar a produced in association with either a pair of top quarks or a single top and a W boson in the dilepton decay channel is performed using the full Run 2 pp data sample collected by the ATLAS detector at the LHC. The search targets the dominant decay channel of the pseudoscalar mass probed in this analysis: $a \rightarrow b\bar{b}$. The search covers the pseudoscalar boson mass between 12 and 100 GeV, involving both the kinematic regime where the decay products of the pseudoscalar merge into large B -jets

and the regime where the b -tagged jets are resolved. Limits on the signal production cross section times the branching ratio of the decay into a pair of bottom quarks are extracted. Assuming $\text{BR}(a \rightarrow b\bar{b}) = 100\%$, the mass region between 50 and 80 GeV is excluded for a coupling of the pseudoscalar to the top quark of 0.5, while a coupling of 1.0 is excluded for all masses. These model independent results are the first limits of their kind and complement previous searches by ATLAS [17] and CMS [16] exploring leptonic decays of the pseudoscalar.

Acknowledgements We thank CERN for the very successful operation of the LHC and its injectors, as well as the support staff at CERN and at our institutions worldwide without whom ATLAS could not be operated efficiently. The crucial computing support from all WLCG partners is acknowledged gratefully, in particular from CERN, the ATLAS Tier-1 facilities at TRIUMF/SFU (Canada), NDGF (Denmark, Norway, Sweden), CC-IN2P3 (France), KIT/GridKA (Germany), INFN-CNAF (Italy), NL-T1 (Netherlands), PIC (Spain), RAL (UK) and BNL (USA), the Tier-2 facilities worldwide and large non-WLCG resource providers. Major contributors of computing resources are listed in Ref. [99]. We gratefully acknowledge the support of ANPCyT, Argentina; YerPhI, Armenia; ARC, Australia; BMWFW and FWF, Austria; ANAS, Azerbaijan; CNPq and FAPESP, Brazil; NSERC, NRC and CFI, Canada; CERN; ANID, Chile; CAS, MOST and NSFC, China; Minciencias, Colombia; MEYS CR, Czech Republic; DNR and DNSRC, Denmark; IN2P3-CNRS and CEA-DRF/IRFU, France; SRNSFG, Georgia; BMBF, HGF and MPG, Germany; GSI, Greece; RGC and Hong Kong SAR, China; ICHEP and Academy of Sciences and Humanities, Israel; INFN, Italy; MEXT and JSPS, Japan; CNRST, Morocco; NWO, Netherlands; RCN, Norway; MNiSW, Poland; FCT, Portugal; MNE/IFA, Romania; MSTDI, Serbia; MSSR, Slovakia; ARIS and MVZL, Slovenia; DSI/NRF, South Africa; MICIU/AEI, Spain; SRC and Wallenberg Foundation, Sweden; SERI, SNSF and Cantons of Bern and Geneva, Switzerland; NSTC, Taipei; TENMAK, Türkiye; STFC/UKRI, United Kingdom; DOE and NSF, United States of America. Individual groups and members have received support from BCKDF, CANARIE, CRC and DRAC, Canada; CERN-CZ, FORTE and PRIMUS, Czech Republic; COST, ERC, ERDF, Horizon 2020, ICSC-NextGenerationEU and Marie Skłodowska-Curie Actions, European Union; Investissements d'Avenir Labex, Investissements d'Avenir Idex and ANR, France; DFG and AvH Foundation, Germany; Herakleitos, Thales and Aristeia programmes co-financed by EU-ESF and the Greek NSRF, Greece; BSF-NSF and MINERVA, Israel; NCN and NAWA, Poland; La Caixa Banking Foundation, CERCA Programme Generalitat de Catalunya and PROMETEO and GenT Programmes Generalitat Valenciana, Spain; Göran Gustafssons Stiftelse, Sweden; The Royal Society and Leverhulme Trust, United Kingdom. In addition, individual members wish to acknowledge support from Armenia: Yerevan Physics Institute (FAPERJ); CERN: European Organization for Nuclear Research (CERN DOCT); Chile: Agencia Nacional de Investigación y Desarrollo (FONDECYT 1230812, FONDECYT 1230987, FONDECYT 1240864); China: Chinese Ministry of Science and Technology (MOST-2023YFA1605700, MOST-2023YFA1609300), National Natural Science Foundation of China (NSFC - 12175119, NSFC 12275265); Czech Republic: Czech Science Foundation (GACR - 24-11373S), Ministry of Education Youth and Sports (ERC-CZ-LL2327, FORTE CZ.02.01.01/00/22_008/0004632), PRIMUS Research Programme (PRIMUS/21/SCI/017); EU: H2020 European Research Council (ERC - 101002463); European Union: European Research Council (ERC - 948254, ERC 101089007, ERC, BARD, 101116429), European Regional Development Fund (SMASH COFUND 101081355, SLO ERDF), Horizon 2020 Framework Programme (MUCCA -

CHIST-ERA-19-XAI-00), European Union, Future Artificial Intelligence Research (FAIR-NextGenerationEU PE00000013), Horizon 2020 (EuroHPC - EHPC-DEV-2024D11-051), Italian Center for High Performance Computing, Big Data and Quantum Computing (ICSC, NextGenerationEU); France: Agence Nationale de la Recherche (ANR-21-CE31-0022, ANR-22-EDIR-0002); Germany: Baden-Württemberg Stiftung (BW Stiftung-Postdoc Eliteprogramme), Deutsche Forschungsgemeinschaft (DFG - 469666862, DFG - CR 312/5-2); China: Research Grants Council (GRF); Italy: Istituto Nazionale di Fisica Nucleare (ICSC, NextGenerationEU), Ministero dell'Università e della Ricerca (NextGenEU I53D23000820006 M4C2.1.1); Japan: Japan Society for the Promotion of Science (JSPS KAKENHI JP22H01227, JSPS KAKENHI JP22H04944, JSPS KAKENHI JP22KK0227, JSPS KAKENHI JP23KK0245, JSPS KAKENHI JP24K23939); Norway: Research Council of Norway (RCN-314472); Poland: Ministry of Science and Higher Education (IDUB AGH, POB8, D4 no 9722), Polish National Science Centre (NCN 2021/42/E/ST2/00350, NCN OPUS 2023/51/B/ST2/02507, NCN OPUS nr 2022/47/B/ST2/03059, NCN UMO-2019/34/E/ST2/00393, UMO-2022/47/O/ST2/00148, UMO-2023/49/B/ST2/04085, UMO-2023/51/B/ST2/00920, UMO-2024/53/N/ST2/00869); Portugal: Foundation for Science and Technology (FCT); Spain: Generalitat Valenciana (Artemisa, FEDER, IDIFEDER/2018/048), Ministry of Science and Innovation (MCIN & NextGenEU PCI2022-135018-2, MICIN & FEDER PID2021-125273NB, RYC2019-028510-I, RYC2020-030254-I, RYC2021-031273-I, RYC2022-038164-I); Sweden: Carl Trygger Foundation (Carl Trygger Foundation CTS 22:2312), Swedish Research Council (Swedish Research Council 2023-04654, VR 2021-03651, VR 2022-03845, VR 2022-04683, VR 2023-03403, VR 2024-05451), Knut and Alice Wallenberg Foundation (KAW 2018.0458, KAW 2022.0358, KAW 2023.0366); Switzerland: Swiss National Science Foundation (SNSF- PCEFP2_194658); United Kingdom: Leverhulme Trust (Leverhulme Trust RPG-2020-004), Royal Society (NIF-R1-231091); United States of America: U.S. Department of Energy (ECA DE-AC02-76SF00515), Neubauer Family Foundation.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Funded by SCOAP³.

References

- ATLAS Collaboration, Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC. *Phys. Lett. B* **716**, 1 (2012). <https://doi.org/10.1016/j.physletb.2012.08.020>. arXiv:1207.7214 [hep-ex]
- C.M.S. Collaboration, Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC. *Phys. Lett. B* **716**, 30 (2012). <https://doi.org/10.1016/j.physletb.2012.08.021>. arXiv:1207.7235 [hep-ex]
- L. Evans, P. Bryant, L.H.C. Machine, *JINST* **3**, S08001 (2008). <https://doi.org/10.1088/1748-0221/3/08/S08001>
- G.C. Branco et al., Theory and phenomenology of two-Higgs-doublet models. *Phys. Rep.* **516**, 1 (2012). <https://doi.org/10.1016/j.physrep.2012.02.002>. arXiv:1106.0034 [hep-ph]
- M. Aoki, S. Kanemura, K. Tsumura, K. Yagyu, Models of Yukawa interaction in the two Higgs doublet model, and their collider phenomenology. *Phys. Rev. D* **80**, 015017 (2009). <https://doi.org/10.1103/PhysRevD.80.015017>. arXiv:0902.4665 [hep-ph]
- Y.A. Golfand, E.P. Likhtman, Extension of the algebra of Poincare group generators and violation of p invariance. *JETP Lett.* **13**, 323 (1971). https://doi.org/10.1142/9789814542340_0001
- D.V. Volkov, V.P. Akulov, Is the neutrino a goldstone particle? *Phys. Lett. B* **46**, 109 (1973). [https://doi.org/10.1016/0370-2693\(73\)90490-5](https://doi.org/10.1016/0370-2693(73)90490-5)
- J. Wess, B. Zumino, Supergauge transformations in four dimensions. *Nucl. Phys. B* **70**, 39 (1974). [https://doi.org/10.1016/0550-3213\(74\)90355-1](https://doi.org/10.1016/0550-3213(74)90355-1)
- J. Wess, B. Zumino, Supergauge invariant extension of quantum electrodynamics. *Nucl. Phys. B* **78**, 1 (1974). [https://doi.org/10.1016/0550-3213\(74\)90112-6](https://doi.org/10.1016/0550-3213(74)90112-6)
- S. Ferrara, B. Zumino, Supergauge invariant Yang-Mills theories. *Nucl. Phys. B* **79**, 413 (1974). [https://doi.org/10.1016/0550-3213\(74\)90559-8](https://doi.org/10.1016/0550-3213(74)90559-8)
- A. Salam, J. Strathdee, Super-symmetry and non-Abelian gauges. *Phys. Lett. B* **51**, 353 (1974). [https://doi.org/10.1016/0370-2693\(74\)90226-3](https://doi.org/10.1016/0370-2693(74)90226-3)
- M.R. Buckley, D. Feld, D. Gonçalves, Scalar simplified models for dark matter. *Phys. Rev. D* **91** (2015). <https://doi.org/10.1103/physrevd.91.015017>. arXiv:1410.6497 [hep-ph]
- M. Bauer, U. Haisch, F. Kahlhoefer, Simplified dark matter models with two Higgs doublets: I. Pseudoscalar mediators. *JHEP* **05**, 138 (2017). [https://doi.org/10.1007/jhep05\(2017\)138](https://doi.org/10.1007/jhep05(2017)138). arXiv:1701.07427 [hep-ph]
- J.E. Kim, Light pseudoscalars, particle physics and cosmology. *Phys. Rep.* **150**, 1 (1987). [https://doi.org/10.1016/0370-1573\(87\)90017-2](https://doi.org/10.1016/0370-1573(87)90017-2)
- M. Casolino, T. Farooque, A. Juste, T. Liu, M. Spannowsky, Probing a light CP-odd scalar in di-top-associated production at the LHC. *Eur. Phys. J. C* **75**, 498 (2015). <https://doi.org/10.1140/epjc/s10052-015-3708-y>. arXiv:1507.07004 [hep-ph]
- CMS Collaboration, Search for a scalar or pseudoscalar dilepton resonance produced in association with a massive vector boson or top quark-antiquark pair in multilepton events at $\sqrt{s} = 13$ TeV. *Phys. Rev. D* **110**, 012013 (2024). <https://doi.org/10.1103/PhysRevD.110.012013>. arXiv:2402.11098 [hep-ex]
- ATLAS Collaboration, Search for a new pseudoscalar decaying into a pair of muons in events with a top-quark pair at $\sqrt{s} = 13$ TeV with the ATLAS detector. *Phys. Rev. D* **108**, 092007 (2023). <https://doi.org/10.1103/PhysRevD.108.092007>. arXiv:2304.14247 [hep-ex]
- ATLAS Collaboration, The ATLAS Experiment at the CERN Large Hadron Collider. *JINST* **3**, S08003 (2008). <https://doi.org/10.1088/1748-0221/3/08/S08003>
- ATLAS Collaboration, ATLAS Insertable B-Layer: Technical Design Report, ATLAS-TDR-19; CERN-LHCC-2010-013 (2010). <https://cds.cern.ch/record/1291633>. Addendum: ATLAS-TDR-19-ADD-1; CERN-LHCC-2012-009 (2012). <https://cds.cern.ch/record/1451888>
- B. Abbott et al., Production and integration of the ATLAS Insertable B-Layer. *JINST* **13**, T05008 (2018). <https://doi.org/10.1088/1748-0221/13/05/T05008>. arXiv:1803.00844 [physics.ins-det]
- G. Avoni et al., The new LUCID-2 detector for luminosity measurement and monitoring in ATLAS. *JINST* **13**, P07017 (2018). <https://doi.org/10.1088/1748-0221/13/07/P07017>

22. ATLAS Collaboration, Performance of the ATLAS trigger system in 2015. *Eur. Phys. J. C* **77**, 317 (2017). <https://doi.org/10.1140/epjc/s10052-017-4852-3>. [arXiv:1611.09661](https://arxiv.org/abs/1611.09661) [hep-ex]
23. ATLAS Collaboration, Software and computing for Run 3 of the ATLAS experiment at the LHC. *Eur. Phys. J. C* **85**, 234 (2025). <https://doi.org/10.1140/epjc/s10052-024-13701-w>. [arXiv:2404.06335](https://arxiv.org/abs/2404.06335) [hep-ex]
24. ATLAS Collaboration, Luminosity determination in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector at the LHC. *Eur. Phys. J. C* **83**, 982 (2023). <https://doi.org/10.1140/epjc/s10052-023-11747-w>. [arXiv:2212.09379](https://arxiv.org/abs/2212.09379) [hep-ex]
25. T. Sjöstrand, S. Mrenna, P. Skands, A brief introduction to PYTHIA 8.1. *Comput. Phys. Commun.* **178**, 852 (2008). <https://doi.org/10.1016/j.cpc.2008.01.036>. [arXiv:0710.3820](https://arxiv.org/abs/0710.3820) [hep-ph]
26. ATLAS Collaboration, The Pythia 8 A3 tune description of ATLAS minimum bias and inelastic measurements incorporating the Donnachie–Landshoff diffractive model, ATL-PHYS-PUB-2016-017 (2016). <https://cds.cern.ch/record/2206965>
27. S. Agostinelli et al., Geant4—a simulation toolkit. *Nucl. Instrum. Methods A* **506**, 250 (2003). [https://doi.org/10.1016/S0168-9002\(03\)01368-8](https://doi.org/10.1016/S0168-9002(03)01368-8)
28. ATLAS Collaboration, The ATLAS simulation infrastructure. *Eur. Phys. J. C* **70**, 823 (2010). <https://doi.org/10.1140/epjc/s10052-010-1429-9>. [arXiv:1005.4568](https://arxiv.org/abs/1005.4568) [physics.ins-det]
29. J. Alwall et al., The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations. *JHEP* **07**, 079 (2014). [https://doi.org/10.1007/JHEP07\(2014\)079](https://doi.org/10.1007/JHEP07(2014)079). [arXiv:1405.0301](https://arxiv.org/abs/1405.0301) [hep-ph]
30. ATLAS Collaboration, Studies of Monte Carlo predictions for the $t\bar{t}b\bar{b}$ process, ATL-PHYS-PUB-2022-006 (2022). <https://cds.cern.ch/record/2802806>
31. S. Frixione, G. Ridolfi, P. Nason, A positive-weight next-to-leading-order Monte Carlo for heavy flavour hadroproduction. *JHEP* **09**, 126 (2007). <https://doi.org/10.1088/1126-6708/2007/09/126>. [arXiv:0707.3088](https://arxiv.org/abs/0707.3088) [hep-ph]
32. P. Nason, A new method for combining NLO QCD with shower Monte Carlo algorithms. *JHEP* **11**, 040 (2004). <https://doi.org/10.1088/1126-6708/2004/11/040>. [arXiv:hep-ph/0409146](https://arxiv.org/abs/hep-ph/0409146)
33. S. Frixione, P. Nason, C. Oleari, Matching NLO QCD computations with parton shower simulations: the POWHEG method. *JHEP* **11**, 070 (2007). <https://doi.org/10.1088/1126-6708/2007/11/070>. [arXiv:0709.2092](https://arxiv.org/abs/0709.2092) [hep-ph]
34. S. Alioli, P. Nason, C. Oleari, E. Re, A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX. *JHEP* **06**, 043 (2010). [https://doi.org/10.1007/JHEP06\(2010\)043](https://doi.org/10.1007/JHEP06(2010)043). [arXiv:1002.2581](https://arxiv.org/abs/1002.2581) [hep-ph]
35. NNPDF Collaboration, R.D. Ball et al., Parton distributions for the LHC run II. *JHEP* **04**, 040 (2015). [https://doi.org/10.1007/JHEP04\(2015\)040](https://doi.org/10.1007/JHEP04(2015)040). [arXiv:1410.8849](https://arxiv.org/abs/1410.8849) [hep-ph]
36. T. Sjöstrand et al., An introduction to PYTHIA 8.2. *Comput. Phys. Commun.* **191**, 159 (2015). <https://doi.org/10.1016/j.cpc.2015.01.024>. [arXiv:1410.3012](https://arxiv.org/abs/1410.3012) [hep-ph]
37. ATLAS Collaboration, ATLAS Pythia 8 tunes to 7 TeV data, ATL-PHYS-PUB-2014-021 (2014). <https://cds.cern.ch/record/1966419>
38. NNPDF Collaboration, R.D. Ball et al., Parton distributions with LHC data. *Nucl. Phys. B* **867**, 244 (2013). <https://doi.org/10.1016/j.nuclphysb.2012.10.003>. [arXiv:1207.1303](https://arxiv.org/abs/1207.1303) [hep-ph]
39. J. Pumplin et al., New generation of parton distributions with uncertainties from global QCD analysis. *JHEP* **07**, 012 (2002). <https://doi.org/10.1088/1126-6708/2002/07/012>. [arXiv:hep-ph/0201195](https://arxiv.org/abs/hep-ph/0201195)
40. E. Bothmann et al., Event generation with Sherpa 2.2. *SciPost Phys.* **7**, 034 (2019). <https://doi.org/10.21468/SciPostPhys.7.3.034>. [arXiv:1905.09127](https://arxiv.org/abs/1905.09127) [hep-ph]
41. S. Schumann, F. Krauss, A parton shower algorithm based on Catani–Seymour dipole factorisation. *JHEP* **03**, 038 (2008). <https://doi.org/10.1088/1126-6708/2008/03/038>. [arXiv:0709.1027](https://arxiv.org/abs/0709.1027) [hep-ph]
42. S. Höche, F. Krauss, M. Schönherr, F. Siegert, A critical appraisal of NLO+PS matching methods. *JHEP* **09**, 049 (2012). [https://doi.org/10.1007/JHEP09\(2012\)049](https://doi.org/10.1007/JHEP09(2012)049). [arXiv:1111.1220](https://arxiv.org/abs/1111.1220) [hep-ph]
43. S. Höche, F. Krauss, M. Schönherr, F. Siegert, QCD matrix elements + parton showers. The NLO case. *JHEP* **04**, 027 (2013). [https://doi.org/10.1007/JHEP04\(2013\)027](https://doi.org/10.1007/JHEP04(2013)027). [arXiv:1207.5030](https://arxiv.org/abs/1207.5030) [hep-ph]
44. S. Catani, F. Krauss, B.R. Webber, R. Kuhn, QCD matrix elements + parton showers. *JHEP* **11**, 063 (2001). <https://doi.org/10.1088/1126-6708/2001/11/063>. [arXiv:hep-ph/0109231](https://arxiv.org/abs/hep-ph/0109231)
45. S. Höche, F. Krauss, S. Schumann, F. Siegert, QCD matrix elements and truncated showers. *JHEP* **05**, 053 (2009). <https://doi.org/10.1088/1126-6708/2009/05/053>. [arXiv:0903.1219](https://arxiv.org/abs/0903.1219) [hep-ph]
46. D.J. Lange, The EvtGen particle decay simulation package. *Nucl. Instrum. Methods A* **462**, 152 (2001). [https://doi.org/10.1016/S0168-9002\(01\)00089-4](https://doi.org/10.1016/S0168-9002(01)00089-4)
47. M. Beneke, P. Falgari, S. Klein, C. Schwinn, Hadronic top-quark pair production with NNLL threshold resummation. *Nucl. Phys. B* **855**, 695 (2012). <https://doi.org/10.1016/j.nuclphysb.2011.10.021>. [arXiv:1109.1536](https://arxiv.org/abs/1109.1536) [hep-ph]
48. M. Cacciari, M. Czakon, M. Mangano, A. Mitov, P. Nason, Top-pair production at hadron colliders with next-to-next-to-leading logarithmic soft-gluon resummation. *Phys. Lett. B* **710**, 612 (2012). <https://doi.org/10.1016/j.physletb.2012.03.013>. [arXiv:1111.5869](https://arxiv.org/abs/1111.5869) [hep-ph]
49. P. Bärnreuther, M. Czakon, A. Mitov, Percent-level-precision physics at the LHC: next-to-next-to-leading order QCD corrections to $q\bar{q} \rightarrow t\bar{t} + X$. *Phys. Rev. Lett.* **109**, 132001 (2012). <https://doi.org/10.1103/PhysRevLett.109.132001>. [arXiv:1204.5201](https://arxiv.org/abs/1204.5201) [hep-ph]
50. M. Czakon, A. Mitov, NNLO corrections to top-pair production at hadron colliders: the all-fermionic scattering channels. *JHEP* **12**, 054 (2012). [https://doi.org/10.1007/JHEP12\(2012\)054](https://doi.org/10.1007/JHEP12(2012)054). [arXiv:1207.0236](https://arxiv.org/abs/1207.0236) [hep-ph]
51. M. Czakon, A. Mitov, NNLO corrections to top pair production at hadron colliders: the quark-gluon reaction. *JHEP* **01**, 080 (2013). [https://doi.org/10.1007/JHEP01\(2013\)080](https://doi.org/10.1007/JHEP01(2013)080). [arXiv:1210.6832](https://arxiv.org/abs/1210.6832) [hep-ph]
52. M. Czakon, P. Fiedler, A. Mitov, Total top-quark pair-production cross section at hadron colliders through $\mathcal{O}(\alpha_s^3)$. *Phys. Rev. Lett.* **110**, 252004 (2013). <https://doi.org/10.1103/PhysRevLett.110.252004>. [arXiv:1303.6254](https://arxiv.org/abs/1303.6254) [hep-ph]
53. M. Czakon, A. Mitov, Top++: a program for the calculation of the top-pair cross-section at hadron colliders. *Comput. Phys. Commun.* **185**, 2930 (2014). <https://doi.org/10.1016/j.cpc.2014.06.021>. [arXiv:1112.5675](https://arxiv.org/abs/1112.5675) [hep-ph]
54. N. Kidonakis, Two-loop soft anomalous dimensions for single top quark associated production with a W^- or H^- . *Phys. Rev. D* **82**, 054018 (2010). <https://doi.org/10.1103/PhysRevD.82.054018>. [arXiv:1005.4451](https://arxiv.org/abs/1005.4451) [hep-ph]
55. N. Kidonakis, ‘Top Quark Production’, in *Proceedings, Helmholtz International Summer School on Physics of Heavy Quarks and Hadrons (HQ2013) (JINR, Dubna, Russia, 15th–28th July 2013)*, p. 139. <https://doi.org/10.3204/DESY-PROC-2013-03/Kidonakis>. [arXiv:1311.0283](https://arxiv.org/abs/1311.0283) [hep-ph]
56. D. de Florian et al., Handbook of LHC Higgs Cross Sections: 4. Deciphering the Nature of the Higgs Sector (2017). <https://doi.org/10.23731/CYRM-2017-002>. [arXiv:1610.07922](https://arxiv.org/abs/1610.07922) [hep-ph]
57. C. Anastasiou, L. Dixon, K. Melnikov, F. Petriello, High-precision QCD at hadron colliders: electroweak gauge boson rapidity distributions at next-to-next-to leading order. *Phys. Rev. D* **69**, 094008 (2004). <https://doi.org/10.1103/PhysRevD.69.094008>. [arXiv:hep-ph/0312266](https://arxiv.org/abs/hep-ph/0312266)

58. ATLAS Collaboration, Electron and photon performance measurements with the ATLAS detector using the 2015–2017 LHC proton–proton collision data. JINST **14**, P12006 (2019). <https://doi.org/10.1088/1748-0221/14/12/P12006>. arXiv:1908.00005 [hep-ex]
59. ATLAS Collaboration, Evidence for the associated production of the Higgs boson and a top quark pair with the ATLAS detector. Phys. Rev. D **97**, 072003 (2018). <https://doi.org/10.1103/PhysRevD.97.072003>. arXiv:1712.08891 [hep-ex]
60. ATLAS Collaboration, Muon reconstruction and identification efficiency in ATLAS using the full Run 2 pp collision data set at $\sqrt{s} = 13$ TeV. Eur. Phys. J. C **81**, 578 (2021). <https://doi.org/10.1140/epjc/s10052-021-09233-2>. arXiv:2012.00578 [hep-ex]
61. ATLAS Collaboration, Jet reconstruction and performance using particle flow with the ATLAS detector. Eur. Phys. J. C **77**, 466 (2017). <https://doi.org/10.1140/epjc/s10052-017-5031-2>. arXiv:1703.10485 [hep-ex]
62. M. Cacciari, G.P. Salam, G. Soyez, The anti- k_t jet clustering algorithm. JHEP **04**, 063 (2008). <https://doi.org/10.1088/1126-6708/2008/04/063>. arXiv:0802.1189 [hep-ph]
63. M. Cacciari, G.P. Salam, G. Soyez, FastJet user manual. Eur. Phys. J. C **72**, 1896 (2012). <https://doi.org/10.1140/epjc/s10052-012-1896-2>. arXiv:1111.6097 [hep-ph]
64. ATLAS Collaboration, Jet energy scale and resolution measured in proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. Eur. Phys. J. C **81**, 689 (2021). <https://doi.org/10.1140/epjc/s10052-021-09402-3>. arXiv:2007.02645 [hep-ex]
65. ATLAS Collaboration, Performance of pile-up mitigation techniques for jets in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector. Eur. Phys. J. C **76**, 581 (2016). <https://doi.org/10.1140/epjc/s10052-016-4395-z>. arXiv:1510.03823 [hep-ex]
66. ATLAS Collaboration, ATLAS flavour-tagging algorithms for the LHC Run 2 pp collision dataset. Eur. Phys. J. C **83**, 681 (2023). <https://doi.org/10.1140/epjc/s10052-023-11699-1>. arXiv:2211.16345 [physics.data-an]
67. ATLAS Collaboration, ATLAS b -jet identification performance and efficiency measurement with $t\bar{t}$ events in pp collisions at $\sqrt{s} = 13$ TeV. Eur. Phys. J. C **79**, 970 (2019). <https://doi.org/10.1140/epjc/s10052-019-7450-8>. arXiv:1907.05120 [hep-ex]
68. ATLAS Collaboration, Measurement of the c -jet mistagging efficiency in $t\bar{t}$ events using pp collision data at $\sqrt{s} = 13$ TeV collected with the ATLAS detector. Eur. Phys. J. C **82**, 95 (2022). <https://doi.org/10.1140/epjc/s10052-021-09843-w>. arXiv:2109.10627 [hep-ex]
69. ATLAS Collaboration, Calibration of the light-flavour jet mistagging efficiency of the b -tagging algorithms with Z+jets events using 139 fb $^{-1}$ of ATLAS proton–proton collision data at $\sqrt{s} = 13$ TeV. Eur. Phys. J. C **83**, 728 (2023). <https://doi.org/10.1140/epjc/s10052-023-11736-z>. arXiv:2301.06319 [hep-ex]
70. M. Cacciari, G.P. Salam, Pileup subtraction using jet areas. Phys. Lett. B **659**, 119 (2008). <https://doi.org/10.1016/j.physletb.2007.09.077>. arXiv:0707.1378 [hep-ph]
71. M. Cacciari, G.P. Salam, G. Soyez, The catchment area of jets. JHEP **04**, 005 (2008). <https://doi.org/10.1088/1126-6708/2008/04/005>. arXiv:0802.1188 [hep-ph]
72. ATLAS Collaboration, Deep Sets based Neural Networks for Impact Parameter Flavour Tagging in ATLAS, ATL-PHYS-PUB-2020-014 (2020). <https://cds.cern.ch/record/2718948>
73. ATLAS Collaboration, Variable Radius, Exclusive- k_T , and Center-of-Mass Subjet Reconstruction for Higgs($\rightarrow b\bar{b}$) Tagging in ATLAS, ATL-PHYS-PUB-2017-010 (2017). <https://cds.cern.ch/record/2268678>
74. ATLAS Collaboration, Calibration of a soft secondary vertex tagger using proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. Phys. Rev. D **110**, 032015 (2024). <https://doi.org/10.1103/PhysRevD.110.032015>. arXiv:2405.03253 [hep-ex]
75. V. Kostyukhin, Secondary vertex based b -tagging, tech. rep. ATL-PHYS-2003-033, revised version number 1 submitted on 2003-09-22 11:21:01: CERN (2003). <https://cds.cern.ch/record/685550>
76. ATLAS Collaboration, Performance of the reconstruction of large impact parameter tracks in the inner detector of ATLAS. Eur. Phys. J. C **83**, 1081 (2023). <https://doi.org/10.1140/epjc/s10052-023-12024-6>. arXiv:2304.12867 [hep-ex]
77. ATLAS Collaboration, DeXTer: Deep Sets based Neural Networks for Low- p_T $X \rightarrow b\bar{b}$ Identification in ATLAS, ATL-PHYS-PUB-2022-042 (2022). <https://cds.cern.ch/record/2825434>
78. ATLAS Collaboration, The performance of missing transverse momentum reconstruction and its significance with the ATLAS detector using 140 fb $^{-1}$ of $\sqrt{s} = 13$ TeV pp collisions (2024). arXiv:2402.05858 [hep-ex]
79. B. Nachman, P. Nef, A. Schwartzman, M. Swiatkowski, C. Wanotayaroj, Jets from Jets: re-clustering as a tool for large radius jet reconstruction and grooming at the LHC. JHEP **02**, 075 (2015). [https://doi.org/10.1007/JHEP02\(2015\)075](https://doi.org/10.1007/JHEP02(2015)075). arXiv:1407.2922 [hep-ph]
80. ATLAS Collaboration, Performance of electron and photon triggers in ATLAS during LHC Run 2. Eur. Phys. J. C **80**, 47 (2020). <https://doi.org/10.1140/epjc/s10052-019-7500-2>. arXiv:1909.00761 [hep-ex]
81. ATLAS Collaboration, Performance of the ATLAS muon triggers in Run 2. JINST **15**, P09015 (2020). <https://doi.org/10.1088/1748-0221/15/09/p09015>. arXiv:2004.13447 [physics.ins-det]
82. ATLAS Collaboration, ATLAS data quality operations and performance for 2015–2018 data-taking. JINST **15**, P04003 (2020). <https://doi.org/10.1088/1748-0221/15/04/P04003>. arXiv:1911.04632 [physics.ins-det]
83. ATLAS Collaboration, Search for charged Higgs bosons decaying into a top quark and a bottom quark at $\sqrt{s} = 13$ TeV with the ATLAS detector. JHEP **06**, 145 (2021). [https://doi.org/10.1007/JHEP06\(2021\)145](https://doi.org/10.1007/JHEP06(2021)145). arXiv:2102.10076 [hep-ex]
84. ATLAS Collaboration, Measurement of the $t\bar{t}t\bar{t}$ production cross section in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. JHEP **11**, 118 (2021). [https://doi.org/10.1007/JHEP11\(2021\)118](https://doi.org/10.1007/JHEP11(2021)118). arXiv:2106.11683 [hep-ex]
85. ATLAS Collaboration, Search for a new scalar resonance in flavour-changing neutral-current top-quark decays $t \rightarrow qX$ ($q = u, c$), with $X \rightarrow b\bar{b}$, in proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. JHEP **07**, 199 (2023). [https://doi.org/10.1007/JHEP07\(2023\)199](https://doi.org/10.1007/JHEP07(2023)199). arXiv:2301.03902 [hep-ex]
86. CMS Collaboration, Measurement of the $t\bar{t}$ production cross section in the all-jet final state in pp collisions at $\sqrt{s} = 7$ TeV. JHEP **05**, 065 (2013). [https://doi.org/10.1007/JHEP05\(2013\)065](https://doi.org/10.1007/JHEP05(2013)065). arXiv:1302.0508 [hep-ex]
87. CMS Collaboration, Measurement of the cross section for $t\bar{t}$ production with additional jets and b jets in pp collisions at $\sqrt{s} = 13$ TeV. JHEP **07**, 125 (2020). [https://doi.org/10.1007/JHEP07\(2020\)125](https://doi.org/10.1007/JHEP07(2020)125). arXiv:2003.06467 [hep-ex]
88. ATLAS Collaboration, Measurements of inclusive and differential fiducial cross-sections of $t\bar{t}$ production with additional heavy-flavour jets in proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. JHEP **04**, 046 (2019). [https://doi.org/10.1007/JHEP04\(2019\)046](https://doi.org/10.1007/JHEP04(2019)046). arXiv:1811.12113 [hep-ex]
89. A. Hoecker et al., TMVA—Toolkit for Multivariate Data Analysis (2009). arXiv:physics/0703039 [physics.data-an]
90. A. Paszke et al., PyTorch: an imperative style, high-performance deep learning library (2019). <https://doi.org/10.48550/arXiv.1912.01703>. arXiv:1912.01703 [cs.LG]
91. G. Cowan, K. Cranmer, E. Gross, O. Vitells, Asymptotic formulae for likelihood-based tests of new physics. Eur. Phys. J. C **71**, 1554 (2011). <https://doi.org/10.1140/epjc/s10052-011-1554-0>. arXiv:1007.1727 [physics.data-an]. Erratum: Eur. Phys. J. C **73**, 2501 (2013)

98. ATLAS Collaboration, Measurement of the associated production of a top-antitop-quark pair and a Higgs boson decaying into a $b\bar{b}$ pair in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector at the LHC. *Eur. Phys. J. C* **85**, 210 (2025). <https://doi.org/10.1140/epjc/s10052-025-13740-x>. [arXiv:2407.10904](https://arxiv.org/abs/2407.10904) [hep-ex]
99. ATLAS Collaboration, ATLAS Computing Acknowledgements, ATL-SOFT-PUB-2025-001 (2025). <https://cds.cern.ch/record/2922210>

ATLAS Collaboration*

G. Aad¹⁰⁴, E. Aakvaag¹⁷, B. Abbott¹²³, S. Abdelhameed^{119a}, K. Abeling⁵⁵, N.J. Abicht⁴⁹, S. H. Abidi³⁰, M. Aboelela⁴⁵, A. Aboulhorma^{36c}, H. Abramowicz¹⁵⁷, Y. Abulaiti¹²⁰, B. S. Acharya^{69a,69b,n}, A. Ackermann^{63a}, C. Adam Bourdarios⁴, L. Adamczyk^{86a}, S. V. Addepalli¹⁴⁹, M. J. Addison¹⁰³, J. Adelman¹¹⁸, A. Adiguzel^{22c}, T. Adye¹³⁷, A. A. Affolder¹³⁹, Y. Afik⁴⁰, M. N. Agaras¹³, A. Aggarwal¹⁰², C. Agheorghiesei^{28c}, F. Ahmadov^{39,ae}, S. Ahuja⁹⁷, X. Ai^{143b}, G. Aielli^{76a,76b}, A. Aikot¹⁶⁹, M. Ait Tamlihat^{36e}, B. Aitbenkikh^{36a}, M. Akbiyik¹⁰², T. P. A. Åkesson¹⁰⁰, A. V. Akimov¹⁵¹, D. Akiyama¹⁷⁴, N. N. Akolkar²⁵, S. Aktas^{22a}, G. L. Alberghi^{24b}, J. Albert¹⁷¹, P. Albicocco⁵³, G. L. Albouy⁶⁰, S. Alderweireldt⁵², Z. L. Alegria¹²⁴, M. Aleksa³⁷, I. N. Aleksandrov³⁹, C. Alexa^{28b}, T. Alexopoulos¹⁰, F. Alfonsi^{24b}, M. Algren⁵⁶, M. Alhroob¹⁷³, B. Ali¹³⁵, H. M. J. Ali^{93,x}, S. Ali³², S. W. Alibocus⁹⁴, M. Aliev^{34c}, G. Alimonti^{71a}, W. Alkahi⁵⁵, C. Allaire⁶⁶, B. M. M. Allbrooke¹⁵², J. S. Allen¹⁰³, J. F. Allen⁵², P. P. Allport²¹, A. Aloisio^{72a,72b}, F. Alonso⁹², C. Alpigiani¹⁴², Z. M. K. Alsolami⁹³, A. Alvarez Fernandez¹⁰², M. Alves Cardoso⁵⁶, M. G. Alviggi^{72a,72b}, M. Aly¹⁰³, Y. Amaral Coutinho^{83b}, A. Ambler¹⁰⁶, C. Amelung³⁷, M. Ameri¹⁰³, C. G. Ames¹¹¹, T. Amezza¹³⁰, D. Amidei¹⁰⁸, B. Amini⁵⁴, K. Amirie¹⁶¹, A. Amirkhanov³⁹, S. P. Amor Dos Santos^{133a}, K. R. Amos¹⁶⁹, D. Amperiadou¹⁵⁸, S. An⁸⁴, C. Anastopoulos¹⁴⁵, T. Andeen¹¹, J. K. Anders⁹⁴, A. C. Anderson⁵⁹, A. Andreazza^{71a,71b}, S. Angelidakis⁹, A. Angerami⁴², A. V. Anisenkov³⁹, A. Annovi^{74a}, C. Antel⁵⁶, E. Antipov¹⁵¹, M. Antonelli⁵³, F. Anulli^{75a}, M. Aoki⁸⁴, T. Aoki¹⁵⁹, M. A. Aparo¹⁵², L. Aperio Bella⁴⁸, M. Apicella³¹, C. Appelt¹⁵⁷, A. Apyan²⁷, S. J. Arbiol Val⁸⁷, C. Arcangeletti⁵³, A. T. H. Arce⁵¹, J-F. Arguin¹¹⁰, S. Argyropoulos¹⁵⁸, J.-H. Arling⁴⁸, O. Arnaez⁴, H. Arnold¹⁵¹, G. Artoni^{75a,75b}, H. Asada¹¹³, K. Asai¹²¹, S. Asai¹⁵⁹, S. Asatryan¹⁷⁹, N. A. Asbah³⁷, R. A. Ashby Pickering¹⁷³, A. M. Aslam⁹⁷, K. Assamagan³⁰, R. Astalos^{29a}, K. S. V. Astrand¹⁰⁰, S. Atashi¹⁶⁵, R. J. Atkin^{34a}, H. Atmani^{36f}, P. A. Atmasiddha¹³¹, K. Augsten¹³⁵, A. D. Aurio⁴¹, V. A. Austrup¹⁰³, G. Avolio³⁷, K. Axiotis⁵⁶, G. Azuelos^{110,ai}, D. Babal^{29b}, H. Bachacou¹³⁸, K. Bachas^{158,r}, A. Bachi³⁵, E. Bachmann⁵⁰, M. J. Backes^{63a}, A. Badea⁴⁰, T. M. Baer¹⁰⁸, P. Bagnaia^{75a,75b}, M. Bahmani¹⁹, D. Bahner⁵⁴, K. Bai¹²⁶, J. T. Baines¹³⁷, L. Baines⁹⁶, O. K. Baker¹⁷⁸, E. Bakos¹⁶, D. Bakshi Gupta⁸, L. E. Balabram Filho^{83b}, V. Balakrishnan¹²³, R. Balasubramanian⁴, E. M. Baldin³⁸, P. Balek^{86a}, E. Ballabene^{24a,24b}, F. Balli¹³⁸, L. M. Baltes^{63a}, W. K. Balunas³³, J. Balz¹⁰², I. Bamwidhi^{119b}, E. Banas⁸⁷, M. Bandieramonte¹³², A. Bandyopadhyay²⁵, S. Bansal²⁵, L. Barak¹⁵⁷, M. Barakat⁴⁸, E. L. Barberio¹⁰⁷, D. Barberis^{18b}, M. Barbero¹⁰⁴, M. Z. Barel¹¹⁷, T. Barillari¹¹², M-S. Barisits³⁷, T. Barklow¹⁴⁹, P. Baron¹³⁶, D. A. Baron Moreno¹⁰³, A. Baroncelli⁶², A. J. Barr¹²⁹, J. D. Barr⁹⁸, F. Barreiro¹⁰¹, J. Barreiro Guimarães da Costa¹⁴, M. G. Barros Teixeira^{133a}, S. Barsov³⁸, F. Bartels^{63a}, R. Bartoldus¹⁴⁹, A. E. Barton⁹³, P. Bartos^{29a}, A. Basan¹⁰², M. Baselga⁴⁹, S. Bashiri⁸⁷, A. Bassalat^{66,b}, M. J. Basso^{162a}, S. Bataju⁴⁵, R. Bate¹⁷⁰, R. L. Bates⁵⁹, S. Batlamous¹⁰¹, M. Battaglia¹³⁹, D. Battulga¹⁹, M. Bause^{75a,75b}, M. Bauer⁷⁹, P. Bauer²⁵, L. T. Bayer⁴⁸, L. T. Bazzano Hurrell³¹, J. B. Beacham¹¹², T. Beau¹³⁰, J. Y. Beauchamp⁹², P. H. Beauchemin¹⁶⁴, P. Bechtle²⁵, H. P. Beck^{20,q}, K. Becker¹⁷³, A. J. Beddall⁸², V. A. Bednyakov³⁹, C. P. Bee¹⁵¹, L. J. Beemster¹⁶, M. Begalli^{83d}, M. Begel³⁰, J. K. Behr⁴⁸, J. F. Beirer³⁷, F. Beisiegel²⁵, M. Belfkir^{119b}, G. Bella¹⁵⁷, L. Bellagamba^{24b}, A. Bellerive³⁵, C. D. Bellgraph⁶⁸, P. Bellos²¹, K. Beloborodov³⁸, D. Benchechroun^{36a}, F. Bendebba^{36a}, Y. Benhammou¹⁵⁷, K. C. Benkendorfer⁶¹, L. Beresford⁴⁸, M. Beretta⁵³, E. Bergeas Kuutmann¹⁶⁷, N. Berger⁴, B. Bergmann¹³⁵, J. Beringer^{18a}, G. Bernardi⁵, C. Bernius¹⁴⁹, F. U. Bernlochner²⁵, F. Bernon³⁷, A. Berrocal Guardia¹³, T. Berry⁹⁷, P. Berta¹³⁶, A. Berthold⁵⁰, A. Berti^{133a}, R. Bertrand¹⁰⁴, S. Bethke¹¹², A. Betti^{75a,75b}, A. J. Bevan⁹⁶, L. Bezio⁵⁶, N. K. Bhalla⁵⁴, S. Bharthuar¹¹², S. Bhatta¹⁵¹, P. Bhattacharai¹⁴⁹, Z. M. Bhatti¹²⁰, K. D. Bhide⁵⁴, V. S. Bhopatkar¹²⁴, R. M. Bianchi¹³², G. Bianco^{24b,24a}, O. Biebel¹¹¹, M. Biglietti^{77a}, C. S. Billingsley⁴⁵, Y. Bimgdi^{36f}, M. Bindi⁵⁵, A. Bingham¹⁷⁷, A. Bingul^{22b}, C. Bini^{75a,75b}, G. A. Bird³³, M. Birman¹⁷⁵, M. Biros¹³⁶, S. Biryukov¹⁵², T. Bisanz⁴⁹, E. Bisceglie^{24b,24a}, J. P. Biswal¹³⁷, D. Biswas¹⁴⁷,

I. Bloch⁴⁸, A. Blue⁵⁹, U. Blumenschein⁹⁶, J. Blumenthal¹⁰², V. S. Bobrovnikov³⁹, M. Boehler⁵⁴, B. Boehm¹⁷², D. Bogavac¹³, A. G. Bogdanchikov³⁸, L. S. Boggia¹³⁰, V. Boisvert⁹⁷, P. Bokan³⁷, T. Bold^{86a}, M. Bomben⁵, M. Bona⁹⁶, M. Boonekamp¹³⁸, A. G. Borbély⁵⁹, I. S. Bordulev³⁸, G. Borissov⁹³, D. Bortoletto¹²⁹, D. Boscherini^{24b}, M. Bosman¹³, K. Bouaouda^{36a}, N. Bouchhar¹⁶⁹, L. Boudet⁴, J. Boudreau¹³², E. V. Bouhova-Thacker⁹³, D. Boumediene⁴¹, R. Bouquet^{57b,57a}, A. Boveia¹²², J. Boyd³⁷, D. Boye³⁰, I. R. Boyko³⁹, L. Bozianu⁵⁶, J. Bracinik²¹, N. Brahimi⁴, G. Brandt¹⁷⁷, O. Brandt³³, B. Brau¹⁰⁵, J. E. Brau¹²⁶, R. Brenner¹⁷⁵, L. Brenner¹¹⁷, R. Brenner¹⁶⁷, S. Bressler¹⁷⁵, G. Brianti^{78a,78b}, D. Britton⁵⁹, D. Britzger¹¹², I. Brock²⁵, R. Brock¹⁰⁹, G. Brooijmans⁴², A. J. Brooks⁶⁸, E. M.
Brooks^{162b}, E. Brost³⁰, L. M. Brown^{171,162a}, L. E. Bruce⁶¹, T. L. Bruckler¹²⁹, P. A. Bruckman de Renstrom⁸⁷, B. Brüers⁴⁸, A. Bruni^{24b}, G. Bruni^{24b}, D. Brunner^{47a,47b}, M. Bruschi^{24b}, N. Bruscino^{75a,75b}, T. Buanes¹⁷, Q. Buat¹⁴², D. Buchin¹¹², A. G. Buckley⁵⁹, O. Bulekov⁸², B. A. Bullard¹⁴⁹, S. Burdin⁹⁴, C. D. Burgard⁴⁹, A. M. Burger⁹¹, B. Burghgrave⁸, O. Burlayenko⁵⁴, J. Burleson¹⁶⁸, J. C. Burzynski¹⁴⁸, E. L. Busch⁴², V. Büscher¹⁰², P. J. Bussey⁵⁹, J. M. Butler²⁶, C. M. Buttar⁵⁹, J. M. Butterworth⁹⁸, W. Buttinger¹³⁷, C. J. Buxo Vazquez¹⁰⁹, A. R. Buzykaev³⁹, S. Cabrera Urbán¹⁶⁹, L. Cadamuro⁶⁶, D. Caforio⁵⁸, H. Cai¹³², Y. Cai^{24b,114c,24a}, Y. Cai^{114a}, V. M. M. Cairo³⁷, O. Cakir^{3a}, N. Calace³⁷, P. Calafiura^{18a}, G. Calderini¹³⁰, P. Calfayan³⁵, G. Callea⁵⁹, L. P. Caloba^{83b}, D. Calvet⁴¹, S. Calvet⁴¹, R. Camacho Toro¹³⁰, S.
Camarda³⁷, D. Camarero Munoz²⁷, P. Camarri^{76a,76b}, C. Camincher¹⁷¹, M. Campanelli⁹⁸, A. Camplani⁴³, V. Canale^{72a,72b}, A. C. Canbay^{3a}, E. Canonero⁹⁷, J. Cantero¹⁶⁹, Y. Cao¹⁶⁸, F. Capocasa²⁷, M. Capua^{44b,44a}, A. Carbone^{71a,71b}, R. Cardarelli^{76a}, J. C. J. Cardenas⁸, M. P. Cardiff²⁷, G. Carducci^{44b,44a}, T. Carli³⁷, G. Carlino^{72a}, J. I. Carlotto¹³, B. T. Carlson^{132,s}, E. M. Carlson¹⁷¹, J. Carmignani⁹⁴, L. Carminati^{71a,71b}, A. Carnelli⁴, M. Carnesale³⁷, S. Caron¹¹⁶, E. Carquin^{140f}, I. B. Carr¹⁰⁷, S. Carrá^{73a,73b}, G. Carratta^{24b,24a}, A. M. Carroll¹²⁶, M. P. Casado^{13,i}, M. Caspar⁴⁸, F. L. Castillo⁴, L. Castillo Garcia¹³, V. Castillo Gimenez¹⁶⁹, N. F. Castro^{133a,133e}, A. Catinaccio³⁷, J. R. Catmore¹²⁸, T. Cavaliere⁴, V. Cavaliere³⁰, L. J. Caviedes Betancourt^{23b}, E. Celebi⁸², S. Cella³⁷, V. Cepaitis⁵⁶, K. Cerny¹²⁵, A. S. Cerqueira^{83a}, A. Cerri^{74a,74b,al}, L.
Cerrito^{76a,76b}, F. Cerutti^{18a}, B. Cervato^{71a,71b}, A. Cervelli^{24b}, G. Cesarini⁵³, S. A. Cetin⁸², P. M. Chabrilat¹³⁰, S. Chakraborty¹⁷³, J. Chan^{18a}, W. Y. Chan¹⁵⁹, J. D. Chapman³³, E. Chapon¹³⁸, B. Chargeishvili^{155b}, D. G. Charlton²¹, C. Chauhan¹³⁶, Y. Che^{114a}, S. Chekanov⁶, S. V. Chekulaev^{162a}, G. A. Chelkov^{39,a}, B. Chen¹⁵⁷, B. Chen¹⁷¹, H. Chen^{114a}, H. Chen³⁰, J. Chen^{144a}, J. Chen¹⁴⁸, M. Chen¹²⁹, S. Chen⁸⁹, S. J. Chen^{114a}, X. Chen^{144a}, X. Chen^{15,ah}, Z. Chen⁶², C. L. Cheng¹⁷⁶, H. C. Cheng^{64a}, S. Cheong¹⁴⁹, A. Cheplakov³⁹, E. Cherepanova¹¹⁷, R. Cherkaoui El Moursli^{36e}, E. Cheu⁷, K. Cheung⁶⁵, L. Chevalier¹³⁸, V. Chiarella⁵³, G. Chiarelli^{74a}, G. Chiodini^{70a}, A. S. Chisholm²¹, A. Chitan^{28b}, M. Chitishvili¹⁶⁹, M. V. Chizhov^{39,t}, K. Choi¹¹, Y. Chou¹⁴², E. Y. S. Chow¹¹⁶, K. L. Chu¹⁷⁵, M. C.
Chu^{64a}, X. Chu^{14,114c}, Z. Chubinidze⁵³, J. Chudoba¹³⁴, J. J. Chwastowski⁸⁷, D. Cieri¹¹², K. M. Ciesla^{86a}, V. Cindro⁹⁵, A. Ciochio^{18a}, F. Ciotto^{72a,72b}, Z. H. Citron¹⁷⁵, M. Citterio^{71a}, D. A. Ciubotaru^{28b}, A. Clark⁵⁶, P. J. Clark⁵², N. Clarke Hall⁹⁸, C. Clarry¹⁶¹, S. E. Clawson⁴⁸, C. Clement^{47a,47b}, Y. Coadou¹⁰⁴, M. Cobal^{69a,69c}, A. Coccaro^{57b}, R. F. Coelho Barrue^{133a}, R. Coelho Lopes De Sa¹⁰⁵, S. Coelli^{71a}, L. S. Colangeli¹⁶¹, B. Cole⁴², P. Collado Soto¹⁰¹, J. Collot⁶⁰, R. Coluccia^{70a,70b}, P. Conde Muño^{133a,133g}, M. P. Connell^{34c}, S. H. Connell^{34c}, E. I. Conroy¹²⁹, F. Conventi^{72a,aj}, H. G. Cooke²¹, A. M. Cooper-Sarkar¹²⁹, L. Corazzina^{75a,75b}, F. A. Corchia^{24b,24a}, A. Cordeiro Oudot Choi¹⁴², L. D. Corpe⁴¹, M. Corradi^{75a,75b}, F. Corriveau^{106,ac}, A. Cortes-Gonzalez¹⁵⁹, M. J. Costa¹⁶⁹, F. Costanza⁴, D. Costanzo¹⁴⁵, B. M. Cote¹²², J. Couthures⁴, G. Cowan⁹⁷, K.
Cranmer¹⁷⁶, L. Cremer⁴⁹, D. Cremonini^{24b,24a}, S. Crépe-Renaudin⁶⁰, F. Crescioli¹³⁰, T. Cresta^{73a,73b}, M. Cristinziani¹⁴⁷, M. Cristoforetti^{78a,78b}, V. Croft¹¹⁷, J. E. Crosby¹²⁴, G. Crosetti^{44b,44a}, A. Cueto¹⁰¹, H. Cui⁹⁸, Z. Cui⁷, W. R. Cunningham⁵⁹, F. Curcio¹⁶⁹, J. R. Curran⁵², M. J. Da Cunha Sargedas De Sousa^{57b,57a}, J. V. Da Fonseca Pinto^{83b}, C. Da Via¹⁰³, W. Dabrowski^{86a}, T. Dado³⁷, S. Dahbi¹⁵⁴, T. Dai¹⁰⁸, D. Dal Santo²⁰, C. Dallapiccola¹⁰⁵, M. Dam⁴³, G. D'amen³⁰, V. D'Amico¹¹¹, J. Damp¹⁰², J. R. Dandoy³⁵, D. Dannheim³⁷, G. D'anniballe^{74a,74b}, M. Danninger¹⁴⁸, V. Dao¹⁵¹, G. Darbo^{57b}, S. J. Das³⁰, F. Dattola⁴⁸

S. Demers¹⁷⁸, M. Demichev³⁹, S. P. Denisov³⁸, H. Denizli^{22a,m}, L. D'Eramo⁴¹, D. Derendarz⁸⁷, F. Derue¹³⁰, P. Dervan⁹⁴, K. Desch²⁵, F. A. Di Bello^{57b,57a}, A. Di Ciccio^{76a,76b}, L. Di Ciccio⁴, A. Di Domenico^{75a,75b}, C. Di Donato^{72a,72b}, A. Di Girolamo³⁷, G. Di Gregorio³⁷, A. Di Luca^{78a,78b}, B. Di Micco^{77a,77b}, R. Di Nardo^{77a,77b}, K. F. Di Petrillo⁴⁰, M. Diamantopoulou³⁵, F. A. Dias¹¹⁷, M. A. Diaz^{140a,140b}, A. R. Didenko³⁹, M. Didenko¹⁶⁹, S. D. Diefenbacher^{18a}, E. B. Diehl¹⁰⁸, S. Díez Cornell⁴⁸, C. Díez Pardos¹⁴⁷, C. Dimitriadi¹⁵⁰, A. Dimitrievska²¹, A. Dimri¹⁵¹, J. Dingfelder²⁵, T. Dingley¹²⁹, I-M. Dinu^{28b}, S. J. Dittmeier^{63b}, F. Dittus³⁷, M. Divisek¹³⁶, B. Dixit⁹⁴, F. Djama¹⁰⁴, T. Djobava^{155b}, C. Doglioni^{103,100}, A. Dohnalova^{29a}, Z. Dolezal¹³⁶, K. Domijan^{86a}, K. M. Dona⁴⁰, M. Donadelli^{83d}, B. Dong¹⁰⁹, J. Donini⁴¹, A. D'Onofrio^{72a,72b}, M. D'Onofrio⁹⁴, J. Dopke¹³⁷, A. Doria^{72a}, N. Dos Santos Fernandes^{133a}, P. Dougan¹⁰³, M. T. Dova⁹², A. T. Doyle⁵⁹, M. A. Draguet¹²⁹, M. P. Drescher⁵⁵, E. Dreyer¹⁷⁵, I. Drivas-koulouris¹⁰, M. Drnevich¹²⁰, M. Drozdova⁵⁶, D. Du⁶², T. A. du Pree¹¹⁷, Z. Duan^{114a}, F. Dubinin³⁹, M. Dubovsky^{29a}, E. Duchovni¹⁷⁵, G. Duckeck¹¹¹, P. K. Duckett⁹⁸, O. A. Ducu^{28b}, D. Duda⁵², A. Dudarev³⁷, E. R. Duden²⁷, M. D'uffizi¹⁰³, L. Duflo⁶⁶, M. Dührssen³⁷, I. Duminica^{28g}, A. E. Dumitriu^{28b}, M. Dunford^{63a}, S. Dungs⁴⁹, K. Dunne^{47a,47b}, A. Duperrin¹⁰⁴, H. Duran Yildiz^{3a}, M. Düren⁵⁸, A. Durglishvili^{155b}, D. Duvnjak³⁵, B. L. Dwyer¹¹⁸, G. I. Dyckes^{18a}, M. Dyndal^{86a}, B. S. Dziedzic³⁷, Z. O. Earnshaw¹⁵², G. H. Eberwein¹²⁹, B. Eckerova^{29a}, S. Eggebrecht⁵⁵, E. Egidio Purcino De Souza^{83e}, G. Eigen¹⁷, K. Einsweiler^{18a}, T. Ekelof¹⁶⁷, P. A. Ekman¹⁰⁰, S. El Farkh^{36b}, Y. El Ghazali⁶², H. El Jarrari³⁷, A. El Moussaouy^{36a}, V. Ellajosyula¹⁶⁷, M. Ellert¹⁶⁷, F. Ellinghaus¹⁷⁷, N. Ellis³⁷, J. Elmsheuser³⁰, M. Elsayy^{119a}, M. Elsing³⁷, D. Emelianov¹³⁷, Y. Enari⁸⁴, I. Ene^{18a}, S. Epari¹¹⁰, D. Ernani Martins Neto⁸⁷, F. Ernst³⁷, M. Errenst¹⁷⁷, M. Escalier⁶⁶, C. Escobar¹⁶⁹, E. Etzion¹⁵⁷, G. Evans^{133a,133b}, H. Evans⁶⁸, L. S. Evans⁹⁷, A. Ezhilov³⁸, S. Ezzarqtouni^{36a}, F. Fabbri^{24b,24a}, L. Fabbri^{24b,24a}, G. Facini⁹⁸, V. Fadeyev¹³⁹, R. M. Fakhruddinov³⁸, D. Fakoudis¹⁰², S. Falciano^{75a}, L. F. Falda Ulhoa Coelho^{133a}, F. Fallavollita¹¹², G. Falsetti^{44b,44a}, J. Faltova¹³⁶, C. Fan¹⁶⁸, K. Y. Fan^{64b}, Y. Fan¹⁴, Y. Fang^{14,114c}, M. Fanti^{71a,71b}, M. Faraj^{69a,69b}, Z. Farazpay⁹⁹, A. Farbin⁸, A. Farilla^{77a}, T. Farooque¹⁰⁹, J. N. Farr¹⁷⁸, S. M. Farrington^{137,52}, F. Fassi^{36e}, D. Fassouliotis⁹, L. Fayard⁶⁶, P. Federic¹³⁶, P. Federicova¹³⁴, O. L. Fedin^{38a}, M. Feickert¹⁷⁶, L. Feligioni¹⁰⁴, D. E. Fellers^{18a}, C. Feng^{143a}, Z. Feng¹¹⁷, M. J. Fenton¹⁶⁵, L. Ferencz⁴⁸, B. Fernandez Barbadillo⁹³, P. Fernandez Martinez⁶⁷, M. J. V. Fernoux¹⁰⁴, J. Ferrando⁹³, A. Ferrari¹⁶⁷, P. Ferrari^{117,116}, R. Ferrari^{73a}, D. Ferrere⁵⁶, C. Ferretti¹⁰⁸, M. P. Fewell¹, D. Fiacco^{75a,75b}, F. Fiedler¹⁰², P. Fiedler¹³⁵, S. Filimonov³⁹, M. S. Filip^{28b,u}, A. Filipčič⁹⁵, E. K. Filmer^{162a}, F. Filthaut¹¹⁶, M. C. N. Fiolhais^{133a,133c,c}, L. Fiorini¹⁶⁹, W. C. Fisher¹⁰⁹, T. Fitschen¹⁰³, P. M. Fitzhugh¹³⁸, I. Fleck¹⁴⁷, P. Fleischmann¹⁰⁸, T. Flick¹⁷⁷, M. Flores^{34d,ag}, L. R. Flores Castillo^{64a}, L. Flores Sanz De Acedo³⁷, F. M. Follega^{78a,78b}, N. Fomin³³, J. H. Foo¹⁶¹, A. Formica¹³⁸, A. C. Forti¹⁰³, E. Fortin³⁷, A. W. Fortman^{18a}, L. Foster^{18a}, L. Fountas^{9,j}, D. Fournier⁶⁶, H. Fox⁹³, P. Francavilla^{74a,74b}, S. Francescato⁶¹, S. Franchellucci⁵⁶, M. Franchini^{24b,24a}, S. Franchino^{63a}, D. Francis³⁷, L. Franco¹¹⁶, V. Franco Lima³⁷, L. Franconi⁴⁸, M. Franklin⁶¹, G. Frattari²⁷, Y. Y. Frid¹⁵⁷, J. Friend⁵⁹, N. Fritzsche³⁷, A. Froch⁵⁶, D. Froidevaux³⁷, J. A. Frost¹²⁹, Y. Fu¹⁰⁹, S. Fuenzalida Garrido^{140f}, M. Fujimoto¹⁰⁴, K. Y. Fung^{64a}, E. Furtado De Simas Filho^{83e}, M. Furukawa¹⁵⁹, J. Fuster¹⁶⁹, A. Gaa⁵⁵, A. Gabrielli^{24b,24a}, A. Gabrielli¹⁶¹, P. Gadow³⁷, G. Gagliardi^{57b,57a}, L. G. Gagnon^{18a}, S. Gaid^{88b}, S. Galantzan¹⁵⁷, J. Gallagher¹, E. J. Gallas¹²⁹, A. L. Gallen¹⁶⁷, B. J. Gallop¹³⁷, K. K. Gan¹²², S. Ganguly¹⁵⁹, Y. Gao⁵², A. Garabaglu¹⁴², F. M. Garay Walls^{140a,140b}, C. García¹⁶⁹, A. Garcia Alonso¹¹⁷, A. G. Garcia Caffaro¹⁷⁸, J. E. García Navarro¹⁶⁹, M. Garcia-Sciveres^{18a}, G. L. Gardner¹³¹, R. W. Gardner⁴⁰, N. Garelli¹⁶⁴, R. B. Garg¹⁴⁹, J. M. Gargan⁵², C. A. Garner¹⁶¹, C. M. Garvey^{34a}, V. K. Gassmann¹⁶⁴, G. Gaudio^{73a}, V. Gautam¹³, P. Gauzzi^{75a,75b}, J. Gavanovic⁹⁵, I. L. Gavrilenko^{133a}, A. Gavriluk³⁸, C. Gay¹⁷⁰, G. Gaycken¹²⁶, E. N. Gazis¹⁰, A. Gekow¹²², C. Gemme^{57b}, M. H. Genest⁶⁰, A. D. Gentry¹¹⁵, S. George⁹⁷, T. Gerasis⁴⁶, A. A. Gerwin¹²³, P. Gessinger-Befurt³⁷, M. E. Geyik¹⁷⁷, M. Ghani¹⁷³, K. Ghorbanian⁹⁶, A. Ghosal¹⁴⁷, A. Ghosh¹⁶⁵, A. Ghosh⁷, B. Giacobbe^{24b}, S. Giagu^{75a,75b}, T. Giani¹¹⁷, A. Giannini⁶², S. M. Gibson⁹⁷, M. Gignac¹³⁹, D. T. Gil^{86b}, A. K. Gilbert^{86a}, B. J. Gilbert⁴², D. Gillberg³⁵, G. Gilles¹¹⁷, D. M. Gingrich^{2,ai}, M. P. Giordani^{69a,69c}, P. F. Giraud¹³⁸, G. Giugliarelli^{69a,69c}, D. Giugni^{71a}, F. Giuli^{76a,76b}, I. Gkialas^{9,j}, L. K. Gladilin³⁸, C. Glasman¹⁰¹, M. Glazewska²⁰, R. M. Gleason¹⁶⁵, G. Glemža⁴⁸, M. Glisic¹²⁶, I. Gnesi^{44b}, Y. Go³⁰, M. Goblirsch-Kolb³⁷, B. Gocke⁴⁹, D. Godin¹¹⁰, B. Gokturk^{22a}, S. Goldfarb¹⁰⁷, T. Golling⁵⁶, M. G. D. Gololo^{34c}, D. Golubkov³⁸, J. P. Gombas¹⁰⁹, A. Gomes^{133a,133b}, G. Gomes Da Silva¹⁴⁷, A. J. Gomez Delegido¹⁶⁹, R. Gonçalves^{133a}, L. Gonella²¹, A. Gongadze^{155c}, F. Gonnella²¹, J. L. Gonski¹⁴⁹, R. Y. González Andana⁵², S. González de la Hoz¹⁶⁹, M. V. Gonzalez Rodrigues⁴⁸, R. Gonzalez Suarez¹⁶⁷, S. Gonzalez-Sevilla⁵⁶, L. Goossens³⁷, B. Gorini³⁷

E. Gorini^{70a,70b} , A. Gorišek⁹⁵ , T. C. Gosart¹³¹ , A. T. Goshaw⁵¹ , M. I. Gostkin³⁹ , S. Goswami¹²⁴ , C. A. Gottardo³⁷ , S. A. Gotz¹¹¹ , M. Goughri^{36b} , A. G. Goussiou¹⁴² , N. Govender^{34c} , R. P. Grabarczyk¹²⁹ , I. Grabowska-Bold^{86a} , K. Graham³⁵ , E. Gramstad¹²⁸ , S. Grancagnolo^{70a,70b} , C. M. Grant¹ , P. M. Gravila^{28f} , F. G. Gravili^{70a,70b} , H. M. Gray^{18a} , M. Greco¹¹² , M. J. Green¹ , C. Grefe²⁵ , A. S. Grefsrud¹⁷ , I. M. Gregor⁴⁸ , K. T. Greif¹⁶⁵ , P. Grenier¹⁴⁹ , S. G. Grewe¹¹² , A. A. Grillo¹³⁹ , K. Grimm³² , S. Grinstein^{13,y} , J.-F. Grivaz⁶⁶ , E. Gross¹⁷⁵ , J. Grosse-Knetter⁵⁵ , L. Guan¹⁰⁸ , G. Guerrieri³⁷ , R. Guevara¹²⁸ , R. Gugel¹⁰² , J. A. M. Guhit¹⁰⁸ , A. Guida¹⁹ , E. Guillon¹⁷³ , S. Guindon³⁷ , F. Guo^{14,114c} , J. Guo^{144a} , L. Guo⁴⁸ , L. Guo^{114b,w} , Y. Guo¹⁰⁸ , A. Gupta⁴⁹ , R. Gupta¹³² , S. Gupta²⁷ , S. Gurbuz²⁵ , S. S. Gurdasani⁴⁸ , G. Gustavino^{75a,75b} , P. Gutierrez¹²³ , L. F. Gutierrez Zagazeta¹³¹ , M. Gutsche⁵⁰ , C. Gutschow⁹⁸ , C. Gwenlan¹²⁹ , C. B. Gwilliam⁹⁴ , E. S. Haaland¹²⁸ , A. Haas¹²⁰ , M. Habedank⁵⁹ , C. Haber^{18a} , H. K. Hadavand⁸ , A. Haddad⁴¹ , A. Hadeef⁵⁰ , A. I. Hagan⁹³ , J. J. Hahn¹⁴⁷ , E. H. Haines⁹⁸ , M. Haleem¹⁷² , J. Haley¹²⁴ , G. D. Hallowell¹⁰⁴ , L. Halser²⁰ , K. Hamano¹⁷¹ , M. Hamer²⁵ , S. E. D. Hammoud⁶⁶ , E. J. Hampshire⁹⁷ , J. Han^{143a} , L. Han^{114a} , L. Han⁶² , S. Han^{18a} , K. Hanagaki⁸⁴ , M. Hance¹³⁹ , D. A. Hangal⁴² , H. Hanif¹⁴⁸ , M. D. Hank¹³¹ , J. B. Hansen⁴³ , P. H. Hansen⁴³ , D. Harada⁵⁶ , T. Harenberg¹⁷⁷ , S. Harkusha¹⁷⁹ , M. L. Harris¹⁰⁵ , Y. T. Harris²⁵ , J. Harrison¹³ , N. M. Harrison¹²² , P. F. Harrison¹⁷³ , M. L. E. Hart⁹⁸ , N. M. Hartman¹¹² , N. M. Hartmann¹¹¹ , R. Z. Hasan^{97,137} , Y. Hasegawa¹⁴⁶ , F. Haslbeck¹²⁹ , S. Hassan¹⁷ , R. Hauser¹⁰⁹ , M. Haviernik¹³⁶ , C. M. Hawkes²¹ , R. J. Hawkins³⁷ , Y. Hayashi¹⁵⁹ , D. Hayden¹⁰⁹ , C. Hayes¹⁰⁸ , R. L. Hayes¹¹⁷ , C. P. Hays¹²⁹ , J. M. Hays⁹⁶ , H. S. Hayward⁹⁴ , M. He^{14,114c} , Y. He⁴⁸ , Y. He⁹⁸ , N. B. Heatley⁹⁶ , V. Hedberg¹⁰⁰ , C. Heidegger⁵⁴ , K. K. Heidegger⁵⁴ , J. Heilman³⁵ , S. Heim⁴⁸ , T. Heim^{18a} , J. G. Heinlein¹³¹ , J. J. Heinrich¹²⁶ , L. Heinrich¹¹² , J. Hejbal¹³⁴ , M. Helbig⁵⁰ , A. Held¹⁷⁶ , S. Hellesund¹⁷ , C. M. Helling¹⁷⁰ , S. Hellman^{47a,47b} , A. M. Henriques Correia³⁷ , H. Herde¹⁰⁰ , Y. Hernández Jiménez¹⁵¹ , L. M. Herrmann²⁵ , T. Herrmann⁵⁰ , G. Herten⁵⁴ , R. Hertenberger¹¹¹ , L. Hervas³⁷ , M. E. Hesping¹⁰² , N. P. Hessey^{162a} , J. Hessler¹¹² , M. Hidaoui^{36b} , N. Hidic¹³⁶ , E. Hill¹⁶¹ , T. S. Hillersoy¹⁷ , S. J. Hillier²¹ , J. R. Hinds¹⁰⁹ , F. Hinterkeuser²⁵ , M. Hirose¹²⁷ , S. Hirose¹⁶³ , D. Hirschbuehl¹⁷⁷ , T. G. Hitchings¹⁰³ , B. Hiti⁹⁵ , J. Hobbs¹⁵¹ , R. Hobincu^{28e} , N. Hod¹⁷⁵ , A. M. Hodges¹⁶⁸ , M. C. Hodgkinson¹⁴⁵ , B. H. Hodgkinson¹²⁹ , A. Hoecker³⁷ , D. D. Hofer¹⁰⁸ , J. Hofer¹⁶⁹ , M. Holzbock³⁷ , L. B. A. H. Hommels³³ , V. Homsak¹²⁹ , B. P. Honan¹⁰³ , J. J. Hong⁶⁸ , T. M. Hong¹³² , B. H. Hooberman¹⁶⁸ , W. H. Hopkins⁶ , M. C. Hoppesch¹⁶⁸ , Y. Horii¹¹³ , M. E. Horstmann¹¹² , S. Hou¹⁵⁴ , M. R. Housenga¹⁶⁸ , A. S. Howard⁹⁵ , J. Howarth⁵⁹ , J. Hoya⁶ , M. Hrabovsky¹²⁵ , T. Hryn'ova⁴ , P. J. Hsu⁶⁵ , S.-C. Hsu¹⁴² , T. Hsu⁶⁶ , M. Hu^{18a} , Q. Hu⁶² , S. Huang³³ , X. Huang^{14,114c} , Y. Huang¹³⁶ , Y. Huang^{114b} , Y. Huang¹⁰² , Y. Huang¹⁴ , Z. Huang⁶⁶ , Z. Hubacek¹³⁵ , M. Huebner²⁵ , F. Hugging²⁵ , T. B. Huffman¹²⁹ , M. Hufnagel Maranha De Faria^{83a} , C. A. Hugli⁴⁸ , M. Huhtinen³⁷ , S. K. Huiberts¹⁷ , R. Hulsken¹⁰⁶ , C. E. Hultquist^{18a} , N. Huseynov^{12,g} , J. Huston¹⁰⁹ , J. Huth⁶¹ , R. Hyneman⁷ , G. Iacobucci⁵⁶ , G. Iakovidis³⁰ , L. Iconomidou-Fayard⁶⁶ , J. P. Iddon³⁷ , P. Iengo^{72a,72b} , R. Iguchi¹⁵⁹ , Y. Iiyama¹⁵⁹ , T. Iizawa¹⁵⁹ , Y. Ikegami⁸⁴ , D. Iliadis¹⁵⁸ , N. Ilic¹⁶¹ , H. Imam^{36a} , G. Inacio Goncalves^{83d} , S. A. Infante Cabanas^{140c} , T. Ingebretsen Carlson^{47a,47b} , J. M. Inglis⁹⁶ , G. Introzzi^{73a,73b} , M. Iodice^{77a} , V. Ippolito^{75a,75b} , R. K. Irwin⁹⁴ , M. Ishino¹⁵⁹ , W. Islam¹⁷⁶ , C. Issever¹⁹ , S. Istin^{22a,an} , K. Itabashi⁸⁴ , H. Ito¹⁷⁴ , R. Iuppa^{78a,78b} , A. Ivina¹⁷⁵ , V. Izzo^{72a} , P. Jacka¹³⁴ , P. Jackson¹ , P. Jain⁴⁸ , K. Jakobs⁵⁴ , T. Jakoubek¹⁷⁵ , J. Jamieson⁵⁹ , W. Jang¹⁵⁹ , S. Jankovych¹³⁶ , M. Javurkova¹⁰⁵ , P. Jawahar¹⁰³ , L. Jeanty¹²⁶ , J. Jejelava^{155a,af} , P. Jenni^{54,f} , C. E. Jessiman³⁵ , C. Jia^{143a} , H. Jia¹⁷⁰ , J. Jia¹⁵¹ , X. Jia^{14,114c} , Z. Jia^{114a} , C. Jiang⁵² , Q. Jiang^{64b} , S. Jiggins⁴⁸ , M. Jimenez Ortega¹⁶⁹ , J. Jimenez Pena¹³ , S. Jin^{114a} , A. Jinaru^{28b} , O. Jinnouchi¹⁴¹ , P. Johansson¹⁴⁵ , K. A. Johns⁷ , J. W. Johnson¹³⁹ , F. A. Jolly⁴⁸ , D. M. Jones¹⁵² , E. Jones⁴⁸ , K. S. Jones⁸ , P. Jones³³ , R. W. L. Jones⁹³ , T. J. Jones⁹⁴ , H. L. Joos^{55,37} , R. Joshi¹²² , J. Jovicevic¹⁶

J. Kirk¹³⁷, A. E. Kiryunin¹¹², S. Kita¹⁶³, O. Kivernyk²⁵, M. Klassen¹⁶⁴, C. Klein³⁵, L. Klein¹⁷², M. H. Klein⁴⁵, S. B. Klein⁵⁶, U. Klein⁹⁴, A. Klimentov³⁰, T. Klioutchnikova³⁷, P. Kluit¹¹⁷, S. Kluth¹¹², E. Kneringer⁷⁹, T. M. Knight¹⁶¹, A. Knue⁴⁹, M. Kobel⁵⁰, D. Kobylanski¹⁷⁵, S. F. Koch¹²⁹, M. Kocian¹⁴⁹, P. Kodyš¹³⁶, D. M. Koeck¹²⁶, T. Koffas³⁵, O. Kolay⁵⁰, I. Koletsou⁴, T. Komarek⁸⁷, K. Köneke⁵⁵, A. X. Y. Kong¹, T. Kono¹²¹, N. Konstantinidis⁹⁸, P. Kontaxakis⁵⁶, B. Konya¹⁰⁰, R. Kopeliansky⁴², S. Koperny^{86a}, K. Korcyl⁸⁷, K. Kordas^{158,d}, A. Korn⁹⁸, S. Korn⁵⁵, I. Korolkov¹³, N. Korotkova³⁸, B. Kortman¹¹⁷, O. Kortner¹¹², S. Kortner¹¹², W. H. Kostecka¹¹⁸, M. Kostov^{29a}, V. V. Kostyukhin¹⁴⁷, A. Kotsokechagia³⁷, A. Kotwal⁵¹, A. Koulouris³⁷, A. Kourkumeli-Charalampidi^{73a,73b}, C. Kourkumelis⁹, E. Kourlitis¹¹², O. Kovanda¹²⁶, R. Kowalewski¹⁷¹, W. Kozanecki¹²⁶, A. S. Kozhin³⁸, V. A. Kramarenko³⁸, G. Kramberger⁹⁵, P. Kramer²⁵, M. W. Krasny¹³⁰, A. Krasznahorkay¹⁰⁵, A. C. Kraus¹¹⁸, J. W. Kraus¹⁷⁷, J. A. Kremer⁴⁸, N. B. Krengel¹⁴⁷, T. Kresse⁵⁰, L. Kretschmann¹⁷⁷, J. Kretzschmar⁹⁴, K. Kreul¹⁹, P. Krieger¹⁶¹, K. Krizka²¹, K. Kroeninger⁴⁹, H. Kroha¹¹², J. Kroll¹³⁴, J. Kroll¹³¹, K. S. Krowpman¹⁰⁹, U. Kruchonak³⁹, H. Krüger²⁵, N. Krumnack⁸¹, M. C. Kruse⁵¹, O. Kuchinskaia³⁹, S. Kuday^{3a}, S. Kuehn³⁷, R. Kuesters⁵⁴, T. Kuhl⁴⁸, V. Kukhtin³⁹, Y. Kulchitsky³⁹, S. Kuleshov^{140d,140b}, J. Kull¹, M. Kumar^{34g}, N. Kumari⁴⁸, P. Kumari^{162b}, A. Kupco¹³⁴, T. Kupfer⁴⁹, A. Kupich³⁸, O. Kuprash⁵⁴, H. Kurashige⁸⁵, L. L. Kurchaninov^{162a}, O. Kurdysh⁴, Y. A. Kurochkin³⁸, A. Kurova³⁸, M. Kuze¹⁴¹, A. K. Kvam¹⁰⁵, J. Kvita¹²⁵, N. G. Kyriacou¹⁰⁸, C. Lacasta¹⁶⁹, F. Lacava^{75a,75b}, H. Lacker¹⁹, D. Lacour¹³⁰, N. N. Lad⁹⁸, E. Ladygin³⁹, A. Lafarge⁴¹, B. Laforge¹³⁰, T. Lagouri¹⁷⁸, F. Z. Lahbabi^{36a}, S. Lai⁵⁵, J. E. Lambert¹⁷¹, S. Lammers⁶⁸, W. Lampl⁷, C. Lampoudis^{158,d}, G. Lamprinoudis¹⁰², A. N. Lancaster¹¹⁸, E. Lançon³⁰, U. Landgraf⁵⁴, M. P. J. Landon⁹⁶, V. S. Lang⁵⁴, O. K. B. Langrekken¹²⁸, A. J. Lankford¹⁶⁵, F. Lanni³⁷, K. Lantzsch²⁵, A. Lanza^{73a}, M. Lanzac Berrocal¹⁶⁹, J. F. Laporte¹³⁸, T. Lari^{71a}, D. Larsen¹⁷, L. Larson¹¹, F. Lasagni Manghi^{24b}, M. Lassnig³⁷, S. D. Lawlor¹⁴⁵, R. Lazaridou¹⁷³, M. Lazzaroni^{71a,71b}, H. D. M. Le¹⁰⁹, E. M. Le Boulicaut¹⁷⁸, L. T. Le Pottier^{18a}, B. Leban^{24b,24a}, F. Ledroit-Guillon⁶⁰, T. F. Lee^{162b}, L. L. Leeuw^{34c}, M. Lefebvre¹⁷¹, C. Leggett^{18a}, G. Lehmann Miotto³⁷, M. Leigh⁵⁶, W. A. Leight¹⁰⁵, W. Leinonen¹¹⁶, A. Leisos^{158,v}, M. A. L. Leite^{83c}, C. E. Leitgeb¹⁹, R. Leitner¹³⁶, K. J. C. Leney⁴⁵, T. Lenz²⁵, S. Leone^{74a}, C. Leonidopoulos⁵², A. Leopold¹⁵⁰, J. H. Lepage Bourbonnais³⁵, R. Les¹⁰⁹, C. G. Lester³³, M. Levchenko³⁸, J. Levêque⁴, L. J. Levinson¹⁷⁵, G. Levri^{24b,24a}, M. P. Lewicki⁸⁷, C. Lewis¹⁴², D. J. Lewis⁴, L. Lewitt¹⁴⁵, A. Li³⁰, B. Li^{143a}, C. Li¹⁰⁸, C-Q. Li¹¹², H. Li^{143a}, H. Li¹⁰³, H. Li¹⁵, H. Li⁶², H. Li^{143a}, J. Li^{144a}, K. Li¹⁴, L. Li^{144a}, R. Li¹⁷⁸, S. Li^{14,114c}, S. Li^{144b,144a}, T. Li⁵, X. Li¹⁰⁶, Z. Li¹⁵⁹, Z. Li^{14,114c}, Z. Li⁶², S. Liang^{14,114c}, Z. Liang¹⁴, M. Liberatore¹³⁸, B. Liberti^{76a}, K. Lie^{64c}, J. Lieber Marin^{83e}, H. Lien⁶⁸, H. Lin¹⁰⁸, S. F. Lin¹⁵¹, L. Linden¹¹¹, R. E. Lindley⁷, J. H. Lindon³⁷, J. Ling⁶¹, E. Lipeles¹³¹, A. Lipniacka¹⁷, A. Lister¹⁷⁰, J. D. Little⁶⁸, B. Liu¹⁴, B. X. Liu^{114b}, D. Liu^{144b,144a}, D. Liu¹³⁹, E. H. L. Liu²¹, J. K. K. Liu¹²⁰, K. Liu^{144b}, K. Liu^{144b,144a}, M. Liu⁶², M. Y. Liu⁶², P. Liu¹⁴, Q. Liu^{144b,142,144a}, X. Liu⁶², X. Liu^{143a}, Y. Liu^{114b,114c}, Y. L. Liu^{143a}, Y. W. Liu⁶², Z. Liu^{66,1}, S. L. Lloyd⁹⁶, E. M. Lobodzinska⁴⁸, P. Loch⁷, E. Lodhi¹⁶¹, T. Lohse¹⁹, K. Lohwasser¹⁴⁵, E. Loiacono⁴⁸, J. D. Lomas²¹, J. D. Long⁴², I. Longarini¹⁶⁵, R. Longo¹⁶⁸, A. Lopez Solis¹³, N. A. Lopez-canelas⁷, N. Lorenzo Martinez⁴, A. M. Lory¹¹¹, M. Losada^{119a}, G. Löschke Centeno¹⁵², X. Lou^{47a,47b}, X. Lou^{14,114c}, A. Lounis⁶⁶, P. A. Love⁹³, M. Lu⁶⁶, S. Lu¹³¹, Y. J. Lu¹⁵⁴, H. J. Lubatti¹⁴², C. Luci^{75a,75b}, F. L. Lucio Alves^{114a}, F. Luehring⁶⁸, B. S. Lunday¹³¹, O. Lundberg¹⁵⁰, J. Lunde³⁷, N. A. Luongo⁶, M. S. Lutz³⁷, A. B. Lux²⁶, D. Lynn³⁰, R. Lysak¹³⁴, V. Lysenko¹³⁵, E. Lytken¹⁰⁰, V. Lyubushkin³⁹, T. Lyubushkina³⁹, M. M. Lyukova¹⁵¹, M. Firdaus M. Soberi⁵², H. Ma³⁰, K. Ma⁶², L. L. Ma^{143a}, W. Ma⁶², Y. Ma¹²⁴, J. C. MacDonald¹⁰², P. C. Machado De Abreu Farias^{83c}, R. Madar⁴¹, T. Madula⁹⁸, J. Maeda⁸⁵, T. Maeno³⁰, P. T. Mafa^{34c,k}, H. Maguire¹⁴⁵, V. Maiboroda⁶⁶, A. Maio^{133a,133b,133d}, K. Maj^{86a}, O. Majersky⁴⁸, S. Majewski¹²⁶, R. Makhmanazarov³⁸, N. Makovec⁶⁶, V. Maksimovic¹⁶, B. Malaescu¹³⁰, J. Malamant¹²⁸, Pa. Malecki⁸⁷, V. P. Maleev³⁸, F. Malek^{60,p}, M. Mali⁹⁵, D. Malito⁹⁷, U. Mallik^{80,*}, A. Maloizel⁵, S. Maltezos¹⁰, A. Malvezzi Lopes^{83d}, S. Malyukov³⁹, J. Mamuzic¹³, G. Mancini⁵³, M. N. Mancini²⁷, G. Manco^{73a,73b}, J. P. Mandalia⁹⁶, S. S. Mandary¹⁵², I. Mandić⁹⁵, L. Manhaes de Andrade Filho^{83a}, I. M. Maniatis¹⁷⁵, J. Manjarres Ramos⁹¹, D. C. Mankad¹⁷⁵, A. Mann¹¹¹, T. Manoussos³⁷, M. N. Mantinan⁴⁰, S. Manzoni³⁷, L. Mao^{144a}, X. Mapekula^{34c}, A. Marantis¹⁵⁸, R. R. Marcelo Gregorio⁹⁶, G. Marchiori⁵, M. Marcisovsky¹³⁴, C. Marcon^{71a}, E. Maricic¹⁶, M. Marinescu⁴⁸, S. Marium⁴⁸, M. Marjanovic¹²³, A. Markhoos⁵⁴, M. Markovitch⁶⁶, M. K. Maroun¹⁰⁵, G. T. Marsden¹⁰³, E. J. Marshall⁹³, Z. Marshall^{18a}, S. Marti-Garcia¹⁶⁹, J. Martin⁹⁸, T. A. Martin¹³⁷, V. J. Martin⁵², B. Martin dit Latour¹⁷, L. Martinelli^{75a,75b}, M. Martinez^{13,y}, P. Martinez Agullo¹⁶⁹,

V. I. Martinez Outschoorn¹⁰⁵, P. Martinez Suarez¹³, S. Martin-Haugh¹³⁷, G. Martinovicova¹³⁶, V. S. Martoiu^{28b}, A. C. Martyniuk⁹⁸, A. Marzin³⁷, D. Mascione^{78a,78b}, L. Masetti¹⁰², J. Masik¹⁰³, A. L. Maslennikov³⁹, S. L. Mason⁴², P. Massarotti^{72a,72b}, P. Mastrandrea^{74a,74b}, A. Mastroberardino^{44b,44a}, T. Masubuchi¹²⁷, T. T. Mathew¹²⁶, J. Matousek¹³⁶, D. M. Mattern⁴⁹, J. Maurer^{28b}, T. Maurin⁵⁹, A. J. Maury⁶⁶, B. Maček⁹⁵, C. Mavungu Tsava¹⁰⁴, D. A. Maximov³⁸, A. E. May¹⁰³, E. Mayer⁴¹, R. Mazini^{34g}, I. Maznas¹¹⁸, S. M. Mazza¹³⁹, E. Mazzeo³⁷, J. P. Mc Gowan¹⁷¹, S. P. Mc Kee¹⁰⁸, C. A. Mc Lean⁶, C. C. McCracken¹⁷⁰, E. F. McDonald¹⁰⁷, A. E. McDougall¹¹⁷, L. F. Mcelhinney⁹³, J. A. Mcfayden¹⁵², R. P. McGovern¹³¹, R. P. Mckenzie^{34g}, T. C. McLachlan⁴⁸, D. J. McLaughlin⁹⁸, S. J. McMahon¹³⁷, C. M. Mcpartland⁹⁴, R. A. McPherson^{171,ac}, S. Mehlhase¹¹¹, A. Mehta⁹⁴, D. Melini¹⁶⁹, B. R. Mellado Garcia^{34g}, A. H. Melo⁵⁵, F. Meloni⁴⁸, A. M. Mendes Jacques Da Costa¹⁰³, L. Meng⁹³, S. Menke¹¹², M. Mentink³⁷, E. Meoni^{44b,44a}, G. Mercado¹¹⁸, S. Merianos¹⁵⁸, C. Merlassino^{69a,69c}, C. Meroni^{71a,71b}, J. Metcalfe⁶, A. S. Mete⁶, E. Meuser¹⁰², C. Meyer⁶⁸, J.-P. Meyer¹³⁸, Y. Miao^{114a}, R. P. Middleton¹³⁷, M. Mihovilovic⁶⁶, L. Mijović⁵², G. Mikenberg¹⁷⁵, M. Mikesstikova¹³⁴, M. Mikuž⁹⁵, H. Mildner¹⁰², A. Milic³⁷, D. W. Miller⁴⁰, E. H. Miller¹⁴⁹, L. S. Miller³⁵, A. Milov¹⁷⁵, D. A. Milstead^{47a,47b}, T. Min^{114a}, A. A. Minaenko³⁸, I. A. Minashvili^{155b}, A. I. Mincer¹²⁰, B. Mindur^{86a}, M. Mineev³⁹, Y. Mino⁸⁹, L. M. Mir¹³, M. Miralles Lopez⁵⁹, M. Mironova^{18a}, M. C. Missio¹¹⁶, A. Mitra¹⁷³, V. A. Mitsou¹⁶⁹, Y. Mitsumori¹¹³, O. Miu¹⁶¹, P. S. Miyagawa⁹⁶, T. Mkrtychyan^{63a}, M. Mlinarevic⁹⁸, T. Mlinarevic⁹⁸, M. Mlynarikova³⁷, S. Mobius²⁰, M. H. Mohamed Farook¹¹⁵, S. Mohapatra⁴², S. Mohiuddin¹²⁴, G. Mokgatitswane^{34g}, L. Moleri¹⁷⁵, U. Molinatti¹²⁹, L. G. Mollier²⁰, B. Mondal¹⁴⁷, S. Mondal¹³⁵, K. Mönig⁴⁸, E. Monnier¹⁰⁴, L. Monsonis Romero¹⁶⁹, J. Montejo Berlingen¹³, A. Montella^{47a,47b}, M. Montella¹²², F. Montereali^{77a,77b}, F. Monticelli⁹², S. Monzani^{69a,69c}, A. Morancho Tarda⁴³, N. Morange⁶⁶, A. L. Moreira De Carvalho⁴⁸, M. Moreno Llácer¹⁶⁹, C. Moreno Martinez⁵⁶, J. M. Moreno Perez^{23b}, P. Morettini^{57b}, S. Morgenstern³⁷, M. Morii⁶¹, M. Morinaga¹⁵⁹, M. Moritsu⁹⁰, F. Morodei^{75a,75b}, P. Moschovakos³⁷, B. Moser⁵⁴, M. Mosidze^{155b}, T. Moskalets⁴⁵, P. Moskvitina¹¹⁶, J. Moss³², P. Moszkowicz^{86a}, A. Moussa^{36d}, Y. Moyal¹⁷⁵, H. Moyano Gomez¹³, E. J. W. Moyse¹⁰⁵, O. Mtintsilana^{34g}, S. Muanza¹⁰⁴, M. Mucha²⁵, J. Mueller¹³², R. Müller³⁷, G. A. Mullier¹⁶⁷, A. J. Mullin³³, J. J. Mullin⁵¹, A. C. Mullins⁴⁵, A. E. Mulski⁶¹, D. P. Mungo¹⁶¹, D. Munoz Perez¹⁶⁹, F. J. Munoz Sanchez¹⁰³, W. J. Murray^{173,137}, M. Muškinja⁹⁵, C. Mwewa⁴⁸, A. G. Myagkov^{38,a}, A. J. Myers⁸, G. Myers¹⁰⁸, M. Myska¹³⁵, B. P. Nachman^{18a}, K. Nagai¹²⁹, K. Nagano⁸⁴, R. Nagasaka¹⁵⁹, J. L. Nagle^{30,ak}, E. Nagy¹⁰⁴, A. M. Nairz³⁷, Y. Nakahama⁸⁴, K. Nakamura⁸⁴, K. Nakkalil⁵, A. Nandi^{63b}, H. Nanjo¹²⁷, E. A. Narayanan⁴⁵, Y. Narukawa¹⁵⁹, I. Naryshkin³⁸, L. Nasella^{71a,71b}, S. Nasri^{119b}, C. Nass²⁵, G. Navarro^{23a}, J. Navarro-Gonzalez¹⁶⁹, A. Nayaz¹⁹, P. Y. Nechaeva³⁸, S. Nechaeva^{24b,24a}, F. Nechansky¹³⁴, L. Nedic¹²⁹, T. J. Neep²¹, A. Negri^{73a,73b}, M. Negrini^{24b}, C. Nellist¹¹⁷, C. Nelson¹⁰⁶, K. Nelson¹⁰⁸, S. Nemecek¹³⁴, M. Nessi^{37,h}, M. S. Neubauer¹⁶⁸, J. Newell⁹⁴, P. R. Newman²¹, Y. W. Y. Ng¹⁶⁸, B. Ngair^{119a}, H. D. N. Nguyen¹¹⁰, J. D. Nichols¹²³, R. B. Nickerson¹²⁹, R. Nicolaidou¹³⁸, J. Nielsen¹³⁹, M. Niemeyer⁵⁵, J. Niermann³⁷, N. Nikiforou³⁷, V. Nikolaenko^{38,a}, I. Nikolic-Audit¹³⁰, P. Nilsson³⁰, I. Ninca⁴⁸, G. Ninio¹⁵⁷, A. Nisati^{75a}, N. Nishu², R. Nisius¹¹², N. Nitika^{69a,69c}, J.-E. Nitschke⁵⁰, E. K. Nkadimeng^{34b}, T. Nobe¹⁵⁹, T. Nommensen¹⁵³, M. B. Norfolk¹⁴⁵, B. J. Norman³⁵, M. Noury^{36a}, J. Novak⁹⁵, T. Novak⁹⁵, R. Novotny¹³⁵, L. Nozka¹²⁵, K. Ntekas¹⁶⁵, N. M. J. Nunes De Moura Junior^{83b}, J. Ocariz¹³⁰, A. Ochi⁸⁵, I. Ochoa^{133a}, S. Oerdek^{48,z}, J. T. Offermann⁴⁰, A. Ogrodnik¹³⁶, A. Oh¹⁰³, C. C. Ohm¹⁵⁰, H. Oide⁸⁴, M. L. Ojeda³⁷, Y. Okumura¹⁵⁹, L. F. Oleiro Seabra^{133a}, I. Oleksiyuk⁵⁶, G. Oliveira Correa¹³, D. Oliveira Damazio³⁰, J. L. Oliver¹⁶⁵, Ö. O. Öncel⁵⁴, A. P. O'Neill²⁰, A. Onofre^{133a,133e}, P. U. E. Onyisi¹¹, M. J. Oreglia⁴⁰, D. Orestano^{77a,77b}, R. Orlandini^{77a,77b}, R. S. Orr¹⁶¹, L. M. Osojnak¹³¹, Y. Osumi¹¹³, G. Otero y Garzon³¹, H. Otono⁹⁰, G. J. Ottino^{18a}, M. Ouchrif^{36d}, F. Ould-Saada¹²⁸, T. Ovsiannikova¹⁴², M. Owen⁵⁹, R. E. Owen¹³⁷, V. E. Ozcan^{22a}, F. Ozturk⁸⁷, N. Ozturk⁸, S. Ozturk⁸², H. A. Pacey¹²⁹, K. Pachal^{162a}, A. Pacheco Pages¹³, C. Padilla Aranda¹³, G. Padovano^{75a,75b}, S. Pagan Griso^{18a}, G. Palacino⁶⁸, A. Palazzo^{70a,70b}, J. Pampel²⁵, J. Pan¹⁷⁸, T. Pan^{64a}, D. K. Panchal¹¹, C. E. Pandini⁶⁰, J. G. Panduro Vazquez¹³⁷, H. D. Pandya¹, H. Pang¹³⁸, P. Pani⁴⁸, G. Panizzo^{69a,69c}, L. Panwar¹³⁰, L. Paolozzi⁵⁶, S. Parajuli¹⁶⁸, A. Paramonov⁶, C. Paraskevopoulos⁵³, D. Paredes Hernandez^{64b}, A. Pareti^{73a,73b}, K. R. Park⁴², T. H. Park¹¹², F. Parodi^{57b,57a}, J. A. Parsons⁴², U. Parzefall⁵⁴, B. Pascual Dias⁴¹, L. Pascual Dominguez¹⁰¹, E. Pasqualucci^{75a}, S. Passaggio^{57b}, F. Pastore⁹⁷, P. Patel⁸⁷, U. M. Patel⁵¹, J. R. Pater¹⁰³, T. Pauly³⁷, F. Pauwels¹³⁶, C. I. Pazos¹⁶⁴, M. Pedersen¹²⁸, R. Pedro^{133a}, S. V. Peleganchuk³⁸, O. Penc³⁷, E. A. Pender⁵², S. Peng¹⁵, G. D. Penn¹⁷⁸, K. E. Penski¹¹¹, M. Penzin³⁸, B. S. Peralva^{83d}, A. P. Pereira Peixoto¹⁴², L. Pereira Sanchez¹⁴⁹,

D. V. Perepelitsa^{30,ak}, G. Perera¹⁰⁵, E. Perez Codina³⁷, M. Perganti¹⁰, H. Pernegger³⁷, S. Perrella^{75a,75b}, O. Perrin⁴¹, K. Peters⁴⁸, R. F. Y. Peters¹⁰³, B. A. Petersen³⁷, T. C. Petersen⁴³, E. Petir¹⁰⁴, V. Petousis¹³⁵, A. R. Petri^{71a,71b}, C. Petridou^{158,d}, T. Petru¹³⁶, A. Petrukhin¹⁴⁷, M. Pettee^{18a}, A. Petukhov⁸², K. Petukhova³⁷, R. Pezoa^{140f}, L. Pezzotti^{24b,24a}, G. Pezzullo¹⁷⁸, L. Pfaffenbichler³⁷, A. J. Pflieger³⁷, T. M. Pham¹⁷⁶, T. Pham¹⁰⁷, P. W. Phillips¹³⁷, G. Piacquadio¹⁵¹, E. Pianori^{18a}, F. Piazza¹²⁶, R. Piegai³¹, D. Pietreanu^{28b}, A. D. Pilkington¹⁰³, M. Pinamonti^{69a,69c}, J. L. Pinfeld², B. C. Pinheiro Pereira^{133a}, J. Pinol Bel¹³, A. E. Pinto Pinoargote¹³⁰, L. Pintucci^{69a,69c}, K. M. Piper¹⁵², A. Pirttikoski⁵⁶, D. A. Pizzi³⁵, L. Pizzimento^{64b}, A. Plebani³³, M.-A. Pleier³⁰, V. Pleskot¹³⁶, E. Plotnikova³⁹, G. Poddar⁹⁶, R. Poettgen¹⁰⁰, L. Poggioli¹³⁰, S. Polacek¹³⁶, G. Polesello^{73a}, A. Poley¹⁴⁸, A. Polini^{24b}, C. S. Pollard¹⁷³, Z. B. Pollock¹²², E. Pompa Pacchi¹²³, N. I. Pond⁹⁸, D. Ponomarenko⁶⁸, L. Pontecorvo³⁷, S. Popa^{28a}, G. A. Popeneciu^{28d}, A. Poreba³⁷, D. M. Portillo Quintero^{162a}, S. Pospisil¹³⁵, M. A. Postill¹⁴⁵, P. Postolache^{28c}, K. Potamianos¹⁷³, P. A. Potepa^{86a}, I. N. Potrap³⁹, C. J. Potter³³, H. Potti¹⁵³, J. Poveda¹⁶⁹, M. E. Pozo Astigarraga³⁷, R. Pozzi³⁷, A. Prades Ibanez^{76a,76b}, J. Pretel¹⁷¹, D. Price¹⁰³, M. Primavera^{70a}, L. Primomo^{69a,69c}, M. A. Principe Martin¹⁰¹, R. Privara¹²⁵, T. Procter^{86b}, M. L. Proffitt¹⁴², N. Proklova¹³¹, K. Prokofiev^{64c}, G. Proto¹¹², J. Proudfoot⁶, M. Przybycien^{86a}, W. W. Przygoda^{86b}, A. Psallidas⁴⁶, J. E. Puddefoot¹⁴⁵, D. Pudzha⁵³, D. Pyatizbyantseva¹¹⁶, J. Qian¹⁰⁸, R. Qian¹⁰⁹, D. Qichen¹⁰³, Y. Qin¹³, T. Qiu⁵², A. Quadri⁵⁵, M. Queitsch-Maitland¹⁰³, G. Quetant⁵⁶, R. P. Quinn¹⁷⁰, G. Rabanal Bolanos⁶¹, D. Rafanoharana¹¹², F. Raffaelli^{76a,76b}, F. Ragusa^{71a,71b}, J. L. Rainbolt⁴⁰, J. A. Raine⁵⁶, S. Rajagopalan³⁰, E. Ramakoti³⁹, L. Rambelli^{57b,57a}, I. A. Ramirez-Berend³⁵, K. Ran^{48,114c}, D. S. Rankin¹³¹, N. P. Rapheeha^{34g}, H. Rasheed^{28b}, D. F. Rassloff^{63a}, A. Rastogi^{18a}, S. Rave¹⁰², S. Ravera^{57b,57a}, B. Ravina³⁷, I. Ravinovich¹⁷⁵, M. Raymond³⁷, A. L. Read¹²⁸, N. P. Readioff¹⁴⁵, D. M. Rebuzzi^{73a,73b}, A. S. Reed¹¹², K. Reeves²⁷, J. A. Reidelsturz¹⁷⁷, D. Reikher¹²⁶, A. Rej⁴⁹, C. Rembser³⁷, H. Ren⁶², M. Renda^{28b}, F. Renner⁴⁸, A. G. Rennie⁵⁹, A. L. Rescia⁴⁸, S. Resconi^{71a}, M. Ressegotti^{57b,57a}, S. Rettie³⁷, W. F. Rettie³⁵, E. Reynolds^{18a}, O. L. Rezanova³⁹, P. Reznicek¹³⁶, H. Riani^{36d}, N. Ribaric⁵¹, E. Ricci^{78a,78b}, R. Richter¹¹², S. Richter^{47a,47b}, E. Richter-Was^{86b}, M. Ridel¹³⁰, S. Ridouani^{36d}, P. Rieck¹²⁰, P. Riedler³⁷, E. M. Riefel^{47a,47b}, J. O. Rieger¹¹⁷, M. Rijssenbeek¹⁵¹, M. Rimoldi³⁷, L. Rinaldi^{24b,24a}, P. Rincke^{167,55}, G. Ripellino¹⁶⁷, I. Riu¹³, J. C. Rivera Vergara¹⁷¹, F. Rizatdinova¹²⁴, E. Rizvi⁹⁶, B. R. Roberts^{18a}, S. S. Roberts¹³⁹, D. Robinson³³, M. Robles Manzano¹⁰², A. Robson⁵⁹, A. Rocchi^{76a,76b}, C. Roda^{74a,74b}, S. Rodriguez Bosca³⁷, Y. Rodriguez Garcia^{23a}, A. M. Rodríguez Vera¹¹⁸, S. Roe³⁷, J. T. Roemer³⁷, O. Røhne¹²⁸, R. A. Rojas³⁷, C. P. A. Roland¹³⁰, A. Romaniouk⁷⁹, E. Romano^{73a,73b}, M. Romano^{24b}, A. C. Romero Hernandez¹⁶⁸, N. Rompotis⁹⁴, L. Roos¹³⁰, S. Rosati^{75a}, B. J. Rosser⁴⁰, E. Rossi¹²⁹, E. Rossi^{72a,72b}, L. P. Rossi⁶¹, L. Rossini⁵⁴, R. Rosten¹²², M. Rotaru^{28b}, B. Rottler⁵⁴, D. Rousseau⁶⁶, D. Rouso⁴⁸, S. Roy-Garand¹⁶¹, A. Rozanov¹⁰⁴, Z. M. A. Rozario⁵⁹, Y. Rozen¹⁵⁶, A. Rubio Jimenez¹⁶⁹, V. H. Ruelas Rivera¹⁹, T. A. Ruggeri¹, A. Ruggiero¹²⁹, A. Ruiz-Martinez¹⁶⁹, A. Rummler³⁷, Z. Rurikova⁵⁴, N. A. Rusakovich³⁹, H. L. Russell¹⁷¹, G. Russo^{75a,75b}, J. P. Rutherford⁷, S. Rutherford Colmenares³³, M. Rybar¹³⁶, P. Rybczynski^{86a}, A. Ryzhov⁴⁵, J. A. Sabater Iglesias⁵⁶, H.F.-W. Sadrozinski¹³⁹, F. Safai Tehrani^{75a}, S. Saha¹, M. Sahinsoy⁸², B. Sahoo¹⁷⁵, A. Saibel¹⁶⁹, B. T. Saifuddin¹²³, M. Saimpert¹³⁸, G. T. Saito^{83c}, M. Saito¹⁵⁹, T. Saito¹⁵⁹, A. Sala^{71a,71b}, A. Salnikov¹⁴⁹, J. Salt¹⁶⁹, A. Salvador Salas¹⁵⁷, F. Salvatore¹⁵², A. Salzburger³⁷, D. Sammel⁵⁴, E. Sampson⁹³, D. Sampsonidis^{158,d}, D. Sampsonidou¹²⁶, J. Sánchez¹⁶⁹, V. Sanchez Sebastian¹⁶⁹, H. Sandaker¹²⁸, C. O. Sander⁴⁸, J. A. Sandesara¹⁷⁶, M. Sandhoff¹⁷⁷, C. Sandoval^{23b}, L. Sanfilippo^{63a}, D. P. C. Sankey¹³⁷, T. Sano⁸⁹, A. Sansoni⁵³, L. Santi³⁷, C. Santoni⁴¹, H. Santos^{133a,133b}, A. Santra¹⁷⁵, E. Sanzani^{24b,24a}, K. A. Saoucha^{88b}, J. G. Saraiva^{133a,133d}, J. Sardain⁷, O. Sasaki⁸⁴, K. Sato¹⁶³, C. Sauer³⁷, E. Sauvan⁴, P. Savard^{161,ai}, R. Sawada¹⁵⁹, C. Sawyer¹³⁷, L. Sawyer⁹⁹, C. Sbarra^{24b}, A. Sbrizzi^{24b,24a}, T. Scanlon⁹⁸, J. Schaarschmidt¹⁴², U. Schäfer¹⁰², A. C. Schaffer^{66,45}, D. Schaile¹¹¹, R. D. Schamberger¹⁵¹, C. Scharf¹⁹, M. M. Schefer²⁰, V. A. Schegelsky³⁸, D. Scheirich¹³⁶, M. Schernau^{140e}, C. Scheulen⁵⁶, C. Schiavi^{57b,57a}, M. Schioppa^{44b,44a}, B. Schlag¹⁴⁹, S. Schlenker³⁷, J. Schmeing¹⁷⁷, E. Schmidt¹¹², M. A. Schmidt¹⁷⁷, K. Schmieden¹⁰², C. Schmitt¹⁰², N. Schmitt¹⁰², S. Schmitt⁴⁸, L. Schoeffel¹³⁸, A. Schoening^{63b}, P. G. Scholer³⁵, E. Schopf¹⁴⁷, M. Schott²⁵, S. Schramm⁵⁶, T. Schroer⁵⁶, H.-C. Schultz-Coulon^{63a}, M. Schumacher⁵⁴, B. A. Schumm¹³⁹, Ph. Schune¹³⁸, H. R. Schwartz¹³⁹, A. Schwartzman¹⁴⁹, T. A. Schwarz¹⁰⁸, Ph. Schwemling¹³⁸, R. Schwiendorst¹⁰⁹, F. G. Sciacca²⁰, A. Sciandra³⁰, G. Sciolla²⁷, F. Scuri^{74a}, C. D. Sebastiani³⁷, K. Sedlaczek¹¹⁸, S. C. Seidel¹¹⁵, A. Seiden¹³⁹, B. D. Seidlitz⁴², C. Seitz⁴⁸, J. M. Seixas^{83b}, G. Sekhniaidze^{72a}, L. Selem⁶⁰, N. Semprini-Cesari^{24b,24a}, A. Semushin¹⁷⁹, D. Sengupta⁵⁶, V. Senthilkumar¹⁶⁹, L. Serin⁶⁶, M. Sessa^{72a,72b}, H. Severini¹²³

F. Sforza^{57b,57a} , A. Sfyrta⁵⁶ , Q. Sha¹⁴ , E. Shabalina⁵⁵ , H. Shaddix¹¹⁸ , A. H. Shah³³ , R. Shaheen¹⁵⁰ , J. D. Shahinian¹³¹ , M. Shamim³⁷ , L. Y. Shan¹⁴ , M. Shapiro^{18a} , A. Sharma³⁷ , A. S. Sharma¹⁷⁰ , P. Sharma³⁰ , P. B. Shatalov³⁸ , K. Shaw¹⁵² , S. M. Shaw¹⁰³ , Q. Shen^{144a} , D. J. Sheppard¹⁴⁸ , P. Sherwood⁹⁸ , L. Shi⁹⁸ , X. Shi¹⁴ , S. Shimizu⁸⁴ , C. O. Shimmin¹⁷⁸ , I. P. J. Shipsey^{129,*} , S. Shirabe⁹⁰ , M. Shiyakova^{39,aa} , M. J. Shochet⁴⁰ , D. R. Shope¹²⁸ , B. Shrestha¹²³ , S. Shrestha^{122,am} , I. Shreyber³⁹ , M. J. Shroff¹⁷¹ , P. Sicho¹³⁴ , A. M. Sickles¹⁶⁸ , E. Sideras Haddad^{34g,166} , A. C. Sidley¹¹⁷ , A. Sidoti^{24b} , F. Siegert⁵⁰ , Dj. Sijacki¹⁶ , F. Sili⁹² , J. M. Silva⁵² , I. Silva Ferreira^{83b} , M. V. Silva Oliveira³⁰ , S. B. Silverstein^{47a} , S. Simion⁶⁶ , R. Simoniello³⁷ , E. L. Simpson¹⁰³ , H. Simpson¹⁵² , L. R. Simpson⁶ , S. Simsek⁸² , S. Sindhu⁵⁵ , P. Sinervo¹⁶¹ , S. N. Singh²⁷ , S. Singh³⁰ , S. Sinha⁴⁸ , S. Sinha¹⁰³ , M. Sioli^{24b,24a} , K. Sioulas⁹ , I. Siral³⁷ , E. Sitnikova⁴⁸ , J. Sjölin^{47a,47b} , A. Skaf⁵⁵ , E. Skorda²¹ , P. Skubic¹²³ , M. Slawinska⁸⁷ , I. Slazyk¹⁷ , I. Sliusar¹²⁸ , V. Smakhtin¹⁷⁵ , B. H. Smart¹³⁷ , S. Yu. Smirnov^{140b} , Y. Smirnov⁸² , L. N. Smirnova^{38,a} , O. Smirnova¹⁰⁰ , A. C. Smith⁴² , D. R. Smith¹⁶⁵ , J. L. Smith¹⁰³ , M. B. Smith³⁵ , R. Smith¹⁴⁹ , H. Smitmanns¹⁰² , M. Smizanska⁹³ , K. Smolek¹³⁵ , P. Smolyanskiy¹³⁵ , A. A. Snesarev³⁹ , H. L. Snoek¹¹⁷ , S. Snyder³⁰ , R. Sobie^{171,ac} , A. Soffer¹⁵⁷ , C. A. Solans Sanchez³⁷ , E. Yu. Soldatov³⁹ , U. Soldevila¹⁶⁹ , A. A. Solodkov^{34g} , S. Solomon²⁷ , A. Soloshenko³⁹ , K. Solovieva⁵⁴ , O. V. Solovyanov⁴¹ , P. Sommer⁵⁰ , A. Sonay¹³ , A. Sopczak¹³⁵ , A. L. Sopio⁵² , F. Sopkova^{29b} , J. D. Sorenson¹¹⁵ , I. R. Sotarriva Alvarez¹⁴¹ , V. Sothilingam^{63a} , O. J. Soto Sandoval^{140c,140b} , S. Sottocornola⁶⁸ , R. Soualah^{88a} , Z. Soumami^{36e} , D. South⁴⁸ , N. Soybelman¹⁷⁵ , S. Spagnolo^{70a,70b} , M. Spalla¹¹² , D. Sperlich⁵⁴ , B. Spisso^{72a,72b} , D. P. Spiteri⁵⁹ , L. Splendori¹⁰⁴ , M. Spousta¹³⁶ , E. J. Staats³⁵ , R. Stamen^{63a} , E. Stanecka⁸⁷ , W. Stanek-Maslouska⁴⁸ , M. V. Stange⁵⁰ , B. Stanislaus^{18a} , M. M. Stanitzki⁴⁸ , B. Stapf⁴⁸ , E. A. Starchenko³⁸ , G. H. Stark¹³⁹ , J. Stark⁹¹ , P. Staroba¹³⁴ , P. Starovoitov^{88b} , R. Staszewski⁸⁷ , G. Stavropoulos⁴⁶ , A. Steff³⁷ , P. Steinberg³⁰ , B. Stelzer^{148,162a} , H. J. Stelzer¹³² , O. Stelzer-Chilton^{162a} , H. Stenzel⁵⁸ , T. J. Stevenson¹⁵² , G. A. Stewart³⁷ , J. R. Stewart¹²⁴ , M. C. Stockton³⁷ , G. Stoicea^{28b} , M. Stolarski^{133a} , S. Stonjek¹¹² , A. Straessner⁵⁰ , J. Strandberg¹⁵⁰ , S. Strandberg^{47a,47b} , M. Stratmann¹⁷⁷ , M. Strauss¹²³ , T. Streblor¹⁰⁴ , P. Strizenec^{29b} , R. Ströhmer¹⁷² , D. M. Strom¹²⁶ , R. Stroynowski⁴⁵ , A. Strubig^{47a,47b} , S. A. Stucci³⁰ , B. Stugu¹⁷ , J. Stupak¹²³ , N. A. Styles⁴⁸ , D. Su¹⁴⁹ , S. Su⁶² , X. Su⁶² , D. Suchy^{29a} , K. Sugizaki¹³¹ , V. V. Sulin³⁸ , M. J. Sullivan⁹⁴ , D. M. S. Sultan¹²⁹ , L. Sultanaliev³⁸ , S. Sultansoy^{3b} , S. Sun¹⁷⁶ , W. Sun¹⁴ , O. Sunneborn Gudnadottir¹⁶⁷ , N. Sur¹⁰⁰ , M. R. Sutton¹⁵² , H. Suzuki¹⁶³ , M. Svatos¹³⁴ , P. N. Swallow³³ , M. Swiatkowski^{162a} , T. Swirski¹⁷² , I. Sykora^{29a} , M. Sykora¹³⁶ , T. Sykora¹³⁶ , D. Ta¹⁰² , K. Tackmann^{48,z} , A. Taffard¹⁶⁵ , R. Tafirout^{162a} , Y. Takubo⁸⁴ , M. Talby¹⁰⁴ , A. A. Talyshv³⁸ , K. C. Tam^{64b} , N. M. Tamir¹⁵⁷ , A. Tanaka¹⁵⁹ , J. Tanaka¹⁵⁹ , R. Tanaka⁶⁶ , M. Tanasini¹⁵¹ , Z. Tao¹⁷⁰ , S. Tapia Araya^{140f} , S. Tapprogge¹⁰² , A. Tarek Abouelfadl Mohamed¹⁰⁹ , S. Tarem¹⁵⁶ , K. Tariq¹⁴ , G. Tarna^{28b} , G. F. Tartarelli^{71a} , M. J. Tartarin⁹¹ , P. Tas¹³⁶ , M. Tasevsky¹³⁴ , E. Tassi^{44b,44a} , A. C. Tate¹⁶⁸ , G. Tateno¹⁵⁹ , Y. Tayalati^{36e,ab} , G. N. Taylor¹⁰⁷ , W. Taylor^{162b} , A. S. Tegetmeier⁹¹ , P. Teixeira-Dias⁹⁷ , J. J. Teoh¹⁶¹ , K. Terashi¹⁵⁹ , J. Terron¹⁰¹ , S. Terzo¹³ , M. Testa⁵³ , R. J. Teuscher^{161,ac} , A. Thaler⁷⁹ , O. Theiner⁵⁶ , T. Theveniaux-Pelzer¹⁰⁴ , D. W. Thomas⁹⁷ , J. P. Thomas²¹ , E. A. Thompson^{18a} , P. D. Thompson²¹ , E. Thomson¹³¹ , R. E. Thornberry⁴⁵ , C. Tian⁶² , Y. Tian⁵⁶ , V. Tikhomirov⁸² , Yu. A. Tikhonov³⁹ , S. Timoshenko³⁸ , D. Timoshyn¹³⁶ , E. X. L. Ting¹ , P. Tipton¹⁷⁸ , A. Tishelman-Charny³⁰ , K. Todome¹⁴¹ , S. Todorova-Nova¹³⁶ , S. Todt⁵⁰ , L. Toffolin^{69a,69c} , M. Togawa⁸⁴ , J. Tojo⁹⁰ , S. Tokár^{29a} , O. Toldaiev⁶⁸ , G. Tolkachev¹⁰⁴ , M. Tomoto^{84,113} , L. Tompkins^{149,o} , E. Torrence¹²⁶ , H. Torres⁹¹ , E. Torró Pastor¹⁶⁹ , M. Toscani³¹ , C. Toscir⁴⁰ , M. Tost¹¹ , D. R. Tovey¹⁴⁵ , T. Trefzger¹⁷² , P. M. Tricarico¹³ , A. Tricoli³⁰ , I. M. Trigger^{162a} , S. Trincaz-Duvold¹³⁰ , D. A. Trischuk²⁷ , A. Tropina³⁹ , L. Truong^{34c} , M. Trzebinski⁸⁷ , A. Trzupek⁸⁷ , F. Tsai¹⁵¹ , M. Tsai¹⁰⁸ , A. Tsiamis¹⁵⁸ , P. V. Tsiarashka³⁹ , S. Tsigaridas^{162a} , A. Tsirigotis^{158,v} , V. Tsiskaridze¹⁶¹ , E. G. Tskhadadze^{155a} , M. Tsopoulou¹⁵⁸ , Y. Tsujikawa⁸⁹ , I. I. Tsukerman³⁸ , V. Tsulaia^{18a} , S. Tsuno⁸⁴ , K. Tsur¹²¹ , D. Tsybychev¹⁵¹ , Y. Tu^{64b} , A. Tudorache^{28b} , V. Tudorache^{28b} , S. B. Tuncay¹²⁹ , S.

E. W. Varnes⁷, C. Varni^{18b}, D. Varouchas⁶⁶, L. Varriale¹⁶⁹, K. E. Varvell¹⁵³, M. E. Vasile^{28b}, L. Vaslin⁸⁴, M. D. Vassilev¹⁴⁹, A. Vasyukov³⁹, L. M. Vaughan¹²⁴, R. Vavricka¹³⁶, T. VazquezSchroeder¹³, J. Veatch³², V. Vecchio¹⁰³, M. J. Veen¹⁰⁵, I. Veliscek³⁰, I. Velkovska⁹⁵, L. M. Veloce¹⁶¹, F. Veloso^{133a,133c}, S. Veneziano^{75a}, A. Ventura^{70a,70b}, S. Ventura Gonzalez¹³⁸, A. Verbytskyi¹¹², M. Verducci^{74a,74b}, C. Vergis⁹⁶, M. Verissimo De Araujo^{83b}, W. Verkerke¹¹⁷, J. C. Vermeulen¹¹⁷, C. Vernieri¹⁴⁹, M. Vessella¹⁶⁵, M. C. Vetterli^{148,ai}, A. Vgenopoulos¹⁰², N. Viaux Maira^{140f}, T. Vickey¹⁴⁵, O. E. Vickey Boeriu¹⁴⁵, G. H. A. Viehhauser¹²⁹, L. Vigani^{63b}, M. Vigi¹¹², M. Villa^{24b,24a}, M. Villaplana Perez¹⁶⁹, E. M. Villhauer⁴⁰, E. Vilucchi⁵³, M. Vincent¹⁶⁹, M. G. Vinciter³⁵, A. Visibile¹¹⁷, C. Vittori³⁷, I. Vivarelli^{24b,24a}, E. Voevodina¹¹², F. Vogel¹¹¹, J. C. Voigt⁵⁰, P.
Vokac¹³⁵, Yu. Volkotrub^{86b}, E. Von Toerne²⁵, B. Vormwald³⁷, K. Vorobev⁵¹, M. Vos¹⁶⁹, K. Voss¹⁴⁷, M. Vozak³⁷, L. Vozdecky¹²³, N. Vranjes¹⁶, M. Vranjes Milosavljevic¹⁶, M. Vreeswijk¹¹⁷, N. K. Vu^{144b,144a}, R. Vuillermet³⁷, O. Vujanovic¹⁰², I. Vukotic⁴⁰, I. K. Vyas³⁵, J. F. Wack³³, S. Wada¹⁶³, C. Wagner¹⁴⁹, J. M. Wagner^{18a}, W. Wagner¹⁷⁷, S. Wahdan¹⁷⁷, H. Wahlberg⁹², C. H. Waits¹²³, J. Walder¹³⁷, R. Walker¹¹¹, K. Walkingshaw Pass⁵⁹, W. Walkowiak¹⁴⁷, A. Wall¹³¹, E. J. Wallin¹⁰⁰, T. Wamorkar^{18a}, A. Wang⁶², A. Z. Wang¹³⁹, C. Wang¹⁰², C. Wang¹¹, H. Wang^{18a}, J. Wang^{64c}, P. Wang¹⁰³, P. Wang⁹⁸, R. Wang⁶¹, R. Wang⁶, S. M. Wang¹⁵⁴, S. Wang¹⁴, T. Wang⁶², T. Wang⁶², W. T. Wang⁸⁰, W. Wang¹⁴, X. Wang¹⁶⁸, X. Wang^{144a}, X. Wang⁴⁸, Y.
Wang^{114a}, Y. Wang⁶², Z. Wang¹⁰⁸, Z. Wang^{144b}, Z. Wang¹⁰⁸, C. Wanotayaroj⁸⁴, A. Warburton¹⁰⁶, A. L. Warnerbring¹⁴⁷, N. Warrack⁵⁹, S. Waterhouse⁹⁷, A. T. Watson²¹, H. Watson⁵², M. F. Watson²¹, E. Watton⁵⁹, G. Watts¹⁴², B. M. Waugh⁹⁸, J. M. Webb⁵⁴, C. Weber³⁰, H. A. Weber¹⁹, M. S. Weber²⁰, S. M. Weber^{63a}, C. Wei⁶², Y. Wei⁵⁴, A. R. Weidberg¹²⁹, E. J. Weik¹²⁰, J. Weingarten⁴⁹, C. Weiser⁵⁴, C. J. Wells⁴⁸, T. Wenaus³⁰, B. Wendland⁴⁹, T. Wengler³⁷, N. S. Wenke¹¹², N. Wermes²⁵, M. Wessels^{63a}, A. M. Wharton⁹³, A. S. White⁶¹, A. White⁸, M. J. White¹, D. Whiteson¹⁶⁵, L. Wickremasinghe¹²⁷, W. Wiedenmann¹⁷⁶, M. Wielers¹³⁷, R. Wierda¹⁵⁰, C. Wiglesworth⁴³, H. G. Wilkens³⁷, J. J. H. Wilkinson³³, D. M. Williams⁴², H. H. Williams¹³¹, S. Williams³³, S. Willocq¹⁰⁵, B. J. Wilson¹⁰³, D. J.
Wilson¹⁰³, P. J. Windischhofer⁴⁰, F. I. Winkel³¹, F. Winklmeier¹²⁶, B. T. Winter⁵⁴, M. Wittgen¹⁴⁹, M. Wobisch⁹⁹, T. Wojtkowski⁶⁰, Z. Wolffs¹¹⁷, J. Wollrath³⁷, M. W. Wolter⁸⁷, H. Wolters^{133a,133c}, M. C. Wong¹³⁹, E. L. Woodward⁴², S. D. Worm⁴⁸, B. K. Wosiek⁸⁷, K. W. Woźniak⁸⁷, S. Wozniowski⁵⁵, K. Wraight⁵⁹, C. Wu¹⁶¹, C. Wu²¹, J. Wu¹⁵⁹, M. Wu^{114b}, M. Wu¹¹⁶, S. L. Wu¹⁷⁶, S. Wu¹⁴, X. Wu⁶², Y. Wu⁶², Z. Wu⁴, J. Wuerzinger¹¹², T. R. Wyatt¹⁰³, B. M. Wynne⁵², S. Xella⁴³, L. Xia^{114a}, M. Xia¹⁵, M. Xie⁶², A. Xiong¹²⁶, J. Xiong^{18a}, D. Xu¹⁴, H. Xu⁶², L. Xu⁶², R. Xu¹³¹, T. Xu¹⁰⁸, Y. Xu¹⁴², Z. Xu⁵², Z. Xu^{114a}, B. Yabsley¹⁵³, S. Yacoob^{34a}, Y. Yamaguchi⁸⁴, E. Yamashita¹⁵⁹, H. Yamauchi¹⁶³, T.
Yamazaki^{18a}, Y. Yamazaki⁸⁵, S. Yan⁵⁹, Z. Yan¹⁰⁵, H. J. Yang^{144a,144b}, H. T. Yang⁶², S. Yang⁶², T. Yang^{64c}, X. Yang³⁷, X. Yang¹⁴, Y. Yang¹⁵⁹, Y. Yang⁶², W-M. Yao^{18a}, C. L. Yardley¹⁵², J. Ye¹⁴, S. Ye³⁰, X. Ye⁶², Y. Yeh⁹⁸, I. Yeletsikh³⁹, B. Yeo^{18b}, M. R. Yexley⁹⁸, T. P. Yildirim¹²⁹, P. Yin⁴², K. Yorita¹⁷⁴, C. J. S. Young³⁷, C. Young¹⁴⁹, N. D. Young¹²⁶, Y. Yu⁶², J. Yuan^{14,114c}, M. Yuan¹⁰⁸, R. Yuan^{144b,144a}, L. Yue⁹⁸, M. Zaazoua⁶², B. Zabinski⁸⁷, I. Zahir^{36a}, A. Zaid^{57b,57a}, Z. K. Zak⁸⁷, T. Zakareishvili¹⁶⁹, S. Zambito⁵⁶, J. A. Zamora Saa^{140d}, J. Zang¹⁵⁹, R. Zanzottera^{71a,71b}, O. Zaplatilek¹³⁵, C. Zeitnitz¹⁷⁷, H. Zeng¹⁴, J. C. Zeng¹⁶⁸, D. T. Zenger Jr²⁷, O. Zenin³⁸, T. Ženiš^{29a}, S. Zenz⁹⁶, D. Zerwas⁶⁶, M.
Zhai^{14,114c}, D. F. Zhang¹⁴⁵, G. Zhang¹⁴, J. Zhang^{143a}, J. Zhang⁶, K. Zhang^{14,114c}, L. Zhang⁶², L. Zhang^{114a}, P. Zhang^{14,114c}, R. Zhang^{114a}, S. Zhang⁹¹, T. Zhang¹⁵⁹, Y. Zhang¹⁴², Y. Zhang⁹⁸, Y. Zhang⁶², Y. Zhang^{114a}, Z. Zhang^{143a}, Z. Zhang⁶⁶, H. Zhao¹⁴², T. Zhao^{143a}, Y. Zhao³⁵, Z. Zhao⁶², Z. Zhao⁶², A. Zhemchugov³⁹, J. Zheng^{114a}, K. Zheng¹⁶⁸, X. Zheng⁶², Z. Zheng¹⁴⁹, D. Zhong¹⁶⁸, B. Zhou¹⁰⁸, H. Zhou⁷, N. Zhou^{144a}, Y. Zhou¹⁵, Y. Zhou^{114a}, Y. Zhou⁷, C. G. Zhu¹⁴⁹, J. Zhu¹⁰⁸, X. Zhu^{144b}, Y. Zhu^{144a}, Y. Zhu⁶², X. Zhuang¹⁴, K. Zhukov⁶⁸, N. I. Zimine³⁹

- ⁷ Department of Physics, University of Arizona, Tucson, AZ, USA
- ⁸ Department of Physics, University of Texas at Arlington, Arlington, TX, USA
- ⁹ Physics Department, National and Kapodistrian University of Athens, Athens, Greece
- ¹⁰ Physics Department, National Technical University of Athens, Zografou, Greece
- ¹¹ Department of Physics, University of Texas at Austin, Austin, TX, USA
- ¹² Institute of Physics, Azerbaijan Academy of Sciences, Baku, Azerbaijan
- ¹³ Institut de Física d'Altes Energies (IFAE), Barcelona Institute of Science and Technology, Barcelona, Spain
- ¹⁴ Institute of High Energy Physics, Chinese Academy of Sciences, Beijing, China
- ¹⁵ Physics Department, Tsinghua University, Beijing, China
- ¹⁶ Institute of Physics, University of Belgrade, Belgrade, Serbia
- ¹⁷ Department for Physics and Technology, University of Bergen, Bergen, Norway
- ¹⁸ (a) Physics Division, Lawrence Berkeley National Laboratory, Berkeley, CA, USA; (b) University of California, Berkeley, CA, USA
- ¹⁹ Institut für Physik, Humboldt Universität zu Berlin, Berlin, Germany
- ²⁰ Albert Einstein Center for Fundamental Physics and Laboratory for High Energy Physics, University of Bern, Bern, Switzerland
- ²¹ School of Physics and Astronomy, University of Birmingham, Birmingham, UK
- ²² (a) Department of Physics, Bogazici University, Istanbul, Türkiye; (b) Department of Physics Engineering, Gaziantep University, Gaziantep, Türkiye; (c) Department of Physics, Istanbul University, Istanbul, Türkiye
- ²³ (a) Facultad de Ciencias y Centro de Investigaciones, Universidad Antonio Nariño, Bogotá, Colombia; (b) Departamento de Física, Universidad Nacional de Colombia, Bogotá, Colombia
- ²⁴ (a) Dipartimento di Fisica e Astronomia A. Righi, Università di Bologna, Bologna, Italy; (b) INFN Sezione di Bologna, Bologna, Italy
- ²⁵ Physikalisches Institut, Universität Bonn, Bonn, Germany
- ²⁶ Department of Physics, Boston University, Boston, MA, USA
- ²⁷ Department of Physics, Brandeis University, Waltham, MA, USA
- ²⁸ (a) Transilvania University of Brasov, Brasov, Romania; (b) Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest, Romania; (c) Department of Physics, Alexandru Ioan Cuza University of Iasi, Iasi, Romania; (d) National Institute for Research and Development of Isotopic and Molecular Technologies, Physics Department, Cluj-Napoca, Romania; (e) National University of Science and Technology Politehnica, Bucharest, Romania; (f) West University in Timisoara, Timisoara, Romania; (g) Faculty of Physics, University of Bucharest, Bucharest, Romania
- ²⁹ (a) Faculty of Mathematics, Physics and Informatics, Comenius University, Bratislava, Slovak Republic; (b) Department of Subnuclear Physics, Institute of Experimental Physics of the Slovak Academy of Sciences, Kosice, Slovak Republic
- ³⁰ Physics Department, Brookhaven National Laboratory, Upton, NY, USA
- ³¹ Facultad de Ciencias Exactas y Naturales, Departamento de Física, y CONICET, Instituto de Física de Buenos Aires (IFIBA), Universidad de Buenos Aires, Buenos Aires, Argentina
- ³² California State University, Long Beach, CA, USA
- ³³ Cavendish Laboratory, University of Cambridge, Cambridge, UK
- ³⁴ (a) Department of Physics, University of Cape Town, Cape Town, South Africa; (b) iThemba Labs, Western Cape, South Africa; (c) Department of Mechanical Engineering Science, University of Johannesburg, Johannesburg, South Africa; (d) National Institute of Physics, University of the Philippines Diliman (Philippines), Quezon City, Philippines; (e) University of South Africa, Department of Physics, Pretoria, South Africa; (f) University of Zululand, KwaDlangezwa, South Africa; (g) School of Physics, University of the Witwatersrand, Johannesburg, South Africa
- ³⁵ Department of Physics, Carleton University, Ottawa, ON, Canada
- ³⁶ (a) Faculté des Sciences Ain Chock, Université Hassan II de Casablanca, Casablanca, Morocco; (b) Faculté des Sciences, Université Ibn-Tofail, Kénitra, Morocco; (c) Faculté des Sciences Semlalia, LPHEA-Marrakech, Université Cadi Ayyad, Marrakech, Morocco; (d) LPMR, Faculté des Sciences, Université Mohamed Premier, Oujda, Morocco; (e) Faculté des sciences, Université Mohammed V, Rabat, Morocco; (f) Institute of Applied Physics, Mohammed VI Polytechnic University, Ben Guerir, Morocco
- ³⁷ CERN, Geneva, Switzerland
- ³⁸ Affiliated with an Institute Formerly Covered by a Cooperation Agreement with CERN, Geneva, Switzerland
- ³⁹ Affiliated with an International Laboratory Covered by a Cooperation Agreement with CERN, Geneva, Switzerland

- ⁴⁰ Enrico Fermi Institute, University of Chicago, Chicago, IL, USA
- ⁴¹ LPC, CNRS/IN2P3, Université Clermont Auvergne, Clermont-Ferrand, France
- ⁴² Nevis Laboratory, Columbia University, Irvington, NY, USA
- ⁴³ Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark
- ⁴⁴ ^(a)Dipartimento di Fisica, Università della Calabria, Rende, Italy; ^(b)Laboratori Nazionali di Frascati, INFN Gruppo Collegato di Cosenza, Cosenza, Italy
- ⁴⁵ Physics Department, Southern Methodist University, Dallas, TX, USA
- ⁴⁶ National Centre for Scientific Research “Demokritos”, Agia Paraskevi, Greece
- ⁴⁷ ^(a)Department of Physics, Stockholm University, Stockholm, Sweden; ^(b)Oskar Klein Centre, Stockholm, Sweden
- ⁴⁸ Deutsches Elektronen-Synchrotron DESY, Hamburg and Zeuthen, Germany
- ⁴⁹ Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany
- ⁵⁰ Institut für Kern- und Teilchenphysik, Technische Universität Dresden, Dresden, Germany
- ⁵¹ Department of Physics, Duke University, Durham, NC, USA
- ⁵² SUPA-School of Physics and Astronomy, University of Edinburgh, Edinburgh, UK
- ⁵³ INFN e Laboratori Nazionali di Frascati, Frascati, Italy
- ⁵⁴ Physikalisches Institut, Albert-Ludwigs-Universität Freiburg, Freiburg, Germany
- ⁵⁵ II. Physikalisches Institut, Georg-August-Universität Göttingen, Göttingen, Germany
- ⁵⁶ Département de Physique Nucléaire et Corpusculaire, Université de Genève, Geneva, Switzerland
- ⁵⁷ ^(a)Dipartimento di Fisica, Università di Genova, Genoa, Italy; ^(b)INFN Sezione di Genova, Genoa, Italy
- ⁵⁸ II. Physikalisches Institut, Justus-Liebig-Universität Giessen, Giessen, Germany
- ⁵⁹ SUPA-School of Physics and Astronomy, University of Glasgow, Glasgow, UK
- ⁶⁰ LPSC, Université Grenoble Alpes, CNRS/IN2P3, Grenoble INP, Grenoble, France
- ⁶¹ Laboratory for Particle Physics and Cosmology, Harvard University, Cambridge, MA, USA
- ⁶² Department of Modern Physics and State Key Laboratory of Particle Detection and Electronics, University of Science and Technology of China, Hefei, China
- ⁶³ ^(a)Kirchhoff-Institut für Physik, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany; ^(b)Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany
- ⁶⁴ ^(a)Department of Physics, Chinese University of Hong Kong, Shatin, N.T., Hong Kong, China; ^(b)Department of Physics, University of Hong Kong, Hong Kong, China; ^(c)Department of Physics and Institute for Advanced Study, Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong, China
- ⁶⁵ Department of Physics, National Tsing Hua University, Hsinchu, Taiwan
- ⁶⁶ IJCLab, CNRS/IN2P3, Université Paris-Saclay, 91405 Orsay, France
- ⁶⁷ Centro Nacional de Microelectrónica (IMB-CNM-CSIC), Barcelona, Spain
- ⁶⁸ Department of Physics, Indiana University, Bloomington, IN, USA
- ⁶⁹ ^(a)INFN Gruppo Collegato di Udine, Sezione di Trieste, Udine, Italy; ^(b)ICTP, Trieste, Italy; ^(c)Dipartimento Politecnico di Ingegneria e Architettura, Università di Udine, Udine, Italy
- ⁷⁰ ^(a)INFN Sezione di Lecce, Lecce, Italy; ^(b)Dipartimento di Matematica e Fisica, Università del Salento, Lecce, Italy
- ⁷¹ ^(a)INFN Sezione di Milano, Milan, Italy; ^(b)Dipartimento di Fisica, Università di Milano, Milan, Italy
- ⁷² ^(a)INFN Sezione di Napoli, Naples, Italy; ^(b)Dipartimento di Fisica, Università di Napoli, Naples, Italy
- ⁷³ ^(a)INFN Sezione di Pavia, Pavia, Italy; ^(b)Dipartimento di Fisica, Università di Pavia, Pavia, Italy
- ⁷⁴ ^(a)INFN Sezione di Pisa, Pisa, Italy; ^(b)Dipartimento di Fisica E. Fermi, Università di Pisa, Pisa, Italy
- ⁷⁵ ^(a)INFN Sezione di Roma, Rome, Italy; ^(b)Dipartimento di Fisica, Sapienza Università di Roma, Rome, Italy
- ⁷⁶ ^(a)INFN Sezione di Roma Tor Vergata, Rome, Italy; ^(b)Dipartimento di Fisica, Università di Roma Tor Vergata, Rome, Italy
- ⁷⁷ ^(a)INFN Sezione di Roma Tre, Rome, Italy; ^(b)Dipartimento di Matematica e Fisica, Università Roma Tre, Rome, Italy
- ⁷⁸ ^(a)INFN-TIFPA, Povo, Italy; ^(b)Università degli Studi di Trento, Trento, Italy
- ⁷⁹ Universität Innsbruck, Department of Astro and Particle Physics, Innsbruck, Austria
- ⁸⁰ University of Iowa, Iowa City, IA, USA
- ⁸¹ Department of Physics and Astronomy, Iowa State University, Ames, IA, USA
- ⁸² Istinye University, Sariyer, Istanbul, Türkiye
- ⁸³ ^(a)Departamento de Engenharia Elétrica, Universidade Federal de Juiz de Fora (UFJF), Juiz de Fora, Brazil; ^(b)Universidade Federal do Rio De Janeiro COPPE/EE/IF, Rio de Janeiro, Brazil; ^(c)Instituto de Física,

- Universidade de São Paulo, São Paulo, Brazil; ^(d)Rio de Janeiro State University, Rio de Janeiro, Brazil; ^(e)Federal University of Bahia, Bahia, Brazil
- ⁸⁴ KEK, High Energy Accelerator Research Organization, Tsukuba, Japan
- ⁸⁵ Graduate School of Science, Kobe University, Kobe, Japan
- ⁸⁶ ^(a)AGH University of Krakow, Faculty of Physics and Applied Computer Science, Kraków, Poland; ^(b)Marian Smoluchowski Institute of Physics, Jagiellonian University, Kraków, Poland
- ⁸⁷ Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland
- ⁸⁸ ^(a)Khalifa University of Science and Technology, Abu Dhabi, United Arab Emirates; ^(b)University of Sharjah, Sharjah, United Arab Emirates
- ⁸⁹ Faculty of Science, Kyoto University, Kyoto, Japan
- ⁹⁰ Research Center for Advanced Particle Physics and Department of Physics, Kyushu University, Fukuoka, Japan
- ⁹¹ L2IT, CNRS/IN2P3, UPS, Université de Toulouse, Toulouse, France
- ⁹² Instituto de Física La Plata, Universidad Nacional de La Plata and CONICET, La Plata, Argentina
- ⁹³ Physics Department, Lancaster University, Lancaster, UK
- ⁹⁴ Oliver Lodge Laboratory, University of Liverpool, Liverpool, UK
- ⁹⁵ Department of Experimental Particle Physics, Jožef Stefan Institute and Department of Physics, University of Ljubljana, Ljubljana, Slovenia
- ⁹⁶ Department of Physics and Astronomy, Queen Mary University of London, London, UK
- ⁹⁷ Department of Physics, Royal Holloway University of London, Egham, UK
- ⁹⁸ Department of Physics and Astronomy, University College London, London, UK
- ⁹⁹ Louisiana Tech University, Ruston, LA, USA
- ¹⁰⁰ Fysiska institutionen, Lunds universitet, Lund, Sweden
- ¹⁰¹ Departamento de Física Teórica C-15 and CIAFF, Universidad Autónoma de Madrid, Madrid, Spain
- ¹⁰² Institut für Physik, Universität Mainz, Mainz, Germany
- ¹⁰³ School of Physics and Astronomy, University of Manchester, Manchester, UK
- ¹⁰⁴ CPPM, CNRS/IN2P3, Aix-Marseille Université, Marseille, France
- ¹⁰⁵ Department of Physics, University of Massachusetts, Amherst, MA, USA
- ¹⁰⁶ Department of Physics, McGill University, Montreal, QC, Canada
- ¹⁰⁷ School of Physics, University of Melbourne, Victoria, Australia
- ¹⁰⁸ Department of Physics, University of Michigan, Ann Arbor, MI, USA
- ¹⁰⁹ Department of Physics and Astronomy, Michigan State University, East Lansing, MI, USA
- ¹¹⁰ Group of Particle Physics, University of Montreal, Montreal, QC, Canada
- ¹¹¹ Fakultät für Physik, Ludwig-Maximilians-Universität München, Munich, Germany
- ¹¹² Max-Planck-Institut für Physik (Werner-Heisenberg-Institut), Munich, Germany
- ¹¹³ Graduate School of Science and Kobayashi-Maskawa Institute, Nagoya University, Nagoya, Japan
- ¹¹⁴ ^(a)Department of Physics, Nanjing University, Nanjing, China; ^(b)School of Science, Shenzhen Campus of Sun Yat-sen University, Guangzhou, China; ^(c)University of Chinese Academy of Science (UCAS), Beijing, China
- ¹¹⁵ Department of Physics and Astronomy, University of New Mexico, Albuquerque, NM, USA
- ¹¹⁶ Institute for Mathematics, Astrophysics and Particle Physics, Radboud University/Nikhef, Nijmegen, Netherlands
- ¹¹⁷ Nikhef National Institute for Subatomic Physics and University of Amsterdam, Amsterdam, Netherlands
- ¹¹⁸ Department of Physics, Northern Illinois University, DeKalb, IL, USA
- ¹¹⁹ ^(a)New York University Abu Dhabi, Abu Dhabi, United Arab Emirates; ^(b)United Arab Emirates University, Al Ain, United Arab Emirates
- ¹²⁰ Department of Physics, New York University, New York, NY, USA
- ¹²¹ Ochanomizu University, Otsuka, Bunkyo-ku, Tokyo, Japan
- ¹²² Ohio State University, Columbus, OH, USA
- ¹²³ Homer L. Dodge Department of Physics and Astronomy, University of Oklahoma, Norman, OK, USA
- ¹²⁴ Department of Physics, Oklahoma State University, Stillwater, OK, USA
- ¹²⁵ Palacký University, Joint Laboratory of Optics, Olomouc, Czech Republic
- ¹²⁶ Institute for Fundamental Science, University of Oregon, Eugene, OR, USA
- ¹²⁷ Graduate School of Science, University of Osaka, Osaka, Japan
- ¹²⁸ Department of Physics, University of Oslo, Oslo, Norway
- ¹²⁹ Department of Physics, Oxford University, Oxford, UK

- 130 LPNHE, CNRS/IN2P3, Sorbonne Université, Université Paris Cité, Paris, France
- 131 Department of Physics, University of Pennsylvania, Philadelphia, PA, USA
- 132 Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh, PA, USA
- 133 (a) Laboratório de Instrumentação e Física Experimental de Partículas - LIP, Lisbon, Portugal; (b) Departamento de Física, Faculdade de Ciências, Universidade de Lisboa, Lisbon, Portugal; (c) Departamento de Física, Universidade de Coimbra, Coimbra, Portugal; (d) Centro de Física Nuclear da Universidade de Lisboa, Lisbon, Portugal; (e) Departamento de Física, Escola de Ciências, Universidade do Minho, Braga, Portugal; (f) Departamento de Física Teórica y del Cosmos, Universidad de Granada, Granada, Spain; (g) Departamento de Física, Instituto Superior Técnico, Universidade de Lisboa, Lisbon, Portugal
- 134 Institute of Physics of the Czech Academy of Sciences, Prague, Czech Republic
- 135 Czech Technical University in Prague, Prague, Czech Republic
- 136 Faculty of Mathematics and Physics, Charles University, Prague, Czech Republic
- 137 Particle Physics Department, Rutherford Appleton Laboratory, Didcot, UK
- 138 IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France
- 139 Santa Cruz Institute for Particle Physics, University of California Santa Cruz, Santa Cruz, CA, USA
- 140 (a) Departamento de Física, Pontificia Universidad Católica de Chile, Santiago, Chile; (b) Millennium Institute for Subatomic Physics at High Energy Frontier (SAPHIR), Santiago, Chile; (c) Instituto de Investigación Multidisciplinario en Ciencia y Tecnología y Departamento de Física, Universidad de La Serena, La Serena, Chile; (d) Universidad Andres Bello, Department of Physics, Santiago, Chile; (e) Instituto de Alta Investigación, Universidad de Tarapacá, Arica, Chile; (f) Departamento de Física, Universidad Técnica Federico Santa María, Valparaíso, Chile
- 141 Department of Physics, Institute of Science, Tokyo, Japan
- 142 Department of Physics, University of Washington, Seattle, WA, USA
- 143 (a) Institute of Frontier and Interdisciplinary Science and Key Laboratory of Particle Physics and Particle Irradiation (MOE), Shandong University, Qingdao, China; (b) School of Physics, Zhengzhou University, Zhengzhou, China
- 144 (a) State Key Laboratory of Dark Matter Physics, School of Physics and Astronomy, Shanghai Jiao Tong University, Key Laboratory for Particle Astrophysics and Cosmology (MOE), SKLPPC, Shanghai, China; (b) State Key Laboratory of Dark Matter Physics, Tsung-Dao Lee Institute, Shanghai Jiao Tong University, Shanghai, China
- 145 Department of Physics and Astronomy, University of Sheffield, Sheffield, UK
- 146 Department of Physics, Shinshu University, Nagano, Japan
- 147 Department Physik, Universität Siegen, Siegen, Germany
- 148 Department of Physics, Simon Fraser University, Burnaby, BC, Canada
- 149 SLAC National Accelerator Laboratory, Stanford, CA, USA
- 150 Department of Physics, Royal Institute of Technology, Stockholm, Sweden
- 151 Departments of Physics and Astronomy, Stony Brook University, Stony Brook, NY, USA
- 152 Department of Physics and Astronomy, University of Sussex, Brighton, UK
- 153 School of Physics, University of Sydney, Sydney, Australia
- 154 Institute of Physics, Academia Sinica, Taipei, Taiwan
- 155 (a) E. Andronikashvili Institute of Physics, Iv. Javakhishvili Tbilisi State University, Tbilisi, Georgia; (b) High Energy Physics Institute, Tbilisi State University, Tbilisi, Georgia; (c) University of Georgia, Tbilisi, Georgia
- 156 Department of Physics, Technion, Israel Institute of Technology, Haifa, Israel
- 157 Raymond and Beverly Sackler School of Physics and Astronomy, Tel Aviv University, Tel Aviv, Israel
- 158 Department of Physics, Aristotle University of Thessaloniki, Thessaloníki, Greece
- 159 International Center for Elementary Particle Physics and Department of Physics, University of Tokyo, Tokyo, Japan
- 160 Graduate School of Science and Technology, Tokyo Metropolitan University, Tokyo, Japan
- 161 Department of Physics, University of Toronto, Toronto, ON, Canada
- 162 (a) TRIUMF, Vancouver, BC, Canada; (b) Department of Physics and Astronomy, York University, Toronto, ON, Canada
- 163 Division of Physics and Tomonaga Center for the History of the Universe, Faculty of Pure and Applied Sciences, University of Tsukuba, Tsukuba, Japan
- 164 Department of Physics and Astronomy, Tufts University, Medford, MA, USA
- 165 Department of Physics and Astronomy, University of California Irvine, Irvine, CA, USA
- 166 University of West Attica, Athens, Greece
- 167 Department of Physics and Astronomy, University of Uppsala, Uppsala, Sweden
- 168 Department of Physics, University of Illinois, Urbana, IL, USA

- ¹⁶⁹ Instituto de Física Corpuscular (IFIC), Centro Mixto Universidad de Valencia - CSIC, Valencia, Spain
 - ¹⁷⁰ Department of Physics, University of British Columbia, Vancouver, BC, Canada
 - ¹⁷¹ Department of Physics and Astronomy, University of Victoria, Victoria, BC, Canada
 - ¹⁷² Fakultät für Physik und Astronomie, Julius-Maximilians-Universität Würzburg, Würzburg, Germany
 - ¹⁷³ Department of Physics, University of Warwick, Coventry, UK
 - ¹⁷⁴ Waseda University, Tokyo, Japan
 - ¹⁷⁵ Department of Particle Physics and Astrophysics, Weizmann Institute of Science, Rehovot, Israel
 - ¹⁷⁶ Department of Physics, University of Wisconsin, Madison, WI, USA
 - ¹⁷⁷ Fakultät für Mathematik und Naturwissenschaften, Fachgruppe Physik, Bergische Universität Wuppertal, Wuppertal, Germany
 - ¹⁷⁸ Department of Physics, Yale University, New Haven, CT, USA
 - ¹⁷⁹ Yerevan Physics Institute, Yerevan, Armenia
- ^a Also at Affiliated with an Institute Formerly Covered by a Cooperation Agreement with CERN, Geneva, Switzerland
- ^b Also at An-Najah National University, Nablus, Palestine
- ^c Also at Borough of Manhattan Community College, City University of New York, New York, NY, USA
- ^d Also at Center for Interdisciplinary Research and Innovation (CIRI-AUTH), Thessaloniki, Greece
- ^e Also at Centre of Physics of the Universities of Minho and Porto (CF-UM-UP), Porto, Portugal
- ^f Also at CERN, Geneva, Switzerland
- ^g Also at CMD-AC UNEC Research Center, Azerbaijan State University of Economics (UNEC), Baku, Azerbaijan
- ^h Also at Département de Physique Nucléaire et Corpusculaire, Université de Genève, Geneva, Switzerland
- ⁱ Also at Departament de Física de la Universitat Autònoma de Barcelona, Barcelona, Spain
- ^j Also at Department of Financial and Management Engineering, University of the Aegean, Chios, Greece
- ^k Also at Department of Mathematical Sciences, University of South Africa, Johannesburg, South Africa
- ^l Also at Department of Modern Physics and State Key Laboratory of Particle Detection and Electronics, University of Science and Technology of China, Hefei, China
- ^m Also at Department of Physics, Bolu Abant İzzet Baysal University, Bolu, Türkiye
- ⁿ Also at Department of Physics, King's College London, London, UK
- ^o Also at Department of Physics, Stanford University, Stanford, CA, USA
- ^p Also at Department of Physics, Stellenbosch University, South Africa
- ^q Also at Department of Physics, University of Fribourg, Fribourg, Switzerland
- ^r Also at Department of Physics, University of Thessaly, Volos, Greece
- ^s Also at Department of Physics, Westmont College, Santa Barbara, USA
- ^t Also at Faculty of Physics, Sofia University 'St. Kliment Ohridski', Sofia, Bulgaria
- ^u Also at Faculty of Physics, University of Bucharest, Bucharest, Romania
- ^v Also at Hellenic Open University, Patras, Greece
- ^w Also at Henan University, Henan, China
- ^x Also at Imam Mohammad Ibn Saud Islamic University, Riyadh, Saudi Arabia
- ^y Also at Institutio Catalana de Recerca i Estudis Avancats, ICREA, Barcelona, Spain
- ^z Also at Institut für Experimentalphysik, Universität Hamburg, Hamburg, Germany
- ^{aa} Also at Institute for Nuclear Research and Nuclear Energy (INRNE) of the Bulgarian Academy of Sciences, Sofia, Bulgaria
- ^{ab} Also at Institute of Applied Physics, Mohammed VI Polytechnic University, Ben Guerir, Morocco
- ^{ac} Also at Institute of Particle Physics (IPP), Toronto, Canada
- ^{ad} Also at Institute of Physics and Technology, Mongolian Academy of Sciences, Ulaanbaatar, Mongolia
- ^{ae} Also at Institute of Physics, Azerbaijan Academy of Sciences, Baku, Azerbaijan
- ^{af} Also at Institute of Theoretical Physics, Ilia State University, Tbilisi, Georgia
- ^{ag} Also at National Institute of Physics, University of the Philippines Diliman (Philippines), Quezon City, Philippines
- ^{ah} Also at The Collaborative Innovation Center of Quantum Matter (CICQM), Beijing, China
- ^{ai} Also at TRIUMF, Vancouver, BC, Canada
- ^{aj} Also at Università di Napoli Parthenope, Naples, Italy
- ^{ak} Also at Department of Physics, University of Colorado Boulder, Colorado, USA
- ^{al} Also at University of Sienna, Siena, Italy

^{am} Also at Washington College, Chestertown, MD, USA

^{an} Also at Physics Department, Yeditepe University, Istanbul, Türkiye

* Deceased