

# Current-enhanced excited states in lattice QCD three-point functions

Lorenzo Barca<sup>\*</sup>

*John von Neumann-Institut für Computing (NIC), Deutsches Elektronen-Synchrotron DESY,  
Platanenallee 6, 15738 Zeuthen, Germany*



(Received 2 September 2025; accepted 14 October 2025; published 24 November 2025)

Excited-state contamination remains one of the leading sources of systematic uncertainty in the precise determination of hadron structure observables from lattice QCD. In this letter, we present a general argument, inspired by meson dominance ideas and implemented through the variational method, to identify which excited states are enhanced by the choice of the inserted current and kinematics. The argument is supported by numerical evidence across multiple hadronic channels and provides both a conceptual understanding and practical guidance to account for excited-state effects in hadron three-point function analyses.

DOI: [10.1103/PhysRevD.112.L091503](https://doi.org/10.1103/PhysRevD.112.L091503)

**Introduction.** Lattice quantum chromodynamics (lattice QCD) provides a first-principles framework to compute a wide range of hadronic observables by numerically simulating the strong interactions of quarks and gluons. In particular, hadron three-point correlation functions offer valuable insight into the internal structure of hadrons and their interactions mediated by external currents. These functions encode how a hadron responds to the insertion of a current—probing, for instance, its form factors or charge distributions—while accounting for the full complexity of QCD dynamics. The standard expression for a hadronic three-point function is

$$C_{3pt}(\vec{p}', t; \vec{q}, \tau) = \langle O_H(\vec{p}', t) \mathcal{J}(\vec{q}, \tau) O_H^\dagger(\vec{p}, 0) \rangle, \quad (1)$$

where  $O_H$  and  $\bar{O}_H$  are interpolators that annihilate and create hadrons with the quantum numbers of  $H$ , and  $\mathcal{J}$  is a current operator that mediates the interaction. For simplicity, we consider the case where the same hadron operator appears at both source and sink, and the current is a local bilinear operator. However, the general argument applies more broadly. The spectral decomposition of the three-point function exposes its dependence on the matrix elements of the current between hadronic states. Explicitly, one finds

$$C_{3pt}(\vec{p}', t; \vec{q}, \tau) = Z_H' Z_H^\dagger \langle H' | \mathcal{J} | H \rangle e^{-E_H'(t-\tau)} e^{-E_H \tau} + \sum_f \sum_i Z_f' Z_i^\dagger \langle f' | \mathcal{J} | i \rangle e^{-E_f'(t-\tau)} e^{-E_i \tau}, \quad (2)$$

where the prefactors  $Z$  arise from the overlaps of the interpolating operators with the physical states. The ground-state contribution is shown explicitly, while the double sum accounts for all possible remaining excited initial and/or final states with the same quantum numbers as  $H$ . In the asymptotic regime  $t \gg \tau \gg 0$ , contributions from excited states are exponentially suppressed if  $E_H' < E_f'$  or  $E_H < E_i$ . In this limit, the ground state matrix element can be, in principle, isolated. However, in practice, the signal-to-noise ratio of most hadronic correlators deteriorates exponentially with increasing time separation, complicating the extraction of hadron properties. Consequently, lattice QCD analyses are constrained to intermediate source-sink separations where excited-state contamination remains significant, making its understanding essential. Recently, alternative methods based on the Lanczos algorithm have been proposed to extract matrix elements through an iterative procedure [1].

Notably, long before QCD, current algebra and meson dominance models provided powerful predictions for hadron dynamics. Vector meson dominance (VMD), proposed by Sakurai [2], postulates that the electromagnetic current couples to hadrons predominantly via the  $\rho$ ,  $\omega$ , and  $\phi$  mesons. Based on VMD, quantitative predictions were made for processes such as  $\rho \rightarrow e^+ e^-$  decay [3],  $\omega \rightarrow \pi^0 \gamma$  and  $\phi \rightarrow \eta \gamma$  [4,5], as well as  $e^+ e^- \rightarrow \pi^+ \pi^-$ ,  $\pi^+ \pi^- \pi^0$  and the  $q^2$  dependence of pion form factors [6]; see [7] for a review. Similarly, current algebra and the Weinberg sum rules, together with vector and axial-vector meson dominance [8], yield the empirical relation  $m_{a_1}/m_\rho \approx \sqrt{2}$ , in striking agreement with experiment. The partially conserved axial current (PCAC) relation further explains the strong coupling of axial currents to pions, reflecting their role as pseudo-Goldstone bosons of spontaneously broken chiral symmetry, and underpins soft-pion theorems, pion

<sup>\*</sup>Contact author: [lorenzo.barca@desy.de](mailto:lorenzo.barca@desy.de)

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP<sup>3</sup>.

pole dominance in form factors, and low-energy current-induced pion production [9–12].

Drawing on the intuition from meson dominance models, one can predict which multihadron states are preferentially produced in a given channel, provided the relevant symmetries and kinematics permit it. These *current-enhanced* states can dominate excited-state contamination in lattice QCD correlators, even if their masses exceed those of other states. In this work, we use the variational method to show that these contributions originate from a *volume enhancement* of specific excited states generated by quark-line disconnected diagrams, and that only certain operators can produce them. Combining this with meson dominance arguments yields clear, physically intuitive predictions for which states dominate in a given three-point function. In the nucleon channel and in  $B \rightarrow \pi$  decays, these predictions agree with chiral perturbation theory (ChPT) expectations, and they extend naturally to cases without ChPT guidance. Our framework is general, applicable to any hadron three-point (or higher), and we present numerical evidence confirming these predictions.

*Volume enhancement of current-enhanced states.* To show the current-enhanced excited-state contributions, we employ the variational method with an extended basis of single and multihadron operators that have the quantum numbers of the initial and final operators:  $\mathbb{B}_H = \{O_H, O_{H_1}, O_{H_2}, O_{HM}, O_{H_1M_1}, \dots\}$ . Using this basis, we construct a matrix of two-point correlation functions, which reads

$$C_{2pt}(\vec{p}, t)_{kj} = \langle O_k(\vec{p}, t) O_j^\dagger(\vec{p}, 0) \rangle \quad O_{k,j} \in \mathbb{B}_H. \quad (3)$$

We solve the generalized eigenvalue problem

$$C_{2pt}(\vec{p}, t) V(\vec{p}, t_0) = C_{2pt}(\vec{p}, t_0) \Lambda(\vec{p}, t) V(\vec{p}, t) \quad t > t_0, \quad (4)$$

which yields the matrix of eigenvalues  $\Lambda(\vec{p}, t) = \text{diag}(\lambda^H, \lambda^{H^*}, \lambda^{HM}, \dots)$  and eigenvectors  $V(\vec{p}, t) = (\vec{v}_{O_H}, \vec{v}_{O_{H_1}}, \vec{v}_{O_{H_2}}, \vec{v}_{O_{HM}}, \vec{v}_{O_{H_1M_1}}, \dots)$ . The eigenvectors can be used to diagonalize the system [13,14] and to construct an improved operator  $\tilde{O}_H$  like

$$\tilde{O}_H = \sum_{O_i \in \mathbb{B}_H} v_{O_i}^H O_i. \quad (5)$$

By construction, this operator overlaps only with the state  $H$ , assuming the variational basis spans the full Hilbert space. However, in practice, it is not feasible to include the infinite tower of interpolating operators required to fully reconstruct the spectrum. Due to this limitation, one must work with a truncated basis, and with the right choice of  $t_0$  and  $t$  in Eq. (4), the associated systematic effects are exponentially suppressed by the energy gaps between the omitted states and the resolved ones [15]. Using this diagonalized operator, we can compute the improved three-point functions

$$\tilde{C}_{3pt}(\vec{p}', t; \vec{q}, \tau) = \langle \tilde{O}_H(\vec{p}', t) \mathcal{J}(\vec{q}, \tau) \tilde{O}_H^\dagger(\vec{p}, 0) \rangle. \quad (6)$$

This expression involves a linear combination of three-point correlators built from all operators in the variational basis. Specifically, it requires evaluating correlation functions of the form

$$\langle O_k(\vec{p}', t) \mathcal{J}(\vec{q}, \tau) O_j^\dagger(\vec{p}, 0) \rangle \quad O_{k,j} \in \mathbb{B}_H. \quad (7)$$

Among these are three-point correlators of the type

$$\langle O_H(\vec{p}'_H, t) O_M(\vec{p}'_M, t) \mathcal{J}(\vec{q}, \tau) O_H^\dagger(\vec{p}, 0) \rangle, \quad (8)$$

with  $\vec{p}' = \vec{p}'_H + \vec{p}'_M$ . These terms, as we argue below, can be significantly enhanced for specific choices of current, momenta, and interpolating operators. To illustrate the enhancement mechanism, we rewrite the three-point function in Eq. (8) in terms of the Wick contractions obtained from the quark content of the hadron operators,

$$\begin{aligned} & \langle O_H(\vec{p}'_H, t) O_M(\vec{p}'_M, t) \mathcal{J}(\vec{q}, \tau) O_H^\dagger(\vec{p}, 0) \rangle \\ &= \langle W_C(\vec{p}'_H; \vec{p}'_M; \vec{q}) + W_D(\vec{p}'_H; \vec{p}'_M; \vec{q}) \\ &+ W_{\text{disc}}(\vec{p}'_H; \vec{p}'_M; \vec{q}) \rangle. \end{aligned} \quad (9)$$

In lattice QCD,  $W_C$  denotes the *connected* diagrams,  $W_D$  the *direct* diagrams, and  $W_{\text{disc}}$  the *disconnected* diagrams. Both  $W_D$  and  $W_{\text{disc}}$  are *quark-line disconnected*. Which contributions appear, and with what relative weight, depends on the quark flavor structure of the interpolating operators, the current, and the kinematics. To make these contributions explicit, we write each term in coordinate space, including the appropriate Fourier transforms to project onto the desired momenta,

$$\begin{aligned} W_C(\vec{p}'_H; \vec{p}'_M; \vec{q}) &= \sum_{\vec{x}_H, \vec{x}_M, \vec{z}} e^{-i\vec{p}'_H \cdot \vec{x}_H} e^{-i\vec{p}'_M \cdot \vec{x}_M} e^{i\vec{q} \cdot \vec{z}} \langle O_H(\vec{x}_H, t) \\ &\times O_M(\vec{x}_M, t) \mathcal{J}(\vec{z}, \tau) O_H^\dagger(\vec{x}_0, 0) \rangle_F, \end{aligned} \quad (10)$$

$$\begin{aligned} W_D(\vec{p}'_H; \vec{p}'_M; \vec{q}) &= \sum_{\vec{x}_H, \vec{x}_M, \vec{z}} e^{-i\vec{p}'_H \cdot \vec{x}_H} e^{-i\vec{p}'_M \cdot \vec{x}_M} e^{i\vec{q} \cdot \vec{z}} \langle O_H(\vec{x}_H, t) \\ &\times O_H^\dagger(\vec{x}_0, 0) \rangle_F \langle O_M(\vec{x}_M, t) \mathcal{J}(\vec{z}, \tau) \rangle_F, \end{aligned} \quad (11)$$

$$\begin{aligned} W_{\text{disc}}(\vec{p}'_H; \vec{p}'_M; \vec{q}) &= \sum_{\vec{x}_H, \vec{x}_M, \vec{z}} e^{-i\vec{p}'_H \cdot \vec{x}_H} e^{-i\vec{p}'_M \cdot \vec{x}_M} e^{i\vec{q} \cdot \vec{z}} \langle O_H(\vec{x}_H, t) \\ &\times O_H^\dagger(\vec{x}_0, 0) \rangle_F \langle O_M(\vec{x}_M, t) \rangle_F \langle \mathcal{J}(\vec{z}, \tau) \rangle_F. \end{aligned} \quad (12)$$

Here,  $\langle \cdot \rangle_F$  denotes the Wick contractions, expressed as traces over products of quark propagators. Examples of the connected and direct diagrams are shown in Fig. S1 in Supplemental Material [16], for the case of a baryon-meson

two-hadron operator at the sink and a baryon interpolator at the source.

A central observation of this work is that, when the meson momentum matches the current momentum transfer,  $\vec{p}'_M = \vec{q}$ , the amplitude of the quark-line disconnected contributions—such as those in Eqs. (11)–(12)—is significantly larger than the quark-line connected contribution in Eq. (10). In the noninteracting limit, such contributions are expected to be enhanced by the spatial volume ( $L^3/a^3$ ), since translational invariance can be used for the current-meson two-point functions. Conversely, multiparticle overlaps are suppressed with the inverse spatial volume, and in Eq. (6) this suppression—carried by the eigenvector components—is compensated by the volume enhancement of quark-line disconnected diagrams. In several nucleon channels—axial-vector, vector, and scalar—we find that at  $\tau = t$  the quark-line disconnected contributions are  $\mathcal{O}(100)$  larger than the connected ones on an ensemble with  $L = 24a = 2.4$  fm and  $m_\pi = 429$  MeV. By comparing results from two volumes [18],  $L = 24a = 2.4$  fm and  $L = 48a = 4.2$  fm, we confirm that this enhancement scales approximately with the spatial volume, in agreement with expectations in the noninteracting limit. Our simulations use ensembles generated by the Coordinated Lattice Simulations (CLS) effort with  $\mathcal{O}(a)$ -improved Wilson fermions [19]. This approximate volume scaling is also observed in [20], which adopts the twisted-mass formulation of the action, and which we expect to be a general feature of lattice QCD three-point functions.

Therefore, in the improved three-point functions of Eq. (6) and at the moderate intermediate distances currently accessible ( $\tau, t-\tau \lesssim 0.8$  fm for nucleon observables), operators whose correlation functions include quark-line disconnected contributions dominate over those that do not. This simple but important observation has direct implications: the presence (or absence) of quark-line disconnected diagrams in a given three-point function indicates which excited states are most likely to contribute significantly at accessible distances. This connection becomes especially transparent in a partially quenched framework [21], where individual contractions can be mapped to specific intermediate states in the spectral decomposition.

For instance, the meson–current term appearing in Eq. (11) contains the matrix element  $\langle 0 | \mathcal{J} | M \rangle$ , describing the creation of a meson by the current. The quantum numbers of the meson  $M$  can be inferred from current algebra, providing theoretical guidance on which states couple most strongly to a given current. Importantly, although the quark-line disconnected diagrams are large in amplitude, their time dependence reflects the propagation of multiparticle states, and leads to a faster exponential decay compared to ground-state contributions present in connected diagrams.

This behavior becomes clear in the schematic structure of the connected and direct terms,

$$W_C = Z'_{HM} \langle (HM)' | \mathcal{J} | H \rangle Z_H^\dagger e^{-E'_{HM}(t-\tau)} e^{-E_H \tau} + |Z'_H| \langle H' | \mathcal{J} | H \rangle Z_H^\dagger e^{-E'_H(t-\tau)} e^{-E_H \tau} + \dots, \quad (13)$$

$$W_D = \mathcal{O}(L^3/a^3) |Z_H|^2 |Z_M|^2 e^{-E_H t} e^{-E_M(t-\tau)} + \dots \quad (14)$$

The connected contributions  $W_C$  contain both single- and multiparticle intermediate states, as propagation from source to sink can proceed through either type. In contrast,  $W_D$  is dominated by multiparticle propagation and thus decays more rapidly in Euclidean time. Consequently, although quark-line disconnected terms can be  $\mathcal{O}(100)$  larger in amplitude, they are exponentially suppressed at large time separations relative to the ground-state contribution in  $W_C$ . For the source-sink separations currently accessible in lattice QCD, however, these enhanced contributions dominate the signal and must be properly accounted for. This behavior is supported by numerical evidence reported in [20,22], and is consistent with the qualitative picture from partially quenched theory [21], which associates multihadron contributions with specific diagram topologies.

Importantly, the states created by the current in these diagrams are not limited to stable single hadrons. In lattice QCD, resonances such as the  $\rho$  or  $\sigma$  mesons do not appear as asymptotic states due to their strong decays. Instead, the current often couples directly to their decay products: for instance, the vector current can produce  $\pi\pi$  in P wave, while the scalar current can create  $\pi\pi$  in S wave. It is important to keep in mind that the dominant states can depend sensitively on the ensemble parameters—such as the pion mass and spatial volume—since these determine which multihadron states lie closest to threshold. We provide further discussion and numerical examples of such cases below.

*Strategies to account for current-enhanced states.* The identification of quark-line disconnected diagrams as the dominant contributions at short and intermediate source-sink separations provides a valuable insight: once these diagrams are known to drive the contamination, one can develop targeted strategies to account for the corresponding excited states.

This can be tackled through these proposed strategies:

- (i) *Variational analysis with tailored operators:* Include interpolators with strong overlap onto the enhanced multiparticle states, such as  $N\sigma$  [18],  $N\pi$  [20,23], or  $N\rho$ -like operators (see Supplemental Material [16]). This model-independent approach isolates and subtracts specific excited-state contributions, and enables determination of transition matrix elements, e.g.  $\langle N\pi | \mathcal{J}^\mu | N \rangle$  [22],  $\langle \pi | \mathcal{J}^\mu | \pi\pi \rangle$  [24],  $\langle K | \mathcal{J}^\mu | K\pi \rangle$  [25], timelike meson form factors beyond the elastic region [26],  $\langle \rho | \mathcal{J} | B \rangle$  and  $\langle \pi\pi | \mathcal{J}^\mu | B \rangle$  [27]. Recent results show that including only operators producing current-enhanced states already yields accurate and robust outcomes.

- (ii) *Two-point analysis of meson-current diagrams:* Compute the quark-line disconnected diagrams—or just the current-meson two-point function—to extract the energy of the associated multiparticle state and identify which states are current enhanced. These states can be then incorporated into effective field theory to compute analytic expressions for their contributions, or used as priors in multistate fits, or to directly remove contamination in standard three-point functions [28,29].
- (iii) *Effective field theory estimates:* Use low-energy EFTs, such as ChPT, treating the current-enhanced meson explicitly to compute analytic expressions for their contributions [30–34]. Information from the current-meson two-point analysis guides which couplings to include in the chiral Lagrangian.

While the variational method—particularly with an extended operator basis—offers the most reliable control over current-enhanced states, it is computationally expensive. The other two strategies are significantly cheaper and—when combined—can still yield robust analytic estimates of excited-state contamination.

*Examples: Numerical evidence in nucleon three-point functions.* The expression for the nucleon three-point function reads

$$C_{3pt}(\vec{p}', t; \vec{q}, \tau) = \langle O_N(\vec{p}', t) \mathcal{J}(\vec{q}, \tau) O_N^\dagger(\vec{p}, 0) \rangle, \quad (15)$$

whose spectral decomposition contains the nucleon-current matrix elements  $\langle N(\vec{p}') | \mathcal{J}(\vec{q}) | N(\vec{p}) \rangle$ . In the following, we discuss three examples for which we have numerical evidence of current-enhanced states: axial, scalar, and vector channels. Importantly, the observation of large contamination from certain excited states is not tied to a specific lattice action: different discretizations, such as the Highly Improved Staggered Quark action [33,35], the six stout-smear  $\mathcal{O}(a)$ -improved Wilson-Clover action with Iwasaki gauge action [28], the Twisted Mass action [20], a mixed-action setup with Overlap valence fermions on a Domain-Wall-fermion sea [36,37], and ensembles generated by the CLS effort [32,38], report similar effects, for instance in the nucleon channel, consistent with the general argument presented in this work.

**Nucleon axial matrix elements:** In the case of an isovector intermediate axial current  $\mathcal{J} = \mathcal{A}_\mu = \bar{\psi} \gamma_\mu \gamma_5 \psi$ , there is a nonvanishing direct diagram when the meson operator is a pionlike or  $a_1$ -like interpolator. In the first case, the direct diagram reads

$$W_D = \langle O_N(\vec{p}'_N, t) O_N^\dagger(\vec{p}, 0) \rangle_F \langle O_\pi(\vec{p}'_\pi, t) \mathcal{A}_\mu(\vec{q}, \tau) \rangle_F \quad (16)$$

and the gauge average of the pion-axial term yields

$$\langle O_\pi(\vec{p}_\pi, t) \mathcal{A}_\mu(\vec{q}, \tau) \rangle_F = i q_\mu f_\pi \delta_{\vec{p}_\pi, \vec{q}} e^{-E_\pi(t-\tau)}. \quad (17)$$

Therefore, this diagram is nonzero only when the pion carries the same momentum of the current and along the direction of

its component. Several lattice studies have reported a large excited-state contamination in this channel for the standard nucleon three-point function in Eq. (15), e.g., [39,40]. It was then predicted at leading order in ChPT [31,32] that  $N\pi$  states contribute very largely to such channels, and that their contribution is not suppressed by the volume, in agreement with the argument proposed in this letter.

Furthermore, ChPT predicts that  $N\pi$  states contribute only weakly—compared to other channels—to cases with  $\vec{q} = \vec{p}' = \vec{p} = \vec{0}$  [41,42], which include, for example, the axial charge [43]. This is again in agreement with the fact that the direct diagram in Eq. (16) vanishes in these kinematics. However, in the axial channel  $\mathcal{A}_4$  and forward limit with boosted frames, i.e.  $\vec{p}' = \vec{p}$ , Eq. (16) does not vanish and indeed, both ChPT [32] and lattice results [20,23] show a large contamination from  $N\pi$  states.

As for the excited-state contamination in the forward limit with  $\vec{q} = \vec{p}' = \vec{0}$ , we speculate that the axial meson  $a_1$  plays a prominent role, and in particular we suggest to include  $N(\vec{0})a_1(\vec{0})$  in the variational analysis or account for the excited states as discussed in the previous section. The reason is that by including  $N(\vec{0})a_1(\vec{0})$  in the variational basis, the Wick contractions of  $\langle O_{Na_1}(\vec{0}, t) \mathcal{A}_i(\vec{0}, \tau) O_N^\dagger(\vec{0}, 0) \rangle$  contain nonvanishing quark-line disconnected diagrams, which are volume enhanced. Very likely, the  $Na_1$ -like operator will couple well to the scattering states at sufficiently light pion masses, i.e.,  $N \otimes \text{decays products of } a_1$  [44].

Notice also that the inclusion of the  $\mathcal{O}(a)$ -improvement term  $c_A a \partial_\mu \mathcal{P}$  in the external axial currents with Wilson-Clover actions will include additional contributions from the same current-enhanced states ( $N\pi$ ), as both the axial and pseudoscalar currents can exhibit large excited state contamination for certain channels [31,45], albeit suppressed by  $\mathcal{O}(a)$ . This applies generally to other sectors and external currents as well.

**Nucleon scalar matrix elements:** The nucleon scalar matrix elements can be extracted from nucleon three-point functions with a scalar current insertion  $\mathcal{J} = \mathcal{S}$ . Of particular interest are nucleon isoscalar scalar matrix elements in the forward limit, which are proportional to the nucleon scalar coupling and sigma terms [38,46].

Following the same argument of the previous sections, when  $N\sigma$  operators are included in the variational basis, the isoscalar scalar channel can receive sizable contributions from both the direct and the disconnected terms in Eqs. (11)–(12). In this case, the direct and disconnected terms read

$$W_D(\vec{p}'_N; \vec{p}'_\sigma; \vec{0}) = \langle O_N(\vec{p}'_N, t) O_N^\dagger(\vec{0}, 0) \rangle_F \times \langle O_\sigma(\vec{p}'_\sigma, t) \mathcal{S}(\vec{0}, 0) \rangle_F, \quad (18)$$

$$W_{\text{disc}}(\vec{p}'_N; \vec{p}'_\sigma; \vec{0}) = \langle O_N(\vec{p}'_N, t) O_N^\dagger(\vec{0}, 0) \rangle_F \times \langle O_\sigma(\vec{p}'_\sigma, t) \rangle_F \langle \mathcal{S}(\vec{0}, 0) \rangle_F. \quad (19)$$



The specific intermediate states contributing to these terms depend on the ensemble parameters, such as the pion mass and spatial volume.

In the regime where the pion mass is sufficiently large that the low-lying scalar mesons lie below the multiparticle threshold, the physical interpretation is that  $N\sigma$  states contribute to the disconnected piece, while  $Na_0$  states contribute to the direct piece. In [18], we verify numerically the large  $N\sigma$  contribution in the isoscalar channel at  $m_\pi = 429$  MeV, where the sigma lies at  $m_\sigma = 554(49)$  MeV, below the  $\pi\pi$  threshold.

At lighter pion masses, where the  $\sigma$  is unstable, in the sense that its mass is above the  $\pi\pi$  S-wave threshold, the dominant contribution is instead expected from  $N\pi\pi$  S-wave states or from mixed  $N\sigma$ – $N\pi\pi$  components, due to their coupling. A next-to-next-to-leading order ( $N^2$ LO) ChPT calculation at the physical point [33] predicts large  $N\pi$  and  $N\pi\pi$  contributions. However, in [20] a variational analysis including the lowest  $N\pi$  states finds no effect on this observable—consistent with the absence of quark-line disconnected contributions from  $N\pi$  operators in this channel. We therefore advocate using  $N\sigma$ -like operators even at light pion masses, since they will couple to the relevant  $N\pi\pi$  scattering states [47].

**Nucleon vector matrix elements:** The vector matrix elements can be computed by inserting a vector current  $\mathcal{J} = \mathcal{V}_\mu$  in Eq. (15). Following the same steps of the previous section, in the isovector channel, we expect a large contamination from  $N\rho$  states, in the physical scenario where the  $\rho$  energy is below the  $\pi\pi$  energy threshold. This may be counterintuitive as the  $\rho$  is quite heavy and may be therefore missed in the multistate fit analyses. To prove this, we have carried out a variational analysis with  $N$  and  $N\rho$  operators on an ensemble with  $N_f = 3$  and  $m_\pi = 429$  MeV, where the  $\rho$  lies at  $m_\rho \approx 860$  MeV, well below the  $\pi\pi$  P-wave states lying at  $\approx 1360$  MeV. In Supplemental Material [16], we report numerical evidence that shows that by removing  $N\rho$  states from the nucleon isovector vector three-point functions, the excited-state contamination is exponentially reduced, proving that  $N\rho$  states are the dominant contribution. Notice that on this ensemble, the energy of the  $N\rho$  states with zero and unit total momentum is  $E_{N\rho} \approx 2280$  MeV, and 2170 MeV, respectively—substantially higher than the corresponding energies of  $N\sigma$ ,  $N\pi$ , or  $N\pi\pi$  states.

This analysis delivers an important message: current-enhanced multiparticle states can dominate the excited-state contamination even if they are heavy. While such states might appear to be suppressed due to their large energy, their strong overlap with the current via the quark-line disconnected diagrams allow them to contribute significantly at moderate Euclidean time separations. In particular, the  $N\rho$  contribution—even when heavier—can outcompete lighter states such as  $N\pi$ , due to its stronger

coupling to the vector current. In [46], a variational analysis with  $N$ - and  $N\pi$ -like operators shows that the  $N\pi$  states have little effect in this channel, again in perfect agreement with our argument.

Importantly, at sufficiently light pion masses, where the  $\rho$  becomes unstable, we expect that the dominant contribution is coming from  $N\pi\pi$  with  $\pi\pi$  in P wave in the rest frame, or a mixture of  $N\rho$  and  $N\pi\pi$ . A similar observation was already made in [48], where evidence for large contributions from  $N\pi\pi$  excited states in the vector channel was found, consistent with the vector meson dominance picture.

*Other examples: Semileptonic decays.* The argument presented in this work is general and extends naturally to processes where different hadrons appear at the source and sink, such as in semileptonic decays.

In the case of heavy-light decays, like, for instance,  $B \rightarrow \pi \ell \nu_\ell$ , the expression for the three-point function reads

$$\langle O_B(\vec{0}, t) \mathcal{V}_\mu(\vec{q}, \tau) O_\pi^\dagger(-\vec{q}, 0) \rangle. \quad (20)$$

For such correlation functions, the direct diagrams arise from two-hadron operators of the type  $B_\mu^* \pi$ , which can have the quantum numbers of  $B$ . In particular, the expression is

$$W_D(\vec{0}; \vec{p}'_\pi; \vec{q}) = \langle O_{B_\mu^*}(\vec{q}, t) \mathcal{V}_\mu(\vec{q}, \tau) \rangle_F \times \langle O_\pi(-\vec{q}, t) O_\pi^\dagger(-\vec{q}, 0) \rangle_F \quad (21)$$

which predicts that the current-enhanced  $B_\mu^* \pi$  states are in the momentum configuration where the pions in the final and initial state carry the same momentum, and  $B^*$  carries the same momentum as the current. In fact, in [34], it was predicted in heavy meson ChPT, that such states contribute largely to this channel.

Similar discussions hold for other processes like  $B_s \rightarrow K \ell \nu_\ell$ , investigated, for instance, in [49]. In this case, the largest excited-state contamination is expected to come from  $B^* K$  states. These examples illustrate how the mechanism discussed here applies beyond nucleon observables, and the framework can be naturally extended to a broad class of hadronic transitions.

*Conclusions.* Excited-state contamination remains a leading source of systematic uncertainty in lattice QCD determinations of hadron structure. We have identified a key mechanism behind this effect: external currents can couple strongly to specific multiparticle states whose contributions are not volume suppressed, leading to an effective current enhancement of these states. Such current-enhanced states—often overlooked in standard analyses for lack of direct ChPT predictions—can dominate signals at the source-sink separations accessible today. We present a general framework, supported by numerical evidence, that determines which excited states contribute most

significantly and why. The considerations of this work, which rely on symmetries of correlation functions and volume-scaling arguments, apply to any lattice discretization of the QCD action. For discretizations such as staggered fermions, the presence of additional unphysical species may give rise to further current-enhanced states, but the same reasoning applies. This picture agrees with existing chiral effective theory results, but extends beyond it by providing a diagrammatic criterion based on the Wick contraction structure, applicable to any channel. It clarifies, for example, why  $N\pi$  states are not the dominant excited states in certain channels—such as the scalar channel—and thus why their inclusion in [20] yielded improvements only where current-enhanced diagrams contribute, while having little effect elsewhere. In practical terms, our findings motivate the targeted inclusion of current-induced multihadron states in multistate fits, and provide a clear operator-selection criterion for variational analyses. Adopting these strategies can systematically reduce excited-state systematics, paving the way for higher-precision determinations of hadron structure in future lattice QCD calculations.

Simulations were carried out on the QPACE 3 computer of SFB/TRR-55, using an adapted version of the Chroma [50] software package.

*Acknowledgments.* The author gratefully acknowledges illuminating discussions and collaboration with G. Bali and S. Collins, as well as valuable input from J. Green, R. Gupta, A. Patella, S. Prelovšek, S. Schaefer, and R. Sommer. Special thanks are due to G. Bali, J. Green, and K.-F. Liu for comments on an earlier version of this manuscript, and to C. Alexandrou and Y. Li for sharing their data and further confirming the volume scaling of quark-line disconnected diagrams. This work was supported by the German Research Foundation (DFG) through the research unit FOR5269 “Future methods for studying confined gluons in QCD.”

*Data availability.* The data that support the findings of this article are not publicly available upon publication because it is not technically feasible and/or the cost of preparing, depositing, and hosting the data would be prohibitive within the terms of this research project. The data are available from the authors upon reasonable request.

- 
- [1] D. C. Hackett and M. L. Wagman, Block Lanczos algorithm for lattice QCD spectroscopy and matrix elements, *Phys. Rev. D* **112**, 014514 (2025).
  - [2] J. J. Sakurai, Theory of strong interactions, *Ann. Phys. (N.Y.)* **11**, 1 (1960).
  - [3] J. J. Sakurai, *Currents and Mesons* (University of Chicago Press, Chicago, 1969).
  - [4] N. M. Kroll, T. D. Lee, and B. Zumino, Neutral vector mesons and the hadronic electromagnetic current, *Phys. Rev.* **157**, 1376 (1967).
  - [5] R. J. Oakes and J. J. Sakurai, Spectral-function sum rules, omega-phi mixing, and lepton-pair decays of vector mesons, *Phys. Rev. Lett.* **19**, 1266 (1967).
  - [6] M. Gell-Mann and F. Zachariasen, Form-factors and vector mesons, *Phys. Rev.* **124**, 953 (1961).
  - [7] U. G. Meissner, Low-energy hadron physics from effective chiral Lagrangians with vector mesons, *Phys. Rep.* **161**, 213 (1988).
  - [8] S. Weinberg, Precise relations between the spectra of vector and axial vector mesons, *Phys. Rev. Lett.* **18**, 507 (1967).
  - [9] S. L. Adler, Consistency conditions on the strong interactions implied by a partially conserved axial vector current, *Phys. Rev.* **137**, B1022 (1965).
  - [10] S. L. Adler, Consistency conditions on the strong interactions implied by a partially conserved axial-vector current. II, *Phys. Rev.* **139**, B1638 (1965).
  - [11] G. Altarelli, N. Cabibbo, and L. Maiani, The sigma term and low-energy pi-n scattering, *Nucl. Phys.* **B34**, 621 (1971).
  - [12] V. Bernard, N. Kaiser, and U.-G. Meissner, Chiral dynamics in nucleons and nuclei, *Int. J. Mod. Phys. E* **04**, 193 (1995).
  - [13] B. Blossier, M. Della Morte, G. von Hippel, T. Mendes, and R. Sommer, On the generalized eigenvalue method for energies and matrix elements in lattice field theory, *J. High Energy Phys.* **04** (2009) 094.
  - [14] C. J. Shultz, J. J. Dudek, and R. G. Edwards, Excited meson radiative transitions from lattice QCD using variationally optimized operators, *Phys. Rev. D* **91**, 114501 (2015).
  - [15] J. Bulava, M. Donnellan, and R. Sommer, On the computation of hadron-to-hadron transition matrix elements in lattice QCD, *J. High Energy Phys.* **01** (2012) 140.
  - [16] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/69yc-d74z> for a sketch of the quark-line connected and disconnected diagrams, and results on nucleon-vector matrix elements with nucleon and nucleon-vector interpolating operators. The Supplemental Material includes Ref. [17] for the construction of  $N\rho$  operators.
  - [17] S. Prelovsek, U. Skerbis, and C. B. Lang, Lattice operators for scattering of particles with spin, *J. High Energy Phys.* **01** (2017) 129.
  - [18] L. Barca, G. Bali, and S. Collins, Nucleon sigma terms with a variational analysis from lattice QCD, *Phys. Rev. D* **111**, L031505 (2025).
  - [19] M. Bruno *et al.*, Simulation of QCD with  $N_f = 2 + 1$  flavors of non-perturbatively improved Wilson fermions, *J. High Energy Phys.* **02** (2015) 043.
  - [20] C. Alexandrou, G. Koutsou, Y. Li, M. Petschlies, and F. Pittler, Investigation of pion-nucleon contributions to nucleon matrix elements, *Phys. Rev. D* **110**, 094514 (2024).

- [21] K.-F. Liu, Parton degrees of freedom in pdfs from the hadronic tensor and large momentum effective theory, *Phys. Rev. D* **102**, 074502 (2020).
- [22] L. Barca, G. Bali, and S. Collins, Progress on nucleon transition matrix elements with a lattice QCD variational analysis, *Proc. Sci., EuroPLeX2023* (2024) 002.
- [23] L. Barca, G. Bali, and S. Collins, Toward  $N$  to  $N\pi$  matrix elements from lattice QCD, *Phys. Rev. D* **107**, L051505 (2023).
- [24] R. A. Briceño, J. J. Dudek, R. G. Edwards, C. J. Shultz, C. E. Thomas, and D. J. Wilson, The  $\pi\pi \rightarrow \pi\gamma^*$  amplitude and the resonant  $\rho \rightarrow \pi\gamma^*$  transition from lattice QCD, *Phys. Rev. D* **93**, 114508 (2016); *Phys. Rev. D* **105**, 079902(E) (2022).
- [25] A. Radhakrishnan, J. J. Dudek, and R. G. Edwards, Radiative decay of the resonant  $K^*$  and the  $\gamma K \rightarrow K\pi$  amplitude from lattice QCD, *Phys. Rev. D* **106**, 114513 (2022).
- [26] F. G. Ortega-Gama, J. J. Dudek, and R. G. Edwards, Time-like meson form factors beyond the elastic region from lattice QCD, *Phys. Rev. D* **110**, 094505 (2024).
- [27] L. Leskovec, S. Meinel, M. Petschlies, J. Negele, S. Paul, and A. Pochinsky,  $B \rightarrow \rho\ell\nu^-$  resonance form factors from  $B \rightarrow \pi\pi\ell\nu^-$  in lattice QCD, *Phys. Rev. Lett.* **134**, 161901 (2025).
- [28] Y. Aoki, K.-I. Ishikawa, Y. Kuramashi, S. Sasaki, K. Sato, E. Shintani, R. Tsuji, H. Watanabe, and T. Yamazaki, Method for high-precision determination of the nucleon axial structure using lattice QCD: Removing  $\pi N$ -state contamination, *Phys. Rev. D* **112**, 074510 (2025).
- [29] R. Tsuji, Y. Aoki, K.-I. Ishikawa, Y. Kuramashi, S. Sasaki, K. Sato, E. Shintani, H. Watanabe, and T. Yamazaki, Investigating the axial structure of the nucleon based on large-volume lattice QCD at the physical point, *arXiv*: 2505.10998.
- [30] O. Bar, Nucleon-pion-state contribution to nucleon two-point correlation functions, *Phys. Rev. D* **92**, 074504 (2015).
- [31] O. Bar,  $N\pi$ -state contamination in lattice calculations of the nucleon axial form factors, *Phys. Rev. D* **99**, 054506 (2019).
- [32] G. S. Bali, L. Barca, S. Collins, M. Gruber, M. Löffler, A. Schäfer, W. Söldner, P. Wein, S. Weishäupl, and T. Wurm, Nucleon axial structure from lattice QCD, *J. High Energy Phys.* **05** (2020) 126.
- [33] R. Gupta, S. Park, M. Hoferichter, E. Mereghetti, B. Yoon, and T. Bhattacharya, Pion–nucleon sigma term from lattice QCD, *Phys. Rev. Lett.* **127**, 242002 (2021).
- [34] O. Bar, A. Broll, and R. Sommer,  $B\pi$  excited-state contamination in lattice calculations of  $B$ -meson correlation functions, *Eur. Phys. J. C* **83**, 757 (2023).
- [35] Y.-Chull Jang, R. Gupta, B. Yoon, and T. Bhattacharya, Axial vector form factors from lattice QCD that satisfy the PCAC relation, *Phys. Rev. Lett.* **124**, 072002 (2020).
- [36] J. Liang, Y.-B. Yang, K.-F. Liu, A. Alexandru, T. Draper, and R. S. Sufian, Lattice calculation of nucleon isovector axial charge with improved currents, *Phys. Rev. D* **96**, 034519 (2017).
- [37] L. Liu, T. Chen, T. Draper, J. Liang, K.-F. Liu, G. Wang, and Y.-B. Yang, Nucleon isovector scalar charge from overlap fermions, *Phys. Rev. D* **104**, 094503 (2021).
- [38] A. Agadjanov, D. Djukanovic, G. von Hippel, H. B. Meyer, K. Ottnad, and H. Wittig, Nucleon sigma terms with  $N_f = 2 + 1$  flavors of O(a)-improved Wilson fermions, *Phys. Rev. Lett.* **131**, 261902 (2023).
- [39] R. Gupta, Y.-C. Jang, H.-W. Lin, B. Yoon, and T. Bhattacharya, Axial-vector form factors of the nucleon from lattice QCD, *Phys. Rev. D* **96**, 114503 (2017).
- [40] G. S. Bali, S. Collins, M. Gruber, A. Schäfer, P. Wein, and T. Wurm, Solving the PCAC puzzle for nucleon axial and pseudoscalar form factors, *Phys. Lett. B* **789**, 666 (2019).
- [41] B. C. Tiburzi, Chiral corrections to nucleon two- and three-point correlation functions, *Phys. Rev. D* **91**, 094510 (2015).
- [42] M. T. Hansen and H. B. Meyer, On the effect of excited states in lattice calculations of the nucleon axial charge, *Nucl. Phys.* **B923**, 558 (2017).
- [43] Z. B. Hall *et al.*, Signs of non-monotonic finite-volume corrections to  $g_A$ , *arXiv*:2503.09891.
- [44] Particle Data Group, Review of particle physics, *Prog. Theor. Exp. Phys.* **2020**, 083C01 (2020).
- [45] O. Bar,  $N\pi$ -state contamination in lattice calculations of the nucleon pseudoscalar form factor, *Phys. Rev. D* **100**, 054507 (2019).
- [46] C. Alexandrou, S. Bacchio, J. Finkenrath, C. Iona, G. Koutsou, Y. Li, and G. Spanoudes, Nucleon charges and  $\sigma$ -terms in lattice QCD, *Phys. Rev. D* **111**, 054505 (2025).
- [47] L. Barca, G. Bali, S. Collins, and M. Rodekamp (to be published).
- [48] S. Park, R. Gupta, B. Yoon, S. Mondal, T. Bhattacharya, Y.-C. Jang, B. Joó, and F. Winter, Precision nucleon charges and form factors using  $(2 + 1)$ -flavor lattice QCD, *Nucl. Phys. B, Proc. Suppl.* **105**, 054505 (2022).
- [49] A. Bazavov *et al.*,  $B_s \rightarrow K\ell\nu$  decay from lattice QCD, *Phys. Rev. D* **100**, 034501 (2019).
- [50] R. G. Edwards and B. Joo, The Chroma software system for lattice QCD, *Nucl. Phys. B, Proc. Suppl.* **140**, 832 (2005).