

Search for low-mass hidden-valley dark showers with non-prompt muon pairs in proton-proton collisions at $\sqrt{s} = 13$ TeV



The CMS collaboration

Full author list at the end of the paper

E-mail: cms-publication-committee-chair@cern.ch

ABSTRACT: A search for signatures of a dark analog to quantum chromodynamics is performed. The analysis targets long-lived dark mesons that decay into standard-model particles, with a high branching fraction of the dark mesons decaying into muons. The dark mesons are formed by the hadronisation of dark partons, which are produced by a decay of the Higgs boson. The search is performed using a data set corresponding to an integrated luminosity of 41.6 fb^{-1} , which was collected in proton-proton collisions at $\sqrt{s} = 13$ TeV by the CMS experiment at the CERN LHC in 2018 using non-prompt muon triggers. The search is based on resonant muon pair signatures. Machine-learning techniques are employed in the analysis, utilising boosted decision trees to discriminate between signal and background. No significant excess is observed above the standard model expectation. Upper limits on the branching fraction of the Higgs boson decaying to dark partons are determined to be as low as 10^{-4} at 95% confidence level, surpassing and extending the existing limits on models with dark $\tilde{W}$ mesons for mean proper decay lengths of less than 500 mm and for $\tilde{W}$ masses down to 0.3 GeV. First limits are set for extended dark-shower models with two dark flavours that contain dark photons, probing their masses down to 0.33 GeV.

KEYWORDS: Beyond Standard Model, Dark Matter, Hadron-Hadron Scattering

ARXIV EPRINT: [2511.11888](https://arxiv.org/abs/2511.11888)

Contents

1	Introduction	1
2	The CMS detector	3
3	Data sets, triggers and Monte Carlo simulation	5
4	Search strategy	6
5	Event selection and categorisation	7
5.1	Event selection	7
5.2	Event categorisation	10
6	Parametrisation of the dimuon mass distribution	10
7	Results and interpretation	13
7.1	Systematic uncertainties	13
7.2	Constraints on new physics models	15
8	Summary	23
	The CMS collaboration	29

1 Introduction

The standard model (SM) of particle physics has made many successful predictions; however, there remain a number of open questions to be answered. A major unresolved question concerns the nature of dark matter. From various astrophysical observations [1–3], the existence of a “dark” form of matter is inferred, which does not interact with the electromagnetic force but has mass. About 27% of the energy content of our universe is believed to be made up of dark matter, compared to only 5% being visible matter [1, 3–5].

Many theories beyond the SM try to account for dark matter. For example, hidden-valley models, which extend the SM to incorporate a new, non-Abelian gauge group [6–13]. Dark matter particles are charged under this new, dark gauge group. Hidden-valley models can produce events with high-multiplicity sprays of dark particles, known as dark showers. In hidden-valley models with confined dark charges, the dark particles assemble themselves to form dark hadrons. Dark hadrons can decay into SM particles after a measurable lifetime, creating displaced signatures in detectors. Dark showers containing sprays of dark hadrons are expected to create high-multiplicity displaced signatures [6, 14]. This analysis targets long-lived dark mesons that decay with a high branching fraction into muons, by searching for displaced vertices formed from resonant muon pairs.

Three broad classes of dark-sector models are explored: the vector portal model with a long-lived dark vector meson [14]; a model with a long-lived dark photon, Scenario A [15]; and a model with a long-lived dark pion, Scenario B1 [15]. The nomenclature of the latter

two models follows that in ref. [15]. The production of dark showers from decays of the SM Higgs boson is considered because it has a narrow width and is a sensitive probe for low-mass SM singlet states [16].

In the vector portal model, there are two types of dark mesons ($\tilde{\eta}$ and $\tilde{\omega}$), and it is assumed that only one of them has a measurable lifetime. This dark meson is referred to as the visibly-decaying dark particle. The other dark meson is assumed to either decay promptly to the visibly-decaying dark particle or to escape the detector and thus create an apparent momentum imbalance. Even though this choice does not account for the full range of possible dark-shower topologies, it still encompasses the phenomenology of a broad class of models and describes a wide range of distinct signatures with a minimal number of arbitrary parameters [14]. A few theory assumptions are made when constructing the model. It is assumed that there are no new sources of SM flavour violation. Operators up to dimension-5 in the infrared effective theory are allowed [14]. A diagram for the vector portal model is shown in figure 1. The Higgs boson first decays into a pair of dark partons $\psi\bar{\psi}$, which then hadronise through a dark analog to quantum chromodynamics (QCD) into dark hadrons, among them dark vector mesons $\tilde{\omega}$ and dark pseudoscalar mesons $\tilde{\eta}$. The $\tilde{\omega}$ has a non-prompt decay into SM particles, with a significant branching fraction of 10–30% for the decay into muons.

Scenarios A and B1 consist of extended hidden valley models with more complex decay topologies [15]. These models have two dark flavours instead of one, which leads to a wider spectrum of dark hadrons, including three dark pions ($\tilde{\pi}_1$, $\tilde{\pi}_2$, and $\tilde{\pi}_3$) and a heavier pseudoscalar meson $\tilde{\eta}$. The benchmark models include both pointing and non-pointing decay topologies, which refer to whether the dimuon vertices from the decays point back to the beam axis or not. The analysis is designed to be sensitive to both topologies. Diagrams for the decays in these scenarios are shown in figure 2. In both scenarios, the pseudoscalar $\tilde{\eta}$ is assumed to decay promptly into three dark pions, and the dark pions $\tilde{\pi}_1$ and $\tilde{\pi}_2$ are assumed to be stable. In Scenario A, the dark pion $\tilde{\pi}_3$ decays promptly into two dark photons A' , which can then each undergo a non-prompt decay into muons. The vector sum of the momenta of the two dimuon vertices points back to the beam axis, so it is referred to as the pointing scenario. In Scenario B1, the dark pion $\tilde{\pi}_3$ undergoes a displaced decay into two dark photons A' , which can then decay promptly into muons. The $\tilde{\pi}_3$ is a long-lived particle in Scenario B1, while A' is a long-lived particle in Scenario A. The dark photons are produced in the non-prompt decay of a long-lived parent particle in Scenario B1, and the momentum vectors of the resulting dimuon vertices do not point back to the beam axis, so this is referred to as the non-pointing scenario. As the dark photon decay is taken to be prompt, the dimuon vertices overlap with each other in this scenario. The pointing angle, defined as the angle between the momentum vector of the dimuon system and the displacement vector from the primary vertex to the secondary vertex, is used to characterise the decay topology. The primary vertex is taken to be the vertex corresponding to the hardest scattering in the event, evaluated using tracking information alone, as described in section 9.4.1 of ref. [17].

The search presented in this paper is performed using a data set collected in 2018 by the CMS experiment at the CERN LHC using a “data parking” strategy at a proton-proton (pp) center-of-mass energy of 13 TeV, corresponding to an integrated luminosity of 41.6 fb^{-1} [18, 19]. For the B-parking data set, the thresholds used by the trigger algorithms are lowered with




Figure 1. Diagram for the vector portal model. An SM Higgs boson decays to dark partons $\psi\bar{\psi}$, which then hadronise to form dark hadrons including dark vector mesons $\tilde{\omega}$ and dark pseudoscalar mesons $\tilde{\eta}$. The $\tilde{\omega}$ then undergoes displaced decay into SM fermions $f\bar{f}$.

respect to standard data sets, facilitating the selection of low transverse momentum (p_T) muons that are displaced from the primary vertex. This data set enables searches for low-mass, long-lived dark-sector particle decays into displaced muons. The present analysis probes a dimuon invariant mass range of 0.3 to 20 GeV. Machine-learning techniques are applied to perform background rejection. The B-parking data set includes full event reconstruction, which provides input for machine learning in this analysis.

Compared to previous dimuon searches performed at the CERN LHC [20–23], the study presented in this paper is the first to target hidden-valley dark-shower models with dark-sector particles decaying into pairs of muons. The sensitivities of previous searches in the low-mass regime were limited by dimuon mass restrictions. Ref. [20] requires the four-muon mass of muon pairs from two displaced vertices to be compatible with the Higgs boson mass. In ref. [21] and ref. [22] there are requirements on the dimuon mass to be larger than 10 GeV and 15 GeV respectively. Compared to the long-lived particle search in ref. [24] performed by the CMS experiment in the muon detectors, this analysis makes use of the tracker, which complements the sensitivity in the lower displacement region. This analysis exploits a different signature from those in refs. [25, 26], which search for dark showers through the emerging jet signature [27, 28].

Tabulated results for this analysis are provided in the HEPData record [29].

2 The CMS detector

The CMS apparatus [30, 31] is a multipurpose, nearly hermetic detector, designed to trigger on [32, 33] and identify electrons, muons, photons, and (charged and neutral) hadrons [34–36]. A global “particle-flow” algorithm [37] aims to reconstruct all individual particles in an event, combining information provided by the all-silicon inner tracker and by the crystal electromagnetic and brass-scintillator hadron calorimeters, operating inside a 3.8 T superconducting solenoid, with data from the gas-ionisation muon detectors embedded in




Figure 2. Diagrams for the Scenario A model (left) and the Scenario B1 model (right). In these extended models the dark hadronisation produces a spectrum of dark mesons, including the dark pions $\tilde{\pi}_1$, $\tilde{\pi}_2$ and $\tilde{\pi}_3$. The $\tilde{\pi}_3$ then decays into SM fermions $f\bar{f}$ through the dark photon A' . Green is used to indicate a long-lived particle. The A' is a long-lived particle in Scenario A, while $\tilde{\pi}_3$ is a long-lived particle in Scenario B1.

the flux-return yoke outside the solenoid. The reconstructed particles are used to identify jets and leptons and to measure missing transverse momentum [38–40].

Events of interest are selected using a two-tiered trigger system. The first level (L1), composed of custom hardware processors, uses information from the calorimeters and muon detectors to select events at a rate of around 100 kHz within a fixed latency of $4\mu\text{s}$ [32]. The second level, known as the high-level trigger (HLT), consists of a farm of processors running a version of the full event reconstruction software optimised for fast processing, and reduces the event rate to a few kHz before data storage [33].

In standard data sets, full offline reconstruction proceeds soon after the data are recorded. The data-parking strategy was designed to overcome the limitation in the ability to perform prompt reconstruction [19]. The trigger thresholds are lowered in the parking strategy, leading to a higher trigger rate, which exceeds the computing capacity available for prompt event reconstruction. The data stream is therefore transferred, unprocessed, to tape storage (the data stream is “parked”). It is kept in a raw format until sufficient computing resources become available for the events to be reconstructed, for example during shutdowns between data-taking periods.

Muons are identified as tracks in the inner tracker that are consistent with either a track or a segment made up of several hits in the muon system. The energy of muons is obtained from the corresponding track momentum, which is measured from the curvature of the track. Muons are measured in the range of $|\eta| < 2.4$. The single-muon trigger efficiency exceeds 90% over the full η range. The efficiency to reconstruct and identify muons with $p_T > 20\text{ GeV}$ is greater than 96%. Matching muons to tracks measured in the silicon tracker results in a relative p_T resolution of 1% in the barrel and 3% in the endcaps for muons with p_T up to 100 GeV [35].

3 Data sets, triggers and Monte Carlo simulation

For this analysis, the B-parking data set collected by CMS during the 2018 data-taking period [18, 19] is used. In order to enrich the data set in B hadron decays, the events collected contain at least one muon that satisfies kinematic requirements on the transverse momentum and the impact parameter significance. The latter is defined as $\text{IP}_{\text{sig}} = \text{IP}/\sigma_{\text{IP}}$, where IP is the impact parameter of the muon with respect to the beam axis in the transverse plane, and σ_{IP} is its uncertainty. The impact parameter is defined as the distance of closest approach of the muon track to the beam axis. The minimum requirement on IP_{sig} serves the dual purpose of reducing the prompt muon background while enhancing the potential contribution from new long-lived particles. The p_{T} thresholds in the triggers range from 7 to 12 GeV, and the thresholds on IP_{sig} vary between 3 and 6. The sample contains $\sim 1.3 \times 10^{10}$ events, with $\sim 70\%$ being B hadron decays.

Simulated signal samples are produced with PYTHIA 8.240 [41] for the vector portal model, and with PYTHIA 8.309 [42] for Scenario A and Scenario B1 models. The NNPDF3.1 parton distribution functions (PDFs) at next-to-next-to-leading order are used [43]. The search explores a dimuon mass range between 0.3 and 20 GeV. A variety of mean proper decay lengths ($c\tau$) of long-lived particles in these three models are considered, ranging from 0.1 to 500 mm. The search targets low-mass dimuon vertices, which are highly Lorentz boosted. The range of $c\tau$ considered is constrained by the drop in sensitivity at higher displacement. The gluon-gluon fusion production mode of the Higgs boson is considered, which has a cross section of 43.9 pb. A branching fraction of 0.01 is assumed for the Higgs boson decaying into dark partons to provide an illustration of a possible signal. The parameters of the models are introduced in table 1.

The vector portal model utilises three dark colours, and for simplicity, one dark flavour. The resulting dark hadron spectrum consists of two dark flavour singlet states: a spin-zero meson $\tilde{\eta}$, and a spin-one meson $\tilde{\omega}$. The dark quark mass is set to be approximately half the mass of $\tilde{\eta}$ [44]. In the benchmark models it is assumed that $m_{\tilde{\omega}} = \tilde{\Lambda} = m_{\tilde{\eta}}$, where $\tilde{\Lambda}$ is the dark sector confinement scale, so that the decay of $\tilde{\omega}$ into $\tilde{\eta}$ is kinematically forbidden, preventing the $\tilde{\omega}$ from decaying into other dark-sector states [14]. The branching fractions of the $\tilde{\omega}$ meson decaying into SM fermions are set according to ref. [14], and they depend on the $\tilde{\omega}$ mass.

For Scenario A and Scenario B1, the number of dark flavours is set to two and the number of dark colours is set to three. It is assumed that $m_{\tilde{\eta}} = \tilde{\Lambda}$. There is a mixing angle θ that parametrises the isospin violation in the coupling of the dark photon to dark-sector quarks. The mixing angle is taken to be $\sin \theta = 0.1$ so that the isospin breaking is relatively small, which results in the dark pions having similar masses, i.e. $m_{\tilde{\pi}_1} \approx m_{\tilde{\pi}_2} \approx m_{\tilde{\pi}_3}$. This gives a simpler mass spectrum, and also ensures the simulation from PYTHIA provides a reasonable approximation to the hadronisation process. The branching fraction $\mathcal{B}(\tilde{\pi}_3 \rightarrow A'A')$ is assumed to be 1 [15]. The dark photon A' decays into SM fermions by kinematically mixing with the SM photon, and the branching fractions are given by ref. [45]. The assumptions made for all the models are theoretically well-motivated and offer simplified benchmark models for interpretation.

For the background Monte Carlo (MC) simulations, a set of muon-enriched QCD background samples are produced, which contain events with at least one muon with $p_{\text{T}} > 5$ GeV.

Signal model	Model parameter	Range
Vector portal	$m_{\tilde{\omega}}$: mass of the dark vector meson	0.3–20 GeV
	$c\tau$: mean proper decay length of the dark vector meson	0.1–500 mm
Scenario A/B1	$m_{\tilde{\pi}_3}$: mass of the dark pion $\tilde{\pi}_3$	1–12 GeV
	$m_{A'}$: mass of the dark photon	0.33–2.5 GeV
	$c\tau$: mean proper decay length of the dark photon (Scenario A)	0.1–100 mm
	or the dark pion $\tilde{\pi}_3$ (Scenario B1)	

Table 1. Model parameters of the different classes of signal models interpreted by the analysis.

The samples are generated with PYTHIA 8.240, and they are binned in $\hat{p}_T$, which is the scale of momentum transfer of the QCD interactions.

Simulated minimum bias events are combined with simulated signal and background events to describe the effect of additional proton-proton interactions within the same or neighboring bunch crossings, which are referred to as pileup interactions.

4 Search strategy

The search targets the displaced decays of dark mesons and dark photons into pairs of muons. The dark mesons are produced through hadronisation within dark parton showers that are initiated by the decays of SM Higgs bosons. Final states with multiple non-prompt muon pairs are expected to result from the signal models.

An event-level boosted decision tree (BDT) is used to separate the signal from the heavy-flavour QCD background. Minimal selections are first applied to the events. For simplicity, in the pre-selections for the BDT training, a single non-prompt muon trigger with a muon p_T threshold of 9 GeV and an impact parameter significance threshold of 6 is applied. In addition, at least one dimuon secondary vertex (with two oppositely charged muons originating from a common vertex) that contains a muon with $p_T > 5$ GeV is required. Features that discriminate between signal and background are then studied. Examples of these features are the muon multiplicity shown in figure 3 (upper), and the muon transverse impact parameter d_{xy} shown in figure 3 (lower). It can be seen that there is a much larger number of muons in the signal events as compared with the background events. It is also found that muon variables provide excellent discrimination between signal and background, for example kinematic variables and variables related to the impact parameter. The multivariate input includes variables of muons, secondary vertices (produced from tracks of all types of charged particles), and muon secondary vertices (produced from muon tracks). Vertex variables include the χ^2 , pointing angle and displacement. Altogether there are a maximum of 390 input variables to the BDT. Where the value for a given feature does not exist for a particular event, it is set to a default value. As a result, the average number of input variables to the BDT with physical values is about 100. Jet variables are not included to make the BDT more model-independent. The variables providing the greatest discrimination are those associated with muons and muon secondary vertices. Good agreement between the data and the background Monte Carlo is observed for the BDT input variables.

The high muon multiplicity in the signal comes from the high multiplicity of dark mesons in the events, with up to an average of five dark mesons per event in Scenario A for example. Figure 4 shows the multiplicity of dark mesons for representative model parameters for the vector portal model (upper), as well as for Scenario A and B1 (lower). The multiplicity increases with decreasing mass of the dark meson, as expected, because less energy is required to produce a lighter meson. The multiplicity is found to be very similar between Scenario A and Scenario B1 for the same $\tilde{\pi}_3$ meson mass, as the two models have similar decay processes. Moreover, there is a high branching fraction for the dark mesons to decay into muons [14]. The branching fraction to muons drops with increasing mass of the dark meson because of competing decays to $c\bar{c}$, $b\bar{b}$, and $\tau^+\tau^-$. For example, in the vector portal model, the branching fraction to muons is 25% for $m_{\tilde{\omega}} = 2$ GeV, while it is 15% for $m_{\tilde{\omega}} = 20$ GeV. The high muon multiplicity in the dark-shower signal models allows a dedicated BDT to be trained for each of the three classes of models using a variety of input features to suppress the background.

The XGBOOST package [46] is used for the BDT training. Most of the hyperparameters are fixed as their default values [47], while some are fine-tuned to optimise the performance for each of the three signal models. For example, the number of boost rounds is changed to adapt to the size of the training samples. Other hyperparameters that are tuned are the learning rate, gamma, the maximum tree depth, the maximum delta step, subsampling ratios, and regularisation terms. Signal and QCD background samples are used for the training. The signal and background samples are weighted to give equal contributions to the training by using a binary cross-entropy loss function which is weighted event-by-event. Each of the N signal model samples (where, for example, $N = 25$ for the vector portal) is reweighted to give a contribution of $1/N$ to the total signal class after the preselections have been applied. The BDT is applied after the basic analysis selection criteria, which will be described in section 5.1.

5 Event selection and categorisation

The event sample is selected by requiring the presence of two oppositely charged muons that form a secondary vertex (SV) and by applying a minimum BDT score threshold. Events are categorised according to the number of vertices and in bins of transverse displacement (l_{xy}) and pointing angle to increase sensitivity to a signal.

5.1 Event selection

Selected events are required to pass at least one of the B-parking triggers and contain at least one reconstructed muon candidate satisfying $p_T > 9$ GeV, $\text{IP}_{\text{sig}} > 6$, and $|\eta| < 1.5$ that is matched to the muon that passes the trigger requirement by requesting $\Delta R < 0.1$, where $\Delta R = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2}$. Furthermore, the muon must pass the loose muon identification (ID) requirements [35].

At least one SV is required to be formed by a pair of oppositely charged muons. The vertex is reconstructed with a standard Kalman filter [48] algorithm, with the range in the distance from the beam axis extended [49]. This modification improves the reconstruction efficiency for the present search, which probes a transverse displacement of up to about 15 cm, by more than 40% relative to using the vertexing algorithm with standard parameters for different displacements. Quality criteria are then applied to the muon SVs, which are




Figure 3. Distributions of examples of variables that are used in the BDT training for the QCD background and benchmark signal models: muon multiplicity (upper) and muon transverse impact parameter (lower).

$\Delta R(\mu, \mu) < 1.2$, $\chi^2 < 10$, one muon matching the one that triggered the event, and both muons passing the loose ID [35]. If more than one muon SV passes the selections, only the muon SV with the lowest χ^2 is used for the final fit.

A selection is applied on the BDT scores of the events, with a threshold of 0.997 (0.999 for signal mass hypotheses less than 5 GeV) for the vector portal model. The BDT thresholds for Scenario A and Scenario B1 models are 0.98 and 0.991 respectively. The BDT cut alone gives a signal efficiency of $\approx 30\text{--}60\%$ and a background efficiency of $\approx 10^{-4}$ (or $\approx 10^{-5}$ for signal mass hypotheses less than 5 GeV in the vector portal model). The BDT thresholds are optimised using control regions in data (data sidebands) for representative signal mass




Figure 4. Distributions of the multiplicity of dark vector mesons $\tilde{\omega}$ for representative vector portal models (upper), and the multiplicity of dark mesons $\tilde{\pi}_3$ for representative Scenario A and B1 models (lower).

models. The data sidebands for each representative signal mass hypothesis are regions in the dimuon mass distribution that are outside the signal region, which is located at the centre of the mass window. The BDT selections are varied to find the optimal expected upper limits on the branching fraction of the Higgs decaying to dark partons.

The χ^2 of the muon SVs is found to be one of the most important variables of the BDT. The signal muon SVs are better fitted and have a lower χ^2 compared to the background, which is dominated by muon vertices from random muon combinations. The muon SV's transverse displacement and its significance are also found to be important variables. This

Resonance	Lower bound [GeV]	Upper bound [GeV]
K_S	0.43	0.49
η	0.52	0.58
π/ω	0.73	0.84
ϕ	0.96	1.08
J/ψ	2.91	3.27
$\psi(2S)$	3.47	3.89
$Y(1S)$	8.99	9.87
$Y(1S)$	9.61	10.39
$Y(3S)$	9.87	10.77

Table 2. The SM resonances and the corresponding mass windows that are masked in the analysis.

is expected as the signal models contain non-prompt decays, while the QCD background is dominated by prompt processes. The pointing angle is also a good discriminating variable.

5.2 Event categorisation

Events are categorised based on whether there are at least two vertices with similar dimuon mass. An event is classified as a multi-vertex event if there is at least one pair of muon SVs with dimuon masses within 3% of each other, which is about three times the width of the signal. Otherwise the event is classified as a single-vertex event.

Both the single-vertex and multi-vertex events are further categorised by the l_{xy} and pointing angle of the muon SV with the lowest χ^2 . The transverse displacement is defined as the distance in the transverse plane between the primary vertex and the secondary vertex. At increasing displacement from the primary vertex, the QCD background processes are expected to diminish. Events are categorised in the transverse displacement in order to increase the sensitivity to different proper lifetimes of the signal models. Categorising in transverse displacement is found to be more beneficial to the sensitivity than categorising in transverse displacement significance. The l_{xy} categories are defined to be $[0.0, 1.0]$ cm, $(1.0, 10.0]$ cm and beyond 10 cm. The signal efficiency drops considerably for $l_{xy} \gtrsim 20$ cm.

In each l_{xy} bin, events are further divided into bins of the pointing angle. This is to increase sensitivity to signal models that are more pointing than the background. The pointing angle bins are chosen to be $[0.0, 0.2]$ and $(0.2, \pi]$ to maintain sensitivity to non-pointing signal scenarios. Twelve categories are defined in total. The dimuon mass distributions for all categories are shown in figures 5 and 6 for data and two vector portal model signal benchmarks. The grey bands in the plots correspond to mass regions of known SM resonances, which are masked in the search, as listed in table 2. This is to avoid misidentifying the SM resonances as the signal. It can be seen that some of the categories are background-free, which shows the capability of the BDT in rejecting the background.

6 Parametrisation of the dimuon mass distribution

Using the dimuon mass distribution in each category, a parametric fit is performed for different signal mass and lifetime hypotheses for each of the three models. A Voigtian function (which




Figure 5. Dimuon invariant mass distributions for each single-vertex category for data and two vector portal model signal benchmarks. The shaded regions indicate mass regions of known SM resonances, which are masked in the search. A branching fraction of 0.01 is assumed for the Higgs boson decaying into dark partons for illustrating the signals.




Figure 6. Dimuon invariant mass distributions for each multi-vertex category for data and two vector portal model signal benchmarks. The shaded regions indicate mass regions of known SM resonances, which are masked in the search. A branching fraction of 0.01 is assumed for the Higgs boson decaying into dark partons for illustrating the signals.

is the convolution of a Gaussian function and a Breit-Wigner function) is used to fit the signal dimuon mass distribution. From the fits, it is found that the Gaussian width of the signal, σ , is about 0.75% of the mass for the vector portal, and about 0.85% of the mass for Scenario A and Scenario B1, each with an uncertainty of about 15%.

To perform the search, a sliding fit window of $\pm 5 \times \text{HWHM}$ of the signal around the signal mass hypothesis is chosen, where HWHM is the half width at half maximum of the signal resonance and is estimated to be 1% of the mass. To model the background, data sidebands (which are defined as mass regions in the fit window, outside of the region within $\pm 2 \times \text{HWHM}$ of the signal) are fitted with an envelope of functions, which are the first order polynomial, exponential, and power law functions. At each point in the likelihood scan, the particular fitting function that maximises the likelihood is chosen. Discrete profiling is used to account for the background systematic uncertainty from the choice of the function [50]. The parameters of each function are fixed in the likelihood fit, while the normalisation is allowed to vary. Goodness of fit tests have been performed, which show the functions can model the background appropriately. Bias tests have also been performed, and the fit is found to be sufficiently unbiased in the measurement of a potential signal.

For mass windows with very low background, a flat function is used instead of the envelope of functions to ensure stability of fit, following the fitting strategy used in previous dimuon searches [20, 51]. It was found that fitting with this approach gives upper limits that agree with those obtained from a counting experiment.

The background-only fit is shown in figure 7 for two mass windows in an example category, together with the signal expected from representative signal models.

7 Results and interpretation

Maximum likelihood fits to the data are performed under the background-only or background-plus-signal hypotheses, using a sliding mass window for different signal mass hypotheses. The fits are carried out simultaneously for all the categories which contain non-zero expected signal.

7.1 Systematic uncertainties

Various sources of systematic uncertainties in the signal modelling are considered. The systematic uncertainty in non-prompt muon ID efficiency is measured using the “tag and probe” technique [52] in the J/ψ mass region to study the differences between data and simulation. J/ψ from heavy flavour decay is used as a “proxy” for the signal as it is a known displaced resonance that lies in the mass range of the search. Scale factors are derived to account for the differences, and the systematic uncertainty in the ID efficiency is found by varying the scale factors with respect to the nominal values. The uncertainty is found to be between 1% and 17%, depending on the p_T of the muon and the transverse displacement of the muon SV. For the trigger selection, the same technique is used to compare the trigger efficiencies in data and simulation. The systematic uncertainty in the trigger efficiency typically lies between 1% and 2%, depending on the p_T and the impact parameter significance of the muon. The BDT selection efficiency is measured in the J/ψ mass region for data and simulation. The systematic uncertainty in the BDT selection efficiency is found to be 10%. The systematic uncertainty from pileup is obtained by considering the pileup distribution




Figure 7. Dimuon invariant mass distributions in mass windows around 0.67 GeV (upper) and 1.33 GeV (lower), in the single-vertex category with $1 \text{ cm} < l_{xy} < 10 \text{ cm}$ and pointing angle < 0.2 . The background fit is shown together with the signal expected for a representative Scenario A model (upper) and a representative Scenario B1 model (lower). A branching fraction of 0.01 is assumed for the Higgs boson decaying into dark partons for illustrating the signals. The lower panel in each plot shows the pull distribution, defined as the difference between the data and the background fit in each bin divided by the statistical uncertainty.

Source	Systematic uncertainty [%]
Non-prompt muon identification	$\sim 1\text{--}17$
Trigger	$\sim 1\text{--}2$
BDT	10
Pileup	~ 5
Theory uncertainty in ggF Higgs production	3.9
Strong coupling constant	2.6
PDF	1.9
Luminosity	2.5

Table 3. Summary of the systematic uncertainties in the signal yield expectation.

during the time when the B-parking triggers were active. Pileup weights are applied to Monte Carlo events such that the number of interactions per bunch crossing agrees with that observed in data. The pileup modelling uncertainty is found, by varying the weights, to be about 5%. The systematic uncertainty in the luminosity is 2.5% [53]. Various sources of systematic uncertainties in the gluon-gluon fusion Higgs boson production cross section are considered. These include a theory uncertainty of 3.9% due to missing higher order terms in the QCD calculation, and uncertainties in the factorisation and renormalisation scales. There is also an uncertainty of 1.9% from the choice of PDFs, and an uncertainty of 2.6% from the strong coupling constant value [54]. The various sources of systematic uncertainties for the signal are summarised in table 3.

The background systematic uncertainty is obtained from discrete profiling of the envelope of fitting functions used to model the background, which predominantly originates from QCD processes. The background uncertainty is found to be negligible compared to the other systematic uncertainties.

Uncertainties in the trigger efficiency, non-prompt muon ID efficiency and pileup modelling are treated as uncorrelated across categories. Other uncertainties are treated as correlated. The total systematic uncertainty is in the range 13–21%, which is less than the statistical uncertainty in the background normalisation.

7.2 Constraints on new physics models

No significant excess beyond the standard model expectation is observed, and 95% confidence level (CL) upper limits are set on the branching fraction of the Higgs boson decaying to dark partons. Limits are estimated using a modified frequentist approach with the CL_s criterion [55–57]. The limits are obtained for different mass and lifetime hypotheses of the dark shower signal models. For the vector portal, the upper limits on the branching fraction $\mathcal{B}(\text{H} \rightarrow \psi\bar{\psi})$ as functions of the dark vector meson proper decay length are shown in figure 8 for four representative mass hypotheses. The limits on the branching fraction are in the range 10^{-3} to 10^{-5} throughout most of the parameter space.

The limits for the Scenario A model as functions of the dark photon proper decay length are shown in figure 9 for four representative mass hypotheses. The dark photon mass models




Figure 8. Upper limits at 95% CL on the branching fraction $\mathcal{B}(H \rightarrow \psi\bar{\psi})$ as functions of the $\tilde{\omega}$ meson $c\tau$ for representative $\tilde{\omega}$ mass hypotheses and branching fractions for decaying into a muon pair in the vector portal model. It is assumed that $m_{\tilde{\omega}} = \tilde{\Lambda} = m_{\tilde{\eta}}$, where $m_{\tilde{\omega}}$, $\tilde{\Lambda}$, and $m_{\tilde{\eta}}$ are parameters of the dark sector: the mass of the spin-one meson, confinement scale, and the mass of the spin-zero meson, respectively.

chosen for Scenario A are in the low-mass regime (below 2 GeV). The limits obtained are in the range 10^{-2} to 10^{-4} .

The limits for the Scenario B1 model as functions of the dark meson $\tilde{\pi}_3$ proper decay length are shown in figure 10 for four representative mass hypotheses. The Scenario B1 model features a non-pointing dimuon vertex topology, and the benchmarks chosen are for low mass. The limits are found to be comparable to those for the Scenario A models, which shows sensitivity is achieved for non-pointing models as well as pointing models.

The constraints are more stringent in general for lower lifetime hypotheses owing to the higher SV reconstruction efficiency at smaller displacement. For very short lifetimes, the constraints are less stringent due to smaller signal efficiencies. The limits also tend to be more stringent at higher mass due to the lower background present. For similar dark meson masses, stronger sensitivity is achieved for the vector portal because there is a higher dark-meson multiplicity (as shown in figure 4).




Figure 9. Upper limits at 95% CL on the branching fraction $\mathcal{B}(H \rightarrow \psi\bar{\psi})$ as functions of the A' $c\tau$ for representative $\tilde{\pi}_3$ and A' mass hypotheses, and branching fractions of the dark photon decaying into a muon pair in Scenario A. It is assumed that $m_{\tilde{\eta}} = \tilde{\Lambda} = 4m_{\tilde{\pi}_2}$ and $\sin\theta = 0.1$, where $m_{\tilde{\eta}}$ is the mass of the dark-sector pseudoscalar meson and θ is the mixing angle paramtrising the isospin violation. The branching fraction $\mathcal{B}(\tilde{\pi}_3 \rightarrow A'A')$ is assumed to be 1.

The limits obtained for the vector portal model are complementary to those from a previous search performed by the CMS experiment in the muon detectors [24] studying the same model, for which longer proper decay lengths are covered. This search probes new parameter space for proper decay lengths of the dark $\tilde{\omega}$ meson below ~ 100 mm and masses as low as 0.3 GeV. There is also an improvement in the limits of up to two orders of magnitude with respect to ref. [24], which gave the best LHC limits on the model, in the range of proper decay lengths between 100 and 500 mm. The first constraints to date are imposed on the Scenario A and Scenario B1 models.

Limits are also shown as functions of the dark vector meson mass in the vector portal model (figure 11), and the dark photon mass for Scenario A and Scenario B1 models in figure 12 and figure 13, respectively. The limits are shown for two representative proper decay length hypotheses. For Scenarios A and B1, two different cases are considered for the mass ratio between the dark pion $\tilde{\pi}_3$ and the dark photon A' , which are $m_{\tilde{\pi}_3} = 10m_{A'}$ and $m_{\tilde{\pi}_3} = 3m_{A'}$, respectively. The limits are found to be stronger for the case with $m_{\tilde{\pi}_3} = 10m_{A'}$,




Figure 10. Upper limits at 95% CL on the branching fraction $\mathcal{B}(H \rightarrow \psi\bar{\psi})$ as functions of the $\tilde{\pi}_3$ meson $c\tau$ for representative $\tilde{\pi}_3$ and A' mass hypotheses, and branching fractions of the dark photon decaying into a muon pair in Scenario B1. It is assumed that $m_{\tilde{\eta}} = \tilde{\Lambda} = 4m_{\tilde{\pi}_2}$ and $\sin\theta = 0.1$, where $m_{\tilde{\eta}}$ is the mass of the dark-sector pseudoscalar meson and θ is the mixing angle parametrising the isospin violation. The branching fraction $\mathcal{B}(\tilde{\pi}_3 \rightarrow A'A')$ is assumed to be 1.

because of the lower dark pion multiplicity, leading to a more energetic A' . The final-state dimuon system is more Lorentz boosted and is more displaced as a result. The improvement in sensitivity from the higher displacement outweighs the effect of the lower multiplicity of SVs in this case, leading to an improvement in the sensitivity. The decrease in sensitivity at $m_{A'} \sim 0.7$ GeV is due to a decrease in the branching fraction $\mathcal{B}(A' \rightarrow \mu\mu)$, caused by competing decays of A' to $\pi^+\pi^-$ or $\pi^+\pi^-\pi^0$.

Observed limits at 95% CL are shown in the two-dimensional parameter space of the dark vector meson mass and proper decay length for the vector portal in figure 14, the dark photon mass and proper decay length for Scenario A in figure 15, and the dark photon mass and the dark pion ($\tilde{\pi}_3$) proper decay length for Scenario B1 in figure 16.




Figure 11. Upper limits at 95% CL on the branching fraction $\mathcal{B}(H \rightarrow \psi\bar{\psi})$ as functions of the $\tilde{\omega}$ meson mass for representative $c\tau$ hypotheses in the vector portal model. It is assumed that $m_{\tilde{\omega}} = \tilde{\Lambda} = m_{\tilde{\eta}}$, where $m_{\tilde{\omega}}$, $\tilde{\Lambda}$, and $m_{\tilde{\eta}}$ are parameters of the dark sector: the mass of the spin-one meson, confinement scale, and the mass of the spin-zero meson, respectively. The grey bands correspond to mass regions of known SM resonances, which are masked in the search.




Figure 12. Upper limits at 95% CL on the branching fraction $\mathcal{B}(H \rightarrow \psi\bar{\psi})$ as functions of the A' mass in Scenario A, for representative A' $c\tau$ hypotheses, and different ratios of the $\tilde{\pi}_3$ and A' masses. The limits are shown for the cases where $m_{\tilde{\pi}_3} = 10 m_{A'}$ (left column) and $m_{\tilde{\pi}_3} = 3 m_{A'}$ (right column), respectively. It is assumed that $m_{\tilde{\eta}} = \tilde{\Lambda} = 4 m_{\tilde{\pi}_2}$ and $\sin \theta = 0.1$, where $m_{\tilde{\eta}}$ is the mass of the dark-sector pseudoscalar meson and θ is the mixing angle parametrising the isospin violation. The branching fraction $\mathcal{B}(\tilde{\pi}_3 \rightarrow A'A')$ is assumed to be 1. The grey bands correspond to mass regions of known SM resonances, which are masked in the search.




Figure 13. Upper limits at 95% CL on the branching fraction $\mathcal{B}(H \rightarrow \psi\bar{\psi})$ as functions of the A' mass in Scenario B1, for representative $\tilde{\pi}_3$ $c\tau$ hypotheses, and different ratios of the $\tilde{\pi}_3$ and A' masses. The limits are shown for the cases where $m_{\tilde{\pi}_3} = 10m_{A'}$ (left column) and $m_{\tilde{\pi}_3} = 3m_{A'}$ (right column), respectively. It is assumed that $m_{\tilde{\eta}} = \tilde{\Lambda} = 4m_{\tilde{\pi}_2}$ and $\sin\theta = 0.1$, where $m_{\tilde{\eta}}$ is the mass of the dark-sector pseudoscalar meson and θ is the mixing angle parametrising the isospin violation. The branching fraction $\mathcal{B}(\tilde{\pi}_3 \rightarrow A'A')$ is assumed to be 1. The grey bands correspond to mass regions of known SM resonances, which are masked in the search.




Figure 14. Observed upper limits at 95% CL on the branching fraction $\mathcal{B}(H \rightarrow \psi\bar{\psi})$ as a function of the $\tilde{\omega}$ meson mass and $c\tau$ in the vector portal model. The parameter space region that is omitted is kinematically forbidden in the model. It is assumed that $m_{\tilde{\omega}} = \tilde{\Lambda} = m_{\tilde{\eta}}$, where $m_{\tilde{\omega}}$, $\tilde{\Lambda}$, and $m_{\tilde{\eta}}$ are parameters of the dark sector: the mass of the spin-one meson, confinement scale, and the mass of the spin-zero meson, respectively. The grey bands correspond to mass regions of known SM resonances, which are masked in the search.




Figure 15. Observed upper limits at 95% CL on the branching fraction $\mathcal{B}(H \rightarrow \psi\bar{\psi})$ as functions of the A' mass and $c\tau$ for representative ratios of the $\tilde{\pi}_3$ and A' masses in Scenario A. The limits are shown for the cases where $m_{\tilde{\pi}_3} = 10m_{A'}$ (left) and $m_{\tilde{\pi}_3} = 3m_{A'}$ (right), respectively. It is assumed that $m_{\tilde{\eta}} = \tilde{\Lambda} = 4m_{\tilde{\pi}_2}$ and $\sin\theta = 0.1$, where $m_{\tilde{\eta}}$ is the mass of the dark-sector pseudoscalar meson and θ is the mixing angle parametrising the isospin violation. The branching fraction $\mathcal{B}(\tilde{\pi}_3 \rightarrow A'A')$ is assumed to be 1. The grey bands correspond to mass regions of known SM resonances, which are masked in the search.




Figure 16. Observed upper limits at 95% CL on the branching fraction $\mathcal{B}(H \rightarrow \psi\bar{\psi})$ as functions of the A' mass and the $\tilde{\pi}_3$ $c\tau$ for representative ratios of the $\tilde{\pi}_3$ and A' masses in Scenario B1. The limits are shown for the cases where $m_{\tilde{\pi}_3} = 10m_{A'}$ (left) and $m_{\tilde{\pi}_3} = 3m_{A'}$ (right), respectively. It is assumed that $m_{\tilde{\eta}} = \tilde{\Lambda} = 4m_{\tilde{\pi}_2}$ and $\sin\theta = 0.1$, where $m_{\tilde{\eta}}$ is the mass of the dark sector pseudoscalar meson and θ is the mixing angle parametrising the isospin violation. The branching fraction $\mathcal{B}(\tilde{\pi}_3 \rightarrow A'A')$ is assumed to be 1. The grey bands correspond to mass regions of known SM resonances, which are masked in the search.

8 Summary

A search for dark showers has been performed with non-prompt muon pairs, using proton-proton collisions at the CERN LHC at a centre-of-mass energy of 13 TeV, collected by the CMS experiment in 2018, corresponding to an integrated luminosity of 41.6 fb^{-1} . The data set is recorded using the data parking strategy, resulting in a sample of about 10^{10} recorded events, giving access to masses down to the sub-GeV scale. No significant excess beyond the standard model expectation is observed. Upper limits on the branching fraction of the Higgs boson decaying into dark partons are set as low as 10^{-4} at 95% confidence level, providing the most stringent limits to date on the vector portal model with dark $\tilde{\omega}$ mesons of mean proper decay length below 500 mm and masses between 0.3 and 20 GeV. For the first time, limits have been set for extended dark-shower models with two dark flavours (Scenario A and Scenario B1) that contain dark photons, probing their masses down to 0.33 GeV.

Acknowledgments

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centres and personnel of the Worldwide LHC Computing Grid and other centres for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector, and the supporting computing infrastructure provided by the following funding agencies: SC (Armenia), BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MoST, and NSFC (China); MINCIENCIAS (Colombia); MSES

and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); ERC PRG, TARISTU24-TK10 and MoER TK202 (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); SRNSF (Georgia); BMFTR, DFG, and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIT and NRF (Republic of Korea); MES (Latvia); LMTLT (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MES, NSC, and NAWA (Poland); FCT (Portugal); MESTD (Serbia); MICIU/AEI and PCTI (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHEI (Thailand); TUBITAK and TENMAK (Türkiye); NASU (Ukraine); STFC (U.K.); DOE and NSF (U.S.A.).

Individuals have received support from the Marie-Curie programme and the European Research Council and Horizon 2020 Grant, contract Nos. 675440, 724704, 752730, 758316, 765710, 824093, 101115353, 101002207, 101001205, and COST Action CA16108 (European Union); the Leventis Foundation; the Alfred P. Sloan Foundation; the Alexander von Humboldt Foundation; the Science Committee, project no. 22rl-037 (Armenia); the Fonds pour la Formation à la Recherche dans l’Industrie et dans l’Agriculture (FRIA) and Fonds voor Wetenschappelijk Onderzoek contract No. 1228724N (Belgium); the Beijing Municipal Science & Technology Commission, No. Z191100007219010, the Fundamental Research Funds for the Central Universities, the Ministry of Science and Technology of China under Grant No. 2023YFA1605804, the Natural Science Foundation of China under Grant No. 12061141002, 12535004, and USTC Research Funds of the Double First-Class Initiative No. YD2030002017 (China); the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Shota Rustaveli National Science Foundation, grant FR-22-985 (Georgia); the Deutsche Forschungsgemeinschaft (DFG), among others, under Germany’s Excellence Strategy – EXC 2121 “Quantum Universe” – 390833306, and under project number 400140256 – GRK2497; the Hellenic Foundation for Research and Innovation (HFRI), Project Number 2288 (Greece); the Hungarian Academy of Sciences, the New National Excellence Program – ÚNKP, the NKFIH research grants K 131991, K 133046, K 138136, K 143460, K 143477, K 146913, K 146914, K 147048, 2020-2.2.1-ED-2021-00181, TKP2021-NKTA-64, and 2021-4.1.2-NEMZ_KI-2024-00036 (Hungary); the Council of Science and Industrial Research, India; ICSC – National Research Centre for High Performance Computing, Big Data and Quantum Computing, FAIR – Future Artificial Intelligence Research, and CUP I53D23001070006 (Mission 4 Component 1), funded by the NextGenerationEU program (Italy); the Latvian Council of Science; the Ministry of Education and Science, project no. 2022/WK/14, and the National Science Center, contracts Opus 2021/41/B/ST2/01369, 2021/43/B/ST2/01552, 2023/49/B/ST2/03273, and the NAWA contract BPN/PPO/2021/1/00011 (Poland); the Fundação para a Ciência e a Tecnologia, grant CEECIND/01334/2018 (Portugal); the National Priorities Research Program by Qatar National Research Fund; MICIU/AEI/10.13039/501100011033, ERDF/EU, “European Union NextGenerationEU/PRTR”, and Programa Severo Ochoa del Principado de Asturias (Spain); the Chulalongkorn Academic into Its 2nd Century Project Advancement Project, the National Science, Research and Innovation Fund program IND_FF_68_369_2300_097, and the Program Management Unit for Human Resources & Institutional Development, Research and Innovation, grant B39G680009 (Thailand); the Kavli Foundation; the Nvidia Corporation;

the SuperMicro Corporation; the Welch Foundation, contract C-1845; and the Weston Havens Foundation (U.S.A.).

Data Availability Statement. Release and preservation of data used by the CMS Collaboration as the basis for publications is guided by the [CMS data preservation, re-use and open access policy](#).

Code Availability Statement. Release and preservation of data used by the CMS Collaboration as the basis for publications is guided by the [CMS data preservation, re-use and open access policy](#).

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY4.0](#)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] V.C. Rubin, N. Thonnard and W.K. Ford Jr., *Rotational properties of 21 SC galaxies with a large range of luminosities and radii, from NGC 4605 ($R = 4\text{kpc}$) to UGC 2885 ($R = 122\text{ kpc}$)*, *Astrophys. J.* **238** (1980) 471 [[INSPIRE](#)].
- [2] M. Markevitch et al., *Direct constraints on the dark matter self-interaction cross-section from the merging galaxy cluster 1E0657-56*, *Astrophys. J.* **606** (2004) 819 [[astro-ph/0309303](#)] [[INSPIRE](#)].
- [3] PLANCK collaboration, *Planck 2018 results. VI. Cosmological parameters*, *Astron. Astrophys.* **641** (2020) A6 [*Erratum ibid.* **652** (2021) C4] [[arXiv:1807.06209](#)] [[INSPIRE](#)].
- [4] A. Arbey and F. Mahmoudi, *Dark matter and the early Universe: a review*, *Prog. Part. Nucl. Phys.* **119** (2021) 103865 [[arXiv:2104.11488](#)] [[INSPIRE](#)].
- [5] M. Persic, P. Salucci and F. Stel, *The universal rotation curve of spiral galaxies: 1. The dark matter connection*, *Mon. Not. Roy. Astron. Soc.* **281** (1996) 27 [[astro-ph/9506004](#)] [[INSPIRE](#)].
- [6] M.J. Strassler and K.M. Zurek, *Echoes of a hidden valley at hadron colliders*, *Phys. Lett. B* **651** (2007) 374 [[hep-ph/0604261](#)] [[INSPIRE](#)].
- [7] T. Hur, D.-W. Jung, P. Ko and J.Y. Lee, *Electroweak symmetry breaking and cold dark matter from strongly interacting hidden sector*, *Phys. Lett. B* **696** (2011) 262 [[arXiv:0709.1218](#)] [[INSPIRE](#)].
- [8] G.D. Kribs, T.S. Roy, J. Terning and K.M. Zurek, *Quirky Composite Dark Matter*, *Phys. Rev. D* **81** (2010) 095001 [[arXiv:0909.2034](#)] [[INSPIRE](#)].
- [9] H. Beauchesne, E. Bertuzzo and G. Grilli Di Cortona, *Dark matter in Hidden Valley models with stable and unstable light dark mesons*, *JHEP* **04** (2019) 118 [[arXiv:1809.10152](#)] [[INSPIRE](#)].
- [10] A. Francis, R.J. Hudspith, R. Lewis and S. Tulin, *Dark Matter from Strong Dynamics: The Minimal Theory of Dark Baryons*, *JHEP* **12** (2018) 118 [[arXiv:1809.09117](#)] [[INSPIRE](#)].
- [11] E. Bernreuther, F. Kahlhoefer, M. Krämer and P. Tunney, *Strongly interacting dark sectors in the early Universe and at the LHC through a simplified portal*, *JHEP* **01** (2020) 162 [[arXiv:1907.04346](#)] [[INSPIRE](#)].
- [12] H.-C. Cheng, L. Li and E. Salvioni, *A theory of dark pions*, *JHEP* **01** (2022) 122 [[arXiv:2110.10691](#)] [[INSPIRE](#)].

- [13] H.-C. Cheng, X.-H. Jiang, L. Li and E. Salvioni, *Dark showers from Z-dark Z' mixing*, *JHEP* **04** (2024) 081 [[arXiv:2401.08785](#)] [[INSPIRE](#)].
- [14] S. Knapen, J. Shelton and D. Xu, *Perturbative benchmark models for a dark shower search program*, *Phys. Rev. D* **103** (2021) 115013 [[arXiv:2103.01238](#)] [[INSPIRE](#)].
- [15] S. Born, R. Karur, S. Knapen and J. Shelton, *Scouting for dark showers at CMS and LHCb*, *Phys. Rev. D* **108** (2023) 035034 [[arXiv:2303.04167](#)] [[INSPIRE](#)].
- [16] D. Curtin et al., *Exotic decays of the 125 GeV Higgs boson*, *Phys. Rev. D* **90** (2014) 075004 [[arXiv:1312.4992](#)] [[INSPIRE](#)].
- [17] D. Contardo et al., *Technical Proposal for the Phase-II Upgrade of the CMS Detector*, CERN-LHCC-2015-010 (2015) [[DOI:10.17181/CERN.VU8I.D59J](#)] [[INSPIRE](#)].
- [18] CMS collaboration, *Enriching the physics program of the CMS experiment via data scouting and data parking*, *Phys. Rept.* **1115** (2025) 678 [[arXiv:2403.16134](#)] [[INSPIRE](#)].
- [19] CMS collaboration, *Recording and reconstructing 10 billion unbiased b hadron decays in CMS*, *EPJ Web Conf.* **245** (2020) 01025 [[INSPIRE](#)].
- [20] CMS collaboration, *Search for long-lived particles decaying into muon pairs in proton-proton collisions at $\sqrt{s} = 13$ TeV collected with a dedicated high-rate data stream*, *JHEP* **04** (2022) 062 [[arXiv:2112.13769](#)] [[INSPIRE](#)].
- [21] CMS collaboration, *Search for long-lived particles decaying to final states with a pair of muons in proton-proton collisions at $\sqrt{s} = 13.6$ TeV*, *JHEP* **05** (2024) 047 [[arXiv:2402.14491](#)] [[INSPIRE](#)].
- [22] ATLAS collaboration, *Search for long-lived particles in final states with displaced dimuon vertices in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Phys. Rev. D* **99** (2019) 012001 [[arXiv:1808.03057](#)] [[INSPIRE](#)].
- [23] LHCb collaboration, *Searches for low-mass dimuon resonances*, *JHEP* **10** (2020) 156 [[arXiv:2007.03923](#)] [[INSPIRE](#)].
- [24] CMS collaboration, *Search for long-lived particles decaying in the CMS muon detectors in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *Phys. Rev. D* **110** (2024) 032007 [[arXiv:2402.01898](#)] [[INSPIRE](#)].
- [25] CMS collaboration, *Search for dark QCD with emerging jets in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **07** (2024) 142 [[arXiv:2403.01556](#)] [[INSPIRE](#)].
- [26] CMS collaboration, *Search for new particles decaying to a jet and an emerging jet*, *JHEP* **02** (2019) 179 [[arXiv:1810.10069](#)] [[INSPIRE](#)].
- [27] P. Schwaller, D. Stolarski and A. Weiler, *Emerging Jets*, *JHEP* **05** (2015) 059 [[arXiv:1502.05409](#)] [[INSPIRE](#)].
- [28] S. Renner and P. Schwaller, *A flavoured dark sector*, *JHEP* **08** (2018) 052 [[arXiv:1803.08080](#)] [[INSPIRE](#)].
- [29] *HEPData record for this analysis*, (2025) [DOI:10.17182/hepdata.165888](#).
- [30] CMS collaboration, *The CMS Experiment at the CERN LHC, 2008* *JINST* **3** S08004 [[INSPIRE](#)].
- [31] CMS collaboration, *Development of the CMS detector for the CERN LHC Run 3, 2024* *JINST* **19** P05064 [[arXiv:2309.05466](#)] [[INSPIRE](#)].
- [32] CMS collaboration, *Performance of the CMS Level-1 trigger in proton-proton collisions at $\sqrt{s} = 13$ TeV, 2020* *JINST* **15** P10017 [[arXiv:2006.10165](#)] [[INSPIRE](#)].

- [33] CMS collaboration, *The CMS trigger system*, 2017 *JINST* **12** P01020 [[arXiv:1609.02366](#)] [[INSPIRE](#)].
- [34] CMS collaboration, *Electron and photon reconstruction and identification with the CMS experiment at the CERN LHC*, 2021 *JINST* **16** P05014 [[arXiv:2012.06888](#)] [[INSPIRE](#)].
- [35] CMS collaboration, *Performance of the CMS muon detector and muon reconstruction with proton-proton collisions at $\sqrt{s} = 13$ TeV*, 2018 *JINST* **13** P06015 [[arXiv:1804.04528](#)] [[INSPIRE](#)].
- [36] CMS collaboration, *Description and Performance of Track and Primary-Vertex Reconstruction with the CMS Tracker*, 2014 *JINST* **9** P10009 [[arXiv:1405.6569](#)] [[INSPIRE](#)].
- [37] CMS collaboration, *Particle-flow reconstruction and global event description with the CMS detector*, 2017 *JINST* **12** P10003 [[arXiv:1706.04965](#)] [[INSPIRE](#)].
- [38] CMS collaboration, *Performance of reconstruction and identification of τ leptons decaying to hadrons and ν_τ in pp collisions at $\sqrt{s} = 13$ TeV*, 2018 *JINST* **13** P10005 [[arXiv:1809.02816](#)] [[INSPIRE](#)].
- [39] CMS collaboration, *Jet energy scale and resolution in the CMS experiment in pp collisions at 8 TeV*, 2017 *JINST* **12** P02014 [[arXiv:1607.03663](#)] [[INSPIRE](#)].
- [40] CMS collaboration, *Performance of missing transverse momentum reconstruction in proton-proton collisions at $\sqrt{s} = 13$ TeV using the CMS detector*, 2019 *JINST* **14** P07004 [[arXiv:1903.06078](#)] [[INSPIRE](#)].
- [41] T. Sjöstrand et al., *An introduction to PYTHIA 8.2*, *Comput. Phys. Commun.* **191** (2015) 159 [[arXiv:1410.3012](#)] [[INSPIRE](#)].
- [42] C. Bierlich et al., *A comprehensive guide to the physics and usage of PYTHIA 8.3*, *SciPost Phys. Codeb.* **2022** (2022) 8 [[arXiv:2203.11601](#)] [[INSPIRE](#)].
- [43] NNPDF collaboration, *Parton distributions from high-precision collider data*, *Eur. Phys. J. C* **77** (2017) 663 [[arXiv:1706.00428](#)] [[INSPIRE](#)].
- [44] L. Carloni, J. Rathsman and T. Sjostrand, *Discerning Secluded Sector gauge structures*, *JHEP* **04** (2011) 091 [[arXiv:1102.3795](#)] [[INSPIRE](#)].
- [45] D. Curtin, R. Essig, S. Gori and J. Shelton, *Illuminating Dark Photons with High-Energy Colliders*, *JHEP* **02** (2015) 157 [[arXiv:1412.0018](#)] [[INSPIRE](#)].
- [46] T. Chen and C. Guestrin, *XGBoost: A Scalable Tree Boosting System*, [arXiv:1603.02754](#) [[DOI:10.1145/2939672.2939785](#)] [[INSPIRE](#)].
- [47] T. Chen et al., *xgboost: Extreme gradient boosting*, <https://github.com/dmlc/xgboost> (2024).
- [48] R. Fruhwirth, *Application of Kalman filtering to track and vertex fitting*, *Nucl. Instrum. Meth. A* **262** (1987) 444 [[INSPIRE](#)].
- [49] T. Speer et al., *Vertex fitting in the CMS tracker*, CERN-CMS-NOTE-2006-032 (2006) [[INSPIRE](#)].
- [50] P.D. Dauncey, M. Kenzie, N. Wardle and G.J. Davies, *Handling uncertainties in background shapes: the discrete profiling method*, 2015 *JINST* **10** P04015 [[arXiv:1408.6865](#)] [[INSPIRE](#)].
- [51] CMS collaboration, *Search for a high-mass dimuon resonance produced in association with b quark jets at $\sqrt{s} = 13$ TeV*, *JHEP* **10** (2023) 043 [[arXiv:2307.08708](#)] [[INSPIRE](#)].
- [52] CMS collaboration, *Measurement of the W and Z inclusive production cross sections at $\sqrt{s} = 7$ TeV with the CMS experiment at the LHC*, CMS-PAS-EWK-10-005 (2011) [[INSPIRE](#)].
- [53] CMS collaboration, *CMS luminosity measurement for the 2018 data-taking period at $\sqrt{s} = 13$ TeV*, CMS-PAS-LUM-18-002 (2019) [[INSPIRE](#)].

- [54] LHC HIGGS CROSS SECTION WORKING GROUP collaboration, *Handbook of LHC Higgs Cross Sections: 4. Deciphering the Nature of the Higgs Sector*, *CERN Yellow Rep. Monogr.* **2** (2017) 1 [[arXiv:1610.07922](#)] [[INSPIRE](#)].
- [55] G. Cowan, K. Cranmer, E. Gross and O. Vitells, *Asymptotic formulae for likelihood-based tests of new physics*, *Eur. Phys. J. C* **71** (2011) 1554 [*Erratum ibid.* **73** (2013) 2501] [[arXiv:1007.1727](#)] [[INSPIRE](#)].
- [56] T. Junk, *Confidence level computation for combining searches with small statistics*, *Nucl. Instrum. Meth. A* **434** (1999) 435 [[hep-ex/9902006](#)] [[INSPIRE](#)].
- [57] A.L. Read, *Presentation of search results: The CL_s technique*, *J. Phys. G* **28** (2002) 2693 [[INSPIRE](#)].

The CMS collaboration

A. Hayrapetyan¹, V. Makarenko¹, A. Tumasyan^{1,a}, W. Adam², J.W. Andrejkovic²,
 L. Benato², T. Bergauer², M. Dragicevic², C. Giordano², P.S. Hussain², M. Jeitler^{2,b},
 N. Krammer², A. Li², D. Liko², M. Matthewman², I. Mikulec², J. Schieck^{2,b},
 D. Schwarz², R. Schöffbeck^{2,b}, M. Shooshtari², M. Sonawane², W. Waltenberger²,
 C.-E. Wulz^{2,b}, T. Janssen³, H. Kwon³, D. Ocampo Henao³, T. Van Laer³,
 P. Van Mechelen³, J. Bierkens⁴, N. Breugelmans⁴, J. D'Hondt⁴, S. Dansana⁴,
 A. De Moor⁴, M. Delcourt⁴, F. Heyen⁴, Y. Hong⁴, P. Kashko⁴, S. Lowette⁴,
 I. Makarenko⁴, D. Müller⁴, J. Song⁴, S. Tavernier⁴, M. Tytgat^{4,c}, G.P. Van Onsem⁴,
 S. Van Putte⁴, D. Vannerom⁴, B. Bilin⁵, B. Clerbaux⁵, A.K. Das⁵, I. De Bruyn⁵,
 G. De Lentdecker⁵, H. Evard⁵, L. Favart⁵, P. Gianneios⁵, A. Khalilzadeh⁵, F.A. Khan⁵,
 A. Malara⁵, M.A. Shahzad⁵, L. Thomas⁵, M. Vanden Bemden⁵, C. Vander Velde⁵,
 P. Vanlaer⁵, F. Zhang⁵, M. De Coen⁶, D. Dobur⁶, G. Gokbulut⁶, J. Knolle⁶,
 D. Marckx⁶, K. Skovpen⁶, A.M. Tomaru⁶, N. Van Den Bossche⁶, J. van der Linden⁶,
 J. Vandenbroeck⁶, L. Wezenbeek⁶, S. Bein⁷, A. Benecke⁷, A. Bethani⁷, G. Bruno⁷,
 A. Cappati⁷, J. De Favereau De Jeneret⁷, C. Delaere⁷, F. Gameiro Casalinho⁷,
 A. Giammanco⁷, A.O. Guzel⁷, V. Lemaitre⁷, J. Lidrych⁷, P. Malek⁷, P. Mastrapasqua⁷,
 S. Turkcapar⁷, G.A. Alves⁸, M. Barroso Ferreira Filho⁸, E. Coelho⁸, C. Hensel⁸,
 T. Menezes De Oliveira⁸, C. Mora Herrera⁸, P. Rebello Teles⁸, M. Soeiro⁸,
 E.J. Tonelli Manganote^{8,d}, A. Vilela Pereira^{8,e}, W.L. Aldá Júnior⁹,
 H. Brandao Malbouisson⁹, W. Carvalho⁹, J. Chinellato^{9,f}, M. Costa Reis⁹,
 E.M. Da Costa⁹, G.G. Da Silveira^{9,g}, D. De Jesus Damiao⁹, S. Fonseca De Souza⁹,
 R. Gomes De Souza⁹, S. S. Jesus⁹, T. Laux Kuhn^{9,g}, M. Macedo⁹, K. Mota Amarilo⁹,
 L. Mundim⁹, H. Nogima⁹, J.P. Pinheiro⁹, A. Santoro⁹, A. Sznajder⁹, M. Thiel⁹,
 F. Torres Da Silva De Araujo^{9,h}, C.A. Bernardes^{10,g}, F. Damas¹⁰, E.M. Gregores¹⁰,
 B. Lopes Da Costa¹⁰, I. Maietto Silverio¹⁰, P.G. Mercadante¹⁰, S.F. Novaes¹⁰, B. Orzari¹⁰,
 Sandra S. Padula¹⁰, V. Scheurer¹⁰, T.R. Fernandez Perez Tomei¹⁰, A. Aleksandrov¹¹,
 G. Antchev¹¹, P. Danev¹¹, R. Hadjiiska¹¹, P. Iaydjiev¹¹, M. Shopova¹¹, G. Sultanov¹¹,
 A. Dimitrov¹², L. Litov¹², B. Pavlov¹², P. Petkov¹², A. Petrov¹², S. Keshri¹³,
 D. Laroze¹³, S. Thakur¹³, W. Brooks¹⁴, T. Cheng¹⁵, T. Javaid¹⁵, L. Wang¹⁵,
 L. Yuan¹⁵, Z. Hu¹⁶, Z. Liang¹⁶, J. Liu¹⁶, X. Wang¹⁶, H. Yang¹⁶, G.M. Chen^{17,i},
 H.S. Chen^{17,i}, M. Chen^{17,i}, Y. Chen¹⁷, Q. Hou¹⁷, X. Hou¹⁷, F. Iemmi¹⁷, C.H. Jiang¹⁷,
 A. Kapoor^{17,j}, H. Liao¹⁷, G. Liu¹⁷, Z.-A. Liu^{17,k}, J.N. Song^{17,k}, S. Song¹⁷, J. Tao¹⁷,
 C. Wang^{17,i}, J. Wang¹⁷, H. Zhang¹⁷, J. Zhao¹⁷, A. Agapitos¹⁸, Y. Ban¹⁸,
 A. Carvalho Antunes De Oliveira¹⁸, S. Deng¹⁸, B. Guo¹⁸, Q. Guo¹⁸, C. Jiang¹⁸, A. Levin¹⁸,
 C. Li¹⁸, Q. Li¹⁸, Y. Mao¹⁸, S. Qian¹⁸, S.J. Qian¹⁸, X. Qin¹⁸, C. Quaranta¹⁸, X. Sun¹⁸,
 D. Wang¹⁸, J. Wang¹⁸, M. Zhang¹⁸, Y. Zhao¹⁸, C. Zhou¹⁸, S. Yang¹⁹, Z. You²⁰,
 K. Jaffel²¹, N. Lu²¹, G. Bauer^{22,l,m}, Z. Cui^{22,m}, B. Li^{22,n}, H. Wang²², K. Yi^{22,o},
 J. Zhang²², Y. Li²³, Z. Lin²⁴, C. Lu²⁴, M. Xiao^{24,p}, C. Avila²⁵, D.A. Barbosa Trujillo²⁵,
 A. Cabrera²⁵, C. Florez²⁵, J. Fraga²⁵, J.A. Reyes Vega²⁵, C. Rendón²⁶, M. Rodriguez²⁶,
 A.A. Ruales Barbosa²⁶, J.D. Ruiz Alvarez²⁶, N. Godinovic²⁷, D. Lelas²⁷, A. Sculac²⁷,
 M. Kovac²⁸, A. Petkovic²⁸, T. Sculac²⁸, P. Bargassa²⁹, V. Brigljevic²⁹, B.K. Chitroda²⁹,
 D. Ferencek²⁹, K. Jakovcic²⁹, A. Starodumov²⁹, T. Susa²⁹, A. Attikis³⁰, K. Christoforou³⁰,

C. Leonidou³⁰, C. Nicolaou³⁰, L. Paizanos³⁰, F. Ptochos³⁰, P.A. Razis³⁰, H. Rykaczewski³⁰, H. Saka³⁰, A. Steppenov³⁰, M. Finger^{31,†}, M. Finger Jr.³¹, E. Ayala³², E. Carrera Jarrin³³, S. Elgammal^{34,q}, A. Ellithi Kamel^{34,r}, A. Hussein³⁵, H. Mohammed³⁵, K. Ehataht³⁶, M. Kadastik³⁶, T. Lange³⁶, C. Nielsen³⁶, J. Pata³⁶, M. Raidal³⁶, N. Seeba³⁶, L. Tani³⁶, E. Brücken³⁷, A. Milieva³⁷, K. Osterberg³⁷, M. Voutilainen³⁷, F. Garcia³⁸, P. Inkaew³⁸, K.T.S. Kallonen³⁸, R. Kumar Verma³⁸, T. Lampén³⁸, K. Lassila-Perini³⁸, B. Lehtela³⁸, S. Lehti³⁸, T. Lindén³⁸, N.R. Mancilla Xinto³⁸, M. Myllymäki³⁸, M.m. Rantanen³⁸, S. Saariokari³⁸, N.T. Toikka³⁸, J. Tuominiemi³⁸, N. Bin Norjoharuddeen³⁹, H. Kirschenmann³⁹, P. Luukka³⁹, H. Petrow³⁹, M. Besancon⁴⁰, F. Couderc⁴⁰, M. Dejardin⁴⁰, D. Denegri⁴⁰, P. Devoue⁴⁰, J.L. Faure⁴⁰, F. Ferri⁴⁰, P. Gaigne⁴⁰, S. Ganjour⁴⁰, P. Gras⁴⁰, G. Hamel de Monchenault⁴⁰, M. Kumar⁴⁰, V. Lohezic⁴⁰, Y. Maidannyk⁴⁰, J. Malcles⁴⁰, F. Orlandi⁴⁰, L. Portales⁴⁰, S. Ronchi⁴⁰, M.Ö. Sahin⁴⁰, A. Savoy-Navarro^{40,s}, P. Simkina⁴⁰, M. Titov⁴⁰, M. Tornago⁴⁰, R. Amella Ranz⁴¹, F. Beaudette⁴¹, G. Boldrini⁴¹, P. Busson⁴¹, C. Charlot⁴¹, M. Chiusi⁴¹, T.D. Cuisset⁴¹, O. Davignon⁴¹, A. De Wit⁴¹, T. Debnath⁴¹, I.T. Ehle⁴¹, S. Ghosh⁴¹, A. Gilbert⁴¹, R. Granier de Cassagnac⁴¹, L. Kalipoliti⁴¹, M. Manoni⁴¹, M. Nguyen⁴¹, S. Obraztsov⁴¹, C. Ochando⁴¹, R. Salerno⁴¹, J.B. Sauvan⁴¹, Y. Sirois⁴¹, G. Sokmen⁴¹, L. Urda Gómez⁴¹, A. Zabi⁴¹, A. Zghiche⁴¹, J.-L. Agram^{42,t}, J. Andrea⁴², D. Bloch⁴², J.-M. Brom⁴², E.C. Chabert⁴², C. Collard⁴², G. Coulon⁴², S. Falke⁴², U. Goerlach⁴², R. Haeblerle⁴², A.-C. Le Bihan⁴², M. Meena⁴², O. Poncet⁴², G. Saha⁴², P. Vaucelle⁴², A. Di Florio⁴³, D. Amram⁴⁴, S. Beauceron⁴⁴, B. Blancon⁴⁴, G. Boudoul⁴⁴, N. Chanon⁴⁴, D. Contardo⁴⁴, P. Depasse⁴⁴, H. El Mamouni⁴⁴, J. Fay⁴⁴, S. Gascon⁴⁴, M. Gouzevitch⁴⁴, C. Greenberg⁴⁴, G. Grenier⁴⁴, B. Ille⁴⁴, E. Jourd'Huy⁴⁴, M. Lethuillier⁴⁴, B. Massoteau⁴⁴, L. Mirabito⁴⁴, A. Purohit⁴⁴, M. Vander Donckt⁴⁴, J. Xiao⁴⁴, A. Khvedelidze^{45,u}, I. Lomidze⁴⁵, Z. Tsamalaidze^{45,u}, V. Botta⁴⁶, S. Consuegra Rodríguez⁴⁶, L. Feld⁴⁶, K. Klein⁴⁶, M. Lipinski⁴⁶, D. Meuser⁴⁶, P. Nattland⁴⁶, V. Oppenländer⁴⁶, A. Pauls⁴⁶, D. Pérez Adán⁴⁶, N. Röwert⁴⁶, M. Teroerde⁴⁶, C. Daumann⁴⁷, S. Diekmann⁴⁷, A. Dodonova⁴⁷, N. Eich⁴⁷, D. Eliseev⁴⁷, F. Engelke⁴⁷, J. Erdmann⁴⁷, M. Erdmann⁴⁷, B. Fischer⁴⁷, T. Hebbeker⁴⁷, K. Hoepfner⁴⁷, F. Ivone⁴⁷, A. Jung⁴⁷, N. Kumar⁴⁷, M.y. Lee⁴⁷, F. Mausolf⁴⁷, M. Merschmeyer⁴⁷, A. Meyer⁴⁷, F. Nowotny⁴⁷, A. Pozdnyakov⁴⁷, W. Redjeb⁴⁷, H. Reithler⁴⁷, U. Sarkar⁴⁷, V. Sarkisovi⁴⁷, A. Schmidt⁴⁷, C. Seth⁴⁷, A. Sharma⁴⁷, J.L. Spah⁴⁷, V. Vaulin⁴⁷, S. Zaleski⁴⁷, M.R. Beckers⁴⁸, C. Dziwok⁴⁸, G. Flügge⁴⁸, N. Hoefflich⁴⁸, T. Kress⁴⁸, A. Nowack⁴⁸, O. Pooth⁴⁸, A. Stahl⁴⁸, A. Zotz⁴⁸, H. Aarup Petersen⁴⁹, A. Abel⁴⁹, M. Aldaya Martin⁴⁹, J. Alimena⁴⁹, S. Amoroso⁴⁹, Y. An⁴⁹, I. Andreev⁴⁹, J. Bach⁴⁹, S. Baxter⁴⁹, M. Bayatmakou⁴⁹, H. Becerril Gonzalez⁴⁹, O. Behnke⁴⁹, A. Belvedere⁴⁹, F. Blekman^{49,v}, K. Borrás^{49,w}, A. Campbell⁴⁹, S. Chatterjee⁴⁹, L.X. Coll Saravia⁴⁹, G. Eckerlin⁴⁹, D. Eckstein⁴⁹, E. Gallo^{49,v}, A. Geiser⁴⁹, V. Guglielmi⁴⁹, M. Guthoff⁴⁹, A. Hinzmann⁴⁹, L. Jeppe⁴⁹, M. Kasemann⁴⁹, C. Kleinwort⁴⁹, R. Kogler⁴⁹, M. Komm⁴⁹, D. Krücker⁴⁹, W. Lange⁴⁹, D. Leyva Pernia⁴⁹, K.-Y. Lin⁴⁹, K. Lipka^{49,x}, W. Lohmann^{49,y}, J. Malvaso⁴⁹, R. Mankel⁴⁹, I.-A. Melzer-Pellmann⁴⁹, M. Mendizabal Morentin⁴⁹, A.B. Meyer⁴⁹, G. Milella⁴⁹, K. Moral Figueroa⁴⁹, A. Mussgiller⁴⁹, L.P. Nair⁴⁹, J. Niedziela⁴⁹, A. Nürnberg⁴⁹, J. Park⁴⁹, E. Ranken⁴⁹, A. Raspereza⁴⁹, D. Rastorguev⁴⁹, L. Rygaard⁴⁹,

M. Scham^{49,z,aa}, S. Schnake^{49,w}, C. Schwanenberger^{49,v}, P. Schütze⁴⁹, D. Selivanova⁴⁹, K. Sharko⁴⁹, M. Shchedrolosiev⁴⁹, D. Stafford⁴⁹, M. Torkian⁴⁹, F. Vazzoler⁴⁹, A. Ventura Barroso⁴⁹, R. Walsh⁴⁹, D. Wang⁴⁹, Q. Wang⁴⁹, K. Wichmann⁴⁹, L. Wiens^{49,w}, C. Wissing⁴⁹, Y. Yang⁴⁹, S. Zakharov⁴⁹, A. Zimmermann Castro Santos⁴⁹, A.R. Alves Andrade⁵⁰, M. Antonello⁵⁰, S. Bollweg⁵⁰, M. Bonanomi⁵⁰, K. El Morabit⁵⁰, Y. Fischer⁵⁰, M. Frahm⁵⁰, E. Garutti⁵⁰, A. Grohsjean⁵⁰, A.A. Guvenli⁵⁰, J. Haller⁵⁰, D. Hundhausen⁵⁰, G. Kasieczka⁵⁰, P. Keicher⁵⁰, R. Klanner⁵⁰, W. Korcari⁵⁰, T. Kramer⁵⁰, C.c. Kuo⁵⁰, F. Labe⁵⁰, J. Lange⁵⁰, A. Lobanov⁵⁰, L. Moureaux⁵⁰, A. Nigamova⁵⁰, K. Nikolopoulos⁵⁰, A. Paasch⁵⁰, K.J. Pena Rodriguez⁵⁰, N. Prouvost⁵⁰, B. Raciti⁵⁰, M. Rieger⁵⁰, D. Savoie⁵⁰, P. Schleper⁵⁰, M. Schröder⁵⁰, J. Schwandt⁵⁰, M. Sommerhalder⁵⁰, H. Stadie⁵⁰, G. Steinbrück⁵⁰, R. Ward⁵⁰, B. Wiederspan⁵⁰, M. Wolf⁵⁰, S. Brommer⁵¹, E. Butz⁵¹, Y.M. Chen⁵¹, T. Chwalek⁵¹, A. Dierlamm⁵¹, G.G. Dincer⁵¹, U. Elicabuk⁵¹, N. Faltermann⁵¹, M. Giffels⁵¹, A. Gottmann⁵¹, F. Hartmann^{51,ab}, M. Horzela⁵¹, F. Hummer⁵¹, U. Husemann⁵¹, J. Kieseler⁵¹, M. Klute⁵¹, R. Kunnilan Muhammed Rafeek⁵¹, O. Lavoryk⁵¹, J.M. Lawhorn⁵¹, A. Lintuluoto⁵¹, S. Maier⁵¹, M. Mormile⁵¹, Th. Müller⁵¹, E. Pfeffer⁵¹, M. Presilla⁵¹, G. Quast⁵¹, K. Rabbertz⁵¹, B. Regnery⁵¹, R. Schmieder⁵¹, N. Shadskiy⁵¹, I. Shvetsov⁵¹, H.J. Simonis⁵¹, L. Sowa⁵¹, L. Stockmeier⁵¹, K. Tauqeer⁵¹, M. Toms⁵¹, B. Topko⁵¹, N. Trevisani⁵¹, C. Verstege⁵¹, T. Voigtländer⁵¹, R.F. Von Cube⁵¹, J. Von Den Driesch⁵¹, M. Wassmer⁵¹, R. Wolf⁵¹, W.D. Zeuner⁵¹, X. Zuo⁵¹, G. Anagnostou⁵², G. Daskalakis⁵², A. Kyriakis⁵², G. Melachroinos⁵³, Z. Painesis⁵³, I. Paraskevas⁵³, N. Saoulidou⁵³, K. Theofilatos⁵³, E. Tziaferi⁵³, E. Tzovara⁵³, K. Vellidis⁵³, I. Zisopoulos⁵³, T. Chatzistavrou⁵⁴, G. Karapostoli⁵⁴, K. Kousouris⁵⁴, E. Siamarkou⁵⁴, G. Tsipolitis⁵⁴, I. Bestintzanos⁵⁵, I. Evangelou⁵⁵, C. Foudas⁵⁵, P. Katsoulis⁵⁵, P. Kokkas⁵⁵, P.G. Kosmoglou Kioseoglou⁵⁵, N. Manthos⁵⁵, I. Papadopoulos⁵⁵, J. Strologas⁵⁵, D. Druzhkin⁵⁶, C. Hajdu⁵⁶, D. Horvath^{56,ac,ad}, K. Márton⁵⁶, A.J. Rádl^{56,ae}, F. Sikler⁵⁶, V. Veszpremi⁵⁶, M. Csanád⁵⁷, K. Farkas⁵⁷, A. Fehérkuti^{57,af}, M.M.A. Gadallah^{57,ag}, Á. Kadlecik⁵⁷, M. León Coello⁵⁷, G. Pásztor⁵⁷, G.I. Veres⁵⁷, B. Ujvari⁵⁸, G. Zilizi⁵⁸, G. Bencze⁵⁹, S. Czellar⁵⁹, J. Molnar⁵⁹, Z. Szillasi⁵⁹, T. Csorgo^{60,af}, F. Nemes^{60,af}, T. Novak⁶⁰, I. Szanyi^{60,ah}, S. Bansal⁶¹, S.B. Beri⁶¹, V. Bhatnagar⁶¹, G. Chaudhary⁶¹, S. Chauhan⁶¹, N. Dhingra^{61,ai}, A. Kaur⁶¹, A. Kaur⁶¹, H. Kaur⁶¹, M. Kaur⁶¹, S. Kumar⁶¹, T. Sheokand⁶¹, J.B. Singh⁶¹, A. Singla⁶¹, A. Bhardwaj⁶², A. Chhetri⁶², B.C. Choudhary⁶², A. Kumar⁶², A. Kumar⁶², M. Naimuddin⁶², S. Phor⁶², K. Ranjan⁶², M.K. Saini⁶², P. Palni⁶³, S. Acharya^{64,aj}, B. Gomber⁶⁴, B. Sahu^{64,aj}, S. Mukherjee⁶⁵, S. Bhattacharya⁶⁶, S. Das Gupta⁶⁶, S. Dutta⁶⁶, S. Dutta⁶⁶, S. Sarkar⁶⁶, M.M. Ameen⁶⁷, P.K. Behera⁶⁷, S. Chatterjee⁶⁷, G. Dash⁶⁷, A. Dattamunsi⁶⁷, P. Jana⁶⁷, P. Kalbhor⁶⁷, S. Kamble⁶⁷, J.R. Komaragiri^{67,ak}, T. Mishra⁶⁷, P.R. Pujahari⁶⁷, A.K. Sikdar⁶⁷, R.K. Singh⁶⁷, P. Verma⁶⁷, S. Verma⁶⁷, A. Vijay⁶⁷, B.K. Sirasva⁶⁸, L. Bhatt⁶⁹, S. Dugad⁶⁹, G.B. Mohanty⁶⁹, M. Shelake⁶⁹, P. Suryadevara⁶⁹, A. Bala⁷⁰, S. Banerjee⁷⁰, S. Barman^{70,al}, R.M. Chatterjee⁷⁰, M. Guchait⁷⁰, Sh. Jain⁷⁰, A. Jaiswal⁷⁰, B.M. Joshi⁷⁰, S. Kumar⁷⁰, M. Maity^{70,al}, G. Majumder⁷⁰, K. Mazumdar⁷⁰, S. Parolia⁷⁰, R. Saxena⁷⁰, A. Thachayath⁷⁰, S. Bahinipati^{71,am}, D. Maity^{71,an}, P. Mal⁷¹, K. Naskar^{71,an}, A. Nayak^{71,an}, S. Nayak⁷¹, K. Pal⁷¹, R. Raturi⁷¹, P. Sadangi⁷¹, S.K. Swain⁷¹,

S. Varghese ^{71,an}, D. Vats ^{71,an}, A. Alpina ⁷², S. Dube ⁷², P. Hazarika ⁷², B. Kansal ⁷², A. Laha ⁷², R. Sharma ⁷², S. Sharma ⁷², K.Y. Vaish ⁷², S. Ghosh ⁷³, H. Bakhshiansohi ^{74,ao}, A. Jafari ^{74,ap}, V. Sedighzadeh Dalavi ⁷⁴, M. Zeinali ^{74,aq}, S. Bashiri ⁷⁵, S. Chenarani ^{75,ar}, S.M. Etesami ⁷⁵, Y. Hosseini ⁷⁵, M. Khakzad ⁷⁵, E. Khazaie ⁷⁵, M. Mohammadi Najafabadi ⁷⁵, S. Tizchang ^{75,as}, M. Felcini ⁷⁶, M. Grunewald ⁷⁶, M. Abbrescia ^{77a,77b}, M. Barbieri ^{77a,77b}, M. Buonsante ^{77a,77b}, A. Colaleo ^{77a,77b}, D. Creanza ^{77a,77c}, N. De Filippis ^{77a,77c}, M. De Palma ^{77a,77b}, W. Elmetenawee ^{77a,77b,at}, N. Ferrara ^{77a,77c}, L. Fiore ^{77a}, L. Longo ^{77a}, M. Louka ^{77a,77b}, G. Maggi ^{77a,77c}, M. Maggi ^{77a}, I. Margjeka ^{77a}, V. Mastrapasqua ^{77a,77b}, S. My ^{77a,77b}, F. Nenna ^{77a,77b}, S. Nuzzo ^{77a,77b}, A. Pellecchia ^{77a,77b}, A. Pompili ^{77a,77b}, G. Pugliese ^{77a,77c}, R. Radogna ^{77a,77b}, D. Ramos ^{77a}, A. Ranieri ^{77a}, L. Silvestris ^{77a}, F.M. Simone ^{77a,77c}, A. Stamerra ^{77a,77b}, Ü. Sözbilir ^{77a}, D. Troiano ^{77a,77b}, R. Venditti ^{77a,77b}, P. Verwilligen ^{77a}, A. Zaza ^{77a,77b}, C. Battilana ^{78a,78b}, D. Bonacorsi ^{78a,78b}, P. Capiluppi ^{78a,78b}, F.R. Cavallo ^{78a}, M. Cuffiani ^{78a,78b}, G.M. Dallavalle ^{78a}, T. Diotallevi ^{78a,78b}, F. Fabbri ^{78a}, A. Fanfani ^{78a,78b}, R. Farinelli ^{78a}, D. Fasanella ^{78a}, P. Giacomelli ^{78a}, C. Grandi ^{78a}, L. Guiducci ^{78a,78b}, S. Lo Meo ^{78a,au}, M. Lorusso ^{78a,78b}, L. Lunerti ^{78a}, S. Marcellini ^{78a}, G. Masetti ^{78a}, F.L. Navarra ^{78a,78b}, G. Paggi ^{78a,78b}, A. Perrotta ^{78a}, F. Primavera ^{78a,78b}, A.M. Rossi ^{78a,78b}, S. Rossi Tisbeni ^{78a,78b}, G.P. Siroli ^{78a,78b}, S. Costa ^{79a,79b,av}, A. Di Mattia ^{79a}, A. Lapertosa ^{79a}, R. Potenza ^{79a,79b}, A. Tricoli ^{79a,79b,av}, J. Altork ^{80a,80b}, P. Assiouras ^{80a}, G. Barbagli ^{80a}, G. Bardelli ^{80a}, M. Bartolini ^{80a,80b}, A. Calandri ^{80a,80b}, B. Camaiani ^{80a,80b}, A. Cassese ^{80a}, R. Ceccarelli ^{80a}, V. Ciulli ^{80a,80b}, C. Civinini ^{80a}, R. D'Alessandro ^{80a,80b}, L. Damenti ^{80a,80b}, E. Focardi ^{80a,80b}, T. Kello ^{80a}, G. Latino ^{80a,80b}, P. Lenzi ^{80a,80b}, M. Lizzo ^{80a}, M. Meschini ^{80a}, S. Paoletti ^{80a}, A. Papanastassiou ^{80a,80b}, G. Sguazzoni ^{80a}, L. Viliani ^{80a}, L. Benussi ⁸¹, S. Bianco ⁸¹, S. Meola ^{81,aw}, D. Piccolo ⁸¹, M. Alves Gallo Pereira ^{82a}, F. Ferro ^{82a}, E. Robutti ^{82a}, S. Tosi ^{82a,82b}, A. Benaglia ^{83a}, F. Brivio ^{83a}, V. Camagni ^{83a,83b}, F. Cetorelli ^{83a,83b}, F. De Guio ^{83a,83b}, M.E. Dinardo ^{83a,83b}, P. Dini ^{83a}, S. Gennai ^{83a}, R. Gerosa ^{83a,83b}, A. Ghezzi ^{83a,83b}, P. Govoni ^{83a,83b}, L. Guzzi ^{83a}, M.R. Kim ^{83a}, G. Lavizzari ^{83a,83b}, M.T. Lucchini ^{83a,83b}, M. Malberti ^{83a}, S. Malvezzi ^{83a}, A. Massironi ^{83a}, D. Menasce ^{83a}, L. Moroni ^{83a}, M. Paganoni ^{83a,83b}, S. Palluotto ^{83a,83b}, D. Pedrini ^{83a}, A. Perego ^{83a,83b}, G. Pizzati ^{83a,83b}, S. Ragazzi ^{83a,83b}, T. Tabarelli de Fatis ^{83a,83b}, S. Buontempo ^{84a}, C. Di Fraia ^{84a,84b}, F. Fabozzi ^{84a,84c}, L. Favilla ^{84a,84d}, A.O.M. Iorio ^{84a,84b}, L. Lista ^{84a,84b,ax}, P. Paolucci ^{84a,ab}, B. Rossi ^{84a}, P. Azzi ^{85a}, N. Bacchetta ^{85a,ay}, M. Benettoni ^{85a}, P. Bortignon ^{85a,85c}, G. Bortolato ^{85a,85b}, A.C.M. Bulla ^{85a,85c}, R. Carlin ^{85a,85b}, P. Checchia ^{85a}, T. Dorigo ^{85a,az}, F. Gasparini ^{85a,85b}, U. Gasparini ^{85a,85b}, S. Giorgetti ^{85a}, E. Lusiani ^{85a}, M. Margoni ^{85a,85b}, A.T. Meneguzzo ^{85a,85b}, J. Pazzini ^{85a,85b}, P. Ronchese ^{85a,85b}, R. Rossin ^{85a,85b}, M. Tosi ^{85a,85b}, A. Triossi ^{85a,85b}, S. Ventura ^{85a}, M. Zanetti ^{85a,85b}, P. Zotto ^{85a,85b}, A. Zucchetta ^{85a,85b}, G. Zumerle ^{85a,85b}, A. Braghieri ^{86a}, S. Calzaferri ^{86a}, P. Montagna ^{86a,86b}, M. Pelliccioni ^{86a}, V. Re ^{86a}, C. Riccardi ^{86a,86b}, P. Salvini ^{86a}, I. Vai ^{86a,86b}, P. Vitulo ^{86a,86b}, S. Ajmal ^{87a,87b}, M.E. Ascoti ^{87a,87b}, G.M. Bilei ^{87a}, C. Carrivale ^{87a,87b}, D. Ciangottini ^{87a,87b}, L. Della Penna ^{87a,87b}, L. Fanò ^{87a,87b,†}, V. Mariani ^{87a,87b}, M. Menichelli ^{87a}, F. Moscatelli ^{87a,ba}, A. Rossi ^{87a,87b}, A. Santocchia ^{87a,87b}, D. Spiga ^{87a}, T. Tedeschi ^{87a,87b}, C. Aimè ^{88a,88b}, C.A. Alexe ^{88a,88c}, P. Asenov ^{88a,88b}, P. Azzurri ^{88a}, G. Bagliesi ^{88a}, L. Bianchini ^{88a,88b}, T. Boccali ^{88a},

E. Bossini ^{88a}, D. Bruschini ^{88a,88c}, L. Calligaris ^{88a,88b}, R. Castaldi ^{88a}, F. Cattafesta ^{88a,88c},
 M.A. Ciocci ^{88a,88d}, M. Cipriani ^{88a,88b}, R. Dell’Orso ^{88a}, S. Donato ^{88a,88b}, R. Forti ^{88a,88b},
 A. Giassi ^{88a}, F. Ligabue ^{88a,88c}, A.C. Marini ^{88a,88b}, D. Matos Figueiredo ^{88a},
 A. Messineo ^{88a,88b}, S. Mishra ^{88a}, V.K. Muraleedharan Nair Bindhu ^{88a,88b}, S. Nandan ^{88a},
 F. Palla ^{88a}, M. Riggirello ^{88a,88c}, A. Rizzi ^{88a,88b}, G. Rolandi ^{88a,88c},
 S. Roy Chowdhury ^{88a,bb}, T. Sarkar ^{88a}, A. Scribano ^{88a}, P. Solanki ^{88a,88b}, P. Spagnolo ^{88a},
 F. Tenchini ^{88a,88b}, R. Tenchini ^{88a}, G. Tonelli ^{88a,88b}, N. Turini ^{88a,88d}, F. Vaselli ^{88a,88c},
 A. Venturi ^{88a}, P.G. Verdini ^{88a}, P. Akrap ^{89a,89b}, C. Basile ^{89a,89b}, S.C. Behera ^{89a},
 F. Cavallari ^{89a}, L. Cunqueiro Mendez ^{89a,89b}, F. De Raggi ^{89a,89b}, D. Del Re ^{89a,89b},
 E. Di Marco ^{89a}, M. Diemoz ^{89a}, F. Errico ^{89a}, L. Frosina ^{89a,89b}, R. Gargiulo ^{89a,89b},
 B. Harikrishnan ^{89a,89b}, F. Lombardi ^{89a,89b}, E. Longo ^{89a,89b}, L. Martikainen ^{89a,89b},
 J. Mijuskovic ^{89a,89b}, G. Organtini ^{89a,89b}, N. Palmeri ^{89a,89b}, R. Paramatti ^{89a,89b},
 S. Rahatlou ^{89a,89b}, C. Rovelli ^{89a}, F. Santanastasio ^{89a,89b}, L. Soffi ^{89a}, V. Vladimirov ^{89a,89b},
 N. Amapane ^{90a,90b}, R. Arcidiacono ^{90a,90c}, S. Argiro ^{90a,90b}, M. Arneodo ^{90a,90c},
 N. Bartosik ^{90a,90c}, R. Bellan ^{90a,90b}, A. Bellora ^{90a,90b}, C. Biino ^{90a}, C. Borca ^{90a,90b},
 N. Cartiglia ^{90a}, M. Costa ^{90a,90b}, G. Cotto ^{90a,90b}, R. Covarelli ^{90a,90b}, N. Demaria ^{90a},
 M. Ferrero ^{90a}, L. Finco ^{90a}, M. Grippo ^{90a,90b}, B. Kiani ^{90a,90b}, L. Lanteri ^{90a,90b},
 F. Legger ^{90a}, F. Luongo ^{90a,90b}, C. Mariotti ^{90a}, S. Maselli ^{90a}, A. Mecca ^{90a,90b},
 L. Menzio ^{90a,90b}, P. Meridiani ^{90a}, E. Migliore ^{90a,90b}, M. Monteno ^{90a}, M.M. Obertino ^{90a,90b},
 G. Ortona ^{90a}, L. Pacher ^{90a,90b}, N. Pastrone ^{90a}, M. Ruspa ^{90a,90c}, F. Siviero ^{90a,90b},
 V. Sola ^{90a,90b}, A. Solano ^{90a,90b}, C. Tarricone ^{90a,90b}, D. Trocino ^{90a}, G. Umoret ^{90a,90b},
 R. White ^{90a,90b}, J. Babbar ^{91a,91b}, S. Belforte ^{91a}, V. Candelise ^{91a,91b}, M. Casarsa ^{91a},
 F. Cossutti ^{91a}, K. De Leo ^{91a}, G. Della Ricca ^{91a,91b}, R. Delli Gatti ^{91a,91b}, S. Dogra ⁹²,
 J. Hong ⁹², J. Kim ⁹², T. Kim ⁹², D. Lee ⁹², H. Lee ⁹², J. Lee ⁹², S.W. Lee ⁹², C.S. Moon ⁹²,
 Y.D. Oh ⁹², S. Sekmen ⁹², B. Tae ⁹², Y.C. Yang ⁹², M.S. Kim ⁹³, G. Bak ⁹⁴, P. Gwak ⁹⁴,
 H. Kim ⁹⁴, D.H. Moon ⁹⁴, J. Seo ⁹⁴, E. Asilar ⁹⁵, F. Carnevali ⁹⁵, J. Choi ^{95,bc}, T.J. Kim ⁹⁵,
 Y. Ryou ⁹⁵, S. Ha ⁹⁶, S. Han ⁹⁶, B. Hong ⁹⁶, J. Kim ⁹⁶, K. Lee ⁹⁶, K.S. Lee ⁹⁶, S. Lee ⁹⁶,
 J. Yoo ⁹⁶, J. Goh ⁹⁷, J. Shin ⁹⁷, S. Yang ⁹⁷, Y. Kang ⁹⁸, H. S. Kim ⁹⁸, Y. Kim ⁹⁸,
 S. Lee ⁹⁸, J. Almond ⁹⁹, J.H. Bhyun ⁹⁹, J. Choi ⁹⁹, J. Choi ⁹⁹, W. Jun ⁹⁹, H. Kim ⁹⁹, J. Kim ⁹⁹,
 T. Kim ⁹⁹, Y. Kim ⁹⁹, Y.W. Kim ⁹⁹, S. Ko ⁹⁹, H. Lee ⁹⁹, J. Lee ⁹⁹, J. Lee ⁹⁹, B.H. Oh ⁹⁹,
 S.B. Oh ⁹⁹, J. Shin ⁹⁹, U.K. Yang ⁹⁹, I. Yoon ⁹⁹, W. Jang ¹⁰⁰, D.Y. Kang ¹⁰⁰, D. Kim ¹⁰⁰,
 S. Kim ¹⁰⁰, B. Ko ¹⁰⁰, J.S.H. Lee ¹⁰⁰, Y. Lee ¹⁰⁰, I.C. Park ¹⁰⁰, Y. Roh ¹⁰⁰, I.J. Watson ¹⁰⁰,
 G. Cho ¹⁰¹, K. Hwang ¹⁰¹, B. Kim ¹⁰¹, S. Kim ¹⁰¹, K. Lee ¹⁰¹, H.D. Yoo ¹⁰¹, Y. Lee ¹⁰²,
 I. Yu ¹⁰², T. Beyrouthy ¹⁰³, Y. Gharbia ¹⁰³, F. Alazemi ¹⁰⁴, K. Dreimanis ¹⁰⁵,
 O.M. Eberlins ¹⁰⁵, A. Gaile ¹⁰⁵, C. Munoz Diaz ¹⁰⁵, D. Osite ¹⁰⁵, G. Pikurs ¹⁰⁵, R. Plese ¹⁰⁵,
 A. Potrebko ¹⁰⁵, M. Seidel ¹⁰⁵, D. Sidiropoulos Kontos ¹⁰⁵, N.R. Strautnieks ¹⁰⁶,
 M. Ambrozias ¹⁰⁷, A. Juodagalvis ¹⁰⁷, S. Nargelas ¹⁰⁷, A. Rinkevicius ¹⁰⁷, G. Tamulaitis ¹⁰⁷,
 I. Yusuff ^{108,bd}, Z. Zolkapli ¹⁰⁸, J.F. Benitez ¹⁰⁹, A. Castaneda Hernandez ¹⁰⁹,
 A. Cota Rodriguez ¹⁰⁹, L.E. Cuevas Picos ¹⁰⁹, H.A. Encinas Acosta ¹⁰⁹, L.G. Gallegos Maríñez ¹⁰⁹,
 J.A. Murillo Quijada ¹⁰⁹, L. Valencia Palomo ¹⁰⁹, G. Ayala ¹¹⁰, H. Castilla-Valdez ¹¹⁰,
 H. Crotte Ledesma ¹¹⁰, R. Lopez-Fernandez ¹¹⁰, J. Mejia Guisao ¹¹⁰, R. Reyes-Almanza ¹¹⁰,
 A. Sánchez Hernández ¹¹⁰, C. Oropeza Barrera ¹¹¹, D.L. Ramirez Guadarrama ¹¹¹,
 M. Ramírez García ¹¹¹, I. Bautista ¹¹², F.E. Neri Huerta ¹¹², I. Pedraza ¹¹²,

H.A. Salazar Ibarguen¹¹², C. Uribe Estrada¹¹², I. Bujanja¹¹³, N. Raicevic¹¹³,
P.H. Butler¹¹⁴, A. Ahmad¹¹⁵, M.I. Asghar¹¹⁵, A. Awais¹¹⁵, M.I.M. Awan¹¹⁵,
W.A. Khan¹¹⁵, V. Avati¹¹⁶, L. Forthomme¹¹⁶, L. Grzanka¹¹⁶, M. Malawski¹¹⁶,
K. Piotrkowski¹¹⁶, M. Bluj¹¹⁷, M. Górski¹¹⁷, M. Kazana¹¹⁷, M. Szleper¹¹⁷,
P. Zalewski¹¹⁷, K. Bunkowski¹¹⁸, K. Doroba¹¹⁸, A. Kalinowski¹¹⁸, M. Konecki¹¹⁸,
J. Krolikowski¹¹⁸, A. Muhammad¹¹⁸, P. Fokow¹¹⁹, K. Pozniak¹¹⁹, W. Zabolotny¹¹⁹,
M. Araujo¹²⁰, D. Bastos¹²⁰, C. Beirão Da Cruz E Silva¹²⁰, A. Boletti¹²⁰, M. Bozzo¹²⁰,
T. Camporesi¹²⁰, G. Da Molin¹²⁰, M. Gallinaro¹²⁰, J. Hollar¹²⁰, N. Leonardo¹²⁰,
G.B. Marozzo¹²⁰, A. Petrilli¹²⁰, M. Pisano¹²⁰, J. Seixas¹²⁰, J. Varela¹²⁰, J.W. Wulff¹²⁰,
P. Adzic¹²¹, L. Markovic¹²¹, P. Milenovic¹²¹, V. Milosevic¹²¹, D. Devetak¹²²,
M. Dordevic¹²², J. Milosevic¹²², L. Nadder¹²², V. Rekovic¹²², M. Stojanovic¹²²,
M. Alcalde Martinez¹²³, J. Alcaraz Maestre¹²³, J.A. Brochero Cifuentes¹²³, M. Cepeda¹²³,
M. Cerrada¹²³, N. Colino¹²³, B. De La Cruz¹²³, A. Delgado Peris¹²³,
A. Escalante Del Valle¹²³, Cristina F. Bedoya¹²³, D. Fernández Del Val¹²³,
J.P. Fernández Ramos¹²³, J. Flix¹²³, M.C. Fouz¹²³, M. Gonzalez Hernandez¹²³,
O. Gonzalez Lopez¹²³, S. Goy Lopez¹²³, J.M. Hernandez¹²³, M.I. Josa¹²³,
J. Llorente Merino¹²³, Oliver M. Carretero¹²³, C. Martin Perez¹²³, E. Martin Viscasillas¹²³,
D. Moran¹²³, C. M. Morcillo Perez¹²³, Á. Navarro Tobar¹²³, R. Paz Herrera¹²³,
C. Perez Dengra¹²³, J. Puerta Pelayo¹²³, A. Pérez-Calero Yzquierdo¹²³, I. Redondo¹²³,
J. Vazquez Escobar¹²³, J.F. de Trocóniz¹²⁴, B. Alvarez Gonzalez¹²⁵, J. Ayllon Torresano¹²⁵,
A. Cardini¹²⁵, J. Cuevas¹²⁵, J. Del Riego Badas¹²⁵, D. Estrada Acevedo¹²⁵,
J. Fernandez Menendez¹²⁵, S. Folgueras¹²⁵, I. Gonzalez Caballero¹²⁵, P. Leguina¹²⁵,
M. Obeso Menendez¹²⁵, E. Palencia Cortezon¹²⁵, J. Prado Pico¹²⁵, A. Soto Rodríguez¹²⁵,
C. Vico Villalba¹²⁵, P. Vischia¹²⁵, S. Blanco Fernández¹²⁶, I.J. Cabrillo¹²⁶, A. Calderon¹²⁶,
J. Duarte Campderros¹²⁶, M. Fernandez¹²⁶, G. Gomez¹²⁶, C. Lasasoa García¹²⁶,
R. Lopez Ruiz¹²⁶, C. Martinez Rivero¹²⁶, P. Martinez Ruiz del Arbol¹²⁶, F. Matorras¹²⁶,
P. Matorras Cuevas¹²⁶, E. Navarrete Ramos¹²⁶, J. Piedra Gomez¹²⁶,
C. Quintana San Emeterio¹²⁶, L. Scodellaro¹²⁶, I. Vila¹²⁶, R. Vilar Cortabitarte¹²⁶,
J.M. Vizán García¹²⁶, D.D.C. Wickramaratna¹²⁷, B. Kailasapathy^{127,be},
W.G.D. Dharmaratna^{128,bf}, K. Liyanage¹²⁸, N. Perera¹²⁸, D. Abbaneo¹²⁹, C. Amendola¹²⁹,
R. Ardino¹²⁹, E. Auffray¹²⁹, J. Baechler¹²⁹, D. Barney¹²⁹, J. Bendavid¹²⁹, M. Bianco¹²⁹,
A. Bocci¹²⁹, L. Borgonovi¹²⁹, C. Botta¹²⁹, A. Bragagnolo¹²⁹, C.E. Brown¹²⁹,
C. Caillol¹²⁹, G. Cerminara¹²⁹, P. Connor¹²⁹, D. d’Enterria¹²⁹, A. Dabrowski¹²⁹,
A. David¹²⁹, A. De Roeck¹²⁹, M.M. Defranchis¹²⁹, M. Deile¹²⁹, M. Dobson¹²⁹,
P.J. Fernández Manteca¹²⁹, B.A. Fontana Santos Alves¹²⁹, W. Funk¹²⁹, A. Gaddi¹²⁹,
S. Giani¹²⁹, D. Gigi¹²⁹, K. Gill¹²⁹, F. Glege¹²⁹, M. Glowacki¹²⁹, A. Gruber¹²⁹,
J. Hegeman¹²⁹, J.K. Heikkilä¹²⁹, R. Hofsaess¹²⁹, B. Huber¹²⁹, T. James¹²⁹, P. Janot¹²⁹,
O. Kaluzinska¹²⁹, O. Karacheban^{129,y}, G. Karathanasis¹²⁹, S. Laurila¹²⁹, P. Lecoq¹²⁹,
E. Leutgeb¹²⁹, C. Lourenço¹²⁹, A.-M. Lyon¹²⁹, M. Magherini¹²⁹, L. Malgeri¹²⁹,
M. Mannelli¹²⁹, A. Mehta¹²⁹, F. Meijers¹²⁹, J.A. Merlin¹²⁹, S. Mersi¹²⁹, E. Meschi¹²⁹,
M. Migliorini¹²⁹, F. Monti¹²⁹, F. Moortgat¹²⁹, M. Mulders¹²⁹, M. Musich¹²⁹,
I. Neutelings¹²⁹, S. Orfanelli¹²⁹, F. Pantaleo¹²⁹, M. Pari¹²⁹, G. Petrucciani¹²⁹,
A. Pfeiffer¹²⁹, M. Pierini¹²⁹, M. Pitt¹²⁹, H. Qu¹²⁹, D. Rabady¹²⁹, A. Reimers¹²⁹,

B. Ribeiro Lopes ¹²⁹, F. Riti ¹²⁹, P. Rosado ¹²⁹, M. Rovere ¹²⁹, H. Sakulin ¹²⁹,
 R. Salvatico ¹²⁹, S. Sanchez Cruz ¹²⁹, S. Scarfi ¹²⁹, M. Selvaggi ¹²⁹, A. Sharma ¹²⁹,
 K. Shchelina ¹²⁹, P. Silva ¹²⁹, P. Sphicas ^{129,bg}, A.G. Stahl Leiton ¹²⁹, A. Steen ¹²⁹,
 S. Summers ¹²⁹, D. Treille ¹²⁹, P. Tropea ¹²⁹, E. Vernazza ¹²⁹, J. Wanczyk ^{129,bh},
 S. Wuchterl ¹²⁹, M. Zarucki ¹²⁹, P. Zehetner ¹²⁹, P. Zejdl ¹²⁹, G. Zevi Della Porta ¹²⁹,
 T. Bevilacqua ^{130,bi}, L. Caminada ^{130,bi}, W. Erdmann ¹³⁰, R. Horisberger ¹³⁰, Q. Ingram ¹³⁰,
 H.C. Kaestli ¹³⁰, D. Kotlinski ¹³⁰, C. Lange ¹³⁰, U. Langenegger ¹³⁰, L. Nochte ^{130,bi},
 T. Rohe ¹³⁰, A. Samalan ¹³⁰, T.K. Aarrestad ¹³¹, M. Backhaus ¹³¹, G. Bonomelli ¹³¹,
 C. Cazzaniga ¹³¹, K. Datta ¹³¹, P. De Bryas Dexmiers D'Archiacchiac ^{131,bh}, A. De Cosa ¹³¹,
 G. Dissertori ¹³¹, M. Dittmar ¹³¹, M. Donegà ¹³¹, F. Eble ¹³¹, K. Gedia ¹³¹, F. Glessgen ¹³¹,
 C. Grab ¹³¹, T.G. Harte ¹³¹, N. Härringer ¹³¹, W. Lustermann ¹³¹, M. Malucchi ¹³¹,
 R.A. Manzoni ¹³¹, L. Marchese ¹³¹, A. Mascellani ^{131,bh}, F. Nessi-Tedaldi ¹³¹, F. Pauss ¹³¹,
 V. Perovic ¹³¹, B. Ristic ¹³¹, R. Seidita ¹³¹, J. Steggemann ^{131,bh}, A. Tarabini ¹³¹,
 D. Valsecchi ¹³¹, R. Wallny ¹³¹, C. Amsler ^{132,bj}, F. Bilandzija ¹³², P. Bärtzchi ¹³²,
 M.F. Canelli ¹³², G. Celotto ¹³², K. Cormier ¹³², M. Huwiler ¹³², W. Jin ¹³², A. Jofrehei ¹³²,
 B. Kilminster ¹³², T.H. Kwok ¹³², S. Leontsinis ¹³², V. Lukashenko ¹³², A. Macchiolo ¹³²,
 F. Meng ¹³², M. Missiroli ¹³², J. Motta ¹³², P. Robmann ¹³², M. Senger ¹³², E. Shokr ¹³²,
 F. Stäger ¹³², R. Tramontano ¹³², P. Viscone ¹³², D. Bhowmik ¹³³, C.M. Kuo ¹³³, P.K. Rout ¹³³,
 S. Taj ¹³³, P.C. Tiwari ^{133,ak}, L. Ceard ¹³⁴, K.F. Chen ¹³⁴, Z.g. Chen ¹³⁴, A. De Iorio ¹³⁴,
 W.-S. Hou ¹³⁴, T.h. Hsu ¹³⁴, Y.w. Kao ¹³⁴, S. Karmakar ¹³⁴, G. Kole ¹³⁴, Y.y. Li ¹³⁴,
 R.-S. Lu ¹³⁴, E. Paganis ¹³⁴, X.f. Su ¹³⁴, J. Thomas-Wilsker ¹³⁴, L.s. Tsai ¹³⁴, D. Tsionou ¹³⁴,
 H.y. Wu ¹³⁴, E. Yazgan ¹³⁴, C. Asawatangtrakuldee ¹³⁵, N. Srimanobhas ¹³⁵, Y. Maghrbi ¹³⁶,
 D. Agyel ¹³⁷, F. Dolek ¹³⁷, I. Dumanoglu ^{137,bk}, Y. Guler ^{137,bl}, E. Gurpinar Guler ^{137,bl},
 C. Isik ¹³⁷, O. Kara ¹³⁷, A. Kayis Topaksu ¹³⁷, Y. Komurcu ¹³⁷, G. Onengut ¹³⁷,
 K. Ozdemir ^{137,bm}, B. Tali ^{137,bn}, U.G. Tok ¹³⁷, E. Uslan ¹³⁷, I.S. Zorbakir ¹³⁷, S. Sen ¹³⁸,
 M. Yalvac ^{139,bo}, B. Akgun ¹⁴⁰, I.O. Atakisi ^{140,bp}, E. Gülmez ¹⁴⁰, M. Kaya ^{140,bq},
 O. Kaya ^{140,br}, M.A. Sarkisla ^{140,bs}, S. Tekten ^{140,bt}, D. Boncukcu ¹⁴¹, A. Cakir ¹⁴¹,
 K. Cankocak ^{141,bk,bu}, B. Haciosahinoglu ¹⁴², I. Hos ^{142,bv}, B. Kaynak ¹⁴², S. Ozkorucuklu ¹⁴²,
 O. Potok ¹⁴², H. Sert ¹⁴², C. Simsek ¹⁴², C. Zorbilmez ¹⁴², S. Cerci ¹⁴³, C. Dozen ^{143,bw},
 B. Isildak ^{143,bx}, E. Simsek ¹⁴³, D. Sunar Cerci ¹⁴³, T. Yetkin ^{143,bw}, A. Boyaryntsev ¹⁴⁴,
 O. Dadazhanova ¹⁴⁴, B. Grynyov ¹⁴⁴, L. Levchuk ¹⁴⁵, J.J. Brooke ¹⁴⁶, A. Bundock ¹⁴⁶,
 F. Bury ¹⁴⁶, E. Clement ¹⁴⁶, D. Cussans ¹⁴⁶, D. Dharmender ¹⁴⁶, H. Flacher ¹⁴⁶,
 J. Goldstein ¹⁴⁶, H.F. Heath ¹⁴⁶, M.-L. Holmberg ¹⁴⁶, L. Kreczko ¹⁴⁶, S. Paramesvaran ¹⁴⁶,
 L. Robertshaw ¹⁴⁶, M.S. Sanjrani ^{146,ao}, J. Segal ¹⁴⁶, V.J. Smith ¹⁴⁶, A.H. Ball ¹⁴⁷, K.W. Bell ¹⁴⁷,
 A. Belyaev ^{147,by}, C. Brew ¹⁴⁷, R.M. Brown ¹⁴⁷, D.J.A. Cockerill ¹⁴⁷, A. Elliot ¹⁴⁷,
 K.V. Ellis ¹⁴⁷, J. Gajownik ¹⁴⁷, K. Harder ¹⁴⁷, S. Harper ¹⁴⁷, J. Linacre ¹⁴⁷, K. Manolopoulos ¹⁴⁷,
 M. Moallemi ¹⁴⁷, D.M. Newbold ¹⁴⁷, E. Olaiya ¹⁴⁷, D. Petyt ¹⁴⁷, T. Reis ¹⁴⁷,
 A.R. Sahasransu ¹⁴⁷, G. Salvi ¹⁴⁷, T. Schuh ¹⁴⁷, C.H. Shepherd-Themistocleous ¹⁴⁷,
 I.R. Tomalin ¹⁴⁷, K.C. Whalen ¹⁴⁷, T. Williams ¹⁴⁷, I. Andreou ¹⁴⁸, R. Bainbridge ¹⁴⁸,
 P. Bloch ¹⁴⁸, O. Buchmuller ¹⁴⁸, C.A. Carrillo Montoya ¹⁴⁸, D. Colling ¹⁴⁸, I. Das ¹⁴⁸,
 P. Dauncey ¹⁴⁸, G. Davies ¹⁴⁸, M. Della Negra ¹⁴⁸, S. Fayer ¹⁴⁸, G. Fedi ¹⁴⁸, G. Hall ¹⁴⁸,
 H.R. Hoorani ¹⁴⁸, A. Howard ¹⁴⁸, G. Iles ¹⁴⁸, C.R. Knight ¹⁴⁸, P. Krueper ¹⁴⁸, J. Langford ¹⁴⁸,
 K.H. Law ¹⁴⁸, J. León Holgado ¹⁴⁸, L. Lyons ¹⁴⁸, A.-M. Magnan ¹⁴⁸, B. Maier ¹⁴⁸,

S. Mallios¹⁴⁸, A. Mastronikolis¹⁴⁸, M. Mieskolainen¹⁴⁸, J. Nash^{148,bz}, M. Pesaresi¹⁴⁸,
 P.B. Pradeep¹⁴⁸, B.C. Radburn-Smith¹⁴⁸, A. Richards¹⁴⁸, A. Rose¹⁴⁸, L. Russell¹⁴⁸,
 K. Savva¹⁴⁸, C. Seez¹⁴⁸, R. Shukla¹⁴⁸, A. Tapper¹⁴⁸, K. Uchida¹⁴⁸, G.P. Uttley¹⁴⁸,
 T. Virdee^{148,ab}, M. Vojinovic¹⁴⁸, N. Wardle¹⁴⁸, D. Winterbottom¹⁴⁸, J.E. Cole¹⁴⁹,
 A. Khan¹⁴⁹, P. Kyberd¹⁴⁹, I.D. Reid¹⁴⁹, S. Abdullin¹⁵⁰, A. Brinkerhoff¹⁵⁰, E. Collins¹⁵⁰,
 M.R. Darwish¹⁵⁰, J. Dittmann¹⁵⁰, K. Hatakeyama¹⁵⁰, V. Hegde¹⁵⁰, J. Hiltbrand¹⁵⁰,
 B. McMaster¹⁵⁰, J. Samudio¹⁵⁰, S. Sawant¹⁵⁰, C. Sutantawibul¹⁵⁰, J. Wilson¹⁵⁰,
 J.M. Hogan¹⁵¹, R. Bartek¹⁵², A. Dominguez¹⁵², S. Raj¹⁵², A.E. Simsek¹⁵², S.S. Yu¹⁵²,
 B. Bam¹⁵³, A. Buchot Perraguin¹⁵³, S. Campbell¹⁵³, R. Chudasama¹⁵³, S.I. Cooper¹⁵³,
 C. Crovella¹⁵³, G. Fidalgo¹⁵³, S.V. Gleyzer¹⁵³, A. Khukhunaishvili¹⁵³, K. Matchev¹⁵³,
 E. Pearson¹⁵³, C.U. Perez¹⁵³, P. Rumerio^{153,ca}, E. Usai¹⁵³, R. Yi¹⁵³, S. Cholak¹⁵⁴,
 G. De Castro¹⁵⁴, Z. Demiragli¹⁵⁴, C. Erice¹⁵⁴, C. Fangmeier¹⁵⁴, C. Fernandez Madrazo¹⁵⁴,
 E. Fontanesi¹⁵⁴, J. Fulcher¹⁵⁴, F. Golf¹⁵⁴, S. Jeon¹⁵⁴, J. O’Cain¹⁵⁴, I. Reed¹⁵⁴,
 J. Rohlf¹⁵⁴, K. Salyer¹⁵⁴, D. Sperka¹⁵⁴, D. Spitzbart¹⁵⁴, I. Suarez¹⁵⁴, A. Tsatsos¹⁵⁴,
 E. Wurtz¹⁵⁴, A.G. Zecchinelli¹⁵⁴, G. Barone¹⁵⁵, G. Benelli¹⁵⁵, D. Cutts¹⁵⁵, S. Ellis¹⁵⁵,
 L. Gouskos¹⁵⁵, M. Hadley¹⁵⁵, U. Heintz¹⁵⁵, K.W. Ho¹⁵⁵, T. Kwon¹⁵⁵, L. Lambrecht¹⁵⁵,
 G. Landsberg¹⁵⁵, K.T. Lau¹⁵⁵, J. Luo¹⁵⁵, S. Mondal¹⁵⁵, J. Roloff¹⁵⁵, T. Russell¹⁵⁵,
 S. Sagir^{155,cb}, X. Shen¹⁵⁵, M. Stamenkovic¹⁵⁵, N. Venkatasubramanian¹⁵⁵, S. Abbott¹⁵⁶,
 S. Baradia¹⁵⁶, B. Barton¹⁵⁶, R. Breedon¹⁵⁶, H. Cai¹⁵⁶,
 M. Calderon De La Barca Sanchez¹⁵⁶, E. Cannaert¹⁵⁶, M. Chertok¹⁵⁶, M. Citron¹⁵⁶,
 J. Conway¹⁵⁶, P.T. Cox¹⁵⁶, R. Erbacher¹⁵⁶, O. Kukral¹⁵⁶, G. Mocellin¹⁵⁶, S. Ostrom¹⁵⁶,
 I. Salazar Segovia¹⁵⁶, J.S. Tafoya Vargas¹⁵⁶, W. Wei¹⁵⁶, S. Yoo¹⁵⁶, K. Adamidis¹⁵⁷,
 M. Bachtis¹⁵⁷, D. Campos¹⁵⁷, R. Cousins¹⁵⁷, A. Datta¹⁵⁷, G. Flores Avila¹⁵⁷, J. Hauser¹⁵⁷,
 M. Ignatenko¹⁵⁷, M.A. Iqbal¹⁵⁷, T. Lam¹⁵⁷, Y.f. Lo¹⁵⁷, E. Manca¹⁵⁷,
 A. Nunez Del Prado¹⁵⁷, D. Saltzberg¹⁵⁷, V. Valuev¹⁵⁷, R. Clare¹⁵⁸, J.W. Gary¹⁵⁸,
 G. Hanson¹⁵⁸, A. Aportela¹⁵⁹, A. Arora¹⁵⁹, J.G. Branson¹⁵⁹, S. Cittolin¹⁵⁹,
 S. Cooperstein¹⁵⁹, B. D’Anzi¹⁵⁹, D. Diaz¹⁵⁹, J. Duarte¹⁵⁹, L. Giannini¹⁵⁹, Y. Gu¹⁵⁹,
 J. Guiang¹⁵⁹, V. Krutelyov¹⁵⁹, R. Lee¹⁵⁹, J. Letts¹⁵⁹, H. Li¹⁵⁹, M. Masciovecchio¹⁵⁹,
 F. Mokhtar¹⁵⁹, S. Mukherjee¹⁵⁹, M. Pieri¹⁵⁹, D. Primosch¹⁵⁹, M. Quinnan¹⁵⁹,
 V. Sharma¹⁵⁹, M. Tadel¹⁵⁹, E. Vourliotis¹⁵⁹, F. Würthwein¹⁵⁹, A. Yagil¹⁵⁹, Z. Zhao¹⁵⁹,
 A. Barzdukas¹⁶⁰, L. Brennan¹⁶⁰, C. Campagnari¹⁶⁰, S. Carron Montero^{160,cc},
 K. Downham¹⁶⁰, C. Grieco¹⁶⁰, M.M. Hussain¹⁶⁰, J. Incandela¹⁶⁰, M.W.K. Lai¹⁶⁰, A.J. Li¹⁶⁰,
 P. Masterson¹⁶⁰, J. Richman¹⁶⁰, S.N. Santpur¹⁶⁰, U. Sarica¹⁶⁰, R. Schmitz¹⁶⁰,
 F. Setti¹⁶⁰, J. Sheplock¹⁶⁰, D. Stuart¹⁶⁰, T.Á. Vámi¹⁶⁰, X. Yan¹⁶⁰, D. Zhang¹⁶⁰,
 A. Albert¹⁶¹, S. Bhattacharya¹⁶¹, A. Bornheim¹⁶¹, O. Cerri¹⁶¹, R. Kansal¹⁶¹, J. Mao¹⁶¹,
 H.B. Newman¹⁶¹, G. Reales Gutiérrez¹⁶¹, T. Sievert¹⁶¹, M. Spiropulu¹⁶¹, J.R. Vlimant¹⁶¹,
 R.A. Wynne¹⁶¹, S. Xie¹⁶¹, J. Alison¹⁶², S. An¹⁶², M. Cremonesi¹⁶², V. Dutta¹⁶²,
 E.Y. Ertorer¹⁶², T. Ferguson¹⁶², T.A. Gómez Espinosa¹⁶², A. Harilal¹⁶²,
 A. Kallil Tharayil¹⁶², M. Kanemura¹⁶², C. Liu¹⁶², M. Marchegiani¹⁶², P. Meiring¹⁶²,
 T. Mudholkar¹⁶², S. Murthy¹⁶², P. Palit¹⁶², K. Park¹⁶², M. Paulini¹⁶², A. Roberts¹⁶²,
 A. Sanchez¹⁶², W. Terrill¹⁶², J.P. Cumalat¹⁶³, W.T. Ford¹⁶³, A. Hart¹⁶³, S. Kwan¹⁶³,
 J. Pearkes¹⁶³, C. Savard¹⁶³, N. Schonbeck¹⁶³, K. Stenson¹⁶³, K.A. Ulmer¹⁶³,
 S.R. Wagner¹⁶³, N. Zipper¹⁶³, D. Zuolo¹⁶³, J. Alexander¹⁶⁴, X. Chen¹⁶⁴, J. Dickinson¹⁶⁴,

A. Duquette¹⁶⁴, J. Fan¹⁶⁴, X. Fan¹⁶⁴, J. Grassi¹⁶⁴, S. Hogan¹⁶⁴, P. Kotamnives¹⁶⁴,
 J. Monroy¹⁶⁴, G. Niendorf¹⁶⁴, M. Oshiro¹⁶⁴, J.R. Patterson¹⁶⁴, A. Ryd¹⁶⁴, J. Thom¹⁶⁴,
 P. Wittich¹⁶⁴, R. Zou¹⁶⁴, L. Zygala¹⁶⁴, M. Albrow¹⁶⁵, M. Alyari¹⁶⁵, O. Amram¹⁶⁵,
 G. Apollinari¹⁶⁵, A. Apresyan¹⁶⁵, L.A.T. Bauerdick¹⁶⁵, D. Berry¹⁶⁵, J. Berryhill¹⁶⁵,
 P.C. Bhat¹⁶⁵, K. Burkett¹⁶⁵, J.N. Butler¹⁶⁵, A. Canepa¹⁶⁵, G.B. Cerati¹⁶⁵,
 H.W.K. Cheung¹⁶⁵, F. Chlebana¹⁶⁵, C. Cosby¹⁶⁵, G. Cummings¹⁶⁵, I. Dutta¹⁶⁵,
 V.D. Elvira¹⁶⁵, J. Freeman¹⁶⁵, A. Gandrakota¹⁶⁵, Z. Gecse¹⁶⁵, L. Gray¹⁶⁵, D. Green¹⁶⁵,
 A. Grummer¹⁶⁵, S. Grünendahl¹⁶⁵, D. Guerrero¹⁶⁵, O. Gutsche¹⁶⁵, R.M. Harris¹⁶⁵,
 T.C. Herwig¹⁶⁵, J. Hirschauer¹⁶⁵, V. Innocente¹⁶⁵, B. Jayatilaka¹⁶⁵, S. Jindariani¹⁶⁵,
 M. Johnson¹⁶⁵, U. Joshi¹⁶⁵, B. Klima¹⁶⁵, K.H.M. Kwok¹⁶⁵, S. Lammel¹⁶⁵, C. Lee¹⁶⁵,
 D. Lincoln¹⁶⁵, R. Lipton¹⁶⁵, T. Liu¹⁶⁵, K. Maeshima¹⁶⁵, D. Mason¹⁶⁵, P. McBride¹⁶⁵,
 P. Merkel¹⁶⁵, S. Mrenna¹⁶⁵, S. Nahn¹⁶⁵, J. Ngadiuba¹⁶⁵, D. Noonan¹⁶⁵, S. Norberg¹⁶⁵,
 V. Papadimitriou¹⁶⁵, N. Pastika¹⁶⁵, K. Pedro¹⁶⁵, C. Pena^{165,cd}, C.E. Perez Lara¹⁶⁵,
 F. Ravera¹⁶⁵, A. Reinsvold Hall^{165,ce}, L. Ristori¹⁶⁵, M. Safdari¹⁶⁵, E. Sexton-Kennedy¹⁶⁵,
 N. Smith¹⁶⁵, A. Soha¹⁶⁵, L. Spiegel¹⁶⁵, S. Stoynev¹⁶⁵, J. Strait¹⁶⁵, L. Taylor¹⁶⁵,
 S. Tkaczyk¹⁶⁵, N.V. Tran¹⁶⁵, L. Uplegger¹⁶⁵, E.W. Vaandering¹⁶⁵, C. Wang¹⁶⁵, I. Zoi¹⁶⁵,
 C. Aruta¹⁶⁶, P. Avery¹⁶⁶, D. Bourilov¹⁶⁶, P. Chang¹⁶⁶, V. Cherepanov¹⁶⁶, R.D. Field¹⁶⁶,
 C. Huh¹⁶⁶, E. Koenig¹⁶⁶, M. Kolosova¹⁶⁶, J. Konigsberg¹⁶⁶, A. Korytov¹⁶⁶,
 G. Mitselmakher¹⁶⁶, K. Mohrman¹⁶⁶, A. Muthirakalayil Madhu¹⁶⁶, N. Rawal¹⁶⁶,
 S. Rosenzweig¹⁶⁶, V. Sulimov¹⁶⁶, Y. Takahashi¹⁶⁶, J. Wang¹⁶⁶, T. Adams¹⁶⁷,
 A. Al Kadhim¹⁶⁷, A. Askew¹⁶⁷, S. Bower¹⁶⁷, R. Goff¹⁶⁷, R. Hashmi¹⁶⁷, A. Hassani¹⁶⁷,
 R.S. Kim¹⁶⁷, T. Kolberg¹⁶⁷, G. Martinez¹⁶⁷, M. Mazza¹⁶⁷, H. Prosper¹⁶⁷, P.R. Prova¹⁶⁷,
 R. Yohay¹⁶⁷, B. Alsufyani¹⁶⁸, S. Butalla¹⁶⁸, S. Das¹⁶⁸, M. Hohlmann¹⁶⁸, M. Lavinsky¹⁶⁸,
 E. Yanes¹⁶⁸, M.R. Adams¹⁶⁹, N. Barnett¹⁶⁹, A. Baty¹⁶⁹, C. Bennett¹⁶⁹, R. Cavanaugh¹⁶⁹,
 R. Escobar Franco¹⁶⁹, O. Evdokimov¹⁶⁹, C.E. Gerber¹⁶⁹, H. Gupta¹⁶⁹, M. Hawksworth¹⁶⁹,
 A. Hingrajiya¹⁶⁹, D.J. Hofman¹⁶⁹, Z. Huang¹⁶⁹, J.h. Lee¹⁶⁹, C. Mills¹⁶⁹, S. Nanda¹⁶⁹,
 G. Nigmatkulov¹⁶⁹, B. Ozek¹⁶⁹, T. Phan¹⁶⁹, D. Pilipovic¹⁶⁹, R. Pradhan¹⁶⁹, E. Prifti¹⁶⁹,
 P. Roy¹⁶⁹, T. Roy¹⁶⁹, N. Singh¹⁶⁹, M.B. Tonjes¹⁶⁹, N. Varelas¹⁶⁹, M.A. Wadud¹⁶⁹,
 J. Yoo¹⁶⁹, M. Alhusseini¹⁷⁰, D. Blend¹⁷⁰, K. Dilsiz^{170,cf}, O.K. Köseyan¹⁷⁰,
 A. Mestvirishvili^{170,cg}, O. Neogi¹⁷⁰, H. Ogul^{170,ch}, Y. Onel¹⁷⁰, A. Penzo¹⁷⁰, C. Snyder¹⁷⁰,
 E. Tiras^{170,ci}, B. Blumenfeld¹⁷¹, J. Davis¹⁷¹, A.V. Gritsan¹⁷¹, L. Kang¹⁷¹,
 S. Kyriacou¹⁷¹, P. Maksimovic¹⁷¹, M. Roguljic¹⁷¹, S. Sekhar¹⁷¹, M.V. Srivastav¹⁷¹,
 M. Swartz¹⁷¹, A. Abreu¹⁷², L.F. Alcerro Alcerro¹⁷², J. Anguiano¹⁷², S. Arteaga Escatel¹⁷²,
 P. Baringer¹⁷², A. Bean¹⁷², R. Bhattacharya¹⁷², Z. Flowers¹⁷², D. Grove¹⁷², J. King¹⁷²,
 G. Krintiras¹⁷², M. Lazarovits¹⁷², C. Le Mahieu¹⁷², J. Marquez¹⁷², M. Murray¹⁷²,
 M. Nickel¹⁷², S. Popescu^{172,cj}, C. Rogan¹⁷², C. Royon¹⁷², S. Rudrabhatla¹⁷²,
 S. Sanders¹⁷², C. Smith¹⁷², G. Wilson¹⁷², B. Allmond¹⁷³, N. Islam¹⁷³, A. Ivanov¹⁷³,
 K. Kaadze¹⁷³, Y. Maravin¹⁷³, J. Natoli¹⁷³, G.G. Reddy¹⁷³, D. Roy¹⁷³, G. Sorrentino¹⁷³,
 A. Baden¹⁷⁴, A. Belloni¹⁷⁴, J. Bistany-riebman¹⁷⁴, S.C. Eno¹⁷⁴, N.J. Hadley¹⁷⁴,
 S. Jabeen¹⁷⁴, R.G. Kellogg¹⁷⁴, T. Koeth¹⁷⁴, B. Kronheim¹⁷⁴, S. Lascio¹⁷⁴, P. Major¹⁷⁴,
 A.C. Mignerey¹⁷⁴, C. Palmer¹⁷⁴, C. Papageorgakis¹⁷⁴, M.M. Paranjpe¹⁷⁴, E. Popova^{174,ck},
 A. Shevelev¹⁷⁴, L. Zhang¹⁷⁴, C. Baldenegro Barrera¹⁷⁵, H. Bossi¹⁷⁵, S. Bright-Thonney¹⁷⁵,
 I.A. Cali¹⁷⁵, Y.c. Chen¹⁷⁵, P.c. Chou¹⁷⁵, M. D'Alfonso¹⁷⁵, J. Eysermans¹⁷⁵, C. Freer¹⁷⁵,

G. Gomez-Ceballos¹⁷⁵, M. Goncharov¹⁷⁵, G. Grosso¹⁷⁵, P. Harris¹⁷⁵, D. Hoang¹⁷⁵,
G.M. Innocenti¹⁷⁵, K. Ivanov¹⁷⁵, D. Kovalskyi¹⁷⁵, J. Krupa¹⁷⁵, L. Lavezzo¹⁷⁵,
Y.-J. Lee¹⁷⁵, K. Long¹⁷⁵, C. McGinn¹⁷⁵, A. Novak¹⁷⁵, M.I. Park¹⁷⁵, C. Paus¹⁷⁵,
C. Reissel¹⁷⁵, C. Roland¹⁷⁵, G. Roland¹⁷⁵, S. Rothman¹⁷⁵, T.a. Sheng¹⁷⁵,
G.S.F. Stephans¹⁷⁵, D. Walter¹⁷⁵, J. Wang¹⁷⁵, Z. Wang¹⁷⁵, B. Wyslouch¹⁷⁵, T. J. Yang¹⁷⁵,
B. Crossman¹⁷⁶, W.J. Jackson¹⁷⁶, C. Kapsiak¹⁷⁶, M. Krohn¹⁷⁶, D. Mahon¹⁷⁶, J. Mans¹⁷⁶,
B. Marzocchi¹⁷⁶, R. Rusack¹⁷⁶, O. Sancar¹⁷⁶, R. Saradhy¹⁷⁶, N. Strobbe¹⁷⁶,
K. Bloom¹⁷⁷, D.R. Claes¹⁷⁷, G. Haza¹⁷⁷, J. Hossain¹⁷⁷, C. Joo¹⁷⁷, I. Kravchenko¹⁷⁷,
A. Rohilla¹⁷⁷, J.E. Siado¹⁷⁷, W. Tabb¹⁷⁷, A. Vagnerini¹⁷⁷, A. Wightman¹⁷⁷, F. Yan¹⁷⁷,
H. Bandyopadhyay¹⁷⁸, L. Hay¹⁷⁸, H.w. Hsia¹⁷⁸, I. Iashvili¹⁷⁸, A. Kalogeropoulos¹⁷⁸,
A. Kharchilava¹⁷⁸, A. Mandal¹⁷⁸, M. Morris¹⁷⁸, D. Nguyen¹⁷⁸, S. Rappoccio¹⁷⁸,
H. Rejeb Sfar¹⁷⁸, A. Williams¹⁷⁸, P. Young¹⁷⁸, D. Yu¹⁷⁸, G. Alverson¹⁷⁹, E. Barberis¹⁷⁹,
J. Bonilla¹⁷⁹, B. Bylsma¹⁷⁹, M. Campana¹⁷⁹, J. Dervan¹⁷⁹, Y. Haddad¹⁷⁹, Y. Han¹⁷⁹,
I. Israr¹⁷⁹, A. Krishna¹⁷⁹, M. Lu¹⁷⁹, N. Manganelli¹⁷⁹, R. McCarthy¹⁷⁹, D.M. Morse¹⁷⁹,
T. Orimoto¹⁷⁹, L. Skinnari¹⁷⁹, C.S. Thoreson¹⁷⁹, E. Tsai¹⁷⁹, D. Wood¹⁷⁹, S. Dittmer¹⁸⁰,
K.A. Hahn¹⁸⁰, M. McGinnis¹⁸⁰, Y. Miao¹⁸⁰, D.G. Monk¹⁸⁰, M.H. Schmitt¹⁸⁰,
A. Taliencio¹⁸⁰, M. Velasco¹⁸⁰, J. Wang¹⁸⁰, G. Agarwal¹⁸¹, R. Band¹⁸¹, R. Bucci¹⁸¹,
S. Castells¹⁸¹, A. Das¹⁸¹, A. Ehnis¹⁸¹, R. Goldouzian¹⁸¹, M. Hildreth¹⁸¹,
K. Hurtado Anampa¹⁸¹, T. Ivanov¹⁸¹, C. Jessop¹⁸¹, A. Karneyeu¹⁸¹, K. Lannon¹⁸¹,
J. Lawrence¹⁸¹, N. Loukas¹⁸¹, L. Lutton¹⁸¹, J. Mariano¹⁸¹, N. Marinelli¹⁸¹, I. Mcalister¹⁸¹,
T. McCauley¹⁸¹, C. Mcgrady¹⁸¹, C. Moore¹⁸¹, Y. Musienko^{181,cl}, H. Nelson¹⁸¹,
M. Osherson¹⁸¹, A. Piccinelli¹⁸¹, R. Ruchti¹⁸¹, A. Townsend¹⁸¹, Y. Wan¹⁸¹, M. Wayne¹⁸¹,
H. Yockey¹⁸¹, A. Basnet¹⁸², M. Carrigan¹⁸², R. De Los Santos¹⁸², L.S. Durkin¹⁸²,
C. Hill¹⁸², M. Joyce¹⁸², M. Nunez Ornelas¹⁸², D.A. Wenzl¹⁸², B.L. Winer¹⁸²,
B. R. Yates¹⁸², H. Bouchamaoui¹⁸³, G. Dezoort¹⁸³, P. Elmer¹⁸³, A. Frankenthal¹⁸³,
M. Galli¹⁸³, B. Greenberg¹⁸³, N. Haubrich¹⁸³, K. Kennedy¹⁸³, G. Kopp¹⁸³, Y. Lai¹⁸³,
D. Lange¹⁸³, A. Loeliger¹⁸³, D. Marlow¹⁸³, I. Ojalvo¹⁸³, J. Olsen¹⁸³, F. Simpson¹⁸³,
D. Stickland¹⁸³, C. Tully¹⁸³, S. Malik¹⁸⁴, R. Sharma¹⁸⁴, S. Chandra¹⁸⁵, R. Chawla¹⁸⁵,
A. Gu¹⁸⁵, L. Gutay¹⁸⁵, M. Jones¹⁸⁵, A.W. Jung¹⁸⁵, D. Kondratyev¹⁸⁵, M. Liu¹⁸⁵,
G. Negro¹⁸⁵, N. Neumeister¹⁸⁵, G. Paspalaki¹⁸⁵, S. Piperov¹⁸⁵, N.R. Saha¹⁸⁵,
J.F. Schulte¹⁸⁵, F. Wang¹⁸⁵, A. Wildridge¹⁸⁵, W. Xie¹⁸⁵, Y. Yao¹⁸⁵, Y. Zhong¹⁸⁵,
N. Parashar¹⁸⁶, A. Pathak¹⁸⁶, E. Shumka¹⁸⁶, D. Acosta¹⁸⁷, A. Agrawal¹⁸⁷,
C. Arbour¹⁸⁷, T. Carnahan¹⁸⁷, P. Das¹⁸⁷, K.M. Ecklund¹⁸⁷, S. Freed¹⁸⁷, F.J.M. Geurts¹⁸⁷,
T. Huang¹⁸⁷, I. Krommydas¹⁸⁷, N. Lewis¹⁸⁷, W. Li¹⁸⁷, J. Lin¹⁸⁷, O. Miguel Colin¹⁸⁷,
B.P. Padley¹⁸⁷, R. Redjimi¹⁸⁷, J. Rotter¹⁸⁷, M. Wulansatiti¹⁸⁷, E. Yigitbasi¹⁸⁷,
Y. Zhang¹⁸⁷, O. Bessidskaia Bylund¹⁸⁸, A. Bodek¹⁸⁸, P. de Barbaro^{188,†}, R. Demina¹⁸⁸,
A. Garcia-Bellido¹⁸⁸, H.S. Hare¹⁸⁸, O. Hindrichs¹⁸⁸, N. Parmar¹⁸⁸, P. Parygin^{188,ck},
H. Seo¹⁸⁸, R. Taus¹⁸⁸, B. Chiarito¹⁸⁹, J.P. Chou¹⁸⁹, S.V. Clark¹⁸⁹, S. Donnelly¹⁸⁹,
D. Gadkari¹⁸⁹, Y. Gershtein¹⁸⁹, E. Halkiadakis¹⁸⁹, C. Houghton¹⁸⁹, D. Jaroslawski¹⁸⁹,
A. Kobert¹⁸⁹, S. Konstantinou¹⁸⁹, I. Laflotte¹⁸⁹, A. Lath¹⁸⁹, J. Martins¹⁸⁹,
M. Perez Prada¹⁸⁹, B. Rand¹⁸⁹, J. Reichert¹⁸⁹, P. Saha¹⁸⁹, S. Salur¹⁸⁹, S. Schnetzer¹⁸⁹,
S. Somalwar¹⁸⁹, R. Stone¹⁸⁹, S.A. Thayil¹⁸⁹, S. Thomas¹⁸⁹, J. Vora¹⁸⁹, D. Ally¹⁹⁰,
A.G. Delannoy¹⁹⁰, S. Fiorendi¹⁹⁰, J. Harris¹⁹⁰, T. Holmes¹⁹⁰, A.R. Kanuganti¹⁹⁰,

N. Karunaratna¹⁹⁰, J. Lawless¹⁹⁰, L. Lee¹⁹⁰, E. Nibigira¹⁹⁰, B. Skipworth¹⁹⁰, S. Spanier¹⁹⁰, D. Aebi¹⁹¹, M. Ahmad¹⁹¹, T. Akhter¹⁹¹, K. Androsov¹⁹¹, A. Bolshov¹⁹¹, O. Bouhali^{191,cm}, A. Cagnotta¹⁹¹, V. D'Amante¹⁹¹, R. Eusebi¹⁹¹, P. Flanagan¹⁹¹, J. Gilmore¹⁹¹, Y. Guo¹⁹¹, T. Kamon¹⁹¹, S. Luo¹⁹¹, R. Mueller¹⁹¹, A. Safonov¹⁹¹, N. Akchurin¹⁹², J. Damgov¹⁹², Y. Feng¹⁹², N. Gogate¹⁹², Y. Kazhykarim¹⁹², K. Lamichhane¹⁹², S.W. Lee¹⁹², C. Madrid¹⁹², A. Mankel¹⁹², T. Peltola¹⁹², I. Volobouev¹⁹², E. Appelt¹⁹³, Y. Chen¹⁹³, S. Greene¹⁹³, A. Gurrola¹⁹³, W. Johns¹⁹³, R. Kunnawalkam Elayavalli¹⁹³, A. Melo¹⁹³, D. Rathjens¹⁹³, F. Romeo¹⁹³, P. Sheldon¹⁹³, S. Tuo¹⁹³, J. Velkovska¹⁹³, J. Viinikainen¹⁹³, J. Zhang¹⁹³, B. Cardwell¹⁹⁴, H. Chung¹⁹⁴, B. Cox¹⁹⁴, J. Hakala¹⁹⁴, G. Hamilton Ilha Machado¹⁹⁴, R. Hirosky¹⁹⁴, M. Jose¹⁹⁴, A. Ledovsky¹⁹⁴, C. Mantilla¹⁹⁴, C. Neu¹⁹⁴, C. Ramón Álvarez¹⁹⁴, Z. Wu¹⁹⁴, S. Bhattacharya¹⁹⁵, P.E. Karchin¹⁹⁵, A. Aravind¹⁹⁶, S. Banerjee¹⁹⁶, K. Black¹⁹⁶, T. Bose¹⁹⁶, E. Chavez¹⁹⁶, S. Dasu¹⁹⁶, P. Everaerts¹⁹⁶, C. Galloni¹⁹⁶, H. He¹⁹⁶, M. Herndon¹⁹⁶, A. Herve¹⁹⁶, C.K. Koraka¹⁹⁶, S. Lomte¹⁹⁶, R. Loveless¹⁹⁶, A. Mallampalli¹⁹⁶, A. Mohammadi¹⁹⁶, S. Mondal¹⁹⁶, T. Nelson¹⁹⁶, G. Parida¹⁹⁶, D. Pinna¹⁹⁶, L. Pétré¹⁹⁶, A. Savin¹⁹⁶, V. Shang¹⁹⁶, V. Sharma¹⁹⁶, W.H. Smith¹⁹⁶, D. Teague¹⁹⁶, H.F. Tsoi¹⁹⁶, W. Vetens¹⁹⁶, A. Warden¹⁹⁶, S. Afanasiev¹⁹⁷, V. Alexakhin¹⁹⁷, Yu. Andreev¹⁹⁷, T. Aushev¹⁹⁷, D. Budkouski¹⁹⁷, R. Chistov¹⁹⁷, M. Danilov¹⁹⁷, T. Dimova¹⁹⁷, A. Ershov¹⁹⁷, S. Gninenko¹⁹⁷, I. Gorbunov¹⁹⁷, A. Gribushin¹⁹⁷, A. Kamenev¹⁹⁷, V. Karjavine¹⁹⁷, M. Kirsanov¹⁹⁷, V. Klyukhin¹⁹⁷, O. Kodolova^{197,cn}, V. Korenkov¹⁹⁷, I. Korsakov¹⁹⁷, A. Kozyrev¹⁹⁷, N. Krasnikov¹⁹⁷, A. Lanev¹⁹⁷, A. Malakhov¹⁹⁷, V. Matveev¹⁹⁷, A. Nikitenko^{197,co,cp}, V. Palichik¹⁹⁷, V. Perelygin¹⁹⁷, S. Petrushanko¹⁹⁷, S. Polikarpov¹⁹⁷, O. Radchenko¹⁹⁷, M. Savina¹⁹⁷, V. Shalaev¹⁹⁷, S. Shmatov¹⁹⁷, S. Shulha¹⁹⁷, Y. Skovpen¹⁹⁷, K. Slizhevskiy¹⁹⁷, V. Smirnov¹⁹⁷, O. Teryaev¹⁹⁷, I. Tisova¹⁹⁷, A. Toropin¹⁹⁷, N. Voytishin¹⁹⁷, A. Zarubin¹⁹⁷, I. Zhizhin¹⁹⁷, E. Boos¹⁹⁸, V. Bunichev¹⁹⁸, M. Dubinin^{198,cd}, L. Dudko¹⁹⁸, V. Kim^{198,cl}, V. Murzin¹⁹⁸, V. Oreshkin¹⁹⁸, V. Savrin¹⁹⁸, A. Snigirev¹⁹⁸, D. Sosnov¹⁹⁸

¹ Yerevan Physics Institute, Yerevan, Armenia

² Institut für Hochenergiephysik, Vienna, Austria

³ Universiteit Antwerpen, Antwerpen, Belgium

⁴ Vrije Universiteit Brussel, Brussel, Belgium

⁵ Université Libre de Bruxelles, Bruxelles, Belgium

⁶ Ghent University, Ghent, Belgium

⁷ Université Catholique de Louvain, Louvain-la-Neuve, Belgium

⁸ Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil

⁹ Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

¹⁰ Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo, Brazil

¹¹ Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

¹² University of Sofia, Sofia, Bulgaria

¹³ Instituto De Alta Investigación, Universidad de Tarapacá, Casilla 7 D, Arica, Chile

¹⁴ Universidad Técnica Federico Santa María, Valparaíso, Chile

¹⁵ Beihang University, Beijing, China

¹⁶ Department of Physics, Tsinghua University, Beijing, China

¹⁷ Institute of High Energy Physics, Beijing, China

¹⁸ State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

- ¹⁹ *State Key Laboratory of Nuclear Physics and Technology, Institute of Quantum Matter, South China Normal University, Guangzhou, China*
- ²⁰ *Sun Yat-Sen University, Guangzhou, China*
- ²¹ *University of Science and Technology of China, Hefei, China*
- ²² *Nanjing Normal University, Nanjing, China*
- ²³ *Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) — Fudan University, Shanghai, China*
- ²⁴ *Zhejiang University, Hangzhou, Zhejiang, China*
- ²⁵ *Universidad de Los Andes, Bogota, Colombia*
- ²⁶ *Universidad de Antioquia, Medellin, Colombia*
- ²⁷ *University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia*
- ²⁸ *University of Split, Faculty of Science, Split, Croatia*
- ²⁹ *Institute Rudjer Boskovic, Zagreb, Croatia*
- ³⁰ *University of Cyprus, Nicosia, Cyprus*
- ³¹ *Charles University, Prague, Czech Republic*
- ³² *Escuela Politecnica Nacional, Quito, Ecuador*
- ³³ *Universidad San Francisco de Quito, Quito, Ecuador*
- ³⁴ *Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt*
- ³⁵ *Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum, Egypt*
- ³⁶ *National Institute of Chemical Physics and Biophysics, Tallinn, Estonia*
- ³⁷ *Department of Physics, University of Helsinki, Helsinki, Finland*
- ³⁸ *Helsinki Institute of Physics, Helsinki, Finland*
- ³⁹ *Lappeenranta-Lahti University of Technology, Lappeenranta, Finland*
- ⁴⁰ *IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France*
- ⁴¹ *Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France*
- ⁴² *Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France*
- ⁴³ *Centre de Calcul de l'Institut National de Physique Nucleaire et de Physique des Particules, CNRS/IN2P3, Villeurbanne, France*
- ⁴⁴ *Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France*
- ⁴⁵ *Georgian Technical University, Tbilisi, Georgia*
- ⁴⁶ *RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany*
- ⁴⁷ *RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany*
- ⁴⁸ *RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany*
- ⁴⁹ *Deutsches Elektronen-Synchrotron, Hamburg, Germany*
- ⁵⁰ *University of Hamburg, Hamburg, Germany*
- ⁵¹ *Karlsruher Institut fuer Technologie, Karlsruhe, Germany*
- ⁵² *Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece*
- ⁵³ *National and Kapodistrian University of Athens, Athens, Greece*
- ⁵⁴ *National Technical University of Athens, Athens, Greece*
- ⁵⁵ *University of Ioánnina, Ioánnina, Greece*
- ⁵⁶ *HUN-REN Wigner Research Centre for Physics, Budapest, Hungary*
- ⁵⁷ *MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary*
- ⁵⁸ *Faculty of Informatics, University of Debrecen, Debrecen, Hungary*
- ⁵⁹ *HUN-REN ATOMKI — Institute of Nuclear Research, Debrecen, Hungary*
- ⁶⁰ *Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary*
- ⁶¹ *Panjab University, Chandigarh, India*
- ⁶² *University of Delhi, Delhi, India*
- ⁶³ *Indian Institute of Technology Mandi (IIT-Mandi), Himachal Pradesh, India*
- ⁶⁴ *University of Hyderabad, Hyderabad, India*

- ⁶⁵ *Indian Institute of Technology Kanpur, Kanpur, India*
- ⁶⁶ *Saha Institute of Nuclear Physics, HBNI, Kolkata, India*
- ⁶⁷ *Indian Institute of Technology Madras, Madras, India*
- ⁶⁸ *IISER Mohali, India, Mohali, India*
- ⁶⁹ *Tata Institute of Fundamental Research-A, Mumbai, India*
- ⁷⁰ *Tata Institute of Fundamental Research-B, Mumbai, India*
- ⁷¹ *National Institute of Science Education and Research, An OCC of Homi Bhabha National Institute, Bhubaneswar, Odisha, India*
- ⁷² *Indian Institute of Science Education and Research (IISER), Pune, India*
- ⁷³ *Indian Institute of Technology Hyderabad, Telangana, India*
- ⁷⁴ *Isfahan University of Technology, Isfahan, Iran*
- ⁷⁵ *Institute for Research in Fundamental Sciences (IPM), Tehran, Iran*
- ⁷⁶ *University College Dublin, Dublin, Ireland*
- ^{77a} *INFN Sezione di Bari, Bari, Italy*
- ^{77b} *Università di Bari, Bari, Italy*
- ^{77c} *Politecnico di Bari, Bari, Italy*
- ^{78a} *INFN Sezione di Bologna, Bologna, Italy*
- ^{78b} *Università di Bologna, Bologna, Italy*
- ^{79a} *INFN Sezione di Catania, Catania, Italy*
- ^{79b} *Università di Catania, Catania, Italy*
- ^{80a} *INFN Sezione di Firenze, Firenze, Italy*
- ^{80b} *Università di Firenze, Firenze, Italy*
- ⁸¹ *INFN Laboratori Nazionali di Frascati, Frascati, Italy*
- ^{82a} *INFN Sezione di Genova, Genova, Italy*
- ^{82b} *Università di Genova, Genova, Italy*
- ^{83a} *INFN Sezione di Milano-Bicocca, Milano, Italy*
- ^{83b} *Università di Milano-Bicocca, Milano, Italy*
- ^{84a} *INFN Sezione di Napoli, Napoli, Italy*
- ^{84b} *Università di Napoli ‘Federico II’, Napoli, Italy*
- ^{84c} *Università della Basilicata, Potenza, Italy*
- ^{84d} *Scuola Superiore Meridionale (SSM), Napoli, Italy*
- ^{85a} *INFN Sezione di Padova, Padova, Italy*
- ^{85b} *Università di Padova, Padova, Italy*
- ^{85c} *Università degli Studi di Cagliari, Cagliari, Italy*
- ^{86a} *INFN Sezione di Pavia, Pavia, Italy*
- ^{86b} *Università di Pavia, Pavia, Italy*
- ^{87a} *INFN Sezione di Perugia, Perugia, Italy*
- ^{87b} *Università di Perugia, Perugia, Italy*
- ^{88a} *INFN Sezione di Pisa, Pisa, Italy*
- ^{88b} *Università di Pisa, Pisa, Italy*
- ^{88c} *Scuola Normale Superiore di Pisa, Pisa, Italy*
- ^{88d} *Università di Siena, Siena, Italy*
- ^{89a} *INFN Sezione di Roma, Roma, Italy*
- ^{89b} *Sapienza Università di Roma, Roma, Italy*
- ^{90a} *INFN Sezione di Torino, Torino, Italy*
- ^{90b} *Università di Torino, Torino, Italy*
- ^{90c} *Università del Piemonte Orientale, Novara, Italy*
- ^{91a} *INFN Sezione di Trieste, Trieste, Italy*
- ^{91b} *Università di Trieste, Trieste, Italy*
- ⁹² *Kyungpook National University, Daegu, Korea*
- ⁹³ *Department of Mathematics and Physics — GWNNU, Gangneung, Korea*
- ⁹⁴ *Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea*
- ⁹⁵ *Hanyang University, Seoul, Korea*

- ⁹⁶ *Korea University, Seoul, Korea*
- ⁹⁷ *Kyung Hee University, Department of Physics, Seoul, Korea*
- ⁹⁸ *Sejong University, Seoul, Korea*
- ⁹⁹ *Seoul National University, Seoul, Korea*
- ¹⁰⁰ *University of Seoul, Seoul, Korea*
- ¹⁰¹ *Yonsei University, Department of Physics, Seoul, Korea*
- ¹⁰² *Sungkyunkwan University, Suwon, Korea*
- ¹⁰³ *College of Engineering and Technology, American University of the Middle East (AUM), Dasman, Kuwait*
- ¹⁰⁴ *Kuwait University — College of Science — Department of Physics, Safat, Kuwait*
- ¹⁰⁵ *Riga Technical University, Riga, Latvia*
- ¹⁰⁶ *University of Latvia (LU), Riga, Latvia*
- ¹⁰⁷ *Vilnius University, Vilnius, Lithuania*
- ¹⁰⁸ *National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia*
- ¹⁰⁹ *Universidad de Sonora (UNISON), Hermosillo, Mexico*
- ¹¹⁰ *Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico*
- ¹¹¹ *Universidad Iberoamericana, Mexico City, Mexico*
- ¹¹² *Benemerita Universidad Autonoma de Puebla, Puebla, Mexico*
- ¹¹³ *University of Montenegro, Podgorica, Montenegro*
- ¹¹⁴ *University of Canterbury, Christchurch, New Zealand*
- ¹¹⁵ *National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan*
- ¹¹⁶ *AGH University of Krakow, Krakow, Poland*
- ¹¹⁷ *National Centre for Nuclear Research, Swierk, Poland*
- ¹¹⁸ *Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland*
- ¹¹⁹ *Warsaw University of Technology, Warsaw, Poland*
- ¹²⁰ *Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal*
- ¹²¹ *Faculty of Physics, University of Belgrade, Belgrade, Serbia*
- ¹²² *VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia*
- ¹²³ *Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain*
- ¹²⁴ *Universidad Autónoma de Madrid, Madrid, Spain*
- ¹²⁵ *Universidad de Oviedo, Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Oviedo, Spain*
- ¹²⁶ *Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain*
- ¹²⁷ *University of Colombo, Colombo, Sri Lanka*
- ¹²⁸ *University of Ruhuna, Department of Physics, Matara, Sri Lanka*
- ¹²⁹ *CERN, European Organization for Nuclear Research, Geneva, Switzerland*
- ¹³⁰ *PSI Center for Neutron and Muon Sciences, Villigen, Switzerland*
- ¹³¹ *ETH Zurich — Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland*
- ¹³² *Universität Zürich, Zurich, Switzerland*
- ¹³³ *National Central University, Chung-Li, Taiwan*
- ¹³⁴ *National Taiwan University (NTU), Taipei, Taiwan*
- ¹³⁵ *High Energy Physics Research Unit, Department of Physics, Faculty of Science, Chulalongkorn University, Bangkok, Thailand*
- ¹³⁶ *Tunis El Manar University, Tunis, Tunisia*
- ¹³⁷ *Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey*
- ¹³⁸ *Hacettepe University, Ankara, Turkey*
- ¹³⁹ *Middle East Technical University, Physics Department, Ankara, Turkey*
- ¹⁴⁰ *Bogazici University, Istanbul, Turkey*
- ¹⁴¹ *Istanbul Technical University, Istanbul, Turkey*
- ¹⁴² *Istanbul University, Istanbul, Turkey*
- ¹⁴³ *Yildiz Technical University, Istanbul, Turkey*
- ¹⁴⁴ *Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkiv, Ukraine*
- ¹⁴⁵ *National Science Centre, Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine*
- ¹⁴⁶ *University of Bristol, Bristol, U.K.*

- 147 *Rutherford Appleton Laboratory, Didcot, U.K.*
- 148 *Imperial College, London, U.K.*
- 149 *Brunel University, Uxbridge, U.K.*
- 150 *Baylor University, Waco, Texas, U.S.A.*
- 151 *Bethel University, St. Paul, Minnesota, U.S.A.*
- 152 *Catholic University of America, Washington, DC, U.S.A.*
- 153 *The University of Alabama, Tuscaloosa, Alabama, U.S.A.*
- 154 *Boston University, Boston, Massachusetts, U.S.A.*
- 155 *Brown University, Providence, Rhode Island, U.S.A.*
- 156 *University of California, Davis, Davis, California, U.S.A.*
- 157 *University of California, Los Angeles, California, U.S.A.*
- 158 *University of California, Riverside, Riverside, California, U.S.A.*
- 159 *University of California, San Diego, La Jolla, California, U.S.A.*
- 160 *University of California, Santa Barbara — Department of Physics, Santa Barbara, California, U.S.A.*
- 161 *California Institute of Technology, Pasadena, California, U.S.A.*
- 162 *Carnegie Mellon University, Pittsburgh, Pennsylvania, U.S.A.*
- 163 *University of Colorado Boulder, Boulder, Colorado, U.S.A.*
- 164 *Cornell University, Ithaca, New York, U.S.A.*
- 165 *Fermi National Accelerator Laboratory, Batavia, Illinois, U.S.A.*
- 166 *University of Florida, Gainesville, Florida, U.S.A.*
- 167 *Florida State University, Tallahassee, Florida, U.S.A.*
- 168 *Florida Institute of Technology, Melbourne, Florida, U.S.A.*
- 169 *University of Illinois Chicago, Chicago, Illinois, U.S.A.*
- 170 *The University of Iowa, Iowa City, Iowa, U.S.A.*
- 171 *Johns Hopkins University, Baltimore, Maryland, U.S.A.*
- 172 *The University of Kansas, Lawrence, Kansas, U.S.A.*
- 173 *Kansas State University, Manhattan, Kansas, U.S.A.*
- 174 *University of Maryland, College Park, Maryland, U.S.A.*
- 175 *Massachusetts Institute of Technology, Cambridge, Massachusetts, U.S.A.*
- 176 *University of Minnesota, Minneapolis, Minnesota, U.S.A.*
- 177 *University of Nebraska-Lincoln, Lincoln, Nebraska, U.S.A.*
- 178 *State University of New York at Buffalo, Buffalo, New York, U.S.A.*
- 179 *Northeastern University, Boston, Massachusetts, U.S.A.*
- 180 *Northwestern University, Evanston, Illinois, U.S.A.*
- 181 *University of Notre Dame, Notre Dame, Indiana, U.S.A.*
- 182 *The Ohio State University, Columbus, Ohio, U.S.A.*
- 183 *Princeton University, Princeton, New Jersey, U.S.A.*
- 184 *University of Puerto Rico, Mayaguez, Puerto Rico, U.S.A.*
- 185 *Purdue University, West Lafayette, Indiana, U.S.A.*
- 186 *Purdue University Northwest, Hammond, Indiana, U.S.A.*
- 187 *Rice University, Houston, Texas, U.S.A.*
- 188 *University of Rochester, Rochester, New York, U.S.A.*
- 189 *Rutgers, The State University of New Jersey, Piscataway, New Jersey, U.S.A.*
- 190 *University of Tennessee, Knoxville, Tennessee, U.S.A.*
- 191 *Texas A&M University, College Station, Texas, U.S.A.*
- 192 *Texas Tech University, Lubbock, Texas, U.S.A.*
- 193 *Vanderbilt University, Nashville, Tennessee, U.S.A.*
- 194 *University of Virginia, Charlottesville, Virginia, U.S.A.*
- 195 *Wayne State University, Detroit, Michigan, U.S.A.*
- 196 *University of Wisconsin — Madison, Madison, Wisconsin, U.S.A.*
- 197 *An institute or international laboratory covered by a cooperation agreement with CERN*
- 198 *An institute formerly covered by a cooperation agreement with CERN*

- ^a Also at Yerevan State University, Yerevan, Armenia
- ^b Also at TU Wien, Vienna, Austria
- ^c Also at Ghent University, Ghent, Belgium
- ^d Also at FACAMP — Faculdades de Campinas, Sao Paulo, Brazil
- ^e Also at Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil
- ^f Also at Universidade Estadual de Campinas, Campinas, Brazil
- ^g Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil
- ^h Also at The University of the State of Amazonas, Manaus, Brazil
- ⁱ Also at University of Chinese Academy of Sciences, Beijing, China
- ^j Also at China Center of Advanced Science and Technology, Beijing, China
- ^k Also at University of Chinese Academy of Sciences, Beijing, China
- ^l Also at School of Physics, Zhengzhou University, Zhengzhou, China
- ^m Now at Henan Normal University, Xinxiang, China
- ⁿ Also at University of Shanghai for Science and Technology, Shanghai, China
- ^o Now at The University of Iowa, Iowa City, Iowa, U.S.A.
- ^p Also at Center for High Energy Physics, Peking University, Beijing, China
- ^q Now at British University in Egypt, Cairo, Egypt
- ^r Now at Cairo University, Cairo, Egypt
- ^s Also at Purdue University, West Lafayette, Indiana, U.S.A.
- ^t Also at Université de Haute Alsace, Mulhouse, France
- ^u Also at Another institute or international laboratory covered by a cooperation agreement with CERN
- ^v Also at University of Hamburg, Hamburg, Germany
- ^w Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany
- ^x Also at Bergische University Wuppertal (BUW), Wuppertal, Germany
- ^y Also at Brandenburg University of Technology, Cottbus, Germany
- ^z Also at Forschungszentrum Jülich, Jülich, Germany
- ^{aa} Now at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany
- ^{ab} Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland
- ^{ac} Also at HUN-REN ATOMKI — Institute of Nuclear Research, Debrecen, Hungary
- ^{ad} Now at Universitatea Babes-Bolyai — Facultatea de Fizica, Cluj-Napoca, Romania
- ^{ae} Also at MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary
- ^{af} Also at HUN-REN Wigner Research Centre for Physics, Budapest, Hungary
- ^{ag} Also at Physics Department, Faculty of Science, Assiut University, Assiut, Egypt
- ^{ah} Also at The University of Kansas, Lawrence, Kansas, U.S.A.
- ^{ai} Also at Punjab Agricultural University, Ludhiana, India
- ^{aj} Also at University of Hyderabad, Hyderabad, India
- ^{ak} Also at Indian Institute of Science (IISc), Bangalore, India
- ^{al} Also at University of Visva-Bharati, Santiniketan, India
- ^{am} Also at IIT Bhubaneswar, Bhubaneswar, India
- ^{an} Also at Institute of Physics, Bhubaneswar, India
- ^{ao} Also at Deutsches Elektronen-Synchrotron, Hamburg, Germany
- ^{ap} Also at Isfahan University of Technology, Isfahan, Iran
- ^{aq} Also at Sharif University of Technology, Tehran, Iran
- ^{ar} Also at Department of Physics, University of Science and Technology of Mazandaran, Behshahr, Iran
- ^{as} Also at Department of Physics, Faculty of Science, Arak University, ARAK, Iran
- ^{at} Also at Helwan University, Cairo, Egypt
- ^{au} Also at Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy
- ^{av} Also at Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy
- ^{aw} Also at Università degli Studi Guglielmo Marconi, Roma, Italy
- ^{ax} Also at Scuola Superiore Meridionale, Università di Napoli ‘Federico II’, Napoli, Italy
- ^{ay} Also at Fermi National Accelerator Laboratory, Batavia, Illinois, U.S.A.

- ^{az} Also at Lulea University of Technology, Lulea, Sweden
- ^{ba} Also at Consiglio Nazionale delle Ricerche — Istituto Officina dei Materiali, Perugia, Italy
- ^{bb} Also at UPES — University of Petroleum and Energy Studies, Dehradun, India
- ^{bc} Also at Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France
- ^{bd} Also at Department of Applied Physics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia
- ^{be} Also at Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka
- ^{bf} Also at Saegis Campus, Nugegoda, Sri Lanka
- ^{bg} Also at National and Kapodistrian University of Athens, Athens, Greece
- ^{bh} Also at Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland
- ^{bi} Also at Universität Zürich, Zurich, Switzerland
- ^{bj} Also at Stefan Meyer Institute for Subatomic Physics, Vienna, Austria
- ^{bk} Also at Near East University, Research Center of Experimental Health Science, Mersin, Turkey
- ^{bl} Also at Konya Technical University, Konya, Turkey
- ^{bm} Also at Izmir Bakircay University, Izmir, Turkey
- ^{bn} Also at Adiyaman University, Adiyaman, Turkey
- ^{bo} Also at Bozok Universititesi Rektörlüğü, Yozgat, Turkey
- ^{bp} Also at Istanbul Sabahattin Zaim University, Istanbul, Turkey
- ^{bq} Also at Marmara University, Istanbul, Turkey
- ^{br} Also at Milli Savunma University, Istanbul, Turkey
- ^{bs} Also at Informatics and Information Security Research Center, Gebze/Kocaeli, Turkey
- ^{bt} Also at Kafkas University, Kars, Turkey
- ^{bu} Now at Istanbul Okan University, Istanbul, Turkey
- ^{bv} Also at Istanbul University — Cerrahpasa, Faculty of Engineering, Istanbul, Turkey
- ^{bw} Also at Istinye University, Istanbul, Turkey
- ^{bx} Also at Yildiz Technical University, Istanbul, Turkey
- ^{by} Also at School of Physics and Astronomy, University of Southampton, Southampton, U.K.
- ^{bz} Also at Monash University, Faculty of Science, Clayton, Australia
- ^{ca} Also at Università di Torino, Torino, Italy
- ^{cb} Also at Karamanoğlu Mehmetbey University, Karaman, Turkey
- ^{cc} Also at California Lutheran University, Thousand Oaks, California, U.S.A.
- ^{cd} Also at California Institute of Technology, Pasadena, California, U.S.A.
- ^{ce} Also at United States Naval Academy, Annapolis, Maryland, U.S.A.
- ^{cf} Also at Bingol University, Bingol, Turkey
- ^{cg} Also at Georgian Technical University, Tbilisi, Georgia
- ^{ch} Also at Sinop University, Sinop, Turkey
- ^{ci} Also at Erciyes University, Kayseri, Turkey
- ^{cj} Also at Horia Hulubei National Institute of Physics and Nuclear Engineering (IFIN-HH), Bucharest, Romania
- ^{ck} Now at Another institute formerly covered by a cooperation agreement with CERN
- ^{cl} Also at Another institute formerly covered by a cooperation agreement with CERN
- ^{cm} Also at Hamad Bin Khalifa University (HBKU), Doha, Qatar
- ^{cn} Also at Yerevan Physics Institute, Yerevan, Armenia
- ^{co} Also at Imperial College, London, U.K.
- ^{cp} Now at Yerevan Physics Institute, Yerevan, Armenia
- [†] Deceased