Illuminating the Hidden Sector of String Theory by Shining Light through a Magnetic Field

Steven A. Abel,¹ Joerg Jaeckel,² Valentin V. Khoze,¹ and Andreas Ringwald^{1,2}

¹*Centre for Particle Theory, Durham University, Durham, DH1 3LE, UK*

²*Deutsches Elektronen-Synchrotron DESY, Notkestraße 85, D-22607 Hamburg, Germany*

It has recently been shown that the observation by the PVLAS collaboration of an anomalously large rotation of the polarization plane of light in the presence of a magnetic field in vacuum may originate from pair production of light, $m_f \sim 0.1$ eV, millicharged, $Q_f \sim 3 \times 10^{-6} e$, fermions. Such millicharges arise generically from kinetic-mixing in theories containing at least two $U(1)$ gauge factors. In this letter, we point out that the required multiple $U(1)$ factors, the size of kinetic-mixing, and suitable matter representations to explain the PVLAS data occur very naturally in the context of realistic extensions of the Standard Model (SM) of elementary particle physics based on string theory.

PACS numbers: 11.25.-w, 11.25.Wx, 12.20.-m, 14.80.-j

The absorption probability and the propagation speed of polarized light propagating in a magnetic field may depend on the relative orientation of the polarization and the magnetic field. These effects are known as vacuum magnetic dichroism and birefringence, respectively.

Recently, it was shown [1] that the observation by the PVLAS collaboration [2] of an anomalously large rotation of the polarization plane of light after its passage through a transverse magnetic field in vacuum may be originating from the dichroism caused by pair production of millicharged fermions of mass

$$m_f \simeq 0.1 \text{ eV} \quad (1)$$

and fractional electric charge

$$\epsilon \equiv Q_f/e \simeq 3 \times 10^{-6}, \quad (2)$$

where e is the electron electric charge.

Particles with such small, unquantized charge arise very naturally in so-called paraphoton [3] models, containing, beyond the usual electromagnetic $U(1)$ gauge factor, at least one additional hidden-sector $U(1)$ factor.

The basic observation is that particles with paracharge get an induced electric charge proportional to some small mixing angle between the kinetic terms of photons and paraphotons [4]. Moreover, in models containing more than one paraphoton with at least one paraphoton being exactly massless and one light, $\text{keV} \gg m_{\tilde{\gamma}} \neq 0$, the prohibitively strong astrophysical bounds on the fractional charge, $\epsilon \lesssim 2 \times 10^{-14}$, for $m_f \lesssim \text{few keV}$, arising from energy loss considerations of stars [5], can be relaxed considerably. In a simple model analysed in [6], there are two paraphotons: one massless and one light, and the fermion transforms in the bifundamental representation of these two $U(1)$ factors. In vacuum, the fermion acquires an electric charge ϵ due to a kinetic-mixing between the photon and the two paraphotons. Importantly, however, this electric charge is reduced in the stellar plasma by a multiplicative factor $m_{\tilde{\gamma}}^2/\omega_p^2$, where $\omega_p \sim \text{few keV}$ is

the plasma frequency. This charge screening mechanism is caused by a partial cancellation between two paraphotons interacting with the bifundamental fermion [6]. The vacuum value (2) is therefore perfectly compatible with astrophysical bounds as long as

$$m_{\tilde{\gamma}} \lesssim 0.1 \text{ eV}. \quad (3)$$

This minimal model can be supplemented by an axion-like spin-zero particle, coupled to the millicharged fermions [6]. A triangle diagram then leads to a coupling of the axion-like particle to two photons. The resulting production of axion-like scalars gives an additional (to the one from millicharged fermions) contribution to the vacuum magnetic dichroism and birefringence. One can then expect to explain PVLAS with even smaller values of ϵ , while still not being in conflict with astrophysics.

In this letter, we point out that the required multiple $U(1)$ factors, the size of kinetic-mixing, and suitable matter representations to explain the PVLAS data occur very naturally within the context of realistic extensions of the standard model (SM) of elementary particle physics based on string theory.

Let us begin by recalling the essentials of gauge kinetic-mixing. It arises generally in theories that have, in addition to some visible $U(1)_a$, at least one other additional $U(1)_b$ factor in a hidden sector. In the basis in which the interaction terms have the canonical form, the pure gauge part of the Lagrangian for an arbitrary $U(1)_a \times U(1)_b$ theory can be written as (generalization to more than one additional $U(1)$ factor is straightforward)

$$\mathcal{L}_{\text{gauge}} = -\frac{1}{4} F_{(a)}^{\mu\nu} F_{(a)\mu\nu} - \frac{1}{4} F_{(b)}^{\mu\nu} F_{(b)\mu\nu} + \frac{\chi}{2} F_{(a)}^{\mu\nu} F_{(b)\mu\nu}, \quad (4)$$

where χ parametrizes the mixing. This parameter is directly linked to charge shifts. Notably, starting for example with two fermion species f_a and f_b with charges $(e, 0)$ and $(0, e)$, respectively, under $U(1)_a \times U(1)_b$, one finds that their charges, after diagonalization of the gauge

FIG. 1: (a) One-loop diagram which contributes to kinetic-mixing in field theory, and (b) its equivalent in open string theory (from Ref. [9]).

kinetic term, are shifted by the amount

$$\epsilon \simeq \chi, \quad (5)$$

to leading order in $\chi \ll 1$ [4]. In other words, the hidden sector fermion f_b picks up a small charge $\epsilon \simeq \chi$ under the visible sector $U(1)_a$.

In view of the millicharged particle interpretation of the PVLAS data, the following question immediately arises in this context:

Are there theoretically appealing and phenomenologically viable theories which naturally lead to a value of $\chi \simeq \epsilon \simeq 3 \times 10^{-6}$ for the kinetic-mixing parameter?

We will argue that the answer is yes. Many extensions of the standard model, and indeed most extensions coming from string theory, predict additional hidden $U(1)$ factors which can give rise to the kinetic-mixing phenomenon (e.g., [7, 8, 9, 11]).

In the context of field theory, a non-zero value of χ can be induced quite generally at the one-loop level when there are states which are simultaneously charged under both the visible $U(1)$ and the hidden $U(1)$ factors [4] (cf. the field theory diagram in Fig. 1(a)). The expected magnitude of χ in the field theory setting may be estimated [4, 7] by considering the contribution of two fermions with charges (e_a, e_b) and $(e_a, -e_b)$ and masses m and m' , respectively. Their joint contribution to χ is given by [4]

$$\chi \simeq \frac{e_a e_b}{6\pi^2} \log\left(\frac{m'}{m}\right) \simeq 1.5 \times 10^{-3} \left(\frac{e_a e_b}{4\pi} \frac{1}{\alpha}\right) \log\left(\frac{m'}{m}\right), \quad (6)$$

and can be of order 3×10^{-6} , for natural values of $e_a e_b / (4\pi) \sim \alpha$, with $\alpha = e^2 / (4\pi)$ being the electromagnetic fine-structure constant, and nearly degenerate masses, $m'/m \sim 1.002$. Kinetic-mixing could be avoided in such theories if the particle spectrum has some particular properties, as discussed in Ref. [7]. For example, if one or both of the $U(1)$ gauge factors sits within an unbroken non-abelian gauge symmetry, then kinetic-mixing is not possible simply because of the tracelessness of the generators.

In the string theory context (e.g., [10]), the story is more subtle and more varied, but quite generally there

FIG. 2: Kinetic-mixing in open string models with SUSY breaking on “hidden” branes. The visible sector consists of a phenomenologically well determined supersymmetric configuration of D3-branes at a fixed point in the 6 dimensional compact manifold, possibly with D7-branes passing through to cancel local tadpoles. Global absence of tadpoles is assumed to require additional branes and/or anti-branes in the bulk. Closed string interactions are mediated from hidden to visible sector by cylinder diagrams, and are equivalent to Fig. 1(b).

are clear hints that string theory naturally leads to a generation of $\chi \neq 0$ [7, 8, 9], and can naturally give values of the right order of magnitude. The expected size of χ in a particular string theory setting can be investigated by performing the equivalent one-loop string theory calculation (cf. Fig. 1(b)).

Weakly coupled heterotic closed string models (in which the string scale, $M_s \approx 5 \times 10^{17}$ GeV, is close to the Planck scale, $M_P = 1.2 \times 10^{19}$ GeV) were treated this way in Ref. [7]. In these models, even if the particular string vacuum does not have the $U(1)$ explicitly embedded in a non-abelian group, there remains a “memory” of the underlying non-abelian structure, and the contributions vanish at leading order. Nevertheless, various effects below the string scale reintroduce kinetic-mixing. In particular, low energy supersymmetry (SUSY) breaking will *always* split the matter multiplets contributing to the kinetic-mixing by an amount of order the supersymmetry breaking scale in the hidden sector. If supersymmetry breaking is mediated by gravity, then the latter is roughly $M_{\text{SUSY}} \sim 10^{11}$ GeV and $\chi \sim 10^{-6}$ can naturally be obtained [7].

Other interesting classes of string models are mostly based on configurations of Dirichlet (D)-branes and/or fluxes. (Dp branes are membrane-like objects with p spatial dimensions, where open strings can be attached, thereby becoming matter and gauge fields.) Particularly interesting are the intermediate scale models with $M_s \sim \sqrt{M_{\text{Pl}} M_P} \sim 10^{11}$ GeV. The so-called bottom-up approach [12] is a good way to ascertain general properties of these models; the idea is to fix the gauge properties of the SM or minimal supersymmetric SM (MSSM) *locally* in the compactified space using D-branes, without fixing the *global* details of the compactification. The latter have

only a minor affect on the visible sector phenomenology and may vary. However, the *global* set-up almost certainly requires anti-D-branes in the bulk which absorb certain types of unwanted (Ramond-Ramond) tadpoles but leave supersymmetry broken [12]. Inevitably, these set-ups predict kinetic-mixing and therefore the existence of millicharged particles [8, 9]. This phenomenon generally arises in these models because the hidden sector of D-branes in the bulk carry additional hidden $U(1)$ factors (possibly emerging from $U(N)$), which interact with the visible sector MSSM branes by exchanging closed string modes through the bulk (cf. Fig. 2). Such a closed string exchange (a cylinder diagram) can also be interpreted in the “open string channel” as a kinetic-mixing diagram as shown in Fig. 1(b); the one-loop open string diagram has a heavy string in the loop stretched between brane and anti-brane. The masses of the modes in the loop are given by their stretching energy proportional to their length (i.e. the distance between brane and anti-brane). The reason, the one-loop contributions do not cancel, is that the presence of anti-branes breaks supersymmetry, which is in fact integral to this particular scenario. Consequently there is a residual contribution to kinetic-mixing and hence χ , which is again given by the amount of supersymmetry breaking.

Consider non-degenerate radii, with our three infinite dimensions, $d + p - 3$ large dimensions of radius $R_{i=4..p+d} = R$, and $9 - d - p$ small space dimensions of radius $R_{i=p+d+1..10} = r$, with the visible and hidden sectors living on stacks of Dp -branes wrapping the small dimensions (where $p \geq 3$), and with the distance between hidden and visible branes being generically of size the compact dimension. The result for the mixing parameter when the hidden $U(1)_b$ is unbroken can be written as [8]

$$\chi \sim \pi \frac{\alpha_p}{N} \text{Tr}(\lambda_a) \text{Tr}(\lambda_b) \times \left(\frac{2^{(8-p)/2}}{\alpha_p} \frac{M_s}{M_P} \right)^{\frac{2(5-p)}{6-p}} \left(\frac{R}{r} \right)^{\frac{d-p+3}{6-p}}, \quad (7)$$

(For the present discussion the spatial dimensions of the branes is assumed to be p for both visible and hidden sectors; for the more general set-up, see Ref. [8].) The integer N is a factor corresponding to the Z_N point-group symmetry of the fixed point, a typical value for three generation models being $N = 3$. The Chan-Paton matrices $\lambda_{a,b}$ come from the vertex operators for photon emission off the open string loop, corresponding to the crosses in Figs. 1(b), 2; typically $\text{Tr}(\lambda_{a,b}) \sim 1$. Finally, α_p is the value of the gauge coupling, for which one may take $\alpha_p \sim 1/24$ (the MSSM unification value). For example if $p = d = 3$, then

$$\chi \simeq 5 \times 10^{-10} \frac{R}{r} \left(\frac{M_s}{10^{11} \text{ GeV}} \right)^{\frac{4}{3}}, \quad (8)$$

FIG. 3: Possible values for the kinetic-mixing parameter χ , as a function of the string scale, in the bottom-up approach discussed in the text. Values in the shaded region above the black (lower) and blue (upper) lines are predicted in models with D3- and D4-branes, respectively, for $R/r \geq 1$. The red (vertical) line gives the largest string scale allowed by phenomenology in these models. The green solid (horizontal) line corresponds to the preferred value $\chi \simeq 3 \times 10^{-6}$ for the millicharged particle interpretation of PVLAS, while the green dashed line gives an idea of the sensitivity of near future searches for millicharged particles.

so that an $R/r \sim 10^4$ would give the right value. Varying $p \geq 3$ and d , there is considerable freedom to reduce the ratio R/r , whilst still having $\chi = 10^{-6}$ (cf. Fig. 3). (Note, that the strong bounds on M_s obtained in Ref. [8] from the astrophysical bounds on ϵ do not apply in general due to the charge screening mechanism proposed in Ref. [6].)

Let us now comment on the fermion sector. A crucial ingredient for the electric charge screening mechanism in the stellar plasma is that there have to be *two* hidden sector $U(1)$ s, and the hidden sector fermions have to have charge $(0, \epsilon, -\epsilon)$ under the visible and the two hidden sector $U(1)$ factors, respectively [6]. Note that this is generic in open string models with hidden D-brane sectors, since hidden sector fields arising from open strings stretched between hidden sector branes naturally fall into the bifundamental representation of the two hidden sector $U(1)$ s.

The remaining question is whether there is any reason to expect the mass of the millicharged particles to be ~ 0.1 eV, and indeed there is. Since one of the $U(1)$ s is by assumption unbroken, it is natural to expect some fermions on the hidden brane to be initially massless. However, as we have seen, the $U(1)_b$ mixes with the visible sector symmetries, and of course here the MSSM requires mass terms, namely the μ -term for the Higgs. Generally, these induce two-loop mass terms in the hidden sector, as shown in Fig. 4. These contributions are diluted by the same volume factors ($V_{||}, V_{\perp}$) that cause the dilution of gravity, given by [13]

$$M_s^2/M_P^2 \sim \alpha_b^2 V_{||}/V_{\perp}; \quad (9)$$

FIG. 4: Two-loop induced fermion masses in the hidden sector are directly related to the supersymmetry breaking. The dashed lines and gauge bosons are all stretched between visible and hidden branes, while the internal propagator is a fermion mass propagator of order 1 TeV.

at one-loop the stretched states get a mass-splitting (the inner loop of Fig. 4) of order $(V_{\parallel}/V_{\perp})M_s^2/\mu$, where $\mu \sim 1$ TeV, and at two-loops the diagram receives another volume factor $V_{\parallel}/V_{\perp}$. In total, therefore, the mass induced in the hidden sector is

$$m_{\text{hidden}} = \alpha_p^{-4} \frac{M_s^6}{M_P^4 \mu} \sim \alpha_p^{-4} \frac{M_W^2}{M_P}, \quad (10)$$

which is roughly of the right order of magnitude for $\alpha_p \sim 1/24$. Note, that no new scales beyond what is assumed for the MSSM have been introduced. The conclusion is quite general: sub eV masses are induced for hidden sector fermions if there are fermions with mass ~ 1 TeV in the visible sector.

Therefore, both closed and open string models not only predict the necessary extra $U(1)$ factors and correct fermion representation, but can also accommodate the required values of the kinetic-mixing parameter and fermion masses.

As far as the gauge boson mass is concerned, the good news is that, naturally, a Higgs appears in the bifundamental representation of the hidden $U(1)$ s, leaving automatically one mixed $U(1)$ massless, as required in the charge screening mechanism of [6]. It remains to be seen whether one can come up with a mechanism, perhaps based on accidental symmetries, to stabilize its small sub-eV scale (cf. Eq. (3)). Finally, there may still be room for an additional light spin-zero particle coupled to the hidden sector fermions which could then play the role of an axion-like particle [6, 14].

There are a number of exciting possibilities to test such a scenario in laboratory experiments, allowing for experimental insights into string theory with less model dependence than astrophysical or cosmological considerations.

The existence of millicharged particles can be tested [1] by improving the sensitivity of instruments for the detection of vacuum magnetic birefringence and dichroism [2, 15, 16, 17, 18, 19]. Another sensitive tool is Schwinger pair production in strong electric fields, as they exist, for example, in accelerator cavities [20]. A classical probe is the search for invisible orthopositronium decays [21]. We expect that all these laboratory experiments will probe into the range $\epsilon \sim 10^{-8} - 10^{-6}$. Ad-

ditional axion-like particles, coupled to the millicharged fermions, may be observed in photon regeneration experiments [17, 18, 22, 23, 24, 25, 26]. These searches are complementary to and presently more sensitive than collider techniques based on the effect of kinetic-mixing on precision electroweak observables [27].

In conclusion, string theory models can provide the necessary ingredients for an explanation of the PVLAS result. Independent of this, intermediate string scales $M_s \sim 10^{11}$ GeV predict the existence of millicharged particles with $\epsilon \gtrsim 0.4 \times 10^{-9}$, testable with near future laboratory experiments.

-
- [1] H. Gies, J. Jaeckel and A. Ringwald, hep-ph/0607118.
 - [2] E. Zavattini *et al.* [PVLAS Collaboration], Phys. Rev. Lett. **96**, 110406 (2006).
 - [3] L. B. Okun, Sov. Phys. JETP **56**, 502 (1982).
 - [4] B. Holdom, Phys. Lett. B **166**, 196 (1986).
 - [5] S. Davidson, S. Hannestad and G. Raffelt, JHEP **0005**, 003 (2000).
 - [6] E. Masso and J. Redondo, hep-ph/0606163.
 - [7] K. R. Dienes, C. F. Kolda and J. March-Russell, Nucl. Phys. B **492**, 104 (1997).
 - [8] S. A. Abel and B. W. Schofield, Nucl. Phys. B **685**, 150 (2004).
 - [9] S. Abel and J. Santiago, J. Phys. G **30**, R83 (2004).
 - [10] J. Polchinski, String Theory, Vol. I and II, Cambridge University Press, Cambridge, 1998.
 - [11] B. Batell and T. Cherguetta, Phys. Rev. D **73**, 045016 (2006).
 - [12] G. Aldazabal, L. E. Ibanez, F. Quevedo and A. M. Uranga, JHEP **0008**, 002 (2000).
 - [13] L. E. Ibanez, C. Munoz and S. Rigolin, Nucl. Phys. B **553**, 43 (1999).
 - [14] J. Jaeckel, E. Masso, J. Redondo, A. Ringwald and F. Takahashi, hep-ph/0605313.
 - [15] R. Cameron *et al.* [BFRT Collaboration], Phys. Rev. D **47**, 3707 (1993).
 - [16] S. J. Chen *et al.* [Q & A Collaboration], hep-ex/0308071.
 - [17] C. Rizzo [BMV Collaboration], 2nd ILIAS-CERN-CAST Axion Academic Training 2006, <http://cast.mppmu.mpg.de/>
 - [18] P. Puginat *et al.*, Czech. J. Phys. **55**, A389 (2005); Czech. J. Phys. **56**, C193 (2006).
 - [19] T. Heinzl *et al.*, hep-ph/0601076.
 - [20] H. Gies, J. Jaeckel and A. Ringwald, hep-ph/0608238.
 - [21] A. Rubbia, Int. J. Mod. Phys. A **19**, 3961 (2004).
 - [22] A. Ringwald, Phys. Lett. B **569**, 51 (2003).
 - [23] R. Rabadan, A. Ringwald and K. Sigurdson, Phys. Rev. Lett. **96**, 110407 (2006).
 - [24] G. Cantatore [PVLAS Collaboration], 2nd ILIAS-CERN-CAST Axion Academic Training 2006, <http://cast.mppmu.mpg.de/>
 - [25] U. Kötz, A. Ringwald and T. Tscheuschner, hep-ex/0606058.
 - [26] K. Baker [LIPSS Collaboration], 2nd ILIAS-CERN-CAST Axion Academic Training 2006, <http://cast.mppmu.mpg.de/>
 - [27] J. Kumar and J. D. Wells, hep-ph/0606183.