



Exclusive processes at HERA

Robert Ciesielski (DESY)
on behalf of H1 and ZEUS Collaborations



$$\gamma^* p \rightarrow V p$$


$$V = \rho, \omega, \varphi, J/\Psi, \Psi', Y, \gamma$$



International Conference on High Energy Physics, ICHEP08
Philadelphia, July 29-August 5, 2008


Exclusive processes @HERA - VM Production, DVCS

HERA (DESY): collisions of 27.5 GeV e with 920 GeV p ($\sqrt{s}=318\text{ GeV}$)



$$\gamma^{(*)} p \rightarrow V p$$

exclusive



$$\gamma^{(*)} p \rightarrow V Y$$

proton dissociative (dominates at higher- t)

VM : vector meson or real photon

$\rho, \omega, \varphi, J/\Psi, \Psi', Y, \gamma$

Q^2 : photon virtuality

$0 < Q^2 < 180\text{ GeV}^2$

W : CM energy of the $\gamma^* p$ system ($x=Q^2/W^2$)

$20 < W < 290\text{ GeV}$


t : (4-mom. transfer) 2 at the proton vertex

$|t| < 30\text{ GeV}^2$


HERA data spans exceptionally wide range of kinematic variables

VM Production Mechanisms @ HERA

soft: VDM + Regge theory (hadron level)



hard: pQCD (parton level)



soft Pomeron exchange: $\alpha_{IP}(t) = \alpha_{IP}(0) + \alpha_{IP}' \cdot t$
 $\alpha_{IP}(0) = 1.08, \alpha_{IP}' = 0.25 \text{ GeV}^{-2}$

$$\frac{d\sigma}{dt} \propto e^{-b|t|} (W/W_0)^{4(\alpha_{IP}(t)-1)}$$

- slow rise with W : $\delta \approx 0.22$

$$\sigma(W) \propto W^\delta$$

- shrinkage of the diffractive peak with W :

$$b(W) = b(W_0) + 4\alpha_{IP}' \ln(W/W_0)$$

$$b(W_0) \approx 10 \text{ GeV}^{-2}$$

$$\frac{d\sigma}{dt} \propto e^{-b|t|}$$

$$\sigma_L \propto \alpha_s(Q^2) [xg(x, Q^2)]^2 / Q^6$$


- fast rise with W : $\delta \approx 0.7$
 $(x = Q^2/W^2, \text{ gluon density rises at low-}x)$

- universal t -dependence:

$$b_{2g} \approx 4 - 5 \text{ GeV}^{-2} \quad \text{and} \quad \alpha_{IP}' \approx 0$$

Change of regime expected with rising Q^2 , M_{VM}^2 or t (hard scales of the process)

DVCS – Deeply Virtual Compton Scattering



$$\gamma^* p \rightarrow \gamma p$$



Similar to VM production, but with real photon in the final state

- no VM wave-function uncertainty (non-perturbative part)
- easier access to GPDs - Generalised (skewed) PDFs

GPDs describe the correlations between two partons (x_1, x_2) which differ by

longitudinal, $x = x_1 - x_2$, and transverse, t , momentum at a given Q^2 .

(~3D picture of the proton) Important for exclusive Higgs production @LHC




Irreducible QED background from $ep \rightarrow ep\gamma$ Bethe-Heitler process
Sensitive to the real part of the amplitude via the QCD-QED interference

$$\sigma = \sigma^{DVCS} + \sigma^{BH} \pm \sigma^{interf.}$$

Vector Mesons in Photoproduction [Q²=0]


$\gamma p \rightarrow V p$



$$\sigma(W) \propto W^\delta$$

soft physics: $\delta \approx 0.22$

$\gamma p \rightarrow Y p$




pQCD describes the steep rise of the cross section with M_Y as a hard scale, sensitivity to gluon GPDs

Process becomes harder (steeper W dependence) as M_{VM} becomes larger


VM mass sets hard scale in photoproduction

ρ, φ mesons, W-dependence vs Q^2

ZEUS



$$\sigma(W) \propto W^\delta \text{ in bins of } Q^2$$






Steeper W dependence as Q^2 becomes larger
 Q^2 sets hard scale for light vector mesons

ρ, φ mesons, t-dependence vs Q^2

$$\frac{d\sigma}{dt} \propto e^{-b|t|} \quad \text{in bins of } Q^2$$

b describes the transverse size of the interaction region $b \propto R_p^2 + R_{q\bar{q}}^2$



b slope decreases with Q^2 : $b = 10 \rightarrow 5 \text{ GeV}^{-2}$
Transverse size of dipole decreases with Q^2


DVCS vs Q^2

$$\gamma^* p \rightarrow \gamma p$$


W dependence:

$$\sigma(W) \propto W^\delta$$

t dependence:




First direct measurement of t-dependence using the Leading Proton Spectrometer (proton tag, only elastic contribution)





Steep rise with W,
no significant dependence of δ on Q^2 (within errors)

$$\frac{d\sigma}{dt} \propto e^{-b|t|}$$



First measurement of
 Q^2 -dependence of the b slope

All VMs and DVCS, W and t-dependence vs Q^2+M^2




Similar behaviour of δ and b with Q^2+M^2 for all VM (ρ , φ , J/ψ) and DVCS
 Transition from *soft* to *hard* regime with increasing of hard scale

At higher Q^2+M^2 : point-like dipole probes gluon cloud of the proton (pQCD region)

Comparison to theoretical models, ρ

High precision of data → improved understanding of non-perturbative
quantities: VM wave-function, PDFs (low-x gluons)




$$A \propto \psi_{q\bar{q}}^{Y^*} \otimes \sigma_{q\bar{q}-p} \otimes \psi_{q\bar{q}}^V$$

Models differ for $\sigma_{q\bar{q}-p}$ and $\psi_{q\bar{q}}^V$

KMW - Kowalski-Motyka-Watt

FSS - Forshaw-Sandapen-Shaw


MRT - Martin-Ryskin-Teubner



Large differences between the models and PDF input. HERA data provide constraints

Comparison to theoretical models, DVCS

GPD model – A. Freund et al. (NLO QCD) with
GPD parametrisation by J. Pumplin et al.



Two dimensionless observables:


$$S = \sqrt{\frac{\sigma_{DVCS} Q^4 b(Q^2)}{1 + \rho^2}}$$

S measures the Q^2 evolution of GPD



$$R = \frac{\text{Im}A(\gamma^* p \rightarrow \gamma p)_{t=0}}{\text{Im}A(\gamma^* p \rightarrow \gamma^* p)_{t=0}} = \frac{4\sqrt{\pi} \sigma_{DVCS} b(Q^2)}{\sigma_T(\gamma^* p \rightarrow X) \sqrt{(1 + \rho^2)}}$$

R measures the ratio of GPDs to PDFs ie. skewing effect ($x_1 - x_2$),
R=1 if no skewing ($x_1 = x_2$, GPDs \rightarrow PDFs)

NLO pQCD model based on GPDs describes the $S(Q^2)$ and $R(Q^2)$.
Data has proven its potential to constrain gluon (and sea) GPDs.



DVCS – Beam Charge Asymmetry



The QCD-QED interference term is sensitive to the real part of the QCD amplitude,
It changes sign with lepton beam charge:

$$\sigma = \sigma^{DVCS} + \sigma^{BH} \pm \sigma^{interf.}$$

+ for incoming e^+ beam
- for incoming e^- beam


Beam Charge Asymmetry (BCA) vs ϕ

ϕ is the angle between two planes defined by incoming and outgoing electron and γ^* and outgoing proton

$$BCA = \frac{\sigma^+ - \sigma^-}{\sigma^+ + \sigma^-} = p_1 \cdot \cos(\phi) + \dots$$


$$p_1 = 0.17 \pm 0.03 \pm 0.05 \quad \text{sensitive to GPDs}$$

First measurement in the low-x region at HERA



ρ and φ mesons, Helicity Structure

Angular distributions \rightarrow 15 Spin Density Matrix Elements $r_{kl}^{ij} \rightarrow$ helicity amplitudes $T_{\lambda_\gamma \lambda_{VM}}$




15 combinations of SDME measured in bins of Q^2 and t (not shown).

Test of *s*-channel helicity conservation (SCHC, observed in soft processes):

VM retains helicity of the photon, the only allowed transitions are: $T_{11}: \gamma_T \rightarrow V_T$, $T_{00}: \gamma_L \rightarrow V_L$




$r_{00}^5 \propto T_{01}$, T_{0-1} violates SCHC
(it measures single helicity flip, $\propto \sqrt{|t|}$)



ρ and φ mesons, Polarised Cross Sections

$$\sigma = \sigma_T + \epsilon \cdot \sigma_L \quad \epsilon \approx 0.996$$

Vector Meson production processes → unique opportunity to extract $R = \sigma_L / \sigma_T$
 R measured from 1DIM angular distributions $f(\cos \theta_h, r_{00}^{04})$, in SCHC approximation





R rises with Q^2 . σ_L dominates at higher Q^2 .
 Not shown: R independent of W and t ,
 R decreases with M_ρ

σ_L and σ_T have different $Q^2 + M^2$ dependence.
 Models based on pQCD describe well σ_L , but not σ_T .

ρ and φ mesons, proton vertex factorisation

H1 test of proton vertex factorisation in DIS regime (shown already by ZEUS for $Q^2=0$)



Ratio of p-diss to elastic cross sections consistent with no dependence on Q^2

Similar values for ρ and φ mesons (within errors)


Probability of proton dissociation is independent of the projectile

Summary


- New high statistics measurements of ρ , φ mesons and DVCS
- All observed features are compatible with the expectations of pQCD
 - The cross section is rising with W and the rise becomes steeper as Q^2 or M_v increases
 - The exponential slope of the t distribution is decreasing with Q^2 and levels off at about 5 GeV^{-2}
 - The ratio of cross sections induced by longitudinally and transversely polarised virtual photons increases with Q^2 , but is independent of W and t
 - The violation of SCHC is observed for light vector mesons
 - Proton vertex factorisation is observed in DIS regime
- DVCS process is well described by pQCD+GPD model
- Non of the models is able to describe all the features of the data for light vector mesons

ρ and φ mesons, Polarised Cross Sections


ZEUS



ZEUS



ZEUS



Significant dependence of R on M_V


R consistent with no dependence on W and t .

σ_L and σ_T have the same W and t dependence

- the same transverse size of the interaction region ($b_L = b_T$)
- the same rise of the cross section with energy
- ... the large spatial configurations of a dipole for σ_T are suppressed in VM production?

ρ and φ mesons, Helicity Structure

Angular distributions \rightarrow 15 Spin Density Matrix Elements r_{kl}^{ij} \rightarrow helicity amplitudes $T_{\lambda_y \lambda_{VM}}$
 15 combinations of SDME measured in bins Q^2 and t . Test of *s-channel helicity conservation*, SCHC.
 (SCHC: VM retains helicity of the photon, the only allowed transitions are: $T_{11}: \gamma_T \rightarrow V_T$, $T_{00}: \gamma_L \rightarrow V_L$)




H1 ρ electroproduction (preliminary)

- H1 HERA-1 prel.
- ZEUS
- GK (GPD)

$|t| < 3 \text{ GeV}^2$

$r_{00}^5 \propto T_{01}, T_{0-1}$ violates SCHC (it measures single helicity flip, $\propto \sqrt{|t|}$)

18



H1 ρ electroproduction (preliminary)

- H1 HERA-1 prel.
- ZEUS

@ $Q^2 = 5 \text{ GeV}^2$

Increased precision of recent HERA data allows to:

- Study the VM and DVCS dynamics within QCD
- Test QCD in the transition region $soft \rightarrow hard$
- Given factorisation, test pQCD and constrain non-perturbative quantites