$F_2^{c\bar{c}}$ measurements at HERA

Marcello Bindi¹, on behalf of the ZEUS and H1 Collaborations

1- University of Bologna and INFN - Department of Physics
via Irnerio, 46 - 40127 - Bologna -Italy

The study of charm production at the ep collider HERA is particularly interesting since it provides sensitivity to the gluon density in the proton and it allows different models and schemes of pQCD to be tested. New measurements of the charm contribution $F_2^{c\bar{c}}$ to the proton structure function F_2 for deep inelastic scattering in $e^\pm p$ collisions at HERA are presented. These results have been achieved by the two experiments, ZEUS and H1, using different techniques: tagging of D mesons (D^* , D^0 , $D^\pm$..) in the final states or measuring the displacement of the decay products of charm decays, exploiting their long lifetime. The data are compared to next-to-leading order pQCD calculations.

1 Introduction

At the ep collider HERA, in deep inelastic scattering (DIS) events heavy quarks are mainly produced via boson-gluon fusion in which the virtual photon interacts with a gluon from the proton. From the total cross section for heavy quark production, the charm contribution to the proton structure function F_2 can be extracted and compared to NLO QCD predictions. Since these predictions are sensitive to the gluon density of the proton, the measurement can help in constraining it.

2 D meson tagging

The golden mode is the reconstruction of a $D^*(2010)$ meson through its decay $D^{*+} \rightarrow D^0 \pi_S^+ \rightarrow K^- \pi^+ \pi_S^+$ and the corresponding antiparticle decay. The production of D^* mesons has been measured in the kinematic region $1.5 < Q^2 < 1000 \text{ GeV}^2$, $0.02 < y < 0.7$, $1.5 < p_T(D^*) < 15 \text{ GeV}$ and $|\eta(D^*)| < 1.5$. The double-differential cross section in y and Q^2 has been measured and used to extract the open-charm contribution to F_2 , by using an NLO QCD calculation to extrapolate outside the measured $p_T(D^*)$ and $\eta(D^*)$ region. Both the ZEUS and H1 Collaborations have used this method to extract $F_2^{c\bar{c}}$ from D mesons cross sections.

Other charmed mesons can also be reconstructed: the most copiously produced are D^0 and $D^\pm$. In contrast to the D^* , these mesons have a sizeable lifetime; the reconstruction of a displaced secondary vertices can be used to enhance the signal to background ratio. A cut on the decay length significance has been applied, for example in the production of D^+ mesons (Fig. 1). Differential cross sections in Q^2, x (see Fig. 2 and 3 for D^0 mesons), $p_T(D)$ and $\eta(D)$ have been measured together with double differential cross sections to extract $F_2^{c\bar{c}}$.

Charm tagging with mesons gives a clean signature of charm production, but the extrapolation to the total charm cross section can be large, especially in the low Q^2 region [2].

Figure 1: The D^+ mass peak, reconstructed from the decay $D^+ \rightarrow K^- \pi^+ \pi^-$, before (left) and after (right) the three sigma cut on the significance of the displace vertex reconstructed from the D^+ decay products.

Figure 2: Single differential D^0 cross sections as a function of Q^2 compared to the NLO calculation of HVQDIS. Preliminary ZEUS data from HERA II (points) are compared to the published values from HERA I (triangles).

Figure 3: Single differential D^0 cross sections as a function of x (solid points) compared to the NLO calculation of HVQDIS. The contribution from beauty, as predicted by RAPGAP program, is shown as a dashed line.

3 Inclusive lifetime tagging

An alternative method to tag charm production takes advantage of the long lifetime of the charmed particles, by identifying tracks having impact parameter, δ , significantly displaced from the event vertex, or by reconstructing secondary vertices from the meson decay products. The distributions of the impact parameter and of its significance in an inclusive DIS data sample are shown in Fig. 4, together with the contributions coming separately from beauty, charm and light flavour background. A clear asymmetry of the significance distribution for heavy flavours can be seen, and the fraction of charm and beauty in the data can be extracted by fitting the distribution with the different flavour components.

This method has been used by the H1 Collaboration to obtain the results presented here [9]. The advantage of this technique is that the kinematic region for the measurement of charm production is significantly enlarged, and therefore the extrapolation needed for the measurement of $F_2^{c\bar{c}}$ is strongly reduced.

Figure 4: The distribution of the signed impact parameter, δ , of the track in the $x-y$ plane (left) and of the impact parameter significance, $\delta/\sigma(\delta)$, of the track with second highest absolute significance (S_2). H1 preliminary points from HERA II are compared with the expectation from the Monte Carlo simulation for the various quark, after the fit to the data.

4 Theoretical predictions

The program HVQDIS [10] is used to calculate theoretical predictions for D meson production cross sections at NLO in pQCD. It was used by the ZEUS and H1 Collaborations to extrapolate the measured cross section for a particular D meson final state to $F_2^{c\bar{c}}$. In this program, the production of heavy flavours is described using the fixed flavour number scheme (FFNS), in which the charm quark is not an active flavour in the proton PDF, but is generated dynamically in the hard subprocess. The ZEUS and H1 measurements of extracted $F_2^{c\bar{c}}$ in this way should therefore be compared with NLO QCD predictions evaluated in the FFNS.

5 Results

The charm contribution to the proton structure function, $F_2^{c\bar{c}}$, has been measured at HERA by the ZEUS [2, 3, 4, 5] and H1 [6, 7, 8] collaborations in a wide kinematic region of x ($0.00002 < x < 0.03$) and of the photon virtuality, Q^2 ($1 < Q^2 < 1000 \text{ GeV}^2$). The results for the reduced cross section $\tilde{\sigma}^{c\bar{c}}(x)$ in 5 different Q^2 bins and for $F_2^{c\bar{c}}(x, Q^2)$ are shown in Fig. 5 and 6.

Charm has been tagged by reconstructing a particular charmed meson in the final state or using the long lifetime of the charm decay products. The ZEUS and H1 measurements, obtained from charmed meson reconstruction, are compared with the H1 results from inclusive lifetime measurements. The agreement between the data of the two experiments is good, validating the two different analysis procedures. The data rise with increasing Q^2 , with the rise becoming steeper at lower x , demonstrating the property of scaling violation in charm production.

The data are also compared with pQCD predictions at NLO. Two different parametrisations of the proton PDF have been used for the NLO QCD calculations, in order to check the sensitivity of the predictions to different gluon densities. The PDFs used were CTEQ5F3 [11] and MRST2004FF3 [12], extracted mainly from inclusive scattering data.

The data are in general well described by NLO QCD. The theoretical predictions based on different PDFs show differences in the low x region (Fig. 6), demonstrating that the data

Figure 5: The measured HERA I and II averaged charm reduced cross section $\sigma_{c\bar{c}}$ as a function of x for 5 different Q^2 values compared with predictions of QCD.

Figure 6: The charm contribution to the proton structure function $F_2, F_2^{c\bar{c}}$, at x values between 0.00003 and 0.03 as a function of Q^2 .

have the power to constrain the gluon density of the proton.

References

- [1] Slides:
<http://indico.cern.ch/contributionDisplay.py?contribId=244&sessionId=30&confId=24657>
- [2] ZEUS Coll., Phys. Rev. **D 69**, 012004 (2004).
- [3] ZEUS Coll., JHEP **07**, 074 (2007).
- [4] ZEUS Coll., *Proceedings of the XV International Workshop on Deep Inelastic Scattering and Related Subjects*, Munich, Germany, April 16-20, 2007.
- [5] ZEUS Coll., *Submitted to the Europhysics Conference on High Energy Physics, EPS HEP 2007*, Manchester, England, July 19-25, 2007.
- [6] H1 Coll., Phys. Lett. **B 528**, 199 (2002).
- [7] H1 Coll., Eur. Phys. J. **C 40**, 349 (2005).
- [8] H1 Coll., Eur. Phys. J. **C 45**, 23 (2006).
- [9] A. Mehta [H1 Coll.], *these proceedings*.
- [10] B. W. Harris and J. Smith, Phys. Rev. **D 57**, 2806 (1998).
- [11] CTEQ Coll., Eur. Phys. J. **C 12**, 375 (2000).
- [12] A. D. Martin, W. J. Stirling and R. S. Thorne, Phys. Lett. **B 636**, 259 (2006).