Analytic results for the one-loop NMHV $H\bar{q}qgg$ amplitude

Simon Badger,^a John M. Campbell,^b R. Keith Ellis^c and Ciaran Williams^d

^a*Deutsches Elektronen-Synchrotron DESY,
Platanenallee 6, D-15738 Zeuthen, Germany*

^b*Department of Physics and Astronomy, University of Glasgow,
Glasgow, G12 8QQ, U.K.*

^c*Fermilab,
Batavia, IL 60510, U.S.A.*

^d*Department of Physics, University of Durham,
Durham, DH1 3LE, U.K.*

E-mail: simon.badger@desy.de, j.campbell@physics.gla.ac.uk,
ellis@fnal.gov, ciaran.williams@durham.ac.uk

ABSTRACT: We compute the one-loop amplitude for a Higgs boson, a quark-antiquark pair and a pair of gluons of negative helicity, i.e. for the next-to-maximally helicity violating (NMHV) case, $\mathcal{A}(H, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-)$. The calculation is performed using an effective Lagrangian which is valid in the limit of very large top quark mass. As a result of this paper all amplitudes for the transition of a Higgs boson into 4 partons are now known analytically at one-loop order.

KEYWORDS: Higgs Physics, NLO Computations, Hadronic Colliders, QCD

ARXIV EPRINT: [0910.4481](https://arxiv.org/abs/0910.4481)

Contents

1	Introduction	1
2	Effective Lagrangian	3
3	Definition of colour ordered amplitudes	4
4	Known analytic results for Higgs + 4 parton amplitudes	4
4.1	Tree graph results	5
4.2	One-loop results	6
4.2.1	$H\bar{q}q\bar{q}q$ amplitudes	6
4.2.2	$Hgggg$ amplitudes	6
4.2.3	$H\bar{q}qgg$ amplitudes	6
5	One-loop results	8
5.1	Results for $A_{4;1}(\phi, 1_{\bar{q}}, 2_q, 3_g^-, 4_g^-)$	8
5.1.1	$A_4^L(\phi, 1_{\bar{q}}, 2_q^+, 3_g^-, 4_g^-)$	8
5.1.2	$A_4^R(\phi, 1_{\bar{q}}, 2_q^+, 3_g^-, 4_g^-)$	11
5.1.3	$A_4^f(\phi, 1_{\bar{q}}, 2_q^+, 3_g^-, 4_g^-)$	12
5.1.4	Relation for rational terms	12
5.2	Results for $A_{4;3}(\phi, 1_{\bar{q}}, 2_q, 3_g^-, 4_g^-)$	13
5.2.1	Box-related terms for $A_4^L(\phi, 1_{\bar{q}}, 2_g^-, 3_q^+, 4_g^-)$	13
5.2.2	Full result for $A_{4;3}$	14
6	Numerical results	15
7	Conclusions	15
A	Spinor notation	16
B	Definitions of special functions	17

1 Introduction

The hunt for the standard model Higgs boson is about to enter the endgame phase. The lower and upper limits coming from direct searches at LEP [1] and indirect constraints from precision electroweak data from the Tevatron and LEP [2] are now supplemented by the first direct limit from a hadron collider [3, 4]. With the increase of luminosity at the Tevatron and the advent of running at the LHC, a discovery or a more stringent set of limits is to be expected.

An important search channel for the Higgs boson, in the mass range $115 < m_H < 160$ GeV, is production via weak boson fusion [5]. A Higgs boson produced in this channel is expected to be produced relatively centrally, in association with two hard forward jets. These striking kinematic features are expected to enable a search for such events despite the otherwise overwhelming QCD backgrounds. Confidence in the theoretical prediction for the Higgs signal process is based upon knowledge of next-to-leading order corrections in both QCD [6–8] and in the electroweak sector [9, 10].

However, in addition to the weak process, a significant number of such events may also be produced via the strong interaction. In order to accurately predict the signal and, in particular, to simulate faithfully the expected significance in a given Higgs model, a fully differential NLO calculation of QCD production of a Higgs and two hard jets is also required.

In the Standard Model the Higgs couples to two gluons via a top-quark loop. Calculations which involve the full dependence on m_t are difficult and a drastic simplification can be achieved if one works in an effective theory in which the mass of the top quark is large [11–13]. For inclusive Higgs production this approximation is valid over a wide range of Higgs masses and, for processes with additional jets, the approximation is justified provided that the transverse momentum of each jet is smaller than m_t [14]. Tree-level calculations have been performed in both the large- m_t limit [15, 16] and with the exact- m_t [14] dependence.

Results for the one-loop corrections to all of the Higgs + 4 parton processes have been published in 2005 [17]. Although analytic results were provided for the Higgs $\bar{q}q\bar{q}q$ processes, the bulk of this calculation was performed using a semi-numerical method. In this approach the loop integrals were calculated analytically whereas the coefficients with which they appear in the loop amplitudes were computed numerically using a recursive method. Although some phenomenology was performed using this calculation [18], the implementation of fully analytic formulae will lead to a faster code and permit more extensive phenomenological investigations.

In recent years enormous progress has been made in solving the problem of evaluating virtual corrections to NLO scattering processes. Building upon the remarkable work of Bern, Dixon, Dunbar and Kosower during the mid-nineties [19, 20], unitarity constructions for these virtual corrections have developed into an efficient algebraic technique. The modern generalised unitarity method utilises quadruple cuts with complex momenta to freeze four dimensional loop momenta and uniquely determine the box coefficients [22]. The computation of triangle and bubble coefficients is also reduced to an algebraic procedure by application of an OPP style integrand reduction [23–25] or using direct analytic extraction [26]. Further developments employing D-dimensional cutting techniques [27–30] extend the method to compute full one-loop amplitudes. The procedure is well suited to numerical implementations and a number of automated approaches have been developed to the point of phenomenological applications [31–39].

In this paper we derive a compact analytic formula for the Higgs NMHV amplitude with a quark-antiquark pair and two like-helicity gluons. This is achieved by splitting the real Higgs scalar into two complex scalars (ϕ and $\phi^\dagger$) such that the Higgs amplitude is recovered in the sum [40]. Coefficients of box, triangle and bubble integrals are computed

by applying the generalised unitarity method in four-dimensions. The rational terms are extracted from a Feynman diagram computation which is simplified using the knowledge of unphysical singularities in the cut-constructible terms.

The paper is organised as follows. In section 2 we describe the large top-mass approximation and the decomposition of the Higgs into self-dual (ϕ) and anti-self-dual ($\phi^\dagger$) components. Section 3 recalls the colour decomposition into primitive amplitudes and section 4 provides a guide to the current literature on Higgs + 4 parton amplitudes and recalls the known analytic results for $\phi\bar{q}qgg$ amplitudes that are needed to construct results for the Higgs boson. In section 5 we present analytic results that are sufficient for a complete description of the NMHV amplitude. We numerically evaluate the obtained expressions for the one-loop colour ordered amplitudes in section 6 before drawing our conclusions.

2 Effective Lagrangian

Our calculation is performed using an effective Lagrangian to express the coupling of gluons to the Higgs field [11],

$$\mathcal{L}_H^{\text{int}} = \frac{C}{2} H \text{tr} G_{\mu\nu} G^{\mu\nu}. \quad (2.1)$$

This Lagrangian is obtained by replacing the full one-loop coupling of the Higgs boson to the gluons via an intermediate top quark loop, by an effective local operator. The effective Lagrangian approximation is valid in the limit $m_H < 2m_t$. At the order required in this paper, the coefficient C is given by [12, 13],

$$C = \frac{\alpha_S}{6\pi v} \left(1 + \frac{11}{4\pi} \alpha_S\right) + \mathcal{O}(\alpha_S^3). \quad (2.2)$$

Here v is the vacuum expectation value of the Higgs field, $v = 246 \text{ GeV}$. The trace in eq. (2.1) is over the colour degrees of freedom which, since SU(3) generators in the fundamental representation are normalised such that $\text{tr} T^a T^b = \delta^{ab}$, implies that $\text{tr} G_{\mu\nu} G^{\mu\nu} = G_{\mu\nu}^a G^{a\mu\nu}$.

Following reference [40] we will introduce a complex scalar field,

$$\phi = \frac{1}{2}(H + iA), \quad \phi^\dagger = \frac{1}{2}(H - iA), \quad (2.3)$$

so that the effective Lagrangian, eq. (2.1), can be written as,

$$\begin{aligned} \mathcal{L}_{H,A}^{\text{int}} &= \frac{C}{2} \left[H \text{tr} G_{\mu\nu} G^{\mu\nu} + iA \text{tr} G_{\mu\nu} {}^* G^{\mu\nu} \right] \\ &= C \left[\phi \text{tr} G_{SD\ \mu\nu} G_{SD}^{\mu\nu} + \phi^\dagger \text{tr} G_{ASD\ \mu\nu} G_{ASD}^{\mu\nu} \right], \end{aligned} \quad (2.4)$$

where the gluon field strength has been separated into a self-dual and an anti-self-dual component,

$$G_{SD}^{\mu\nu} = \frac{1}{2}(G^{\mu\nu} + {}^* G^{\mu\nu}), \quad G_{ASD}^{\mu\nu} = \frac{1}{2}(G^{\mu\nu} - {}^* G^{\mu\nu}), \quad {}^* G^{\mu\nu} \equiv \frac{i}{2} \epsilon^{\mu\nu\rho\sigma} G_{\rho\sigma}. \quad (2.5)$$

Calculations performed in terms of the field ϕ are simpler than the calculations for the Higgs boson and, moreover, the amplitudes for $\phi^\dagger$ can be obtained by parity. In the final

stage, the full Higgs boson amplitudes are then written as a combination of ϕ and $\phi^\dagger$ components:

$$\begin{aligned} A(H, \{p_k\}) &= A(\phi, \{p_k\}) + A(\phi^\dagger, \{p_k\}) , \\ A(A, \{p_k\}) &= -i \left(A(\phi, \{p_k\}) - A(\phi^\dagger, \{p_k\}) \right) . \end{aligned} \quad (2.6)$$

3 Definition of colour ordered amplitudes

The colour decomposition of the $H\bar{q}qgg$ amplitudes is exactly the same as for the case $\bar{q}qgg$ which was written down in ref. [21]. For the tree graph there are two colour stripped amplitudes,

$$\mathcal{A}_4^{(0)}(\phi, 1_{\bar{q}}, 2_q, 3_g, 4_g) = Cg^2 \sum_{\sigma \in S_2} (T^{a_{\sigma(3)}} T^{a_{\sigma(4)}})_{i_2}^{\bar{i}_1} A_4^{(0)}(\phi, 1_{\bar{q}}, 2_q, \sigma(3), \sigma(4)) . \quad (3.1)$$

At one-loop level the colour decomposition is,

$$\begin{aligned} \mathcal{A}_4^{(1)}(\phi, 1_{\bar{q}}, 2_q, 3_g, 4_g) &= Cg^4 c_\Gamma \left[N_c \sum_{\sigma \in S_2} (T^{a_{\sigma(3)}} T^{a_{\sigma(4)}})_{i_2}^{\bar{i}_1} A_{4;1}(\phi, 1_{\bar{q}}, 2_q, \sigma(3), \sigma(4)) \right. \\ &\quad \left. + \delta^{a_3 a_4} \delta_{i_2}^{\bar{i}_1} A_{4;3}(\phi, 1_{\bar{q}}, 2_q; 3_g, 4_g) \right] . \end{aligned} \quad (3.2)$$

In these equations g is the strong coupling constant and c_Γ is the ubiquitous one-loop factor,

$$c_\Gamma \equiv \frac{1}{(4\pi)^{2-\epsilon}} \frac{\Gamma(1+\epsilon)\Gamma^2(1-\epsilon)}{\Gamma(1-2\epsilon)} . \quad (3.3)$$

The colour stripped amplitudes $A_{4;1}$ and $A_{4;3}$ can further be decomposed into primitive amplitudes,

$$\begin{aligned} A_{4;1}(\phi, 1_{\bar{q}}, 2_q, 3_g, 4_g) &= A_4^L(\phi, 1_{\bar{q}}, 2_q, 3_g, 4_g) - \frac{1}{N_c^2} A_4^R(\phi, 1_{\bar{q}}, 2_q, 3_g, 4_g) \\ &\quad + \frac{n_f}{N_c} A_4^f(\phi, 1_{\bar{q}}, 2_q, 3_g, 4_g) , \end{aligned} \quad (3.4)$$

and,

$$\begin{aligned} A_{4;3}(\phi, 1_{\bar{q}}, 2_q; 3_g, 4_g) &= A_4^L(\phi, 1_{\bar{q}}, 2_q, 3_g, 4_g) + A_4^R(\phi, 1_{\bar{q}}, 2_q, 3_g, 4_g) + A_4^L(\phi, 1_{\bar{q}}, 3_g, 2_q, 4_g) \\ &\quad + A_4^L(\phi, 1_{\bar{q}}, 2_q, 4_g, 3_g) + A_4^R(\phi, 1_{\bar{q}}, 2_q, 4_g, 3_g) + A_4^L(\phi, 1_{\bar{q}}, 4_g, 2_q, 3_g) . \end{aligned} \quad (3.5)$$

All of these colour decomposition equations, namely eqs. (3.1), (3.2), (3.4), (3.5) are equally valid if the ϕ is replaced by a $\phi^\dagger$ or a Higgs boson H . Sample diagrams contributing to each of the primitive amplitudes are shown in figure 1.

4 Known analytic results for Higgs + 4 parton amplitudes

In this section we review results from the literature and collect formulae, for both tree and one-loop results, that will be useful in constructing the Higgs NMHV amplitude.

Figure 1. Sample diagrams contributing to the primitive amplitudes (a) $A_4^L(\phi, 1_{\bar{q}}, 2_q, 3_g, 4_g)$, (b) $A_4^R(\phi, 1_{\bar{q}}, 2_q, 3_g, 4_g)$, (c) $A_4^L(\phi, 1_{\bar{q}}, 2_g, 3_q, 4_g)$, and (d) $A_4^f(\phi, 1_{\bar{q}}, 2_q, 3_g, 4_g)$. The ϕ field can attach to any gluon line in the diagram.

4.1 Tree graph results

The results for the tree graphs that are primarily of interest here, i.e. $\phi\bar{q}qgg$ amplitudes with gluons of the same helicity, are:

$$-iA_4^{(0)}(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-) = -\frac{\langle 3|(1+4)|2\rangle^2 \langle 41\rangle}{[24] s_{124}} \left[\frac{1}{s_{12}} + \frac{1}{s_{41}} \right] - \frac{\langle 4|(1+3)|2\rangle^2 \langle 13\rangle}{[23] s_{12} s_{123}} + \frac{\langle 1|(3+4)|2\rangle^2}{\langle 12\rangle [24] [23] [34]}, \quad (4.1)$$

$$-iA_4^{(0)}(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^+, 4_g^+) = 0, \quad (4.2)$$

and for the subleading colour piece,

$$-iA_4^{(0)}(\phi, 1_{\bar{q}}^-, 2_g^-, 3_q^+, 4_g^-) = -\frac{\langle 4|(1+2)|3\rangle^2}{[12] [23] s_{123}} - \frac{\langle 2|(1+4)|3\rangle^2}{[34] [41] s_{341}}, \quad (4.3)$$

$$-iA_4^{(0)}(\phi, 1_{\bar{q}}^-, 2_g^+, 3_q^+, 4_g^+) = 0. \quad (4.4)$$

A brief summary of our spinor notation is given in appendix A. Compact analytic expressions for all helicity amplitudes are presented in references [45, 46].

By using parity and charge conjugation [47], we can relate these $\phi\bar{q}qgg$ amplitudes to ones for $\phi^\dagger\bar{q}qgg$ with the same helicity assignments of quark and antiquark. This relation, valid at any order of perturbation theory, n , reads,

$$\mathcal{A}_4^{(n)}(\phi^\dagger, 1_{\bar{q}}^{-h_q}, 2_q^{h_q}, 3_g^{h_3}, 4_g^{h_4}) = -\left[\mathcal{A}_4^{(n)}(\phi, 2_{\bar{q}}^{-h_q}, 1_q^{h_q}, 4_g^{-h_4}, 3_g^{-h_3}) \right] \Big|_{\langle ij \rangle \leftrightarrow [j i]}. \quad (4.5)$$

H amplitude	ϕ amplitude	$\phi^\dagger$ amplitude
$\mathcal{A}(H, 1_g^+, 2_g^+, 3_g^+, 4_g^+)$	$\mathcal{A}(\phi, 1_g^+, 2_g^+, 3_g^+, 4_g^+)$ [44]	$\mathcal{A}(\phi^\dagger, 1_g^+, 2_g^+, 3_g^+, 4_g^+)$ [41]
$\mathcal{A}(H, 1_g^-, 2_g^+, 3_g^+, 4_g^+)$	$\mathcal{A}(\phi, 1_g^-, 2_g^+, 3_g^+, 4_g^+)$ [44]	$\mathcal{A}(\phi^\dagger, 1_g^-, 2_g^+, 3_g^+, 4_g^+)$ [45]
$\mathcal{A}(H, 1_g^-, 2_g^-, 3_g^+, 4_g^+)$	$\mathcal{A}(\phi, 1_g^-, 2_g^-, 3_g^+, 4_g^+)$ [42]	$\mathcal{A}(\phi^\dagger, 1_g^-, 2_g^-, 3_g^+, 4_g^+)$ [42]
$\mathcal{A}(H, 1_g^-, 2_g^+, 3_g^-, 4_g^+)$	$\mathcal{A}(\phi, 1_g^-, 2_g^+, 3_g^-, 4_g^+)$ [43]	$\mathcal{A}(\phi^\dagger, 1_g^-, 2_g^+, 3_g^-, 4_g^+)$ [43]

Table 1. ϕ and $\phi^\dagger$ amplitudes needed to construct a given one-loop $Hg\bar{q}qg$ amplitude, together with the references where they can be obtained. In all cases the $\phi^\dagger$ amplitudes are constructed from the ϕ amplitudes given in the reference using the parity operation. Results for all helicity combinations are also written, in uniform notation, in ref. [45].

We thus see that the $\phi^\dagger$ amplitude in which we are interested is zero,

$$-iA_4^{(0)}(\phi^\dagger, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-) = 0, \quad (4.6)$$

so that, at tree graph level, the NMHV Higgs amplitude in which we will ultimately be interested is simply given by eq. (4.1).

4.2 One-loop results

We begin this section with a brief survey of the literature, to indicate where original calculations of $H\bar{q}q\bar{q}q$ and $Hg\bar{q}qg$ amplitudes may be found. We then turn to the $H\bar{q}qg$ amplitudes, concluding the section by quoting the results for the $\phi^\dagger$ amplitude that must be combined with the new ϕ amplitude calculation that we present here.

4.2.1 $H\bar{q}q\bar{q}q$ amplitudes

The full one-loop results for this process, both for pairs of identical and non-identical quarks, are already available in the literature. The matrix element squared has been computed in ref. [17], with results for the amplitude presented in ref. [46].

4.2.2 $Hg\bar{q}qg$ amplitudes

In principle there are 16 combinations of amplitudes, but this number is reduced to four independent amplitudes by parity and cyclicity. The references to the complete set of needed amplitudes are given in table 1. In addition a nice summary of all the one-loop results for the Higgs + 4 gluon amplitudes is given in ref. [45].

4.2.3 $H\bar{q}qg$ amplitudes

In principle there are 8 combinations of amplitudes, since helicity is conserved on the quark line, but because of parity invariance only four Higgs amplitudes are independent. The references to the amplitudes already calculated in the literature are given in table 2. From this table we see that the Higgs amplitude $\mathcal{A}(H, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-)$ requires, in addition to the calculation of a previously unknown ϕ amplitude, also the results for the corresponding $\phi^\dagger$ amplitude from ref. [46].

H amplitude	ϕ amplitude	$\phi^\dagger$ amplitude
$\mathcal{A}(H, 1_{\bar{q}}^-, 2_q^+, 3_g^+, 4_g^+)$	$\mathcal{A}(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^+, 4_g^+) \text{ [44, 46]}$	$\mathcal{A}(\phi^\dagger, 1_{\bar{q}}^-, 2_q^+, 3_g^+, 4_g^+)$
$\mathcal{A}(H, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-)$	$\mathcal{A}(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-)$	$\mathcal{A}(\phi^\dagger, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-) \text{ [44, 46]}$
$\mathcal{A}(H, 1_{\bar{q}}^-, 2_q^+, 3_g^+, 4_g^-)$	$\mathcal{A}(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^+, 4_g^-) \text{ [46]}$	$\mathcal{A}(\phi^\dagger, 1_{\bar{q}}^-, 2_q^+, 3_g^+, 4_g^-) \text{ [46]}$
$\mathcal{A}(H, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^+)$	$\mathcal{A}(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^+) \text{ [46]}$	$\mathcal{A}(\phi^\dagger, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^+) \text{ [46]}$

Table 2. ϕ and $\phi^\dagger$ amplitudes needed to construct a given one-loop $H\bar{q}qgg$ amplitude, together with the references where they can be obtained. In all cases the $\phi^\dagger$ amplitudes are constructed from the ϕ amplitudes given in the reference, using the parity operation. The cases where the gluons have the same helicity, which have no associated references, are the subject of this paper.

The $\phi^\dagger$ results that we shall need can be derived from the following amplitudes in the case of $A_{4;1}$,

$$-iA_4^L(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^+, 4_g^+) = 2iA_4^{(0)}(\phi^\dagger, 1_{\bar{q}}^-, 2_q^+, 3_g^+, 4_g^+) + \frac{1}{2} \left[\frac{\langle 1|(2+3)|4\rangle}{\langle 23\rangle\langle 34\rangle} + \frac{\langle 12\rangle[23]\langle 31\rangle}{\langle 23\rangle\langle 34\rangle\langle 41\rangle} \right] - \frac{1}{3} \frac{\langle 13\rangle[34]\langle 41\rangle}{\langle 12\rangle\langle 34\rangle^2}, \quad (4.7)$$

$$-iA_4^R(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^+, 4_g^+) = -\frac{1}{2} \left[\frac{\langle 1|(2+3)|4\rangle}{\langle 23\rangle\langle 34\rangle} + \frac{\langle 12\rangle[23]\langle 31\rangle}{\langle 23\rangle\langle 34\rangle\langle 41\rangle} \right], \quad (4.8)$$

$$-iA_4^f(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^+, 4_g^+) = \frac{1}{3} \frac{\langle 13\rangle[34]\langle 41\rangle}{\langle 12\rangle\langle 34\rangle^2}, \quad (4.9)$$

whilst the subleading partial amplitude $A_{4;3}$ also requires the results,

$$-iA_4^L(\phi, 1_{\bar{q}}^-, 2_q^+, 3_q^+, 4_g^+) = -2 \frac{\langle 1|(2+3)|4\rangle^2}{\langle 12\rangle\langle 23\rangle s_{123}} + \frac{1}{2} \left[\frac{\langle 13\rangle\langle 1|(3+4)|2\rangle}{\langle 23\rangle\langle 34\rangle\langle 41\rangle} + \frac{\langle 13\rangle^2[34]}{\langle 12\rangle\langle 23\rangle\langle 34\rangle} \right],$$

$$A_4^f(\phi, 1_{\bar{q}}^-, 2_q^+, 3_q^+, 4_g^+) = 0. \quad (4.10)$$

To obtain the form that is most useful for the calculation of $\mathcal{A}(H, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-)$, we relate the $\phi^\dagger\bar{q}qgg$ amplitudes to the $\phi\bar{q}qgg$ ones by using the relation in eq. (4.5). Thus we obtain the required results by performing the transformation $1 \leftrightarrow 2, 3 \leftrightarrow 4, \langle \rangle \leftrightarrow []$ and reversing the sign. The amplitudes contributing to $A_{4;1}$ are,

$$-iA_4^L(\phi^\dagger, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-) = 2iA_4^{(0)}(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-) + \frac{1}{2} \left[\frac{\langle 3|(1+4)|2\rangle}{[14][34]} + \frac{[21]\langle 14\rangle[24]}{[14][34][23]} \right] - \frac{1}{3} \frac{[24]\langle 34\rangle[23]}{[12][34]^2}. \quad (4.11)$$

$$-iA_4^R(\phi^\dagger, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-) = -\frac{1}{2} \left[\frac{\langle 3|(1+4)|2\rangle}{[14][34]} + \frac{[21]\langle 14\rangle[24]}{[14][34][23]} \right], \quad (4.12)$$

$$-iA_4^f(\phi^\dagger, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-) = \frac{1}{3} \frac{[24]\langle 34\rangle[23]}{[12][34]^2}, \quad (4.13)$$

while the additional subleading contributions become,

$$\begin{aligned}
 -iA_4^L(\phi^\dagger, 1_{\bar{q}}^-, 3_g^-, 2_q^+, 4_g^-) &= 2 \frac{\langle 3|(4+1)|2\rangle^2}{[24][41]s_{124}} - \frac{1}{2} \left[\frac{[21]\langle 4|(1+3)|2\rangle}{[41][13][32]} + \frac{[12]^2\langle 13\rangle}{[24][41][13]} \right] \\
 &= iA_4^{(0)}(\phi, 1_{\bar{q}}^-, 3_g^-, 2_q^+, 4_g^-) + \text{terms antisymmetric in } \{3 \leftrightarrow 4\}.
 \end{aligned} \tag{4.14}$$

We note that all of these amplitudes are finite because of the vanishing of the corresponding tree graph results (see section 4.1).

5 One-loop results

In this section we present analytic expressions for the full one-loop corrections to the process $\mathcal{A}_4^{(1)}(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-)$. All expressions are presented un-renormalised in the four-dimensional helicity (FDH) scheme (setting $\delta_R = 0$) or 't Hooft-Veltman scheme (setting $\delta_R = 1$).

We employ the generalised unitarity method [22, 26, 48–50] to calculate the cut-constructible parts of the left-moving, right-moving and n_f one-loop amplitudes. This relies on the expansion of a one-loop amplitude in terms of scalar basis integrals,

$$A_4^{\text{cut-cons.}}(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-) = \sum_i C_{4;i} \mathcal{I}_{4;i} + \sum_i C_{3;i} \mathcal{I}_{3;i} + \sum_i C_{2;i} \mathcal{I}_{2;i}. \tag{5.1}$$

In this sum each j -point scalar basis integral ($\mathcal{I}_{j;i}$) appears with a coefficient $C_{j;i}$. The sum over i represents the sum over the partitions of the external momenta over the j legs of the basis integral. Multiple cuts isolate different integral functions and allow the construction of a linear system of equations from which the coefficients can be extracted. We use the quadruple cut method [22] which freezes the loop momenta and determines each box coefficient uniquely. Triangle coefficients are determined using the Laurent expansion method [26], whilst the two-point coefficients are determined *via* Stokes' Theorem applied to functions of two complex-conjugated variables [51]. Results were obtained using the QGRAF [52], FORM [53] and S@M [54] packages in order to control the extensive algebra.

5.1 Results for $A_{4;1}(\phi, 1_{\bar{q}}, 2_q, 3_g^-, 4_g^-)$

The partial amplitude $A_{4;1}(\phi, 1_{\bar{q}}, 2_q, 3_g^-, 4_g^-)$ is calculated from three primitive amplitudes according to eq. (3.4). We shall deal with each of these ingredients in turn.

5.1.1 $A_4^L(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-)$

The full result for this primitive amplitude is given by,

$$\begin{aligned}
 -iA_4^L(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-) &= -iA_4^{(0)}(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-) \times V_1^L \\
 &+ \frac{s_{134}^2}{[14][34][2|(1+4)|3]} \left[\text{Ls}_{-1}(s_{14}, s_{34}; s_{134}) + \widetilde{\text{Ls}}_{-1}^{2mh}(s_{12}, s_{134}; s_{34}, m_\phi^2) \right] \\
 &- \frac{\langle 1|(3+4)|2\rangle^2}{\langle 1|(2+3)|4\rangle[23][34]} \left[\text{Ls}_{-1}(s_{34}, s_{23}; s_{234}) + \widetilde{\text{Ls}}_{-1}^{2mh}(s_{12}, s_{234}; s_{34}, m_\phi^2) \right]
 \end{aligned}$$

$$\begin{aligned}
 & + \left[\frac{m_\phi^4 \langle 14 \rangle^2 \langle 24 \rangle}{[12] \langle 2 | (1+4) | 3 \rangle \langle 4 | (1+2) | 3 \rangle s_{124}} - \frac{\langle 3 | (1+4) | 2 \rangle^3}{[12] [24] \langle 3 | (1+2) | 4 \rangle s_{124}} \right] \text{LS}_{-1}(s_{12}, s_{14}; s_{124}) \\
 & + \left[\frac{[23]^2 \langle 4 | (2+3) | 1 \rangle^3}{[12] [13]^3 \langle 4 | (1+2) | 3 \rangle s_{123}} - \frac{m_\phi^4 \langle 13 \rangle^3}{\langle 12 \rangle \langle 1 | (2+3) | 4 \rangle \langle 3 | (1+2) | 4 \rangle s_{123}} \right] \text{LS}_{-1}(s_{12}, s_{23}; s_{123}) \\
 & + \left[\frac{\langle 4 | (1+3) | 2 \rangle^3}{[12] [23] \langle 4 | (1+2) | 3 \rangle s_{123}} - \frac{m_\phi^4 \langle 13 \rangle^3}{\langle 12 \rangle \langle 1 | (2+3) | 4 \rangle \langle 3 | (1+2) | 4 \rangle s_{123}} \right] \\
 & \times \left[\widetilde{\text{LS}}_{-1}^{2mh}(s_{34}, s_{123}; s_{12}, m_\phi^2) + \widetilde{\text{LS}}_{-1}^{2mh}(s_{14}, s_{123}; s_{23}, m_\phi^2) \right] \\
 & + \left[\frac{m_\phi^4 \langle 14 \rangle^2 \langle 24 \rangle}{\langle 12 \rangle \langle 2 | (1+4) | 3 \rangle \langle 4 | (1+2) | 3 \rangle s_{124}} - \frac{\langle 3 | (1+4) | 2 \rangle^2 \langle 3 | (2+4) | 1 \rangle}{[12] [14] \langle 3 | (1+2) | 4 \rangle s_{124}} \right] \\
 & \times \left[\widetilde{\text{LS}}_{-1}^{2mh}(s_{34}, s_{124}; s_{12}, m_\phi^2) + \widetilde{\text{LS}}_{-1}^{2mh}(s_{23}, s_{124}; s_{14}, m_\phi^2) \right] \\
 & - C_{3;\phi|12|34}(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-) \text{I}_3^{\text{m}}(s_{12}, s_{34}, m_\phi^2) \\
 & - C_{3;\phi|41|23}(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-) \text{I}_3^{\text{m}}(s_{23}, s_{14}, m_\phi^2) \\
 & - \frac{2 \langle 13 \rangle^2 \langle 34 \rangle \langle 4 | (1+2) | 3 \rangle [12]}{3} \widehat{\text{L}}_3(s_{123}, s_{12}) \\
 & + \frac{\langle 34 \rangle \langle 31 \rangle (\langle 4 | (1+3) | 2 \rangle [13] - 3 \langle 4 | (2+3) | 1 \rangle [23])}{6 [31]} \widehat{\text{L}}_2(s_{123}, s_{12}) \\
 & + \frac{\langle 13 \rangle \left(16 \langle 4 | (1+3) | 2 \rangle^2 [13]^2 - 3 \langle 4 | (1+3) | 2 \rangle \langle 4 | (1+2) | 3 \rangle [21] [31] + 6 \langle 4 | (2+3) | 1 \rangle^2 [23]^2 \right)}{6 s_{123} [31]^2 [32]} \\
 & \times \widehat{\text{L}}_1(s_{123}, s_{12}) \\
 & - \frac{2 s_{124} \langle 34 \rangle^2 \langle 14 \rangle [42]}{3} \widehat{\text{L}}_3(s_{124}, s_{12}) \\
 & - \langle 34 \rangle \langle 14 \rangle \frac{2 \langle 3 | (1+4) | 2 \rangle [14] - 3 \langle 3 | (1+2) | 4 \rangle [12]}{6 [41]} \widehat{\text{L}}_2(s_{124}, s_{12}) \\
 & - \frac{\langle 3 | (1+4) | 2 \rangle (9 s_{124} \langle 34 \rangle [21] - 22 \langle 3 | (1+4) | 2 \rangle \langle 42 \rangle [12])}{6 s_{124} [41] [21]} \widehat{\text{L}}_1(s_{124}, s_{12}) \\
 & + \frac{\langle 14 \rangle \langle 13 \rangle \langle 4 | (2+3) | 1 \rangle [12]}{2 [31]} \widehat{\text{L}}_2(s_{123}, s_{23}) \\
 & - \langle 13 \rangle \langle 4 | (2+3) | 1 \rangle \frac{3 \langle 4 | (1+3) | 2 \rangle [13] + 2 \langle 4 | (2+3) | 1 \rangle [23]}{2 s_{123} [13]^2} \widehat{\text{L}}_1(s_{123}, s_{23}) \\
 & + \frac{s_{234} \langle 14 \rangle \langle 34 \rangle [42]}{2 [43]} \widehat{\text{L}}_2(s_{234}, s_{23}) + 3 \frac{\langle 34 \rangle \langle 1 | (3+4) | 2 \rangle}{2 [43]} \widehat{\text{L}}_1(s_{234}, s_{23}) \\
 & + R^L \left(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^- \right), \tag{5.2}
 \end{aligned}$$

with,

$$V_1^L = -\frac{1}{\epsilon^2} \left[\left(\frac{\mu^2}{-s_{23}} \right)^\epsilon + \left(\frac{\mu^2}{-s_{34}} \right)^\epsilon + \left(\frac{\mu^2}{-s_{41}} \right)^\epsilon \right] + \frac{13}{6\epsilon} \left(\frac{\mu^2}{-s_{12}} \right)^\epsilon + \frac{119}{18} - \frac{\delta_R}{6}, \tag{5.3}$$

and the remaining rational terms given by,

$$\begin{aligned}
 R^L \left(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^- \right) = & \frac{\langle 34 \rangle \langle 3 | (1+4) | 2 \rangle \left(2 \langle 24 \rangle [42] - \langle 12 \rangle [21] \right)}{12 s_{124} \langle 12 \rangle [21] [41]} \\
 & + \frac{\langle 23 \rangle \langle 4 | (1+3) | 2 \rangle^2 \left(3 \langle 12 \rangle [21] - 2 \langle 23 \rangle [32] \right) - 2 \langle 13 \rangle^2 \langle 24 \rangle \langle 4 | (2+3) | 1 \rangle [21] [32]}{12 s_{123} \langle 12 \rangle \langle 23 \rangle [21] [31] [32]} \\
 & + \frac{5 \langle 34 \rangle^2}{12 \langle 23 \rangle [31]} + \frac{5 \langle 34 \rangle \langle 4 | (1+3) | 2 \rangle}{6 \langle 23 \rangle [31] [32]} + \frac{\langle 4 | (1+3) | 2 \rangle^2}{6 \langle 12 \rangle [21] [31] [32]} \\
 & - \frac{\langle 13 \rangle \langle 14 \rangle \langle 24 \rangle [21]}{3 \langle 12 \rangle \langle 23 \rangle [31] [32]} - \frac{\langle 13 \rangle \langle 34 \rangle}{12 \langle 12 \rangle [41]} - \frac{\langle 34 \rangle^2 [42]}{6 \langle 12 \rangle [21] [41]} + \frac{\langle 13 \rangle \langle 24 \rangle \langle 4 | (1+3) | 4 \rangle}{4 \langle 12 \rangle \langle 23 \rangle [31] [43]} \\
 & - \frac{\langle 13 \rangle \langle 4 | (1+3) | 4 \rangle}{3 \langle 12 \rangle [41] [43]} - \frac{5 \langle 14 \rangle^2 [41]}{12 \langle 12 \rangle [31] [43]} + \frac{\langle 14 \rangle^2 [42]}{6 \langle 12 \rangle [32] [43]}. \quad (5.4)
 \end{aligned}$$

The coefficients of the three mass triangles were calculated using the method of ref. [26],

$$C_{3;\phi|12|34} \left(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^- \right) = \sum_{\gamma=\gamma_{\pm}} \frac{m_{\phi}^4 \langle 34 \rangle^3 \langle 1 K_1^{\flat} \rangle^2}{\gamma \left(\gamma - m_{\phi}^2 \right) \langle 12 \rangle \langle 3 K_1^{\flat} \rangle \langle 4 K_1^{\flat} \rangle}, \quad (5.5)$$

with $K_1 = -p_1 - p_2 - p_3 - p_4$, $K_2 = -p_1 - p_2$ and the massless vector $K_1^{\flat}$ given by,

$$K_1^{\flat \mu} = \gamma \frac{\gamma K_1^{\mu} - K_1^2 K_2^{\mu}}{\gamma^2 - K_1^2 K_2^2}, \quad (5.6)$$

and where γ is given by the two solutions,

$$\gamma_{\pm} = K_1 \cdot K_2 \pm \sqrt{(K_1 \cdot K_2)^2 - K_1^2 K_2^2}. \quad (5.7)$$

The other triangle coefficient is,

$$C_{3;\phi|41|23} \left(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^- \right) = - \sum_{\gamma=\gamma_{\pm}} \frac{m_{\phi}^4 \langle 14 \rangle^2 \langle 3 K_1^{\flat} \rangle^2}{2\gamma \left(\gamma - m_{\phi}^2 \right) \langle 1 K_1^{\flat} \rangle \langle 2 K_1^{\flat} \rangle}, \quad (5.8)$$

with $K_1 = -p_1 - p_2 - p_3 - p_4$, $K_2 = -p_1 - p_4$ and $K_1^{\flat}$ given in terms of these vectors by eq. (5.6).

The definitions of the box integral functions Ls_{-1} and $\widetilde{\text{Ls}}_{-1}^{2mh}$ can be found in appendix B, together with expressions for $\widehat{\text{L}}_1$, $\widehat{\text{L}}_2$ and $\widehat{\text{L}}_3$. In addition to logarithms and polynomial denominators, the latter functions also contain rational terms that protect them from unphysical singularities. Thus, for example,

$$\widehat{\text{L}}_2(s, t) = \frac{\log(s/t)}{(s-t)^2} - \frac{1}{2(s-t)} \left(\frac{1}{s} + \frac{1}{t} \right), \quad (5.9)$$

which is finite in the limit that $s \rightarrow t$.

5.1.2 $A_4^R(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-)$

The result for the right-moving amplitude, $A_4^R(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-)$ is,

$$\begin{aligned}
 -iA_4^R(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-) &= -iA_4^{(0)}(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-) \times V^R \\
 &+ \frac{[1\,2]^2 \langle 4|(1+2)|3\rangle^2}{[1\,3]^3 [2\,3] s_{123}} \text{LS}_{-1}(s_{12}, s_{23}; s_{123}) + \frac{\langle 3|(1+4)|2\rangle^2}{[1\,4][2\,4] s_{124}} \text{LS}_{-1}(s_{14}, s_{12}; s_{124}) \\
 &- \frac{\langle 1|(3+4)|2\rangle^2}{[2\,3][3\,4] \langle 1|(2+3)|4\rangle} \widetilde{\text{LS}}_{-1}^{2mh}(s_{14}, s_{234}; s_{23}, m_\phi^2) \\
 &+ \frac{s_{134}^2}{[1\,4][3\,4] \langle 2|(1+4)|3\rangle} \widetilde{\text{LS}}_{-1}^{2mh}(s_{23}, s_{134}; s_{14}, m_\phi^2) \\
 &- C_{3;\phi|41|23}(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-) \text{I}_3^{\text{m}}(s_{23}, s_{14}, m_\phi^2) - \frac{1}{2} \frac{\langle 1\,4\rangle^2 [1\,2]^2 \langle 3|(1+2)|4\rangle^2}{[1\,4][2\,4] s_{124}} \widehat{\text{L}}_2(s_{124}, s_{12}) \\
 &+ 2 \frac{\langle 3\,4\rangle \langle 3|(1+4)|2\rangle}{[1\,4]} \widehat{\text{L}}_1(s_{124}, s_{12}) + \frac{1}{2} \frac{\langle 3|(1+4)|2\rangle^2}{[1\,4][2\,4] s_{124}} \widehat{\text{L}}_0(s_{124}, s_{12}) \\
 &- \frac{1}{2} \frac{\langle 1\,4\rangle^2 [2\,4]^2 s_{234}^2}{[2\,3][3\,4] \langle 1|(2+3)|4\rangle} \widehat{\text{L}}_2(s_{234}, s_{23}) - 2 \frac{\langle 3\,4\rangle \langle 1|(3+4)|2\rangle}{[3\,4]} \widehat{\text{L}}_1(s_{234}, s_{23}) \\
 &+ \frac{1}{2} \frac{\langle 1|(3+4)|2\rangle^2}{[2\,3][3\,4] \langle 1|(2+3)|4\rangle} \widehat{\text{L}}_0(s_{234}, s_{23}) - \frac{1}{2} \frac{\left(\langle 1\,2\rangle [1\,2] \langle 4|(2+3)|1\rangle\right)^2 [2\,3]}{[1\,3]^3 s_{123}} \widehat{\text{L}}_2(s_{123}, s_{23}) \\
 &+ 2 \frac{\langle 1\,3\rangle [1\,2] \langle 4|(1+2)|3\rangle \langle 4|(2+3)|1\rangle}{\langle 2\,3\rangle [1\,3]^2 [2\,3]} \widehat{\text{L}}_1(s_{123}, s_{23}) \\
 &+ \left[-2 \frac{\langle 1\,3\rangle [1\,2] \langle 4|(1+2)|3\rangle \langle 4|(2+3)|1\rangle}{s_{123} [1\,3]^2 \langle 2\,3\rangle [2\,3]} + \frac{1}{2} \frac{\langle 4|(2+3)|1\rangle^2 [2\,3]}{[1\,3]^3 s_{123}} \right] \widehat{\text{L}}_0(s_{123}, s_{23}) \\
 &- \frac{1}{2} \frac{\left(\langle 1\,3\rangle [1\,2] \langle 4|(1+2)|3\rangle\right)^2}{[1\,3][2\,3] s_{123}} \widehat{\text{L}}_2(s_{123}, s_{12}) \\
 &+ \langle 3\,4\rangle [1\,2] \langle 4|(1+2)|3\rangle \frac{(-2 \langle 1\,3\rangle [1\,3] - \langle 2\,3\rangle [2\,3])}{\langle 2\,3\rangle [1\,3]^2 [2\,3]} \widehat{\text{L}}_1(s_{123}, s_{12}) \\
 &+ [1\,2] \langle 4|(1+2)|3\rangle \frac{\langle 2\,3\rangle \langle 4|(1+3)|2\rangle + 2 \langle 1\,3\rangle \langle 4|(2+3)|1\rangle}{[1\,3]^2 \langle 2\,3\rangle [2\,3] s_{123}} \widehat{\text{L}}_0(s_{123}, s_{12}) \\
 &+ R^R(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-) , \tag{5.10}
 \end{aligned}$$

with

$$V^R = -\frac{1}{\epsilon^2} \left(\frac{\mu^2}{-s_{12}} \right)^\epsilon - \frac{3}{2\epsilon} \left(\frac{\mu^2}{-s_{12}} \right)^\epsilon - \frac{7}{2} - \frac{\delta_R}{2} . \tag{5.11}$$

The remaining rational pieces in eq. (5.10) have the following form:

$$\begin{aligned}
 R^R(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-) = & -\frac{\langle 24 \rangle^2 [21]^2}{2 \langle 23 \rangle [31]^3} + \frac{\langle 4|(1+2)|3 \rangle^2 [21]^2}{2s_{123} [31]^3 [32]} - \frac{\langle 14 \rangle^2 [21]}{2 \langle 12 \rangle [31] [32]} \\
 & + \frac{[21] \left(\langle 13 \rangle^2 \langle 23 \rangle \langle 4|(1+2)|3 \rangle^2 [31]^2 + \langle 12 \rangle^3 \langle 4|(2+3)|1 \rangle^2 [21] [32] \right)}{4s_{123}^2 \langle 12 \rangle \langle 23 \rangle [31]^3 [32]} \\
 & + \frac{\langle 3|(1+4)|2 \rangle^2}{2s_{124} [41] [42]} - \frac{\langle 13 \rangle^2 [21]}{2 \langle 12 \rangle [41] [42]} + \frac{\langle 14 \rangle^2 \langle 3|(1+2)|4 \rangle^2 [21]}{4s_{124}^2 \langle 12 \rangle [41] [42]} \\
 & - \frac{\langle 13 \rangle \langle 14 \rangle [42]}{2 \langle 1|(2+3)|4 \rangle [43]} - \frac{s_{234} \langle 14 \rangle^2 [42]^2}{4 \langle 23 \rangle \langle 1|(2+3)|4 \rangle [32]^2 [43]} - \frac{\langle 14 \rangle^2 [42]^2}{2 \langle 1|(2+3)|4 \rangle [32] [43]}. \quad (5.12)
 \end{aligned}$$

5.1.3 $A_4^f(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-)$

The fermion loop contribution is,

$$\begin{aligned}
 -iA_4^f(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-) = & -iA_4^{(0)}(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-) \times \left[-\frac{2}{3\epsilon} \left(\frac{\mu^2}{-s_{12}} \right)^\epsilon - \frac{10}{9} \right] \\
 & + \frac{2}{3} \langle 13 \rangle^2 \langle 34 \rangle [12] \langle 4|(1+2)|3 \rangle \hat{L}_3(s_{123}, s_{12}) + \frac{2}{3} \langle 14 \rangle^2 \langle 34 \rangle [12] \langle 3|(1+2)|4 \rangle \hat{L}_3(s_{124}, s_{12}) \\
 & + \frac{1}{3} \langle 13 \rangle \langle 34 \rangle \langle 4|(1+3)|2 \rangle \hat{L}_2(s_{123}, s_{12}) + \frac{1}{3} \langle 14 \rangle \langle 34 \rangle \langle 3|(1+4)|2 \rangle \hat{L}_2(s_{124}, s_{12}) \\
 & - \frac{2}{3} \frac{\langle 13 \rangle \langle 4|(1+3)|2 \rangle^2}{\langle 12 \rangle [23] [12]} \hat{L}_1(s_{123}, s_{12}) + \frac{2}{3} \frac{\langle 14 \rangle \langle 3|(1+4)|2 \rangle^2}{\langle 12 \rangle [24] [12]} \hat{L}_1(s_{124}, s_{12}) \\
 & + \frac{2}{3} \frac{\langle 13 \rangle \langle 4|(1+3)|2 \rangle^2}{\langle 12 \rangle [12] [23] s_{123}} \hat{L}_0(s_{123}, s_{12}) + \frac{2}{3} \frac{(s_{12} + s_{14}) \langle 3|(1+4)|2 \rangle^2}{\langle 12 \rangle [14] [24] [12] s_{124}} \hat{L}_0(s_{124}, s_{12}) \\
 & - \frac{\langle 13 \rangle \langle 34 \rangle \langle 4|(1+3)|2 \rangle}{6 \langle 12 \rangle [12] s_{123}} - \frac{\langle 14 \rangle \langle 34 \rangle \langle 3|(1+4)|2 \rangle}{6 \langle 12 \rangle [12] s_{124}} - \frac{1}{3} \frac{\langle 13 \rangle \langle 14 \rangle}{\langle 12 \rangle [34]}. \quad (5.13)
 \end{aligned}$$

5.1.4 Relation for rational terms

We note that the rational terms in the three leading colour primitive amplitudes obey the following relation in the four-dimensional helicity scheme, ($\delta_R = 0$),

$$\begin{aligned}
 \mathcal{R} \Big\{ & A_4^L(\phi, 1_{\bar{q}}, 2_q, 3_g, 4_g) + A_4^R(\phi, 1_{\bar{q}}, 2_q, 3_g, 4_g) + A_4^f(\phi, 1_{\bar{q}}, 2_q, 3_g, 4_g) \Big\} \\
 & + 2A_4^{(0)}(\phi^\dagger, 1_{\bar{q}}, 2_q, 3_g, 4_g) - 2A_4^{(0)}(\phi, 1_{\bar{q}}, 2_q, 3_g, 4_g) = 0, \quad (5.14)
 \end{aligned}$$

a formula analogous to that found in super-symmetric decompositions of QCD amplitudes [21]. This property is helicity independent and has also been checked for the previously known MHV amplitudes [46]. Eq. (5.14) simplifies further for the case of the Higgs field, since the last two terms on the left hand side, proportional to tree graph amplitudes, cancel out. For the NMHV helicity assignment at hand, we note that one of the tree graph amplitudes that appears in eq. (5.14), namely $A_4^{(0)}(\phi^\dagger, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-)$, is zero (c.f. eq. (4.6)). We stress that the $\mathcal{R}$ operation extracts the full rational term, including completion terms from the functions $\hat{L}_3$ and $\hat{L}_2$ and the constants from the V -functions.

Thus it corresponds to dropping all logarithms, box functions and ϵ -dependent terms from the V -functions.

We conclude this section by noting that the three primitive amplitudes for the helicity assignment $(1_{\bar{q}}^-, 2_q^+, 3_g^+, 4_g^+)$ displayed in eqs. (4.7), (4.8) and (4.9), also satisfy eq. (5.14). For these amplitudes, which are purely rational, the $\mathcal{R}$ operation leaves the amplitude unchanged.

5.2 Results for $A_{4;3}(\phi, 1_{\bar{q}}, 2_q, 3_g^-, 4_g^-)$

We can calculate the result for $A_{4;3}$ using eq. (3.5). Given the results for A_L^4 and A_R^4 in the previous section the only missing ingredient is $A_4^L(\phi, 1_{\bar{q}}^-, 2_g^-, 3_q^+, 4_g^-)$.

5.2.1 Box-related terms for $A_4^L(\phi, 1_{\bar{q}}^-, 2_g^-, 3_q^+, 4_g^-)$

The calculation of the box-related terms in $\phi\bar{q}gqg$ ($--+-$) is easily performed using the methods given in ref. [22]. The result is,

$$\begin{aligned}
 -iA_4^{L,\text{box}}(\phi, 1_{\bar{q}}^-, 2_g^-, 3_q^+, 4_g^-) &= -iA_4^{(0)}(\phi, 1_{\bar{q}}^-, 2_g^-, 3_q^+, 4_g^-) \times V_4^L \\
 &+ \frac{\langle 12 \rangle^2 m_\phi^4}{\langle 1|(2+3)|4 \rangle \langle 3|(1+2)|4 \rangle s_{123}} \text{LS}_{-1}(s_{12}, s_{23}; s_{123}) - \frac{\langle 2|(1+4)|3 \rangle^2}{[14][34]s_{134}} \text{LS}_{-1}(s_{14}, s_{34}; s_{134}) \\
 &+ \frac{[34]^2 \langle 1|(3+4)|2 \rangle^2}{[32][24]^3 \langle 1|(2+3)|4 \rangle} \text{LS}_{-1}(s_{23}, s_{34}; s_{234}) + \frac{s_{124}^2}{[12][24]\langle 3|(1+2)|4 \rangle} \text{LS}_{-1}(s_{12}, s_{14}; s_{124}) \\
 &+ \frac{\langle 3|(2+4)|1 \rangle s_{124}^2}{[12][14]\langle 3|(1+2)|4 \rangle \langle 3|(1+4)|2 \rangle} \widetilde{\text{LS}}_{-1}^{2mh}(s_{23}, s_{124}; s_{14}, m_\phi^2) \\
 &- \frac{\langle 1|(2+4)|3 \rangle^3}{[23][34]\langle 1|(2+3)|4 \rangle \langle 1|(3+4)|2 \rangle} \widetilde{\text{LS}}_{-1}^{2mh}(s_{12}, s_{234}; s_{34}, m_\phi^2) \\
 &+ \frac{1}{s_{123}} \left[\frac{m_\phi^4 \langle 12 \rangle^2}{\langle 3|(1+2)|4 \rangle \langle 1|(2+3)|4 \rangle} + \frac{\langle 4|(1+2)|3 \rangle^2}{[12][23]} \right] \\
 &\times \left[\widetilde{\text{LS}}_{-1}^{2mh}(s_{34}, s_{123}; s_{12}, m_\phi^2) + \widetilde{\text{LS}}_{-1}^{2mh}(s_{14}, s_{123}; s_{23}, m_\phi^2) \right], \tag{5.15}
 \end{aligned}$$

with

$$V_4^L = -\frac{1}{\epsilon^2} \left[\left(\frac{\mu^2}{-s_{34}} \right)^\epsilon + \left(\frac{\mu^2}{-s_{41}} \right)^\epsilon \right] + \frac{1}{3\epsilon} \left(\frac{\mu^2}{-s_{123}} \right)^\epsilon + \frac{7}{4} - \frac{\delta_R}{3}. \tag{5.16}$$

As we shall see in the next section, no further information is required for the calculation of the $A_{4;3}$ which is completely determined by box diagrams alone.

5.2.2 Full result for $A_{4;3}$

The full result for the partial amplitude $A_{4;3}$ is,

$$\begin{aligned}
 -iA_{4;3}(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-) &= -iA_4^{(0)}(\phi, 1_{\bar{q}}^-, 2_q^+, 3_g^-, 4_g^-) \times V_5(s_{12}, s_{34}, s_{13}, s_{24}) \\
 &+ \frac{1}{s_{123}} \left[\frac{\langle 4|(1+2)|3\rangle^2 [12]^2}{[13]^3 [23]} + \frac{[23]^2 \langle 4|(2+3)|1\rangle^3}{[13]^3 [12] \langle 4|(1+2)|3\rangle} - \frac{m_\phi^4 \langle 13\rangle^3}{\langle 12\rangle \langle 1|(2+3)|4\rangle \langle 3|(1+2)|4\rangle} \right] \\
 &\quad \times \text{Ls}_{-1}(s_{12}, s_{23}; s_{123}) \\
 &+ \frac{1}{s_{124}} \left[\frac{m_\phi^4 \langle 14\rangle^2 \langle 24\rangle}{\langle 12\rangle \langle 2|(1+4)|3\rangle \langle 4|(1+2)|3\rangle} - \frac{\langle 3|(1+4)|2\rangle^2 \langle 3|(2+4)|1\rangle}{[14] \langle 3|(1+2)|4\rangle [12]} \right] \text{Ls}_{-1}(s_{12}, s_{14}; s_{124}) \\
 &+ \frac{1}{s_{123}} \left[\frac{m_\phi^4 \langle 13\rangle^2}{\langle 1|(2+3)|4\rangle \langle 2|(1+3)|4\rangle} - \frac{\langle 4|(1+3)|2\rangle^2}{[13] [23]} \right] \text{Ls}_{-1}(s_{13}, s_{23}; s_{123}) \\
 &+ \frac{s_{341}^2}{[13] [34] \langle 2|(1+3)|4\rangle} \left[\text{Ls}_{-1}(s_{13}, s_{14}; s_{341}) + \widetilde{\text{Ls}}_{-1}^{2mh}(s_{23}, s_{341}, s_{14}, m_\phi^2) \right] \\
 &+ \frac{s_{341}^2}{[14] [34] \langle 2|(1+4)|3\rangle} \left[\text{Ls}_{-1}(s_{14}, s_{34}; s_{341}) + \widetilde{\text{Ls}}_{-1}^{2mh}(s_{12}, s_{341}, s_{34}, m_\phi^2) \right] \\
 &- \frac{\langle 1|(3+4)|2\rangle^2}{[23] [34] \langle 1|(2+3)|4\rangle} \left[\text{Ls}_{-1}(s_{23}, s_{34}; s_{234}) + \widetilde{\text{Ls}}_{-1}^{2mh}(s_{12}, s_{234}, s_{34}, m_\phi^2) \right] \\
 &+ \frac{[24]^2 \langle 1|(2+4)|3\rangle^2}{[23] [34]^3 \langle 1|(2+3)|4\rangle} \text{Ls}_{-1}(s_{23}, s_{24}; s_{234}) \\
 &- \frac{\langle 1|(3+4)|2\rangle^2}{\langle 1|(2+4)|3\rangle [24] [34]} \widetilde{\text{Ls}}_{-1}^{2mh}(s_{14}, s_{234}, s_{23}, m_\phi^2) \\
 &+ \frac{1}{s_{123}} \left[-\frac{m_\phi^4 \langle 13\rangle^2 \langle 23\rangle}{\langle 12\rangle \langle 2|(1+3)|4\rangle \langle 3|(1+2)|4\rangle} + \frac{\langle 4|(1+3)|2\rangle^2 \langle 4|(2+3)|1\rangle}{\langle 4|(1+2)|3\rangle [12] [13]} \right] \\
 &\quad \times \widetilde{\text{Ls}}_{-1}^{2mh}(s_{14}, s_{123}, s_{23}, m_\phi^2) \\
 &+ \frac{1}{s_{123}} \left[\frac{m_\phi^4 \langle 13\rangle^3}{\langle 12\rangle \langle 3|(1+2)|4\rangle \langle 1|(2+3)|4\rangle} - \frac{\langle 4|(1+3)|2\rangle^3}{\langle 4|(1+2)|3\rangle [12] [23]} \right] \\
 &\quad \times \widetilde{\text{Ls}}_{-1}^{2mh}(s_{24}, s_{123}, s_{13}, m_\phi^2) \\
 &+ \frac{1}{s_{123}} \left[-\frac{m_\phi^4 \langle 13\rangle^2}{\langle 2|(1+3)|4\rangle \langle 1|(2+3)|4\rangle} + \frac{\langle 4|(1+3)|2\rangle^2}{[13] [23]} \right] \widetilde{\text{Ls}}_{-1}^{2mh}(s_{34}, s_{123}, s_{12}, m_\phi^2) \\
 &+ \left\{ 3 \leftrightarrow 4 \right\}, \tag{5.17}
 \end{aligned}$$

where the function containing poles and associated logarithms is conveniently written as,

$$V_5(s_{12}, s_{34}, s_{13}, s_{24}) = -\frac{1}{\epsilon^2} \left[\left(\frac{\mu^2}{-s_{12}} \right)^\epsilon + \left(\frac{\mu^2}{-s_{34}} \right)^\epsilon - \left(\frac{\mu^2}{-s_{13}} \right)^\epsilon - \left(\frac{\mu^2}{-s_{24}} \right)^\epsilon \right]. \tag{5.18}$$

We note that the apparent double pole in ϵ in eq. (5.18) is cancelled upon expanding about $\epsilon = 0$.

This result for the ϕ amplitude is particularly simple, containing neither bubble contributions nor rational terms. This is also true for the helicity amplitude $A_{4;3}(\phi, 1_{\bar{q}}, 2_q, 3_g^-, 4_g^+)$,¹ which can easily be checked using the previously calculated results in ref. [46]. It is therefore more efficient to program the full result for $A_{4;3}$, rather than to program the individual primitive amplitudes using eq. (3.5).

Furthermore, for the case of two negative gluon helicities calculated here one can check using eq. (3.5) and eqs. (4.11), (4.12), (4.15) that the corresponding $\phi^\dagger$ amplitude is zero. Therefore we have,

$$A_{4;3}(H, 1_{\bar{q}}, 2_q^+, 3_g^-, 4_g^-) = iA_{4;3}(A, 1_{\bar{q}}, 2_q^+, 3_g^-, 4_g^-) = A_{4;3}(\phi, 1_{\bar{q}}, 2_q^+, 3_g^-, 4_g^-) . \quad (5.19)$$

6 Numerical results

Here we present evaluations of the new amplitudes at the same kinematic point as used previously in the literature [17, 46]:

$$\begin{aligned} k_\phi &= (-1.0000000000, 0.0000000000, 0.0000000000, 0.0000000000), \\ k_1 &= (0.30674037867, -0.17738694693, -0.01664472021, -0.24969277974), \\ k_2 &= (0.34445032281, 0.14635282800, -0.10707762397, 0.29285022975), \\ k_3 &= (0.22091667641, 0.08911915938, 0.19733901856, 0.04380941793), \\ k_4 &= (0.12789262211, -0.05808504045, -0.07361667438, -0.08696686795). \end{aligned} \quad (6.1)$$

We have used a scale $\mu = m_H$, set $\delta_R = 1$ (corresponding to the 't Hooft-Veltman scheme) and, in assembling the amplitude $A_{4;1}$, have used $n_f = 5$. The results for the final Higgs amplitudes presented in table 3 agree with those from the semi-numerical calculation of ref. [17] to one part in 10^8 . Note that these results depend on an overall phase that can be removed by dividing out by the corresponding Born calculation. Using the analytic expressions for all the $Hg g g g$, $H\bar{q}q g g$ and $H\bar{q}q\bar{q}'q'$ amplitudes that are now available we can also confirm² the numerical values for the matrix elements squared given in ref. [17].

7 Conclusions

In this paper we have computed the last remaining, analytically unknown, helicity amplitude contributing to the NLO corrections to Higgs plus two jet production at hadron colliders. This builds upon the previously known semi-numerical results [17] and completes the set of compact analytic formulae [41–46].

We employed a generalised unitarity approach to calculate the cut-constructible parts of the amplitude. Completion of the logarithmic terms to remove unphysical singularities was used to simplify rational terms extracted from a Feynman diagram calculation. Simplifications in the construction of the subleading colour amplitude, $A_{4;3}$, showed that this component is free from all rational, bubble and triangle terms. Similar relations between

¹The amplitude $A_{4;3}(\phi, 1_{\bar{q}}, 2_q, 3_g^+, 4_g^-)$ is not independent and is obtained by swapping labels 3 and 4.

²Fortran code that calculates all the amplitudes can be downloaded from mcfm.fnal.gov.

	$(\phi, 1_{\bar{q}}, 2_q^+, 3_g^-, 4_g^-)$			$(\phi^\dagger, 1_{\bar{q}}, 2_q^+, 3_g^-, 4_g^-)$
	$1/\epsilon^2$	$1/\epsilon$	ϵ^0	ϵ^0
$A_4^{(0)}$	0	0	+6.4990753590 −2.3930814482 i	0
A_4^L	−19.497226077 +7.1792443445 i	−64.624963049 −45.761120716 i	−31.605583566 −137.56039301 i	−17.355492030 +6.1436166419 i
A_4^R	−6.4990753590 +2.3930814482 i	−23.126318341 −14.670203900 i	−48.741904002 −39.062655529 i	+4.5854677141 −1.3871854529 i
A_4^f	0	−4.3327169060 +1.5953876321 i	−14.980583214 −9.8887497350 i	−0.22812640212 +0.029731707266 i
$A_{4;1}$	−18.775106593 +6.9133464058 i	−69.276566965 −41.472118673 i	−51.157455145 −149.70134751 i	−18.245199113 +6.3473012044 i
$A_{4;3}$	0	+2.6108347714 −0.051191063962 i	+17.757374434 +4.9309701446 i	0

Table 3. Numerical values of $\phi\bar{q}qgg$ and $\phi^\dagger\bar{q}qgg$ primitive amplitudes (above) and the amplitudes multiplying the two different colour structures (below), at the kinematic point defined in eq. (6.1).

the rational terms in the left, right and fermion loop primitive amplitudes were used to find a compact analytic structure.

Our results have been verified against the known numerical results and we envisage that they will provide the means for a faster and more flexible analysis of Higgs phenomenology at the Tevatron and the LHC.

Acknowledgments

We are happy to acknowledge useful discussions with Nigel Glover, Pierpaolo Mastrolia and Giulia Zanderighi. CW acknowledges the award of an STFC studentship and SB acknowledges support from the Helmholtz Gemeinschaft under contract VH-NG-105. Fermilab is operated by Fermi Research Alliance, LLC under Contract No. DE-AC02-07CH11359 with the United States Department of Energy.

A Spinor notation

Our spinor notation is quite standard in the QCD literature, (for a review see refs. [55, 56]) The function $u_\pm(k_i)$ is a massless Weyl spinor of momentum k_i and positive or negative chirality. In terms of these solutions of the Dirac equation, the spinor products are defined by,

$$\langle i j \rangle = \langle i^- | j^+ \rangle = \bar{u}_-(k_i) u_+(k_j), \quad (\text{A.1})$$

$$[i j] = \langle i^+ | j^- \rangle = \bar{u}_+(k_i) u_-(k_j). \quad (\text{A.2})$$

We use the convention $[ij] = \text{sgn}(k_i^0 k_j^0) \langle ji \rangle^*$, so that,

$$\langle ij \rangle [ji] = 2k_i \cdot k_j \equiv s_{ij}. \quad (\text{A.3})$$

We further define,

$$s_{ijl} \equiv (k_i + k_j + k_l)^2 = \langle ij \rangle [ji] + \langle jl \rangle [lj] + \langle il \rangle [li], \quad (\text{A.4})$$

$$\langle a|i|b \rangle = \langle ai \rangle [ib], \quad \langle a|(i+j)|b \rangle = \langle ai \rangle [ib] + \langle aj \rangle [jb], \quad (\text{A.5})$$

$$\langle ji \rangle = -\langle ij \rangle, \quad [ji] = -[ij]. \quad (\text{A.6})$$

Simplification of the formula can sometimes be achieved by using the Schouten identity,

$$\langle ab \rangle \langle cd \rangle = \langle ad \rangle \langle cb \rangle + \langle ac \rangle \langle bd \rangle, \quad (\text{A.7})$$

$$[ab] [cd] = [ad] [cb] + [ac] [bd]. \quad (\text{A.8})$$

B Definitions of special functions

With the definition of the $L_i(s, t)$ basis functions,

$$L_i(s, t) = \frac{\log(s/t)}{(s-t)^i}, \quad (\text{B.1})$$

we can define the completions,

$$\begin{aligned} L_3(s, t) &\rightarrow \widehat{L}_3(s, t) = L_3(s, t) - \frac{1}{2(s-t)^2} \left(\frac{1}{s} + \frac{1}{t} \right), \\ L_2(s, t) &\rightarrow \widehat{L}_2(s, t) = L_2(s, t) - \frac{1}{2(s-t)} \left(\frac{1}{s} + \frac{1}{t} \right), \\ L_1(s, t) &\rightarrow \widehat{L}_1(s, t) = L_1(s, t) \\ L_0(s, t) &\rightarrow \widehat{L}_0(s, t) = L_0(s, t), \end{aligned} \quad (\text{B.2})$$

such that the completed functions are finite in the limit that $s \rightarrow t$. Note that the definition of these completed logarithmic functions is similar in spirit, but different in detail from the definitions in ref. [46].

We also need the box functions from the scalar box with one massive external leg,

$$L_{S-1}(s, t; m^2) = \text{Li}_2\left(1 - \frac{s}{m^2}\right) + \text{Li}_2\left(1 - \frac{t}{m^2}\right) + \log\left(\frac{s}{m^2}\right) \log\left(\frac{t}{m^2}\right) - \frac{\pi^2}{6}, \quad (\text{B.3})$$

and coming from the box with two adjacent massive external legs,

$$\begin{aligned} \widetilde{L}_{S-1}^{2mh}(s, t; m_1^2, m_2^2) &= -\text{Li}_2\left(1 - \frac{m_1^2}{t}\right) - \text{Li}_2\left(1 - \frac{m_2^2}{t}\right) - \frac{1}{2} \log^2\left(\frac{-s}{-t}\right) \\ &\quad + \frac{1}{2} \log\left(\frac{-s}{-m_1^2}\right) \log\left(\frac{-s}{-m_2^2}\right). \end{aligned} \quad (\text{B.4})$$

where the dilogarithm is defined as usual by,

$$\text{Li}_2(x) = -\int_0^x dy \frac{\log(1-y)}{y}. \quad (\text{B.5})$$

I_3^{m} is the three mass triangle function defined, for example, in eq. (II.9) of ref. [57],

$$I_3^{\text{m}}(s_{12}, s_{34}, s_{56}) = \int_0^1 d^3 a_i \delta(1 - a_1 - a_2 - a_3) \frac{1}{[-s_{12}a_1a_2 - s_{34}a_2a_3 - s_{56}a_3a_1]}. \quad (\text{B.6})$$

Explicit results for this integral can be found in refs. [58, 59] and [60].

References

- [1] LEP WORKING GROUP FOR HIGGS BOSON SEARCHES collaboration, R. Barate et al., *Search for the standard model Higgs boson at LEP*, *Phys. Lett. B* **565** (2003) 61 [[hep-ex/0306033](#)] [[SPIRES](#)].
- [2] *Web page of the LEP Electroweak Working Group (LEPEWWG)*, <http://lepewwg.web.cern.ch/LEPEWWG/>.
- [3] CDF collaboration, *Combined CDF and DØ upper limits on standard model Higgs-Boson production with up to 4.2 fb^{-1} of data*, [arXiv:0903.4001](#) [[SPIRES](#)].
- [4] CDF AND D0 collaboration, N. Krumnack, *Combined SM Higgs limits at the Tevatron*, [arXiv:0910.3353](#) [[SPIRES](#)].
- [5] For an extensive review of theoretical aspects of Higgs boson production and prospects for detection, see: A. Djouadi, *The Anatomy of electro-weak symmetry breaking. I: the Higgs boson in the standard model*, *Phys. Rept.* **457** (2008) 1 [[hep-ph/0503172](#)] [[SPIRES](#)] and references therein.
- [6] T. Han, G. Valencia and S. Willenbrock, *Structure function approach to vector boson scattering in pp collisions*, *Phys. Rev. Lett.* **69** (1992) 3274 [[hep-ph/9206246](#)] [[SPIRES](#)].
- [7] T. Figy, C. Oleari and D. Zeppenfeld, *Next-to-leading order jet distributions for Higgs boson production via weak-boson fusion*, *Phys. Rev. D* **68** (2003) 073005 [[hep-ph/0306109](#)] [[SPIRES](#)].
- [8] E.L. Berger and J.M. Campbell, *Higgs boson production in weak boson fusion at next-to-leading order*, *Phys. Rev. D* **70** (2004) 073011 [[hep-ph/0403194](#)] [[SPIRES](#)].
- [9] M. Ciccolini, A. Denner and S. Dittmaier, *Strong and electroweak corrections to the production of Higgs+2 Jets via weak interactions at the LHC*, *Phys. Rev. Lett.* **99** (2007) 161803 [[arXiv:0707.0381](#)] [[SPIRES](#)].
- [10] M. Ciccolini, A. Denner and S. Dittmaier, *Electroweak and QCD corrections to Higgs production via vector-boson fusion at the LHC*, *Phys. Rev. D* **77** (2008) 013002 [[arXiv:0710.4749](#)] [[SPIRES](#)].
- [11] F. Wilczek, *Decays of heavy vector mesons into Higgs particles*, *Phys. Rev. Lett.* **39** (1977) 1304 [[SPIRES](#)].
- [12] A. Djouadi, M. Spira and P.M. Zerwas, *Production of Higgs bosons in proton colliders: QCD corrections*, *Phys. Lett. B* **264** (1991) 440 [[SPIRES](#)].
- [13] S. Dawson, *Radiative corrections to Higgs boson production*, *Nucl. Phys. B* **359** (1991) 283 [[SPIRES](#)].
- [14] V. Del Duca, W. Kilgore, C. Oleari, C. Schmidt and D. Zeppenfeld, *Higgs + 2 jets via gluon fusion*, *Phys. Rev. Lett.* **87** (2001) 122001 [[hep-ph/0105129](#)] [[SPIRES](#)].

- [15] S. Dawson and R.P. Kauffman, *Higgs boson plus multi - jet rates at the SSC*, *Phys. Rev. Lett.* **68** (1992) 2273 [[SPIRES](#)].
- [16] R.P. Kauffman, S.V. Desai and D. Risal, *Production of a Higgs boson plus two jets in hadronic collisions*, *Phys. Rev. D* **55** (1997) 4005 [[hep-ph/9610541](#)] [[SPIRES](#)].
- [17] R.K. Ellis, W.T. Giele and G. Zanderighi, *Virtual QCD corrections to Higgs boson plus four parton processes*, *Phys. Rev. D* **72** (2005) 054018 [Erratum *ibid.* **D 74** (2006) 079902] [[hep-ph/0506196](#)] [[SPIRES](#)].
- [18] J.M. Campbell, R.K. Ellis and G. Zanderighi, *Next-to-leading order Higgs + 2 jet production via gluon fusion*, *JHEP* **10** (2006) 028 [[hep-ph/0608194](#)] [[SPIRES](#)].
- [19] Z. Bern, L.J. Dixon, D.C. Dunbar and D.A. Kosower, *One-Loop n-point gauge theory amplitudes, unitarity and collinear limits*, *Nucl. Phys. B* **425** (1994) 217 [[hep-ph/9403226](#)] [[SPIRES](#)].
- [20] Z. Bern, L.J. Dixon, D.C. Dunbar and D.A. Kosower, *Fusing gauge theory tree amplitudes into loop amplitudes*, *Nucl. Phys. B* **435** (1995) 59 [[hep-ph/9409265](#)] [[SPIRES](#)].
- [21] Z. Bern, L.J. Dixon and D.A. Kosower, *One loop corrections to two quark three gluon amplitudes*, *Nucl. Phys. B* **437** (1995) 259 [[hep-ph/9409393](#)] [[SPIRES](#)].
- [22] R. Britto, F. Cachazo and B. Feng, *Generalized unitarity and one-loop amplitudes in $N = 4$ super-Yang-Mills*, *Nucl. Phys. B* **725** (2005) 275 [[hep-th/0412103](#)] [[SPIRES](#)].
- [23] F. del Aguila and R. Pittau, *Recursive numerical calculus of one-loop tensor integrals*, *JHEP* **07** (2004) 017 [[hep-ph/0404120](#)] [[SPIRES](#)].
- [24] G. Ossola, C.G. Papadopoulos and R. Pittau, *Reducing full one-loop amplitudes to scalar integrals at the integrand level*, *Nucl. Phys. B* **763** (2007) 147 [[hep-ph/0609007](#)] [[SPIRES](#)].
- [25] R.K. Ellis, W.T. Giele and Z. Kunszt, *A numerical unitarity formalism for evaluating one-loop amplitudes*, *JHEP* **03** (2008) 003 [[arXiv:0708.2398](#)] [[SPIRES](#)].
- [26] D. Forde, *Direct extraction of one-loop integral coefficients*, *Phys. Rev. D* **75** (2007) 125019 [[arXiv:0704.1835](#)] [[SPIRES](#)].
- [27] Z. Bern and A.G. Morgan, *Massive loop amplitudes from unitarity*, *Nucl. Phys. B* **467** (1996) 479 [[hep-ph/9511336](#)] [[SPIRES](#)].
- [28] Z. Bern, L.J. Dixon, D.C. Dunbar and D.A. Kosower, *One-loop self-dual and $N = 4$ super Yang-Mills*, *Phys. Lett. B* **394** (1997) 105 [[hep-th/9611127](#)] [[SPIRES](#)].
- [29] C. Anastasiou, R. Britto, B. Feng, Z. Kunszt and P. Mastrolia, *D-dimensional unitarity cut method*, *Phys. Lett. B* **645** (2007) 213 [[hep-ph/0609191](#)] [[SPIRES](#)].
- [30] W.T. Giele, Z. Kunszt and K. Melnikov, *Full one-loop amplitudes from tree amplitudes*, *JHEP* **04** (2008) 049 [[arXiv:0801.2237](#)] [[SPIRES](#)].
- [31] C.F. Berger et al., *An automated implementation of on-shell methods for one-loop amplitudes*, *Phys. Rev. D* **78** (2008) 036003 [[arXiv:0803.4180](#)] [[SPIRES](#)].
- [32] C.F. Berger et al., *One-loop multi-parton amplitudes with a vector boson for the LHC*, [arXiv:0808.0941](#) [[SPIRES](#)].
- [33] C.F. Berger et al., *Precise predictions for $W + 3$ jet production at hadron colliders*, *Phys. Rev. Lett.* **102** (2009) 222001 [[arXiv:0902.2760](#)] [[SPIRES](#)].

- [34] C.F. Berger et al., *Next-to-leading order QCD predictions for $W+3$ -jet distributions at hadron colliders*, *Phys. Rev. D* **80** (2009) 074036 [[arXiv:0907.1984](#)] [[SPIRES](#)].
- [35] R.K. Ellis, K. Melnikov and G. Zanderighi, *Generalized unitarity at work: first NLO QCD results for hadronic $W + 3$ jet production*, *JHEP* **04** (2009) 077 [[arXiv:0901.4101](#)] [[SPIRES](#)].
- [36] R. Keith Ellis, K. Melnikov and G. Zanderighi, *$W+3$ jet production at the Tevatron*, *Phys. Rev. D* **80** (2009) 094002 [[arXiv:0906.1445](#)] [[SPIRES](#)].
- [37] G. Ossola, C.G. Papadopoulos and R. Pittau, *CutTools: a program implementing the OPP reduction method to compute one-loop amplitudes*, *JHEP* **03** (2008) 042 [[arXiv:0711.3596](#)] [[SPIRES](#)].
- [38] G. Bevilacqua, M. Czakon, C.G. Papadopoulos, R. Pittau and M. Worek, *Assault on the NLO Wishlist: $pp \rightarrow t\bar{t}b\bar{b}$* , *JHEP* **09** (2009) 109 [[arXiv:0907.4723](#)] [[SPIRES](#)].
- [39] K. Melnikov and M. Schulze, *NLO QCD corrections to top quark pair production and decay at hadron colliders*, *JHEP* **08** (2009) 049 [[arXiv:0907.3090](#)] [[SPIRES](#)].
- [40] L.J. Dixon, E.W.N. Glover and V.V. Khoze, *MHV rules for Higgs plus multi-gluon amplitudes*, *JHEP* **12** (2004) 015 [[hep-th/0411092](#)] [[SPIRES](#)].
- [41] S.D. Badger and E.W.N. Glover, *One-loop helicity amplitudes for $H \rightarrow$ gluons: the all-minus configuration*, *Nucl. Phys. Proc. Suppl.* **160** (2006) 71 [[hep-ph/0607139](#)] [[SPIRES](#)].
- [42] S.D. Badger, E.W.N. Glover and K. Risager, *One-loop phi-MHV amplitudes using the unitarity bootstrap*, *JHEP* **07** (2007) 066 [[arXiv:0704.3914](#)] [[SPIRES](#)].
- [43] E.W.N. Glover, P. Mastrolia and C. Williams, *One-loop phi-MHV amplitudes using the unitarity bootstrap: the general helicity case*, *JHEP* **08** (2008) 017 [[arXiv:0804.4149](#)] [[SPIRES](#)].
- [44] C.F. Berger, V. Del Duca and L.J. Dixon, *Recursive construction of Higgs+multiparton loop amplitudes: The last of the phi-nite loop amplitudes*, *Phys. Rev. D* **74** (2006) 094021 [*Erratum ibid.* **D 76** (2007) 099901] [[hep-ph/0608180](#)] [[SPIRES](#)].
- [45] S. Badger, E.W.N. Glover, P. Mastrolia and C. Williams, *One-loop Higgs plus four gluon amplitudes: full analytic results*, [arXiv:0909.4475](#) [[SPIRES](#)].
- [46] L.J. Dixon and Y. Sofianatos, *Analytic one-loop amplitudes for a Higgs boson plus four partons*, *JHEP* **08** (2009) 058 [[arXiv:0906.0008](#)] [[SPIRES](#)].
- [47] V. Del Duca, A. Frizzo and F. Maltoni, *Higgs boson production in association with three jets*, *JHEP* **05** (2004) 064 [[hep-ph/0404013](#)] [[SPIRES](#)].
- [48] P. Mastrolia, *On triple-cut of scattering amplitudes*, *Phys. Lett. B* **644** (2007) 272 [[hep-th/0611091](#)] [[SPIRES](#)].
- [49] R. Britto, E. Buchbinder, F. Cachazo and B. Feng, *One-loop amplitudes of gluons in SQCD*, *Phys. Rev. D* **72** (2005) 065012 [[hep-ph/0503132](#)] [[SPIRES](#)].
- [50] R. Britto, B. Feng and P. Mastrolia, *The cut-constructible part of QCD amplitudes*, *Phys. Rev. D* **73** (2006) 105004 [[hep-ph/0602178](#)] [[SPIRES](#)].
- [51] P. Mastrolia, *Double-cut of scattering amplitudes and Stokes' theorem*, *Phys. Lett. B* **678** (2009) 246 [[arXiv:0905.2909](#)] [[SPIRES](#)].

- [52] P. Nogueira, *Automatic Feynman graph generation*, *J. Comput. Phys.* **105** (1993) 279 [[SPIRES](#)].
- [53] J.A.M. Vermaseren, *The FORM project*, *Nucl. Phys. Proc. Suppl.* **183** (2008) 19 [[arXiv:0806.4080](#)] [[SPIRES](#)].
- [54] D. Maître and P. Mastrolia, *S@M, a mathematica implementation of the Spinor-Helicity formalism*, *Comput. Phys. Commun.* **179** (2008) 501 [[arXiv:0710.5559](#)] [[SPIRES](#)].
- [55] M.L. Mangano and S.J. Parke, *Multi-parton amplitudes in gauge theories*, *Phys. Rept.* **200** (1991) 301 [[hep-th/0509223](#)] [[SPIRES](#)].
- [56] L.J. Dixon, *Calculating scattering amplitudes efficiently*, [hep-ph/9601359](#) [[SPIRES](#)].
- [57] Z. Bern, L.J. Dixon and D.A. Kosower, *One-loop amplitudes for e^+e^- to four partons*, *Nucl. Phys. B* **513** (1998) 3 [[hep-ph/9708239](#)] [[SPIRES](#)].
- [58] H.J. Lu and C.A. Perez, *Massless one loop scalar three point integral and associated Clausen, Glaisher and L functions*, SLAC-PUB-5809 [[SPIRES](#)].
- [59] A.I. Davydychev and J.B. Tausk, *A magic connection between massive and massless diagrams*, *Phys. Rev. D* **53** (1996) 7381 [[hep-ph/9504431](#)] [[SPIRES](#)].
- [60] N.I. Usyukina and A.I. Davydychev, *Exact results for three and four point ladder diagrams with an arbitrary number of rungs*, *Phys. Lett. B* **305** (1993) 136 [[SPIRES](#)].