

# Chiral Superfluidity of the Quark-Gluon Plasma

Tigran Kalaydzhyan<sup>1,2</sup>, Ingo Kirsch<sup>1</sup>, Henry Verschelde<sup>3</sup> and Valentin I. Zakharov<sup>2,4</sup>

<sup>1</sup>DESY-Hamburg, <sup>2</sup>ITEP-Moscow, <sup>3</sup>Ghent University, <sup>4</sup>MPI-Munich



## Motivation


### Irregular structure of the QCD vacuum

- fractal distribution of the topological charge density
- low-dimensional defects

### Effective model for sQGP

- effective Lagrangian
- two-component fluid

### Axion-like field within QCD



## Bosonization with a finite cut-off

Euclidean functional integral for  $SU(N_c) \times U_{em}(1)$  is given by

$$\int \mathcal{D}\bar{\psi} \mathcal{D}\psi \exp \left\{ - \int_V d^4x \bar{\psi} (\not{D} - im) \psi + \frac{1}{4} G^{a\mu\nu} G_{\mu\nu}^a + \frac{1}{4} F^{\mu\nu} F_{\mu\nu} \right\},$$

where we define the Dirac operator as

$$\not{D} = -i(\not{\partial} + \mathbf{A} + g\mathbf{G} + \gamma_5 \mathbf{A}_5)$$

- integrate out quarks below a cut-off Dirac eigenvalue  $\Lambda$
- add gauge-invariant terms to the Lagrangian to match the chiral anomaly
- consider a pure gauge  $\mathbf{A}_{5\mu} = \partial_\mu \theta$  for the auxiliary axial field
- and the chiral limit  $m \rightarrow 0$

The **total effective Euclidean Lagrangian** reads as

$$\begin{aligned} \mathcal{L}_E^{(4)} = & \frac{1}{4} G^{a\mu\nu} G_{\mu\nu}^a + \frac{1}{4} F^{\mu\nu} F_{\mu\nu} - j^\mu A_\mu \\ & + \frac{\Lambda^2 N_c}{4\pi^2} \partial^\mu \theta \partial_\mu \theta + \frac{g^2}{16\pi^2} \theta G^{a\mu\nu} \tilde{G}_{\mu\nu}^a + \frac{N_c}{8\pi^2} \theta F^{\mu\nu} \tilde{F}_{\mu\nu} \\ & + \frac{N_c}{24\pi^2} \theta \square^2 \theta - \frac{N_c}{12\pi^2} (\partial^\mu \theta \partial_\mu \theta)^2 \end{aligned}$$

So we get an axion-like field with decay constant  $f = \frac{2\Lambda}{\pi} \sqrt{N_c}$  and a

negligible mass  $m_\theta^2 = \lim_{V \rightarrow \infty} \frac{\langle Q^2 \rangle}{f^2 V} \equiv \chi(T)/f^2$ . We tend to interpret it as a quasiparticle moving along the low-dimensional defects!

## Interpretation of the scale $\Lambda$

From the quartic Lagrangian at  $N_c = N_f = 1$  we get

$$\rho_5 = \frac{1}{2} \left( \frac{\Lambda}{\pi} \right)^2 \mu_5 + \frac{1}{3\pi^2} \mu_5^3$$

- Free quarks (see 0808.3382):  $\Lambda = \pi \sqrt{\frac{2}{3}} \sqrt{T^2 + \frac{\mu^2}{\pi^2}} \propto \rho_{sphaleron}^{-1}$
- Free quarks and strong B-field:  $\Lambda = 2\sqrt{|eB|}$
- Dynamical lattice fermions (1105.0385):  $\Lambda \simeq 3 \text{ GeV} \gg \Lambda_{QCD}$

## Change in entropy and higher order gradient corrections

The terms  $\tau^{\mu\nu}$ ,  $\nu^\mu$  and  $\nu_5^\mu$  denote higher-order gradient corrections and obey the Landau conditions

$$u_\mu \tau^{\mu\nu} = 0, \quad u_\mu \nu^\mu = 0, \quad u_\mu \nu_5^\mu = 0.$$


Using both hydrodynamic equations and constitutive relations one can derive

$$\partial_\mu (s u^\mu - \frac{\mu}{T} \nu^\mu - \frac{\mu_5}{T} \nu_5^\mu) = -\frac{1}{T} (\partial_\mu u_\nu) \tau^{\mu\nu} - \nu^\mu (\partial_\mu \frac{\mu}{T} - \frac{1}{T} E_\mu) - \nu_5^\mu \partial_\mu \frac{\mu_5}{T}.$$


so the entropy production is always nonnegative. This fact tells us that

- we don't need to add any additional terms to the entropy current like in Son and Surowka (0906.5044)
- no additional first-order corrections to the currents.
- in absence of dissipative corrections we obtain  $\partial_\mu (s u^\mu) = 0$ , i.e. only the "normal" component contributes to the entropy current, while the "superfluid" component has zero entropy.

## Fermionic spectrum at finite temperature



- there are two separated parts of the spectrum at intermediate temperatures!
- a strong external magnetic field does not destroy the picture
- all the chiral properties are described by the near zero modes



## Hydrodynamic equations

Equations of motion for the quadratic effective Minkowski Lagrangian

$$\begin{aligned} \partial^\mu \partial_\mu \theta &= \frac{C}{4f^2} F^{\mu\nu} \tilde{F}_{\mu\nu} + \frac{g^2}{32\pi^2 f^2} G_a^{\mu\nu} \tilde{G}_{\mu\nu}^a, \\ \partial_\mu F^{\mu\nu} &= -j^\nu + C(\partial_\sigma \theta) \tilde{F}^{\sigma\nu}, \\ \partial_\mu \tilde{F}^{\mu\nu} &= 0. \end{aligned}$$

Varying the quadratic Lagrangian with respect to axial-vector  $\mathbf{A}_{5\mu} = \partial_\mu \theta$  we obtain the axial current  $j_5^\mu = -f^2 \partial^\mu \theta$  (curl-free!). Conservation law  $\partial_\mu (T^{\mu\nu} + \Theta^{\mu\nu}) = 0$  makes it possible to express divergency of the fluid energy-momentum tensor  $T^{\mu\nu}$  via the one of the electromagnetic stress-energy tensor  $\Theta^{\mu\nu} = F^{\mu\lambda} F_\lambda^\nu + \frac{1}{4} g^{\mu\nu} F_{\alpha\beta} F^{\alpha\beta}$ .

In summary, the hydrodynamic equations are

$$\begin{aligned} \partial_\mu T^{\mu\nu} &= F^{\nu\lambda} (j_\lambda + C \tilde{F}_{\lambda\sigma} \partial^\sigma \theta) \equiv F^{\nu\lambda} (j_\lambda + j_{s\lambda}), \\ \partial_\mu j_5^\mu &= -\frac{C}{4} F^{\mu\nu} \tilde{F}_{\mu\nu} - \frac{g^2}{32\pi^2} G_a^{\mu\nu} \tilde{G}_{\mu\nu}^a, \\ \partial_\mu j^\mu &= 0, \end{aligned}$$

plus the Josephson equation  $u^\mu \partial_\mu \theta + \mu_5 = 0$ .

Corresponding constitutive relations in gradient expansion are

$$\begin{aligned} T^{\mu\nu} &= (\epsilon + P) u^\mu u^\nu + P g^{\mu\nu} + f^2 \partial^\mu \theta \partial^\nu \theta + \tau^{\mu\nu}, \\ j^\mu &= \rho u^\mu + \nu^\mu, \\ j_5^\mu &= -f^2 \partial^\mu \theta + \nu_5^\mu \end{aligned}$$

The stress-energy tensor  $T^{\mu\nu}$  consists of two parts, an ordinary fluid component and a pseudoscalar "superfluid" component. This modifies the equation of state by adding to the r.h.s. a new  $\theta$ -dependent term

$$dP = s dT + \rho d\mu - f^2 d \left[ \frac{1}{2} \partial^\mu \theta \partial_\mu \theta \right],$$

where  $s$  is the entropy density.

## Phenomenological output

Electric and magnetic fields in the fluid rest frame are defined as

$$E^\mu = F^{\mu\nu} u_\nu, \quad B^\mu = \tilde{F}^{\mu\nu} u_\nu \equiv \frac{1}{2} \epsilon^{\mu\nu\alpha\beta} u_\nu F_{\alpha\beta}$$

An additional electric current, induced by  $\theta$ -field

$$j_\lambda^S = C \tilde{F}_{\lambda\sigma} \partial^\sigma \theta = -C \mu_5 B_\lambda + C \epsilon_{\lambda\alpha\sigma\beta} u^\alpha \partial_\sigma \theta E_\beta - u_\lambda (\partial \theta \cdot B)$$

- I term: Chiral Magnetic Effect (electric current along B-field)
- II term: Chiral Electric Effect (electric current transverse to E-field and to both normal and superfluid velocities)
- III term: Chiral Dipole Wave (dipole moment induced by B-field)
- The field  $\theta(\vec{x}, t)$  itself: Chiral Magnetic Wave (propagating imbalance between the number of left- and right-handed quarks)